



UNIVERSITÀ POLITECNICA DELLE MARCHE
Repository ISTITUZIONALE

Altitude affects the quality of the water-extractable organic matter (WEOM) from rhizosphere and bulk soil in European beech forests

This is the peer reviewed version of the following article:

Original

Altitude affects the quality of the water-extractable organic matter (WEOM) from rhizosphere and bulk soil in European beech forests / De Feudis, M; Cardelli, Valeria; Massaccesi, L.; Hofmann, D.; Berns, A. E.; Bol, R.; Cocco, Stefania; Corti, Giuseppe; Agnelli, A.. - In: GEODERMA. - ISSN 0016-7061. - STAMPA. - 302:(2017), pp. 6-13. [10.1016/j.geoderma.2017.04.015]

Availability:

This version is available at: 11566/247498 since: 2022-06-03T16:28:55Z

Publisher:

Published

DOI:10.1016/j.geoderma.2017.04.015

Terms of use:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. The use of copyrighted works requires the consent of the rights' holder (author or publisher). Works made available under a Creative Commons license or a Publisher's custom-made license can be used according to the terms and conditions contained therein. See editor's website for further information and terms and conditions.

This item was downloaded from IRIS Università Politecnica delle Marche (<https://iris.univpm.it>). When citing, please refer to the published version.

(Article begins on next page)

1
2
3 **Altitude affects the quality of the water-extractable organic matter (WEOM) from rhizosphere**
4 **and bulk soil in European beech forests**
5
6

7
8
9 De Feudis, M.^{a*}, Cardelli, V.^b, Massaccesi, L.^a, Hofmann, D.^c, Berns, A.E.^c, Bol, R.^c, Cocco, S.^b,
10
11 Corti, G.^b, Agnelli, A.^a
12
13

14
15
16 ^aDipartimento di Scienze Agrarie, Alimentari e Ambientali, Università degli Studi di Perugia,
17
18 Perugia, Italy
19

20 ^bDipartimento di Scienze Agrarie, Alimentari e Ambientali, Università Politecnica delle Marche,
21
22 Ancona, Italy
23

24 ^cInstitute for Bio- and Geosciences, IBG-3 Agrosphere, Forschungszentrum Jülich GmbH, 52425
25
26 Jülich, Germany
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44

45
46 **Corresponding author:*

47 Mauro De Feudis

48 Dipartimento di Scienze Agrarie, Alimentari ed Ambientali

49 Università degli Studi di Perugia

50 Borgo XX Giugno 72

51 06121 Perugia, Italy

52 e-mail: mauro.defeudis@studenti.unipg.it
53
54
55
56
57
58
59

60
61
62 **Abstract**
63

64 Water-extractable organic matter (WEOM) is the most dynamic and bioavailable fraction of the soil
65 organic matter pool. Although the litter floor is considered the main source of WEOM, roots also
66 release a great amount of labile organic compounds through rhizodeposition processes. This makes
67 the rhizosphere, the small soil volume in proximity to the roots, a soil compartment relatively enriched
68 in WEOM. Since both the rhizosphere and the labile organic C pool are highly sensitive to the
69 environmental conditions we evaluated the characteristics of WEOM from rhizosphere and bulk soil
70 collected from the A horizons of European beech (*Fagus sylvatica* L.) forest soils of Apennines
71 mountains (central Italy) at two altitudes (800 and 1000 m), using elevation as a proxy for temperature
72 change. Specifically, we tested if *i*) the rhizosphere contains higher amounts of WEOM with a greater
73 diversity of compounds with respect to the bulk soil, and *ii*) this effect is more pronounced at higher
74 altitude. At both 800 m and 1000 m above sea level, the main distinction between WEOM from
75 rhizosphere and bulk soil was the larger amounts of sugars in the soil close to the roots. Further, our
76 results indicated an influence of altitude on rhizospheric processes as suggested by the larger
77 concentrations of organic C and soluble phenols, and richness of tannins in the rhizosphere WEOM
78 than in the bulk soil at 1000 m. We attributed this influence to environmental constraints which
79 enhanced the release of labile organics and secondary metabolites by rhizodeposition and
80 humification processes in the rhizosphere. As a whole, our data draw a picture where the roots are
81 able to affect the characteristics of WEOM and environmental constraints enhance the differentiation
82 between rhizosphere and bulk soil. This view confirms the influence of the rhizosphere on the soil C
83 cycle, and the importance of the rhizospheric processes when environmental conditions become
84 harsher.
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108

109 **Keywords:** labile soil C pool; mountain soils; polyphenolic compounds; *ESI FT-ICR MS*; climate
110 change
111
112
113
114

115 **1. Introduction**
116
117
118

119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177

Soil water-extractable organic matter (WEOM) is a mixture of macromolecules and low molecular weight compounds produced by decomposition of fresh (leaf litter, root exudates, decaying fine roots) and old soil organic matter (McDowell, 2003). WEOM is an organic fraction that accounts for a small portion on soil organic matter (SOM), and is composed of easily degradable molecules that represent the main C and energy source for the soil microbial community (Smolander and Kitunen 2002; Kaiser and Kalbitz, 2012). Because of this, WEOM is therefore the most dynamic and bioavailable fraction of the SOM (McDowell, 2003) and it is often considered as an indicator of microbial activity (Gutiérrez-Girón et al., 2015).

Although the decomposition of the litter floor is considered the main source of WEOM (Kalbitz and Kaiser, 2007), roots release a wide range of labile and soluble organic compounds into the soil. In fact, plants can release 10 to 40 % of their total net C assimilation per year from their roots through respiration and rhizodeposition in the form of root exudates, mucilage, sloughed-off cells and decaying roots into the soil environment (Lynch and Whipps, 1990; Bertin et al., 2003). The allocation of labile compounds through rhizodeposition processes makes the rhizosphere, the small soil volume in proximity to the root, a soil compartment enriched with WEOM compared to the bulk soil (e.g., Fujii et al., 2012; Massaccesi et al., 2015; Agnelli et al., 2016). Conversely to the rhizosphere, in the soil far from the roots the labile organics mainly derive from the decomposition of SOM (Jones et al., 2009). The easily decomposable organic compounds released by rhizodeposition processes fuel the rhizosphere heterotrophic microbial community (Boddy et al., 2007; Phillips et al., 2011), which in turn may boost SOM decomposition by the priming effect (Kuzyakov, 2002).

WEOM is involved in several biogeochemical processes, and its molecular composition and concentration in soil are highly sensitive to ecosystem disturbances, such as temperature changes. Many authors reported on the influence of temperature on WEOM properties, although their findings were contrasting possibly because the effect of temperature varied depending on the applied methodologies and the nature of the studied ecosystems. For example, Hassouna et al. (2010)

178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236

recorded a weak impact of seasonal temperature variations on WEOM quantity in cropland soils, while Roth et al. (2015) found that during the warm seasons forest soils showed a lower abundance of labile organic compounds with respect to colder seasons. Williams et al. (2016) detected no seasonal variation in the amount of soluble phenols in pasture soils. Delarue et al. (2014), by using an in-situ experimental warming with open-top chambers, reported that the soluble organic matter increased in the subsurface of *Sphagnum*-dominated peatland at higher temperatures. Further, in laboratory experiments, Ultra et al. (2013) and Yin et al. (2013a, b) found that high temperatures may increase the amount of WEOM by increasing organic C inputs into the soil via enhanced root exudation. The effect of the temperature was also tested by using altitude as a proxy for temperature change (Vincent et al., 2014; Gutiérrez-Girón et al., 2015). In fact, an increase in the contents of more labile C compounds with altitude was reported in both cultivated (e.g., Chen et al., 2006) and natural (e.g., Fernández Sanjurjo et al., 2003; Xu et al., 2015) soils.

The objective of the present study was to evaluate the effect of the altitude-depending temperature on the quantity and quality of the labile and potentially soluble organic pool in the rhizosphere and bulk soil of Mediterranean mountain areas under European beech (*Fagus sylvatica* L.). We compared WEOM properties of rhizosphere and bulk soil of A horizons under European beech forests at two altitudes (800 and 1000 m) on three mountains in central Italy. It is noteworthy to say that the sites at 800 and 1000 m altitude had a mean annual air temperature that differed by 1°C, which is the expected increase of the air temperature for the year 2050 (IPCC, 2013). Through the combination of chemical and spectroscopic analyses we tested the hypotheses that i) the rhizosphere contains higher amounts of WEOM with a greater diversity of compounds with respect to the bulk soil, and ii) this effect is more pronounced at higher altitude since the rhizosphere is more sensitive than the bulk soil to changes of environmental conditions.

2. Materials and methods

2.1. Study areas and soil features

237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295

As study areas, three calcareous massifs were selected on the Apennine chain (central Italy): Mount Terminillo (42°28' N, 12°59' E), Mount San Vicino (43°19' N, 13°03' E), and Mount Acuto (43°28' N, 12°41' E) (Figure S1 of the Supplementary Materials). Two European beech (*Fagus sylvatica* L.) forests were selected at each site at about 800 and 1000 m above sea level, on slopes with an inclination ranging from 25° to 40° and northern aspect. All the forests were coppices in conversion, with the conversion that started from about 20 to about 40 years ago. The mean annual air temperature (MAAT) in the three study areas was 10°C at 800 m and about 9°C at 1000 m with January as the coldest month and July the warmest one. The mean annual precipitation was 1248 mm at Mount Terminillo, 825 mm at Mount San Vicino, and 1430 at Mount Acuto. All the soils had a *mesic* soil temperature regime, and an *udic* soil moisture regime. A detailed description of each site is reported in Table S1 of the Supplementary Materials.

All soils developed from limestone and their morphology, reported in Table S1 of the Supplementary Materials, was obtained per Schoeneberger et al. (2012). Here it suffices to recollect that the mean thickness of the *solum* was 68.8 cm (standard error 15.9) for the soils at 800 m, and 49.3 cm (standard error 9.5) for those at 1000 m. All the soils hosted an *amphimus* type of humus (Baize and Girard, 2008) which is characterized by well-developed O horizons rich in mesofauna that rest on an A horizon with a well-developed crumb structure. According to Soil Survey Staff (2014) the soils at 800 m were classified as Inceptisols (Humudepts) and Mollisols (Hapludolls and Haprendoll), while those at 1000 m were all Mollisols (Haprendoll). Hence, slightly harsher environmental conditions existed at 1000 m because of a lower MAAT and a general reduced soil development than at 800 m.

2.2. Soil sampling

During the late winter 2014 two profiles were opened within a plot of about 100 m² at each altitudinal site (3 study areas x 2 altitudes x 2 profiles). The winter sampling was chosen because European beech is a dormant species, and the low rhizosphere and soil microbial community activities occurring during the cold season allow comparing rhizosphere and bulk soil in a more stable condition rather

296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354

than in the dynamic seasons like spring or summer (De Feudis et al., 2016). The profiles were opened at about 50-60 cm from the stem of selected beech trees with an age ranging between 40 and 65 years old. The influence of the different age of the selected trees on the rhizosphere properties was assumed to be negligible because we considered as rhizosphere only the soil associated with fine roots, the activity of which is rather independent of plant age because of their periodic turnover (Trumbore and Gaudinki, 2003; Agnelli et al., 2014).

A large amount of sample (at least 3 kg) was collected from the A horizon of each soil and stored in a portable refrigerator. The beech rhizosphere of each sample was isolated in the laboratory by picking up the roots with a diameter lesser than 2 mm together with the adhering soil (Cocco et al., 2013; Massaccesi et al., 2015). The roots with a greater diameter than 2 mm were discarded. After a gentle shaking to detach the weakly adhering soil particles, which were then added to the bulk soil (i.e. the soil not strictly adhering to the roots), the soil material firmly adhering to the fine roots was considered as rhizosphere and recovered by further shaking and soft brushing. The root fragments present in the bulk soil were removed by using tweezers under a magnifying lens. Both rhizosphere and bulk soil samples were air-dried and sieved through a 2-mm mesh.

2.3. Soil chemical analysis

The soil pH was determined potentiometrically in water ($\text{pH}_{\text{H}_2\text{O}}$) and in 1 M KCl solution (pH_{KCl}) (solid:liquid ratio of 1:2.5) after 30 minutes of stirring by a combined glass-calomel electrode.

The content of total organic C (TOC) was estimated by K-dichromate digestion, heating the suspension at 180 °C for 30 minutes (Nelson and Sommers, 1996), and total N content was determined by a Carlo Erba EA1110 dry combustion analyzer (Carlo Erba Instruments, Italy).

Available P was estimated according to Olsen et al. (1954).

Total phenol contents were extracted according to Alonso et al. (1998) with a solvent mixture of methanol-distilled water (4:1 v:v ratio) with 2 % trimethylamine. Briefly, 5 mL of solvent mixture were added to 1 g of soil sample, and shaken for 1 h with an orbital shaker (130 rpm). The suspension

355
356
357 was centrifuged at 1400 g for 10 min, and then filtered through a 0.45 µm membrane filter. The
358
359 content of total phenols in the extracts was determined colorimetrically by the Prussian blue method
360
361 (Graham, 1992; Hagerman, 2002). To a mixture composed of 0.1 mL of extract and 3 mL of distilled
362
363 water, 1 mL of 0.016 M $K_3Fe(CN)_6$ (Prussian blue) solution was added, followed immediately by the
364
365 addition of 1 mL of 0.02 M $FeCl_3$ + 0.1 M HCl solution. The mixture was put in a vortex for a while
366
367 and then left to rest for 15 minutes. Subsequently, 5 mL of stabilizer (800 mL of distilled water + 100
368
369 mL of 85 % H_3PO_4 solution + 100 mL of 1 % gum arabic solution) were added and the absorbance
370
371 was read at 700 nm by means of a Lambda EZ 150 UV/VIS Spectrometer (Perkin Elmer, USA)
372
373 against a gallic acid standard. The total phenols concentration was normalized against TOC content
374
375 and expressed as mg gallic acid equivalent g^{-1} organic C.
376
377
378

379 380 381 *2.4. Cross-Polarization Magic-Angle-Spinning Carbon-13 NMR Spectroscopy (CPMAS ^{13}C NMR)*

382
383 Solid-state ^{13}C NMR spectroscopy was employed to assess the main structural components of the
384
385 SOM in the whole soil samples. Solid-state ^{13}C NMR spectra were acquired with the cross-
386
387 polarization magic-angle-spinning (CPMAS) technique performed at 15 kHz, using a 7.05 T Varian
388
389 INOVA™ Unity (Varian Inc., Palo Alto, CA, USA) spectrometer at a frequency of 75.4 MHz. The
390
391 spectra were collected with a sweep width of 25 kHz and an acquisition time of 20 ms. Contact time
392
393 for cross-polarization and recycle delay were 1250 µs and 0.5 s, respectively. The 1H radio frequency
394
395 (RF) field strength was set to 47.0 kHz and the ^{13}C RF field strength to 41.1 kHz. An ascending ramp
396
397 of 15.3 kHz on the 1H RF field was used (Berns and Conte, 2011).
398
399
400

401 402 *2.5. Water-extractable organic matter (WEOM)*

403 404 *2.5.1. WEOM extraction*

405
406 The lack of a standardized method to obtain WEOM makes this organic material an operationally
407
408 defined fraction as its composition depends on the combination of several factors such as, for
409
410 example, extracting solution (water or dilute saline solution), sample pre-treatment (fresh or air-dried)
411
412
413

414
415
416 soil:solution ratio, contact time, and energy of the treatment (Jones and Willett, 2006). Therefore, the
417
418 organic matter obtained by a very mild extraction with a low ionic-strength aqueous solution, a 1:2
419
420 solid:liquid ratio, a short contact time (1 min) and a gentle stirring of the mixture is considered
421
422 representative of the soil solution dissolved organics collected in situ with zero-tension lysimeters
423
424 (Zsolnay 2003; Chantigny et al., 2008). In contrast, a more vigorous extraction using distilled water
425
426 with wider solid : liquid ratio (1:5 or 1:10) and overnight shaking solubilizes greater amount of
427
428 organics because of the partial aggregate disruption and release of organic molecules that are not
429
430 typically in the soil solution (Kaiser et al., 2015). To this regard, since the goal of this study was to
431
432 evaluate the effect of altitude on the whole labile and potentially soluble organic pool of rhizosphere
433
434 and bulk soils, we chose to extract the WEOM by the following procedure (Agnelli et al., 2016): 1 g
435
436 of rhizosphere or bulk soil sample was placed into a plastic container, submerged with distilled water
437
438 (solid:liquid ratio 1:10) and shaken (140 rpm) overnight with an orbital shaker. The suspension was
439
440 centrifuged at 1400 g for 10 min, and then filtered through a 0.45 μm membrane filter. The obtained
441
442 solution was stored in the refrigerator and analyzed within a week.
443
444
445
446
447

448 2.5.2. *Chemical analyses*

449

450 The organic C content of the WEOM was analyzed by a TOC-5000A analyzer (Shimadzu Corp.,
451
452 Tokyo, Japan) on aliquots of the extracted solution after the addition of a few drops of concentrated
453
454 H_3PO_4 to eliminate inorganic C.
455

456 The contents of hexose and pentose sugars in the WEOM were colorimetrically estimated (Hofman
457
458 and Dušek, 2003) by using the anthrone and orcinol methods, respectively. Briefly, for the hexoses,
459
460 5 mL of anthrone reagent (0.2 g of anthrone + 5 ml of ethanol + 95 ml of 75% H_2SO_4) were added to
461
462 1 mL of extracted solution in a shaking ice-water bath. Once the mixture was cooled it was heated in
463
464 a boiling water bath for 10 minutes and, then, cooled again in an ice-bath. Once cooled, the
465
466 absorbance was measured at 625 nm against a glucose standard. For the pentoses, 3 mL of orcinol
467
468 reagent (0.5 g of orcinol + 50 ml of ethanol + 0.18 g $\text{FeCl}_3 \cdot 6\text{H}_2\text{O}$ in 200 ml of concentrated HCl)
469
470
471
472

473
474
475 were added to 1 mL of extracted solution, and the mixture was heated in a boiling water bath for 20
476
477 minutes. Once cooled, the absorbance was read at 672 nm using ribose as standard. The phenols
478
479 concentration in the WEOM was determined colorimetrically by the Prussian blue method.
480
481

482 483 484 *2.5.3. Electrospray Ionization Fourier Transform-Ion Cyclotron Resonance Mass Spectrometry (ESI* 485 486 *FT-ICR MS)*

487
488 WEOM aliquots were analysed by an ESI FT-ICR MS which measures the mass-to-charge ratio of
489
490 organic molecules after their ionisation. The electrospray ionization induces the breaking of the non-
491
492 covalent bonds such as Van der Waals forces and electrostatic interaction, and typically allows the
493
494 detection of molecules with a mass lesser than 1000 Da (Hertkorn et al., 2008; Stubbins and Dittmar,
495
496 2015). The ultra-high resolution in combination with exact mass characterisation (here below 1 ppm)
497
498 of this technology allows identification of thousands of compounds for each sample (Sleighter and
499
500 Hatcher, 2007). However, the FT-ICR MS cannot be used as quantitative analysis (Sipler and
501
502 Seitzinger, 2008) because the polarity of the compounds affects the intensity of the signals: the higher
503
504 the polarity, the higher the intensity. To prevent quenching processes due to soil inherent permanent
505
506 ions a combined desalting and pre-concentration of the WEOM samples was performed by solid
507
508 phase extraction (SPE, C18 hydra cartridges, Machery & Nagel, Düren, Germany), using methanol
509
510 ($\geq 99.98\%$, Ultra LC-MS grade; Carl Roth, Germany) as back eluent. The mean C recovery achieved
511
512 with the SPE cartridges ranged between 66 and 71%, in agreement with the values reported in
513
514 literature (e.g., Stubbins et al., 2012; Stubbins and Dittmar, 2015).
515
516

517
518 The ultra-high-resolution mass spectra were acquired using an ESI-LTQ-FT Ultra (ThermoFisher
519
520 Scientific, San Jose, CA, USA) equipped with a 7 T supra-conducting magnet (Oxford Instruments,
521
522 Abingdon, UK). The mass spectrometer was used in negative ionization mode with 2.9 kV spray
523
524 voltage, -50 V capillary voltage, 275 °C transfer capillary temperature, and nitrogen as sheath gas.
525

526
527 The samples were introduced by a syringe pump providing an infusion rate of 8 $\mu\text{l}/\text{min}$, and full FT-
528
529 ICR MS spectra between 200 and 1000 Da were recorded at a mass resolving power of 400,000 at
530
531

532
533
534 m/z 400 Da. The spectra were averaged out of 7 scans, and each scan was accumulated from 50
535
536 transients. Moreover, the ICR cell filling was adjusted to 500.000 ions per transient to obtain an
537
538 accuracy of 0.4 ppm. The formula assignment was performed by an in-house developed calculation
539
540 program using Scilab routines. The results obtained from the ESI FT-ICR MS were presented by
541
542 plotting van Krevelen diagrams (Kim et al., 2003) of H/C versus O/C molar ratios of the CHO
543
544 compounds (Hofmann et al., 2015). To estimate the proportion of WEOM components four regions
545
546 were delineated within the diagrams according to Sleighter and Hatcher (2007) and Ohno et al.
547
548 (2010), to cluster the detected molecules within the typical ranges of selected biopolymers: lignins
549
550 (H/C: 0.7-1.5, O/C: 0.1-0.67); tannins (H/C: 0.5-1.4, O/C: 0.67-0.9); condensed aromatics (H/C: 0.2-
551
552 0.7, O/C: 0.1-0.67); sugars (H/C: 1.5-2.0, O/C: 0.67-1.0).
553
554
555
556
557

558 *2.6. Statistical analysis*

559
560 Two-way ANOVA was carried out by including rhizosphere and bulk soil (2 levels) and altitudes (2
561
562 levels) as fixed factors. The study area (3 levels) was also included as blocking factor. The normality
563
564 and homoscedasticity of the data were verified by graphical analysis of residuals and transformed if
565
566 necessary. The transformation was selected by the maximum likelihood procedure suggested by Box
567
568 and Cox (1964), as implemented in the boxcox function of the package MASS (Venables and Ripley
569
570 2002). The results presented and discussed are based on mean values and standard error, and the
571
572 comparison of means was assessed by Tukey HSD post-hoc test ($P < 0.05$).
573
574

575
576 Principal component analysis (PCA) was applied only to WEOM properties (contents of water-
577
578 extractable organic C, pentoses, hexoses and phenols, and richness of lignins, tannins and condensed
579
580 aromatic compounds) to identify the variables capable of explaining most of the variability between
581
582 the two altitudes and between rhizosphere and bulk soil. The WEOM dataset was standardized prior
583
584 to PCA by subtracting the mean and dividing by the standard deviation.

585
586 The data were analyzed using R software (R Core Team, 2014).
587
588
589
590

591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649

3. Results

3.1. Rhizosphere and bulk soil properties

The bulk soil showed a higher pH in H₂O and in KCl at 1000 than at 800 m of altitude, while no difference was observed for the pH of the rhizosphere at both altitudes (Table 1). The TOC content was greater in the rhizosphere than in the bulk soil for both altitudes, while the total N content was higher at 1000 than at 800 m, with no difference between rhizosphere and bulk soil (Table 1). The content of available P was similar at the two altitudes, but always with higher contents in the rhizosphere than in the bulk soil (Table 1). Total phenols concentrations, once normalized to the TOC content, showed no difference between rhizosphere and bulk soil, while a greater amount was found at 800 m than at 1000 m (Table 1).

The CPMAS ¹³C NMR spectroscopy detected six functional groups: alkyl-C like lipids and cutin at 0–45 ppm, N-alkyl-C like amino acids at 45–60 ppm, O-alkyl-C such as carbohydrates, hemicellulose, and cellulose at 60–110 ppm, aryl-C such as lignins, tannins, and aromatic C compounds at 110–160 ppm, carboxyl-C like carboxyl acids at 160 – 190 ppm, and carbonyl-C such as aldehydes and ketons at 190–215 ppm. The relative concentration of these organic functional groups in the rhizosphere and bulk soils showed the following general trend: O-alkyl-C > aryl-C ≥ alkyl-C > N-alkyl-C = carboxyl-C > carbonyl-C (Table 2, and Figure S2 of the Supplementary Materials). Rhizosphere and bulk soil showed no difference in the content of the various C forms, but some difference in the SOM composition occurred between the two altitudes: O-alkyl-C and alkyl-C had a larger relative percentage at 800 m, whereas the aryl-C showed greater proportion at 1000 m.

3.2. WEOC extracted from rhizosphere and bulk soil

3.2.1. Chemical and spectroscopic (ESI FT-ICR MS) analyses

The water-extractable organic carbon (WEOC) concentration was similar at 800 and 1000 m above sea level (Table 3). However, whereas for the soils at 800 m no difference occurred between rhizosphere and bulk soil, at 1000 m the rhizosphere had higher WEOC content than the respective

650
651
652 bulk soil. The contents of hexose and pentose sugars were always larger in the rhizosphere than in
653
654 the bulk soil, although no significant difference between the two altitudes occurred (Table 3). The
655
656 concentration of water extractable phenols was greater in the rhizosphere at 1000 m, while it did not
657
658 differ between rhizosphere and bulk soil at 800 m (Table 3).
659

660
661 The analysis of WEOM by ESI FT-ICR MS produced a large data set with thousands of peaks
662
663 corresponding to molecules with different mass-to-charge ratios (figure not shown). In particular, the
664
665 data obtained by the mass analyses showed a similar number of compounds (formulas) in all
666
667 rhizosphere and bulk soil samples from both altitudes (Table 4). Further, the van Krevelen diagrams
668
669 exhibited similar patterns for the different samples with most of the molecules clustered in the regions
670
671 corresponding to lignins and tannins (Figure S3 of the Supplementary Materials, Table 4). However,
672
673 while no difference was found between rhizosphere and bulk soil at 800 m, at 1000 m the rhizosphere
674
675 showed a greater number of formulas that fell in the tannins and condensed aromatics regions of the
676
677 van Krevelen diagram (Table 4). The low proportion of sugars detected in the WEOM of all samples
678
679 was attributed to the fact that the ESI FT-ICR MS weakly ionizes oxygen-rich molecules such as
680
681 carbohydrates without permanent charge (Hertkorn et al., 2007). Furthermore, such high polar
682
683 substances could be partly lost in the SPE process.
684
685
686
687

688 3.2.2. *Principal Component Analysis (PCA)* 689

690
691 The first two components calculated by PCA explained 69.3% of the total variance (Figure 1). The
692
693 variability along the PC1 was mainly affected by the content of pentoses, hexoses, WEOC, and the
694
695 richness of tannins (Table 5). The variability along the second component (PC2) was mainly driven
696
697 by the richness of lignins and condensed aromatic molecules (Table 5).
698

699
700 In the PCA scoring plot (Figure 1) the observations have been grouped (rhizosphere or bulk soil, 800
701
702 or 1000 m above sea level) and the 95% confidence ellipses were drawn for each group. The addition
703
704 of confidence ellipses improved the interpretation of the scoring plot because they showed the degree
705
706 of separation of the groups (Hammer and Harper, 2006; Montanari et al., 2016). The lack of
707
708

709
710
711 overlapping of the confidence ellipses drawn for the WEOM of rhizosphere and bulk soil at 1000 m
712
713 indicated a distinction between the two groups. Conversely, this did not happen between the
714
715 rhizosphere and bulk soil at 800 m. Further, the lack of overlapping of the confidence ellipses between
716
717 the WEOM from the rhizospheres at the two altitudes showed a clear differentiation of the two groups.
718
719

720 721 722 **4. Discussion**

723 724 *4.1. Chemical properties of rhizosphere and bulk soil at 800 and 1000 m of altitude*

725
726 Although literature often reports lower pH in the rhizosphere than in the bulk soil (e.g., Pandey and
727
728 Palni, 2007; Cocco et al., 2013; Massaccesi et al., 2015), in our case the absence of significant
729
730 differences between the two fractions was likely due to the calcareous nature of the parent material,
731
732 as protons released by the roots are neutralized by carbonate dissolution (Richter et al., 2007; Agnelli
733
734 et al., 2016). However, the lower weathering rate occurring at higher altitudes (Riebe et al., 2004)
735
736 particularly in soil developed on alkaline substrates in central Italy (e.g., Corti et al., 2012; Agnelli et
737
738 al., 2016), was responsible for the generally higher pH values at 1000 than at 800 m. Altitude also
739
740 had an effect on organic matter content (Dieleman et al., 2013; Prietzel and Christophel, 2014; Tashi
741
742 et al., 2016). Indeed, the lower temperatures occurring at higher altitudes reduce the net primary
743
744 production and the litterfall input (Zianis and Mencuccini, 2005; Bu et al., 2012), but also soil
745
746 microbial activity (Yimer et al., 2006; Xu et al., 2014). The positive relationship between altitude and
747
748 organic matter content was confirmed by the greater TOC and total N contents at 1000 m than at 800
749
750 m. At both altitudes, the larger amount of TOC and available P in the rhizosphere compared to the
751
752 bulk soil was mainly attributed to C input due to root exudates, border cells and decaying roots
753
754 (Hinsinger et al., 2009, Sokolova, 2015). With regards to P, this element is usually limited in soil
755
756 because it mostly forms stable organic compounds or calcium- or iron-phosphates (Frossard et al.,
757
758 1995; Nelson and Janke, 2007). However, plants are able to mobilize this element by the release of
759
760 phosphatase enzymes (Nannipieri et al., 2011), and through exudation of low molecular weight
761
762 organic acids (LMWOA), sugars, and phenolic compounds (Carvalhais et al., 2011; Bowsher et al.,
763
764
765
766
767

768
769
770 2015). In the same study sites De Feudis et al. (2016) attributed the higher P availability in the
771 rhizosphere to an enhanced organic matter cycling triggered by root exudation.
772

773
774 The chemical structure of the organic matter of rhizosphere and bulk samples, as revealed by the ¹³C
775 NMR spectroscopy, was similar to that found by previous studies on forest floors (e.g., Rumpel et al.,
776 2002; Cepáková and Frouz, 2015), and this similarity derived from the strong influence of the organic
777 horizons on the underlying A horizon. In particular, the input of slightly decomposed organic residues
778 into the mineral soil during forest floor decay could explain the general high proportion of O-alkyl C
779 (Rumpel et al., 2002) which is the most represented constituent in the ¹³C NMR spectra of beech
780 leaves and roots (Angst et al., 2016). Hence, the higher amount of O-alkyl-C at the lower altitude
781 suggested relatively greater litter decomposition due to the higher temperatures which could have
782 favored a greater incorporation of cellulose- and hemicellulose-derived substances in the mineral
783 horizons at 800 m than at 1000 m. In addition to the larger proportion of aliphatic compounds, also
784 the higher concentration of extractable total phenols, which we considered as mostly derived from
785 lignins and humic substances decomposition (Santana et al., 2009), would confirm the higher
786 degradation degree of the organic pool at 800 m than at 1000 m. The results obtained from the ¹³C
787 NMR analysis, together with the TOC contents indicated that temperature affected both the quantity
788 (through mineralization rate) and the quality of organic matter. Indeed, besides reducing the rate of
789 organic matter degradation, the lower temperatures at the higher altitude could have promoted the
790 accumulation of aromatic substances which further fostered a greater organic C storage at 1000 than
791 at 800 m. The NMR analysis did not reveal any difference between the SOM of rhizosphere and bulk
792 soil, although some differences were expected. However, the lack of differences between the two
793 fractions might be masked by the fact that we analyzed the whole organic pool. Indeed, although one
794 of the main roles of the roots is the enrichment of the rhizosphere with labile C compounds (Angst et
795 al., 2016), these latter represent a very small proportion of the whole soil organic carbon pool. In our
796 case, the WEOC was less than 0.5% of TOC.
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826

827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885

4.2. WEOM of rhizosphere and bulk soil at 800 and 1000 m of altitude

The rhizosphere had a larger WEOC content than the bulk soil only at 1000 m where slightly harsher environmental conditions might have 1) affected the quality of root exudation (Chen et al., 2006; Guo et al., 2015), 2) increased the amount of labile C compounds (Fernández Sanjurjo et al., 2003; Xu et al., 2015), and 3) accelerated fine roots production and turnover (Graefe et al., 2008; Girardin et al., 2013). In contrast, rhizospheres at 800 and 1000 m showed no significant difference in WEOC content. These findings contrasted with those of Ultra et al. (2013) and Yin et al. (2013a, b) who reported that warming increases the amount of WEOC in the rhizosphere because of enhanced root exudation. However, these studies were run under controlled laboratory conditions where the only variable was the temperature and no other constraints occurred. Instead, the effect of altitude is more complex than that of an experimental warming because factors other than temperature (e.g., soil development and properties, fine roots turnover, nutrient availability) may control processes in the rhizosphere.

As supported by PCA (Figure 1), WEOM of the rhizosphere at 1000 m was qualitatively different from that of the respective bulk soil and the difference was mostly due to a greater richness of phenols, tannins and condensed aromatic compounds in the rhizosphere WEOM (Tables 3 and 4). Among the phenolic compounds, tannins occur in all the different plant tissues, especially in leaves, roots and bark (Kraus et al., 2003). However, as fine roots are a great part of forest biomass and often decay faster than the leaf litter (Vogt et al. 1996; Stevens et al., 2002), they can be considered a significant contributor of tannins to forest soils (Kraus et al., 2003). Hence, as at higher altitude roots undergo to a faster turnover (Graefe et al., 2008; Girardin et al., 2013), we attributed the larger presence of phenolic compounds in the rhizosphere than in the bulk at 1000 m to the harsher environmental conditions occurring at this altitude.

The richness of lignin and tannin compounds indicated a general presence of recalcitrant organic molecules (e.g., Kögel-Knaber, 2002; Ohno et al., 2010) in the WEOM of both rhizosphere and bulk soil. Indeed, these carboxyl-rich alicyclic molecules are considered as products derived from

886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944

decomposition of organics and important substrate for the formation of humic substances (e.g., Kraus et al., 2003; Hertkorn et al., 2006; Buondonno et al., 2014). At 1000 m, the rhizosphere displayed a greater diversity of compounds belonging to the condensed aromatics class than the bulk soil (Table 4). This fact was interpreted as the result of an increase of the humification rate in the rhizosphere at the higher altitude, with the consequent indirect increase of the number of recalcitrant compounds (Bayer et al., 2002) in the vicinity of the roots. These larger number of condensed aromatic compounds could be partly due to the mobilization of humic substances promoted by the root exudation of LMWOA which are able to break bridges between metals and humic moieties by complexing metal cations (Takeda et al., 2009).

The higher content of sugars in the rhizosphere than in the bulk soil at both altitudes suggested that the major source of these compounds was root exudation (Gunina and Kuzyakov, 2015). The release of sugars, as well as other low molecular weight organic compounds, is considered as a plant strategy to overcome environmental restrictions through the stimulation of the microbial biomass activity (Bowsher et al., 2015; Kaiser et al., 2015). However, we cannot exclude that the larger content of sugars in the rhizosphere was due to the activity of the microbial biomass that, besides using these compounds as an energetic substrate, is able to produce new carbohydrates both directly and following decomposition of plant residues (Gunina and Kuzyakov, 2015).

In addition to the above considerations the different WEOM characteristics between rhizosphere and bulk soil at 1000 m might be the result of the activity of different microbial communities. For example, the meta-analysis conducted by Blankinship et al. (2011) revealed a higher microbial diversity in soils subjected to lower temperatures, while an increased fungal diversity in the rhizosphere of an annual plant (*Eschscholzia californica* Cham.) was found at higher altitude by de Armas-Ricard et al. (2016).

5. Conclusions

945
946
947 We appraised the effect of altitude (800 and 1000 m) on some characteristics of WEOM extracted
948 from the rhizosphere and bulk soil of beech forests in the Apennine mountains (central Italy). The
949 main distinction occurring at both altitudes between WEOM of rhizosphere and bulk soil was due to
950 the larger amounts of sugars in the former. The greater availability of these compounds in the
951 rhizosphere WEOM was attributed to both rhizodeposition and microbial activity. We hypothesized
952 that the influence of altitude on the rhizospheric processes, as revealed by the larger concentrations
953 of organic C and soluble phenols, and richness of tannins in the rhizosphere WEOM than in the bulk
954 soil at 1000 m, was due to environmental constraints that enhanced the release of labile organics and
955 secondary metabolites by rhizodeposition processes. Furthermore, the presence of these easily
956 degradable compounds in the rhizosphere could promote the degradation of more stable soil organic
957 matter leading to an increase in richness of soluble condensed aromatic moieties. Hence,
958 environmental conditions can affect the belowground C allocation with potential impact on the forms
959 and amounts of the rhizosphere labile organic C pool.
960
961 Because of its capability to respond to disturbances, the rhizosphere WEOM could be taken as an
962 index of environmental changes following temperature variations in both time and space. However,
963 more studies on WEOM and its composition are needed to strengthen this aspect.
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982

983 **References**

- 984
985 Agnelli, A., Bol, R., Trumbore, S.E., Dixon, L., Cocco, S., Corti, G., 2014. Carbon and nitrogen in
986 soil and vine roots in harrowed and grass-covered vineyards. *Agric. Ecosyst. Environ.* 193, 70–82.
987
988 Agnelli, A., Massaccesi, L., De Feudis, M., Cocco, S., Courchesne, F., Corti, G., 2016. Holm oak
989 (*Quercus ilex* L.) rhizosphere affects limestone-derived soil under a multi-centennial forest. *Plant*
990 *Soil* 400, 297–314.
991
992 Alonso, M.C., Puig, D., Silgoner, I., Grasserbauer, M., Barceló, D., 1998. Determination of priority
993 phenolic compounds in soil samples by various extraction methods followed by liquid
994
995
996
997
998
999

1000
1001
1002
1003

1004
1005
1006 chromatography-atmospheric pressure chemical ionisation mass spectrometry. J. Chromatogr. A
1007 823, 231–239.
1008
1009
1010 Angst, G., Kögel-Knaber, I, Kirfel, K., Hertel, D., Mueller, C.W., 2016. Spatial distribution and
1011 chemical composition of soil organic matter fractions in rhizosphere and non-rhizosphere soil under
1012 European beech (*Fagus sylvatica* L.). Geoderma 264, 179–187.
1013
1014
1015 Baize, D., Girard, M.-C. (Eds.) 2008. Référentiel Pédologique. Association française pour l'étude du
1016 sol (Afes). Édition Quæ, Versailles, France.
1017
1018
1019 Bayer, C., Martin-Neto, L., Mielniczuk, J., Saab, S.d.C., Milori, D.M.P., Bagnato, V.S., 2002. Tillage
1020 and cropping system effects on soil humic acid characteristics as determined by electron spin
1021 resonance and fluorescence spectroscopies. Geoderma 105, 81–92.
1022
1023
1024 Berns, A.E., Conte, P., 2011. Effect of ramp size and sample spinning speed on CPMAS ¹³C NMR
1025 spectra of soil organic matter. Org. Geochem. 42, 926–935.
1026
1027
1028 Bertin, C., Yang, X., Weston, L.A., 2003. The role of root exudates and allelochemicals in the
1029 rhizosphere. Plant Soil 256, 67–83.
1030
1031
1032 Blankinship, J.C., Niklaus, P.A., Hungate, B.A., 2011. A meta-analysis of responses of soil biota to
1033 global change. Oecologia 165, 553–565.
1034
1035
1036 Boddy, E., Hill, P.W., Farrar, J., Jones, D., 2007. Fast turnover of low molecular weight components
1037 of dissolved organic carbon pool of temperate grassland field soils. Soil Biol. Biochem. 39, 827–
1038 835.
1039
1040
1041 Bowsher, A.W., Ali, R., Harding, S.A., Tsai, C.-J., Donovan, L.A., 2015. Analysis of wild sunflower
1042 (*Helianthus annuus* L.) root exudates using gas chromatography-mass spectrometry. J. Plant Nutr.
1043 Soil Sci. 178, 776–786.
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053 Box, G.E.P., Cox, D.R., 1964. Analysis of transformations. J. R. Stat. Soc. Ser. B Stat. Methodol. 26,
1054 211–252.
1055
1056
1057
1058
1059
1060
1061
1062

1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121

- Bu, X., Ruan, H., Wang, L., Ma, W., Ding, J., Yu, X., 2012. Soil organic matter in density fractions as related to vegetation changes along an altitude gradient in the Wuyi Mountains, southeastern China. *Appl. Soil Ecol.* 52, 42–47.
- Buondonno, A., Capra, G.F., Coppola, E., Dazzi, C., Grilli, E., Odierna, P., Rubino, M., Vacca, S., 2014. Aspects of soil phenolic matter (SPM): An explorative investigation in agricultural, agroforestry, and wood ecosystems. *Geoderma* 213, 235–244.
- Carvalhais, L.C., Dennis, P.G., Fedoseyonko, D., Hajirezaei, M.-R., Borriss, R., von Wirén, N., 2011. Root exudation of sugars, amino acids, and organic acids by maize as affected by nitrogen, phosphorus, potassium, and iron deficiency. *J. Plant Nutr. Soil Sci.* 174, 3–11.
- Cepáková, Š., Frouz, J., 2015. Changes in chemical composition of litter during decomposition: a review of published ¹³C NMR spectra. *J. Soil Sci. Plant Nutr.* 15, 805–815.
- Chantigny, M.H., Angers, D.A., Kaiser, K., Kalbitz, K., 2008. Extraction and characterization of dissolved organic matter. In: Carter, M.R., Gregorich, E.G. (Eds.) *Soil sampling and methods of analysis*, Second edition, CRC Press Taylor & Francis, Boca Raton, FL, Chapter 48.
- Chen, Y.M., Wang, M.K., Zhuang, S.Y., Chiang, P.N., 2006. Chemical and physical properties of rhizosphere and bulk soils of three tea plants cultivated in Ultisols. *Geoderma* 136, 378–387.
- Cocco, S., Agnelli, A., Gobran, G.R., Corti, G., 2013. Changes induced by the roots of *Erica arborea* L. to create a suitable environment in a soil developed from alkaline and fine-texture marine sediments. *Plant Soil* 368, 297–313.
- Corti, G., Cocco, S., Basili, M., Cioci, C., Warburton, J., Agnelli, A., 2012. Soil formation in kettle holes from high altitudes in central Apennines, Italy. *Geoderma* 170, 280–294.

1122
1123
1124 de Armas-Ricard, M., Orlando, J., Bustamante, R., Carù, M., 2016. Microbial communities of bulk
1125 and *Eschscholzia californica* rhizosphere soils at two altitudes in Central Chile. J. Soil Sci. Plant
1126 Nutr. 16, 1–13.
1127
1128
1129
1130 De Feudis, M., Cardelli, V., Massaccesi, L., Bol, R., Willbold, S., Cocco, S., Corti, G., Agnelli, A.,
1131 2016. Effect of beech (*Fagus sylvatica* L.) rhizosphere on phosphorous availability in soils at
1132 different altitudes (central Italy). Geoderma 276, 53–63.
1133
1134
1135
1136
1137 Delarue, F., Gogo, S., Buttler, A., Bragazza, L., Jassey, V.E.J., Bernard, G., Laggoun-Défarage, F.,
1138 2014. Indirect effects of experimental warming on dissolved organic carbon content in subsurface
1139 peat. J. Soils Sediments 14, 1800–1805.
1140
1141
1142
1143 Dieleman, W.I.J., Venter, M., Ramachandra, A., Krockenberger, A.K., Bird, M.I., 2013. Soil carbon
1144 stocks vary predictably with altitude in tropical forests: Implications for soil carbon storage.
1145 Geoderma 204–205, 59–67.
1146
1147
1148
1149 Fernández Sanjurjo, M.J., Corti, G., Certini, G., Ugolini, F.C., 2003. Pedogenesis induced by *Genista*
1150 *aetnensis* (Biv.) DC. on basaltic pyroclastic deposits at different altitudes, Mt. Etna, Italy. Geoderma
1151 115, 223–243.
1152
1153
1154
1155
1156 Frossard, E., Brossard, M., Hedley, M.J., Metherell, A., 1995. Reactions controlling the cycling of P
1157 in soils. In: Tiessen, H. (Ed.) Phosphorus in the Global Environment. John Wiley & Sons Ltd.,
1158 Chichester, pp. 107–137.
1159
1160
1161
1162 Fujii, K., Aoki, M., Kitayama, K., 2012. Biodegradation of low molecular weight organic acids in
1163 rhizosphere soils from a tropical montane rain forest. Soil Biol. Biochem. 47, 142–148.
1164
1165
1166
1167 Girardin, C.A.J., Aragão, L.E.O.C., Malhi, Y., Huaraca Huasco, W., Metcalfe, D.B., Durand, L.,
1168 Mamani, M., Silva-Espejo, J.E., Whittaker, R.J., 2013. Fine root dynamics along an elevational
1169 gradient in tropical Amazonian and Andean forests. Glob. Biogeochem. Cycles 27, 252–264.
1170
1171
1172
1173 Graefe, S., Hertel, D., Leuschner, C., 2008. Fine root dynamics along a 2,000-m elevation transect in
1174 South Ecuadorian mountain rainforests. Plant Soil 313, 155–166.
1175
1176
1177
1178
1179
1180

1181
1182
1183 Graham, H.D., 1992. Stabilization of the Prussian blue color in the determination of polyphenols. J.
1184 Agric. Food Chem. 40, 801–805.
1185
1186
1187 Gunina, A., Kuzyakov, Y., 2015. Sugars in soil and sweets for microorganisms: Review of origin,
1188 content, composition and fate. Soil Biol. Biochem. 90, 87–100.
1189
1190
1191 Guo, H., Cui, H., Jin, H., Yan, Z., Ding, L., Qin, B., 2015. Potential allelochemicals in root zone soils
1192 of *Stellera chamaejasme* L. and variations at different geographical growing sites. Plant Growth
1193 Regul. 77, 335–342.
1194
1195
1196
1197
1198 Gutiérrez-Girón, A., Díaz-Pinés, E., Rubio, A., Gavilán, R.G., 2015. Both altitude and vegetation
1199 affect temperature sensitivity of soil organic matter decomposition in Mediterranean high mountain
1200 soils. Geoderma 237–238, 1–8.
1201
1202
1203
1204 Hagerman, A.E., 2002. The tannin handbook. Miami University, Oxford OH 45056.
1205
1206
1207 Hammer, Ø., Harper, D., 2006. Paleontological data analysis. Blackwell Publishing, Oxford.
1208
1209 Hassouna, M, Massiani, C., Dudal, Y., Pech, N., Theraulaz, F., 2010. Changes in water extractable
1210 organic matter (WEOM) in a calcareous soil under field conditions with time and soil depth.
1211 Geoderma 155, 75–85.
1212
1213
1214
1215 Hertkorn, N., Benner, R., Frommberger, M., Schmitt-Kopplin, P., Witt, M., Kaiser, K., Kettrup, A.,
1216 Hedges, J.I., 2006. Characterization of a major refractory component of marine dissolved organic
1217 matter. Geochim. Cosmochim. Acta 70, 2990–3010.
1218
1219
1220
1221 Hertkorn, N., Frommberger, M., Witt, M., Koch, B.P., Schmitt-Kopplin, Ph., Perdue, E.M., 2008.
1222 Natural organic matter and the event horizon of mass spectrometry. Anal. Chem. 80, 8908–8919.
1223
1224
1225 Hertkorn, N., Ruecker, C., Meringer, M., Gugisch, R., Frommberger, M., Perdue, E.M., Witt, M.,
1226 Schmitt-Kopplin, P., 2007. High-precision frequency measurements: indispensable tools at the core
1227 of the molecular-level analysis of complex systems. Anal. Bioanal. Chem. 389, 1311-1327.
1228
1229
1230
1231 Hinsinger, P., Bengough, G., Vetterlein, D., Young, I.M., 2009. Rhizosphere: biophysics,
1232 biogeochemistry and ecological relevance. Plant Soil 321, 117–152.
1233
1234
1235
1236
1237
1238
1239

1240
1241
1242 Hofman, J., Dušek, L., 2003. Biochemical analysis of soil organic matter and microbial biomass
1243 composition - a pilot study. *Eur. J. Soil Biol.* 39, 217–224.
1244
1245
1246 Hofmann, D., Preuss, G., Mätzler, C., 2015. Evidence for biological shaping of hair ice.
1247 *Biogeosciences* 12, 4261–4273.
1248
1249
1250 IPCC, 2013. *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to*
1251 *the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* [Stocker, T.F., Qin,
1252 D., Plattner, G.-K., Tignor, M., Allen, S.K., Boschung, J., Nauels, A., Xia, Y., Bex, V., Midgley,
1253 P.M. (Eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.
1254
1255
1256
1257
1258
1259 Jones, D.L., Nguyen, C., Finlay, R.D., 2009. Carbon flow in the rhizosphere: carbon trading at the
1260 soil-root interface. *Plant Soil* 321, 5–33.
1261
1262
1263 Jones, D.L., Willett, V.B., 2006. Experimental evaluation of methods to quantify dissolved organic
1264 nitrogen (DON) and dissolved organic carbon (DOC) in soil. *Soil Biol. Biochem.* 38, 991-999.
1265
1266
1267
1268 Kaiser, K., Kalbitz, K., 2012. Cycling downwards - dissolved organic matter in soils. *Soil Biol.*
1269 *Biochem.* 52, 29–32.
1270
1271
1272 Kaiser, M., Kleber, M., Berhe, A.A., 2015. How air-drying and rewetting modify soil organic matter
1273 characteristics: An assessment to improve data interpretation and inference. *Soil Biol. Biochem.* 80,
1274 324–340.
1275
1276
1277
1278 Kalbitz, K., Kaiser, K., 2007. Contribution of dissolved organic matter to carbon storage in forest
1279 mineral soils. *J. Plant Nutr. Soil Sci.* 170, 52–60.
1280
1281
1282
1283 Kim, S., Kramer, R.W., Hatcher, P.G., 2003. Graphical method of analysis of ultrahigh-resolution
1284 broadband mass spectrometry of natural organic matter, the van krevelen diagram. *Anal. Chem.* 75,
1285 5336–5344.
1286
1287
1288
1289 Kögel-Knaber, I., 2002. The macromolecular organic composition of plant and microbial residues as
1290 inputs to soil organic matter. *Soil Biol. Biochem.* 34, 139–162.
1291
1292
1293 Kraus, T.E.C., Dahlgren, R.A., Zasoski, R.J., 2003. Tannins in nutrient dynamics of forest ecosystems
1294 - a review. *Plant Soil* 256, 41–66.
1295
1296
1297
1298

1299
1300
1301 Kuzyakov, Y., 2002. Review: factors affecting rhizosphere priming effects. *J. Plant Nutr. Soil Sci.*
1302 165, 382–396.
1303
1304
1305 Lynch, J.M., Whipps, J.M., 1990. Substrate flow in the rhizosphere. *Plant Soil* 129, 1–10.
1306
1307 Massaccesi, L., Benucci, G.M.N., Gigliotti, G., Cocco, S., Corti, G., Agnelli, A., 2015. Rhizosphere
1308 effect of three plant species of environment under periglacial conditions (Majella Massif, central
1309 Italy). *Soil Biol. Biochem.* 89, 184–195.
1310
1311
1312
1313
1314 McDowell, W.H., 2003. Dissolved organic matter in soils-future directions and unanswered
1315 questions. *Geoderma* 113, 179–186.
1316
1317
1318 Montanari, C., Bargossi, E., Gardini, A., Lanciotti, R., Magnani, R., Gardini, F., Tabanelli, G., 2016.
1319 Correlation between volatile profiles of Italian fermented sausages and their size and starter culture.
1320 *Food Chem.* 192, 736–744.
1321
1322
1323
1324 Nannipieri, P., Giagnoni, L., Landi, L., Renella, G., 2011. Role of phosphatase enzymes in soil. In:
1325 Bunemann, E.K., Obreson, A., Frossard, E. (Eds.) *Phosphorus in action*. Springer, Berlin, pp. 215–
1326 243.
1327
1328
1329
1330
1331 Nelson, D.W., Sommers, L.E., 1996. Total carbon, organic carbon, and organic matter. In: Sparks,
1332 D.L. (Ed.) *Methods of Soil Analysis, Part 3. Chemical Methods*. SSSA and ASA, Madison, pp. 961–
1333 1010.
1334
1335
1336
1337 Nelson, N.O., Janke, R.R., 2007. Phosphorus sources and management in organic production systems.
1338 *HortTechnology* 17, 442–454.
1339
1340
1341
1342 Ohno, T., He, Z.Q., Sleighter, R.L., Honeycutt, C.W., Hatcher, P.G., 2010. Ultrahigh resolution mass
1343 spectrometry and indicator species analysis to identify marker components of soil- and plant
1344 biomass-derived organic matter fractions. *Environ. Sci. Technol.* 44, 8594–8600.
1345
1346
1347
1348 Olsen, S.R., Cole, C.V., Watanabe, F.S., Dean, L.A., 1954. Estimation of available phosphorus in
1349 soils by extraction with sodium bicarbonate. US Department of Agriculture, Circ. 939, Washington.
1350
1351
1352
1353 Pandey, A., Palni, L.M.S., 2007. The rhizosphere effect in trees of the Indian Central Himalaya with
1354 special reference to altitude. *Appl. Ecol. Environ. Res.* 5, 93–102.
1355
1356
1357


1358
1359
1360 Phillips, R.P., Finzi, A.C., Bernhardt, E.S., 2011. Enhanced root exudation induces microbial
1361 feedbacks to N cycling in a pine forest under long-term CO₂ fumigation. *Ecol. Lett.* 14, 187–194.
1362
1363
1364 Prietzel, J., Christophel, D., 2014. Organic carbon stocks in forest soils of German Alps. *Geoderma*
1365 221–222, 28–39.
1366
1367
1368 R Core Team, 2014. R: a language and environment 538 for statistical computing. R Foundation for
1369 Statistical Computing, Vienna, Austria. URL <http://www.R-project.org/>.
1370
1371
1372
1373 Richter, D.B., Oh, N.-H., Fimmen, R., Jackson, J., 2007. The rhizosphere and Soil Formation. In:
1374 Cardon, Z.G., Whitbeck, J.L. (Eds.) *The Rhizosphere: An Ecological Perspective*. Elsevier,
1375 Burlington, pp. 179–200.
1376
1377
1378 Riebe, C.S., Kirchner, J.W., Finkel, R.C., 2004. Sharp decrease in long-term chemical weathering
1379 rates along an altitudinal transect. *Earth Planet. Sci. Lett.* 218, 421–434.
1380
1381
1382
1383 Roth, V.-N., Dittmar, T., Gaupp, R., Gleixner, G., 2015. The molecular composition of dissolved
1384 organic matter in forest soils as a function of pH and temperature. *PLoS ONE* 10, e0119188.
1385
1386
1387 Rumpel, C., Kögel-Knaber, I., Bruhn, F., 2002. Vertical distribution, age, and chemical composition
1388 of organic carbon in two forest soils of different pedogenesis. *Org. Geochem.* 33, 1131–1142.
1389
1390
1391
1392 Santana, C.M., Ferrera, Z.S., Torres Padrón, M.E., Santana Rodríguez, J.J., 2009. Methodologies for
1393 the extraction of phenolic compounds from environmental samples: new approaches. *Molecules* 14,
1394 298–320.
1395
1396
1397
1398 Schoeneberger, P.J., Wysocki, D.A., Benham, E.C., Soil Survey Staff. 2012. Field book for
1399 describing and sampling soils, Version 3.0. Natural Resources Conservation Service, National Soil
1400 Survey Center, Lincoln, NE.
1401
1402
1403
1404 Sipler, R., Seitzinger, S., 2008. Use of electrospray ionization (ESI) mass spectrometry to investigate
1405 complex dissolved organic matter (DOM) and its potential applications in phytoplankton research.
1406
1407
1408
1409 *Harmful Algae* 8, 182–187.
1410
1411
1412
1413
1414
1415
1416

1417
1418
1419 Sleighter, R.L., Hatcher, P.G., 2007. The application of electrospray ionization coupled to ultrahigh
1420 resolution mass spectrometry for the molecular characterization of natural organic matter. *J. Mass*
1421 *Spectrom.* 42, 559–574.
1422
1423
1424
1425 Smolander, A., Kitunen, V., 2002. Soil microbial activities and characteristics of dissolved organic
1426 C and N in relation to tree species. *Soil Biol. Biochem.* 34, 651–660.
1427
1428
1429 Soil Survey Staff, 2014. Keys to soil taxonomy, 12th edition. USDA–Natural Resources Conservation
1430 Service, Washington.
1431
1432
1433
1434 Sokolova, T.A., 2015. Specificity of soil properties in the rhizosphere: Analysis of literature data.
1435 *Eurasian Soil Sci.* 48, 968–980.
1436
1437
1438 Stevens, G.N., Jones, R.H., Mitchell, R.J., 2002. Rapid fine root disappearance in a pine woodland:
1439 a substantial carbon flux. *Can. J. For. Res.* 32, 2225–2230.
1440
1441
1442 Stubbins, A., Dittmar, T., 2015. Illuminating the deep: molecular signatures of photochemical
1443 alteration of dissolved organic matter from North Atlantic deep water. *Mar. Chem.* 177, 318–324.
1444
1445
1446 Stubbins, A., Niggemann, J., Dittmar, T., 2012. Photo-lability of deep ocean dissolved black carbon.
1447 *Biogeosciences* 9, 1661–1670.
1448
1449
1450 Takeda, A., Tsukada, H., Takaku, Y., Hisamatsu, S., 2009. Fractionation of metal complexes with
1451 dissolved organic matter in a rhizosphere soil solution of a humus-rich Andosol using size exclusion
1452 chromatography with inductively coupled plasma-mass spectrometry. *Soil Sci. Plant Nutr.* 55, 349–
1453 357.
1454
1455
1456
1457
1458
1459 Tashi, S., Singh, B., Keitel, C., Adams, M., 2016. Soil carbon and nitrogen stocks in forests along an
1460 altitudinal gradient in the eastern Himalayas and a meta-analysis of global data. *Glob. Change Biol.*
1461 6, 2255–2268.
1462
1463
1464
1465
1466 Trumbore, S.E., Gaudinki, J.B., 2003. The secret lives of roots. *Science* 302, 1344–1345.
1467
1468
1469 Ultra, V.U., Jr., Han, S.-H., Kim, D.-H., 2013. Soil properties and microbial functional structure in
1470 the rhizosphere of *Pinus densiflora* (S. and Z.) exposed to elevated atmospheric temperature and
1471 carbon dioxide. *J. For. Res.* 18, 149–158.
1472
1473
1474
1475

1476
1477
1478 Venables, W.N., Ripley, B.D., 2002. Modern applied statistics with S. 4th Edn. Springer, New York.
1479
1480 Vincent, A.G., Sundqvist, M.K., Wardle, D.A., Giesler, R., 2014. Bioavailable soil phosphorus
1481 decreases with increasing elevation in a Subarctic Tundra Landscape. PLoS ONE 9, e92942.
1482
1483
1484 Vogt, K.A., Vogt, D.J., Palmiotto, P.A., Boon, P., OHara, J., Asbjornsen, H., 1996. Review of root
1485 dynamics in forest ecosystems grouped by climate, climatic forest type and species. Plant Soil 187,
1486 159–219.
1487
1488
1489
1490 Williams, J.S., Dungait, J.A.J., Bol, R., Abbott, G.D., 2016. Contrasting temperature responses of
1491 dissolved organic carbon and phenols leached from soils. Plant Soil 399, 13–27.
1492
1493
1494 Xu, G., Chen, J., Berninger, F., Pumpanen, J., Bai, J., Yu, L., Duan, B., 2015. Labile, recalcitrant,
1495 microbial carbon and nitrogen and the microbial community composition at two *Abies*
1496 *faxoniana* forest elevations under elevated temperatures. Soil Biol. Biochem. 91, 1–13.
1497
1498
1499
1500 Xu, M., Li, X., Cai, X., Gai, J., Li, X., Christie, P., Zhang, J., 2014. Soil microbial community
1501 structure and activity along a montane elevational gradient on the Tibetan Plateau. Eur. J. Soil Biol.
1502 64, 6–14.
1503
1504
1505
1506
1507 Yimer, F., Ledin, S., Abdelkadir, A., 2006. Soil organic carbon and total nitrogen stocks as affected
1508 by topographic aspect and vegetation in the Bale Mountains, Ethiopia. Geoderma 135, 335–344.
1509
1510
1511
1512 Yin, H., Li, Y., Xiao, J., Xu, Z., Cheng, X., Liu, Q., 2013a. Enhanced root exudation stimulates soil
1513 nitrogen transformations in a subalpine coniferous forest under experimental warming. Glob.
1514 Chang. Biol. 19, 2158–2167.
1515
1516
1517
1518 Yin, H., Xiao, J., Li, Y., Chen, Z., Cheng, X., Zhao, C., Liu, Q., 2013b. Warming effects on root
1519 morphological and physiological traits: The potential consequences on soil C dynamics as altered
1520 root exudation. Agric. For. Meteorol. 180, 287–296.
1521
1522
1523
1524 Zianis, D., Mencuccini, M., 2005. Aboveground net primary productivity of beech (*Fagus moesiaca*)
1525 forest: a case study of Naousa forest, northern Greece. Tree Physiol. 25, 713–722.
1526
1527
1528
1529 Zsolnay, A., 2003. Dissolved organic matter: artefacts, definitions, and functions. Geoderma 113,
1530 187–209.
1531
1532
1533
1534

Figure caption

Figure 1. Variation of the properties of the water-extractable organic matter (WEOM) of rhizosphere and bulk from soils under European beech (*Fagus sylvatica* L.) forest at two altitudes (800 and 1000 m), as analysed by principal component analysis (PCA) using standardized data. Central Apennines, Italy. Also the 95% confidence ellipses are drawn.



- Bulk soil 800 m above sea level
- Rhizosphere 800 m above sea level
- ▲ Bulk soil 1000 m above sea level
- ▲ Rhizosphere 1000 m above sea level

Table 1. Main properties of rhizosphere and bulk from the soils under European beech (*Fagus sylvatica* L.) forest at two altitudes (800 and 1000 m). Central Apennines, Italy. Numbers in parentheses are the standard errors (n=3). For each line, mean values with different letters significantly differ for $P < 0.05$.

	800 m above sea level		1000 m above sea level	
	Rhizosphere	Bulk soil	Rhizosphere	Bulk soil
pH _{H₂O}	6.69 (0.29) b	6.77 (0.20) b	6.93 (0.14) ab	7.34 (0.15) a
pH _{KCl}	5.98 (0.27) b	5.99 (0.18) b	6.51 (0.13) ab	6.60 (0.12) a
Total organic C (g kg ⁻¹)	81.11 (5.31) b	68.52 (4.22) c	119.66 (6.25) a	91.69 (8.98) b
Total N (g kg ⁻¹)	5.44 (0.58) b	4.36 (0.48) b	10.75 (0.85) a	8.76 (0.93) a
Available P (mg kg ⁻¹)	26.07 (1.92) a	17.29 (1.39) b	29.38 (0.75) a	20.09 (0.69) b
Total phenols-to-total organic C (mg gallic acid equivalent g ⁻¹ TOC)	4.53 (0.27) a	4.59 (0.26) a	2.87 (0.13) b	3.10 (0.29) b

Table 2. Relative proportions of carbon containing functional groups as detected by CPMAS ¹³C NMR analysis in rhizosphere and bulk from the soils under European beech (*Fagus sylvatica* L.) forest at two altitudes (800 and 1000 m). Central Apennines, Italy. Numbers in parentheses are the standard errors (n=3). Mean values with different letters significantly differ for $P < 0.05$.

Groups	800 m above sea level		1000 m above sea level	
	Rhizosphere	Bulk soil	Rhizosphere	Bulk soil
	% of spectral area			
Alkyl-C	20.3 (0.5) d	20.7 (0.4) d	17.5 (0.4) e	17.0 (0.4) ef
N-Alkyl-C	13.5 (0.6) gh	13.9 (0.4) g	12.4 (0.3) gh	11.9 (0.4) gh
O-Alkyl-C	34.4 (1.2) a	32.7 (0.5) a	30.9 (0.2) b	30.4 (0.4) b
Aryl-C	19.8 (0.4) de	19.8 (0.4) de	24.6 (0.6) c	25.3 (0.7) c
Carboxyl-C	11.0 (1.0) h	11.9 (0.7) gh	12.6 (0.5) gh	13.1 (0.7) gh
Carbonyl-C	1.1 (0.2) k	1.1 (0.2) k	2.0 (0.2) jk	2.4 (0.5) j

Table 3. Concentrations of water soluble organic C (WEOC), hexose and pentose sugars, and phenols of rhizosphere and bulk from the soils under European beech (*Fagus sylvatica* L.) forest at two altitudes (800 and 1000 m). Central Apennines, Italy. Numbers in parentheses are the standard errors (n=3). For each line, mean values with different letters significantly differ for $P < 0.05$.

	800 m above sea level		1000 m above sea level	
	Rhizosphere	Bulk soil	Rhizosphere	Bulk soil
WEOC (g kg ⁻¹)	0.36 (0.04) ab	0.30 (0.02) b	0.44 (0.03) a	0.30 (0.04) b
Hexose sugars (g glucose-C kg ⁻¹ soil)	0.10 (0.02) a	0.04 (0.01) b	0.09 (0.01) a	0.05 (0.01) b
Pentose sugars (g ribose-C kg ⁻¹ soil)	0.05 (0.01) a	0.03 (0.01) b	0.06 (0.00) a	0.02 (0.00) b
Phenols (μg gallic acid equivalent g ⁻¹ soil)	10.81 (0.66) b	11.68 (1.98) b	19.95 (1.94) a	10.12 (1.30) b

Table 4. Number of CHO formulas as detected by ESI FT-ICR MS analysis, and relative proportion of lignins, tannins, condensed aromatics, and sugars of water-extractable organic matter (WEOM) of rhizosphere and bulk from the soils under European beech (*Fagus sylvatica* L.) forest at two altitudes (800 and 1000 m). Central Apennines, Italy. Numbers in parentheses are the standard errors (n=3). For each column, mean values with different letters significantly differ for $P < 0.05$.

	N° of formulas	Lignins	Tannins	Condensed aromatics	Sugars	Others
		%				
<i>800 m above sea level</i>						
Rhizosphere	3403 (824) a	54.7 (6.2) a	26.6 (1.8) ab	6.4 (0.5) a	< 1	12.4
Bulk soil	4114 (926) a	67.5 (6.4) a	20.2 (5.0) b	5.8 (2.3) ab	< 1	6.4
<i>1000 m above sea level</i>						
Rhizosphere	2796 (94) a	56.1 (0.3) a	29.6 (1.5) a	5.7 (0.1) a	< 1	8.6
Bulk soil	3928 (249) a	61.8 (6.2) a	24.6 (1.3) b	2.3 (0.4) b	< 1	11.3

Table 5. Contributes of variables (%) and their correlation coefficients with PC1 and PC2 of the water-extractable organic matter (WEOM) from the soils under European beech (*Fagus sylvatica* L.) forest at two altitudes (800 and 1000 m). Central Apennines, Italy.

Chemical properties	PC1		PC2	
	%	Correlation coefficient	%	Correlation coefficient
Water-extractable organic C	17.96	0.71	3.65	-0.27
Pentoses	29.03	0.91	0.06	0.03
Hexoses	23.99	0.83	5.36	-0.33
Phenols	6.52	0.43	14.41	0.54
Lignins	6.40	-0.43	31.57	0.80
Tannins	13.84	0.63	15.14	0.55
Condensed aromatics	2.27	0.25	29.81	0.78

Supplementary materials

Altitude affects the quality of the water-extractable organic matter (WEOM) from rhizosphere and bulk soil in European beech forests

De Feudis, M.^{a*}, Cardelli, V.^b, Massaccesi, L.^a, Hofmann, D.^c, Berns, A.E.^c, Bol, R.^c, Cocco, S.^b, Corti, G.^b, Agnelli, A.^a

^aDipartimento di Scienze Agrarie, Alimentari e Ambientali, Università degli Studi di Perugia, Perugia, Italy

^bDipartimento di Scienze Agrarie, Alimentari e Ambientali, Università Politecnica delle Marche, Ancona, Italy

^cInstitute for Bio- and Geosciences, IBG-3 Agrosphere, Forschungszentrum Jülich GmbH, 52425 Jülich, Germany

**Corresponding author:*

Mauro De Feudis

Dipartimento di Scienze Agrarie, Alimentari ed Ambientali

Università degli Studi di Perugia

Borgo XX Giugno 72

06121 Perugia, Italy

e-mail: mauro.defeudis@studenti.unipg.it



Figure S1. Map of Italy showing the location of the three study areas (Mt. Terminillo, Mt. San Vicino, and Mt. Acuto). Central Apennines, Italy.




Figure S2. CPMAS ¹³C NMR spectra of rhizosphere and bulk from the soils under European beech (*Fagus sylvatica* L.) forest at two altitudes (800 and 1000 m). Central Apennines, Italy.




Figure S3. van Krevelen diagrams of H/C versus O/C molar ratios for the CHO compounds detected by the ESI FT-ICR MS in the water-extractable organic matter (WEOM) of rhizosphere and bulk from soils under European beech (*Fagus sylvatica* L.) forest at two altitudes (800 and 1000 m). Central Apennines, Italy. Boxes overlain on the plots indicate the major biopolymer classes [lignins (L), tannins (T), condensed aromatics (CA), sugars (S)], whereas colors indicate relative mass peak intensity.

Table S1. General information and features for the soils under European beech (*Fagus sylvatica* L.) forest at two altitudes (800 and 1000 m). Central Apennines, Italy. For symbols see legend.

Horizon	Depth cm	Colour ^a	Structure ^b	Roots ^c	Skeleton %
---------	-------------	---------------------	------------------------	--------------------	---------------

Mount Terminillo. Parent rock: grey limestone with small flintstone layers. Forest management: coppice in conversion to timber forest. Exposure: N. Mean annual precipitation: 1248 mm. Soil temperature regime: *mesic*. Soil moisture regime: *udic*.

Altitude: 800 m

Slope: 40°; Mean annual air temperature: 10.0°C; Mean temperature of the coldest month (January): 1.6°C; Mean temperature of the warmest months (July and August): 19.4°C. Vegetation: *Fagus sylvatica* L. (95%), *Acer* spp., *Laburnum anagyroides* Medik..

Soil: loamy-skeletal, mixed, mesic, Typic Hapludoll (Soil Survey Staff, 2014)

Forest floor	9-0	-	-	0	0
A	0-8	2.5YR 2.5/1	3f,m sbk, fr	3mi,vf,f,m,co	40
AB	8-16	2.5YR	2m pl→2f,m sbk, fr	3mi,vf,f,m,co	10
Bw1	16-23	5YR 3/4	2m pl→2f,m sbk, fr	2mi,vf,f; 3m,co	10
Bw2	23-33	5YR 4/4	2m pl→2f,m sbk, fr	2mi,vf,f,m,co	70
Bw3	33-66	5YR 5/4	1f,m abk, fr	2mi,vf,f,m,co	70
BC	66-77+	10YR 6/6	sg & 1f abk, fr	1mi,vf,f; 3m,co	70

Soil: loamy, mixed, mesic Lithic Haprendoll (Soil Survey Staff, 2014)

Forest floor	4-0	-	-	0	0
A	0-9	2.5YR 2.5/1	3f,m sbk, fr	3mi,vf,f,m,co	10
AB	9-19	5YR 2.5/2	3f,m sbk+abk, fr	2mi,vf,f,m; 3co	40
R	19-22+	-	-	-	-

Altitude: 1000 m

Slope: 40°; Mean annual air temperature: 9.1°C; Mean temperature of the coldest months (January and February): 1.0°C; Mean temperature of the warmest month (July): 18.3°C. Vegetation: *Fagus sylvatica* L. (99%), *Acer* spp..

Soil: loamy-skeletal, mixed, mesic, Lithic Haprendoll (Soil Survey Staff, 2014)

Forest floor	17-0	-	-	0	0
A	0-11	2.5YR 2.5/1	3f cr+sbk, fr	3mi,vf,f,m,co	20
AB	11-20	5YR 2.5/1	3f abk+sbk, fr	2mi,vf,f; 3m,co	40
BA	20-28	5YR 2.5/2	3f abk, fr	1mi,vf,f; 2m; 3co	50
2Bwb	28-38	5YR 4/3	1f,m abk, fr	1mi,vf,f,m,co	20
2Crb	38-49+	10YR 5/4	1f,m abk, fr	1mi,vf,f,m,co	75

Soil: loamy, mixed, mesic, Lithic Haprendoll (Soil Survey Staff, 2014)

Forest floor	16-0	-	-	0	0
A	0-14	2.5YR 2.5/1	3f,m sbk+abk, fr	3mi,vf,f,m; 2co	5
AB	14-24	5YR 2.5/2	3f,m sbk+abk, fr	2mi,vf,f,m,co	30
Cr	24-45+	10YR 6/4	1m abk, fr	2f; 3m,co	75

Mount San Vicino. Parent rock: grey limestone with marl and flintstone layers. Forest management: coppice in conversion to timber forest. Exposure: N. Mean annual precipitation: 825 mm. Soil temperature regime: *mesic*. Soil moisture regime: *udic*.

Altitude: 800 m

Slope: 35°; Mean annual air temperature: 10.0°C; Mean temperature of the coldest month (January): 1.0°C; Mean temperature of the warmest month (July): 21.0°C. Vegetation: *Fagus sylvatica* L. (90%), *Quercus cerris* L., *Castanea sativa* Mill..

Soil: coarse-loamy, mixed, mesic, Typic Hapludoll (Soil Survey Staff, 2014)

Forest floor	9-0	-	-	0	0
A	0-10	7.5YR 5/2	2ccr	3mi,vf,f,m,co	10
AB	10-16	7.5YR 4/2	2f,m abk+sbk, fr	2mi,mf,f	20
Bw1	16-30	7.5YR 5/4	2f sbk, fr	2mi,vf,f	20
Bw2	30-43	10YR 4/3	2f, cr+sbk, fr	1mi,vf,f,m	20
2Bwb	43-68	10YR 4/2	2f cr, fr	3mi,vf,f; 1m	70
3Bwb	68-81	2.5Y 2/0	1f abk, fr	3mi,vf,f; 2m	70
3Crb	81-90+	7.5YR 4/4	1f abk, fr	1vf,f	90

Soil: coarse-loamy, mixed, mesic, Typic Humudept (Soil Survey Staff, 2014)

Forest floor	7-0	-	-	0	0
A	0-9	7.5YR 5/2	2-3f cr, fr	3mi,vf,f,m	20
AB	9-16	10YR3/2	2f cr, fr	2mi,vf,f,m,co	30
Bw1	16-28	10YR5/4	2f sbk, fr	2mi,vf,f; 1m	40
Bw2	28-37	10YR3/4	3m sbk, fr	2mi,vf; 3f,m	20
2Crb	37-40+	10YR3/2	1f abk, fr	1mi,vf,f; 2m,co	80

Altitude: 1000 m

Slope: 45°; Mean annual air temperature: 9.0°C; Mean temperature of the coldest month (January): 0.0°C; Mean temperature of the warmest month (July): 20.0°C. Vegetation: *Fagus sylvatica* L. (80%), *Sorbus aria* Crantz (10%), *Acer* spp., *Quercus cerris* L..

Soil: loamy-skeletal, mixed, mesic, Lithic Haprendoll (Soil Survey Staff, 2014)

Forest floor	6-0	-	-	0	0
A	0-7	5YR 2.5/2	3f cr, fr	3mi,vf,f,m,co	25
AB	7-16	5YR 3/3	3m,f cr+sbk, fr	3mi,vf,f,m,co	50
Bw	16-29	7.5YR 3/3	3f cr+sbk, fr	3mi,vf,f,m,co	70
2Bwb	29-40	2.5YR 3/4	3f sbk, fr	2mi,vf,f; 3m,co	25
2BCb	40-47+	10YR 6/5	1m abk, fr	1mi,vf,f; 2m,co	60

Soil: loamy-skeletal, mixed, mesic, Inceptic Haprendoll (Soil Survey Staff, 2014)

Forest floor	4-0	-	-	0	0
A	0-5	5YR 2.5/2	3f sbk+cr, fr	3mi,vf,f,m	25
AB	5-14	5YR 3/2	3f cr+sbk, fr	3mi,vf,f; 2m	50
Bw1	14-36	5YR 3/3	2-3f,m sbk, fr	3mi,vf,f; 1m	70
Bw2	36-43	5YR 3/4	1m sbk, fr	2mi,vf,f; 1m	70
BC	43-52+	10YR 6/5	1m abk, fr	1mi,vf; 2f,m,co	70

Mount Acuto. Parent rock: white limestone with flintstone layers. Forest management: coppice in conversion to timber forest. Exposure: N-NE. Mean annual precipitation: 1430 mm. Soil temperature regime: *mesic*. Soil moisture regime: *udic*.

Altitude: 800 m

Slope: 40°; Mean annual air temperature: 10.0°C; Mean temperature of the coldest month (January): 0.0°C; Mean temperature of the warmest month (July): 21.0°C. Vegetation: *Fagus sylvatica* L. (95%), *Carpinus betulus* L., *Acer opalus* Mill., *Quercus* spp..

Soil: coarse-loamy, mixed, mesic, Typic Humudept (Soil Survey Staff, 2010)

Forest floor	7-0	-	-	0	0
A	0-3	10YR 2/1	2-3f cr, vfr	2mi,vf,f	5
AB	3-9	10YR 4/4	3f,m sbk, fr	2vf,f,m,co	20
Bw1	9-25	10YR 5/4	1f,m abk+sbk, fr	3mi,vf,f 2m,co	30
Bw2	25-4	10YR 5/4	1m sbk+abk, fr	3mi,vf,f,m,co	30
Bw3	44-76	10YR 5/4	1m sbk, fr	2mi,vf,f; 3m,co	30
Bw4	76-99	2.5YR 5/6	1f,m sbk, fr	2mi,vf; 3f; 2m,co	50
Bw5	99-108	10YR 7/6	1m sbk, fr	1mi,fv,f,m,co	70
BC	108-115+	10YR 7/6	sg & 1f sbk, fr	1mi,fv,f,m,co	70

Soil: coarse-loamy, mixed, mesic, Typic Humudept (Soil Survey Staff, 2014)

Forest floor	7-0	-	-	0	0
A	0-3	10YR 2/1	3f cr, fr	3mi,vf,f,m,co	10
AB	3-8	10YR 2/2	3f cr, fr	3mi,vf,f,m,co	10
Bw1	8-25	10YR 4/4	1f sbk, fr	3mi,vf,f,m,co	20
Bw2	25-38	10YR 4/4	1m,c sbk, fr	2mi,vf,f,m,co	30
Bw3	38-54	10YR 5/6	3f,m sbk, fr	2mi,vf,f,m,co	30
Bw4	54-73	10YR 5/4	1f,m sbk, fr	2mi,vf,f; 1m,co	40
Bw5	73-102	10YR 6/6	1f,m sbk+abk, fr	1mi,vf; 2f	30
BC	102-113+	10YR 7/6	sg & 1m sbk, fr	1mi,fv,f,m,co	50

Altitude: 1000 m

Slope: 25°; Mean annual air temperature: 9.0°C; Mean temperature of the coldest month (January): -1.0°C; Mean temperature of the warmest month (July): 20.0°C. Vegetation: *Fagus sylvatica* L. (100%).

Soil: coarse-loamy, mixed, mesic, Inceptic Haprendoll (Soil Survey Staff, 2014)

Forest floor	5-0	-	-	0	0
A	0-11	7.5YR 3/2	3f,m sbk, fr	2mi,vf; 3f,m	10
AB	11-23	7.5YR 2.5/1	2f,m sbk, fr	2mi,vf,f; 3m,co	30
2Ab	23-33	7.5YR 3/2	3f,m sbk, fr	3mi,vf,f; 3m,co	50
2Bwb	33-54	10YR 7/6	1f abk, fr	2mi,vf,f	80
3Ab	54-63	10YR 4/3	1f sbk, fr	2mi,vf,f	60
3Bwb	63-70	10YR 5/3	1f abk, fr	2mi,vf; 1f	70
4Ab	70-71	7.5YR 3/2	3f abk, fr	1mi,vf,f	70
4Bwb	71-76	10YR 5/3	2f,m abk, fr	1mi,vf,f	75
4BC	76-86+	10YR 6/6	1f,m abk, fr	v1mi,vf,f	80

Soil: loamy-skeletal, mixed, mesic, Inceptic Haprendoll (Soil Survey Staff, 2014)

Forest floor	13-0	-	-	0	0
A	0-13	7.5YR 2.5/1	3f,m cr+sbk, fr	3mi,vf,f,m; 2co	40
AB	13-20	10YR 3/2	1f,m sbk+abk, fr	2mi,vf,f,m,co	50
2Ab	20-36	10YR 2/2	1f abk, fr	2mi,vf,f; 3m,co	50

2Bwb1	36-69	10YR 4/4	1f abk, fr	2mi,vf,f; 1m	60
2Bwb2	69-75	10YR 6/6	1f abk, fr	1mi,vf	60
2BC2	75-85+	10YR 7/6	1f abk, fr	1mi,vf	75

^amoist and crushed, according to the Munsell Soil Color Charts.

^bsg=single grain; 1=weak, 2=moderate, 3=strong; f=fine, m=medium, c=coarse; cr=crumb, abk=angular blocky, sbk=subangular blocky; vfr=very friable, fr=friable.

^c0=absent, v1=very few, 1=few, 2=plentiful, 3=abundant; mi=micro, vf=very fine, f=fine, m=medium, co=coarse.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

HIGHLIGHTS

- Beech rhizosphere effect on WEOM was evaluated at 800 m and 1000 m a.s.l.
- Rhizosphere WEOM had larger amounts of sugars than that of bulk soil
- Organic C and phenols contents were higher in rhizosphere than in bulk soil WEOM at 1000 m
- Higher tannins diversity occurred in rhizosphere than in bulk soil WEOM at 1000 m
- Environmental constraints enhanced the rhizosphere effect on WEOM quality