



UNIVERSITÀ POLITECNICA DELLE MARCHE
Repository ISTITUZIONALE

Oxidative and interactive challenge of cadmium and ocean acidification on the smooth scallop *Flexopecten glaber*.

This is the peer reviewed version of the following article:

Original

Oxidative and interactive challenge of cadmium and ocean acidification on the smooth scallop *Flexopecten glaber* / Nardi, A.; Benedetti, M.; Fattorini, D.; Regoli, F.. - In: AQUATIC TOXICOLOGY. - ISSN 0166-445X. - STAMPA. - 196:(2018), pp. 53-60. [10.1016/j.aquatox.2018.01.008]

Availability:

This version is available at: 11566/253723 since: 2022-06-01T13:00:07Z

Publisher:

Published

DOI:10.1016/j.aquatox.2018.01.008

Terms of use:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. The use of copyrighted works requires the consent of the rights' holder (author or publisher). Works made available under a Creative Commons license or a Publisher's custom-made license can be used according to the terms and conditions contained therein. See editor's website for further information and terms and conditions.

This item was downloaded from IRIS Università Politecnica delle Marche (<https://iris.univpm.it>). When citing, please refer to the published version.

(Article begins on next page)

1 **Oxidative and interactive challenge of cadmium and ocean acidification on the smooth scallop**

2 *Flexopecten glaber*

3

4

5

6 Alessandro Nardi^a, Maura Benedetti^{a,b}, Daniele Fattorini^a and Francesco Regoli^{a,b*}

7

8

9 ^aDepartment of Life and Environmental Sciences (DiSVA), Polytechnic University of Marche,
10 Ancona, Italy

11 ^b CoNISMa, Consorzio Interuniversitario per le Scienze del Mare, Roma, Italy

12

13

14

15

16 *Corresponding author: Prof. Francesco Regoli

17 Dipartimento di Scienze della Vita e dell'Ambiente (DiSVA),

18 Università Politecnica delle Marche,

19 Via Brecce Bianche 60131, Ancona, Italy

20 Phone: +39 0712204613

21 Fax: +39 0712204609;

22 e-mail: f.regoli@univpm.it

23

24

25 **Abstract**

26 Ocean acidification (OA) may affect sensitivity of marine organisms to metal pollution
27 modulating chemical bioavailability, bioaccumulation and biological responsiveness of several
28 cellular pathways. In this study, the smooth scallop *Flexopecten glaber* was exposed to various
29 combinations of reduced pH (pH/pCO₂ 7.4/~3000 µatm) and Cd (20 µg/L). The analyses on cadmium
30 uptake were integrated with those of a wide battery of biomarkers including metallothioneins, single
31 antioxidant defenses and total oxyradical scavenging capacity in digestive gland and gills, lysosomal
32 membrane stability and onset of genotoxic damage in haemocytes. Reduced pH slightly enhanced
33 concentration of Cd in scallops tissues, but no effects were measured in terms of metallothioneins.
34 Induction of some antioxidants by Cd and/or low pH in the digestive gland was not reflected in
35 variations of the total oxyradical scavenging capacity, while the investigated stressors caused a certain
36 inhibition of antioxidants and reduction of the scavenging capacity toward peroxy radical in the gills.
37 Lysosomal membrane stability and onset of genotoxic damages showed high sensitivity with possible
38 synergistic effects of the investigated factors. The overall results suggest that indirect effects of ocean
39 acidification on metals accumulation and toxicity are tissue-specific and modulate oxidative balance
40 through different mechanisms.

41

42 **Keywords:** Oxidative stress; ocean acidification; metal contamination; bioaccumulation; biomarkers;
43 scallops

44 1. Introduction

45 World oceans have absorbed about the 30% of anthropogenic emissions of carbon dioxide (CO₂)
46 in the atmosphere causing changes in the inorganic carbon system equilibrium (Le Quéré *et al.*, 2009).
47 The consequent ocean acidification (OA) is responsible for the continuous reduction of ocean pH,
48 dropped by 0.1 units since the beginning of industrial era (Gattuso and Lavigne, 2009), and expected
49 to further decrease by 0.14 to 0.35 units depending on CO₂ emissions scenarios (Caldeira and Wickett,
50 2005). Scientific literature provides wide evidence that future projections of ocean pH/pCO₂ will
51 affect health status of marine organisms by altering key physiological processes, like calcification
52 rates (Cerrano *et al.*, 2013; Dupont *et al.*, 2010; Gazeau *et al.*, 2007; Jokiel *et al.*, 2008), acid-base
53 balance and ionic homeostasis (Gutowska *et al.*, 2010; Miles *et al.*, 2007; Spicer *et al.*, 2007),
54 metabolism (Lannig *et al.*, 2010; Pan *et al.*, 2015; Stumpp *et al.*, 2012), immune response (Bibby *et*
55 *al.*, 2008; Hernroth *et al.*, 2011, 2012, 2016), larval development (Dupont *et al.*, 2008; Ellis *et al.*,
56 2009; Kurihara *et al.*, 2007; Stumpp *et al.*, 2011) and oxidative stress responses (Benedetti *et al.*,
57 2016; Freitas *et al.*, 2016; Nardi *et al.*, 2017; Pimentel *et al.*, 2015; Rokitta *et al.*, 2012; Soriano-
58 Santiago *et al.*, 2012; Tomanek *et al.*, 2011).

59 Beside the direct effects, there is growing interest for the potential interaction of OA with other
60 environmental stressors, such as the high levels of metal contamination in coastal environments
61 (Ivanina and Sokolova, 2015). In this respect, OA is supposed to increase the ionic and bioavailable
62 fraction of certain metals like copper (Cu²⁺), which typically form strong complexes with carbonate
63 (CO₃²⁻) and hydroxide (OH⁻) ions (Millero *et al.*, 2009). These model predictions have been
64 confirmed by some experimental evidence revealing that high pCO₂/low pH regimes can increase
65 release of metals from polluted sediments (Ardelan *et al.*, 2009; Ardelan and Steinnes, 2010; de Orte
66 *et al.*, 2014a, 2014b) and enhance their bioaccumulation (Duckworth *et al.*, 2017; Götze *et al.*, 2014;
67 Ivanina *et al.*, 2014; Lacoue-Labarthe *et al.*, 2009, 2011; López *et al.*, 2010; Rodríguez-Romero *et*
68 *al.*, 2014). Synergistic effects of high pCO₂/low pH and metal exposure were recently reported on
69 several cellular responses of marine invertebrates (Götze *et al.*, 2014; Ivanina *et al.*, 2013, 2015;

70 Lewis *et al.*, 2013), including the antioxidant status and the onset of oxidative stress (Benedetti *et al.*,
71 2016; Nardi *et al.*, 2017; Ricevuto *et al.*, 2015, 2016; Siddiqui *et al.*, 2015;), which is one of the most
72 relevant pathways by which trace elements exert their toxicity through a sophisticated array of
73 molecular and cellular effects (Regoli and Giuliani, 2014).

74 To provide new insights on the interactions between ocean acidification and metal
75 contamination, this study investigated whether high $p\text{CO}_2$ /low pH may influence bioaccumulation
76 and sub-lethal effects of cadmium in the smooth scallop *Flexopecten glaber*. This species, widely
77 distributed throughout the Mediterranean Sea, has been recently addressed to be a key commercial
78 species, especially in Northwestern Adriatic Sea where it represents about 74% of shellfish fishery
79 (Marčeta *et al.*, 2016, Mazzoldi *et al.*, 2014; Pujolar *et al.*, 2010). Scallops are widely used in
80 ecotoxicological studies, they are typically characterized by high basal concentrations of cadmium in
81 the digestive gland (Bustamante *et al.*, 2002; Mauri *et al.*, 1990; Regoli *et al.*, 1998, 2000, 2002) and
82 they are recently addressed as highly sensitive to ocean acidification (Andersen *et al.*, 2013; Cooley
83 *et al.*, 2015; Schalkhauser *et al.*, 2013; White *et al.*, 2013). The effects of ocean acidification can be
84 exacerbated in shallow coastal and estuarine waters due to freshwaters inputs, which influence
85 carbonate chemistry, nutrients levels, organic matter degradation and pollutants concentrations
86 (Nikinmaa, 2013; Wallace *et al.*, 2014; Wong *et al.*, 2014). In this respect, scallops were exposed to
87 various combinations of Cd and high $p\text{CO}_2$ /low pH, and a complex network of cellular responses was
88 investigated including levels of metallothioneins, variations of antioxidant defenses and total
89 antioxidant capacity in both digestive gland and gills, lysosomal alterations and onset of genotoxic
90 damages in haemocytes. The overall significance of biomarkers responses was synthesized in a
91 cellular hazard index through a quantitative hazard model (Sediqualsoft) which considers the number
92 and magnitude of observed variations, giving a different weight to each biomarker based on the
93 toxicological relevance of biological endpoints (Benedetti *et al.*, 2012; Piva *et al.*, 2011). Results
94 obtained in the present study were expected to contribute to the growing knowledge on the interactive
95 effects of ocean acidification and metals focusing on sensitivity of different tissues in a widely

96 distributed, potentially vulnerable but still poorly investigated species.

97

98 **2. Materials and Methods**

99 *2.1 Animal collection and experimental design*

100 Scallops, *Flexopecten glaber* (4.5 ± 0.5 cm shell length), were obtained in June 2015 from a
101 shellfish farm in an unpolluted area of Venice lagoon, Chioggia, Italy. Organisms were acclimatized
102 for 7 days in aquaria with aerated artificial seawater (ASW; Instant Ocean®) at local environmental
103 conditions of salinity (30 practical salinity units), temperature (20 °C) and pH_{NBS} (8.20).

104 Scallops were then randomly assigned to one of the following treatments, each containing 20
105 organisms in 20 L: 1) control condition (CTRL), at 20°C, pH=8.20/ $p\text{CO}_2 \sim 400$ µatm; 2) cadmium
106 exposure (Cd), 20°C, pH=8.20/ $p\text{CO}_2 \sim 400$ µatm and 20 µg/L cadmium; 3) acidification (A), 20°C,
107 pH=7.40/ $p\text{CO}_2 \sim 3000$ µatm; 4) acidification + Cd (A - Cd), 20°C, pH=7.40/ $p\text{CO}_2 \sim 3000$ µatm and
108 20 µg/L cadmium. Cadmium exposure was representative of a polluted but environmentally realistic
109 scenario (Neff, 2002), while selected target pH was based on scenario RCP 8.5 and the 2014 IPCC
110 WGII AR5 (IPCC, 2014) where future decrease in coastal waters is predicted to be higher than in
111 open ocean; target pH was reached by mixing ASW (pH=8.2) with small amounts of CO₂-saturated
112 ASW as described elsewhere (Nardi *et al.*, 2017). For each experimental condition temperature, pH
113 and salinity were measured daily, while total alkalinity (A_T) was measured every three days during
114 the experiment according to Dickson *et al.*, 2007. Seawater carbonate parameters ($p\text{CO}_2$, and
115 saturation state (Ω) for calcite and aragonite) were calculated in CO2SYS using barometric pressure
116 values (Pierrott *et al.*, 2006); A_T , pH, temperature, salinity values and full seawater chemistry are
117 provided in Table 1. For calculations, we used NBS scale for seawater pH, carbonate constants from
118 Millero (2010), KSO_4^- constant from Dickson *et al.* (2007) and concentration of silicate and phosphate
119 from Instant Ocean® composition (0.21 µmol/kg and 0.05 µmol/kg, respectively). Water was
120 changed every other day, and scallops fed 12 hours prior the water change with a commercial mixture
121 of zooplankton (50-300 µm) for filter-feeding organisms.

122 After ten days, animals were sampled from each tank and tissues collected for chemical and
123 biological analyses. Gills and digestive glands were excised, pooled in 5 samples (each constituted
124 by tissues of 4 individuals), rapidly frozen in liquid nitrogen and maintained at -80°C until analyzed
125 for cadmium content or biomarker responses. Haemolymph was withdrawn from the adductor muscle
126 of 5 specimens and immediately used for the measurement of lysosomal membranes stability and
127 onset of genotoxic damages.

128

129 *2.2 Cadmium determination*

130 Cadmium (Cd) concentrations in scallops were analyzed according to previously described
131 methods (Regoli *et al.*, 2005). For each treatment, digestive glands and gills were dried at 60°C
132 overnight and digested in a microwave digestion system (Mars CEM, CEM Corporation, Matthews
133 NC). Cd was analyzed by atomic absorption spectrophotometry (AAS) using graphite furnace
134 atomization and Zeeman effect (SpectrAA 300 Zeeman, Varian, Mulgrave, VIC, Australia). Quality
135 assurance and quality control was assessed by processing blank samples and reference standard
136 material (Mussel Tissue Standard Reference Material SRM NIST-2977, National Institute of
137 Standards and Technology Gaithersburg, MD, USA). The concentrations obtained for the SRM were
138 always within the 95% confidence interval of certified values. Data are expressed as µg/g dry weight
139 (mean values ± standard deviation, n = 5).

140

141 *2.3 Biomarkers responses*

142 Standardized protocols were used to analyze biomarkers and full methodological details are
143 given in Supplementary Material 1 (SM1). Metallothioneins (MTs), single antioxidant defenses
144 (catalase, glutathione S-transferases, glutathione peroxidases, glutathione reductase activities and
145 total glutathione), total oxyradical scavenging capacity toward peroxy radical (TOSC ROO•) and
146 hydroxyl radical (TOSC HO•) were evaluated in digestive gland and gills. The analysis of the Total
147 Oxyradical Scavenging Capacity (TOSC) is a reliable tool for quantitatively assess the biological

148 resistance to toxicity of different forms of ROS including peroxy radicals, hydroxyl radicals and
149 peroxy nitrite decomposition products (Regoli and Winston, 1998, 1999). The assay is based on the
150 capability of cellular antioxidants to reduce the oxidation of α -keto- γ -methiolbutyric acid (KMBA)
151 in the presence of artificially generated oxyradicals. Compared to individual antioxidants, variations
152 of TOSC have a greater biological relevance and prognostic value, being an impaired capability to
153 neutralize ROS associated to the onset of various forms of oxidative damages like lysosomal
154 dysfunctions, lipid peroxidation and genotoxic effects (Nigro et al., 2002; Camus et al., 2003; Gorbi
155 and Regoli, 2003; Moore et al., 2006). Lysosomal membrane stability (as Neutral Red Retention
156 Time, NRRT) and onset of genotoxic effects as DNA strand breaks (Comet assay) and micronuclei
157 frequency (MN) were analyzed in haemocytes.

158

159 *2.4 Statistical analyses*

160 Analysis of variance (One-way ANOVA) was used to evaluate the effects on all investigated
161 parameters, after ensuring that all data followed the normal distribution (Shapiro-Wilk test) and that
162 variances were homogeneous (Levene's Test). Level of significance was set to $p < 0.05$; *post-hoc*
163 Tukey HSD tests were used to compare group of means. Multivariate principal component analysis
164 (PCA) was applied to visualize the relationships among the different treatments and all statistical
165 analyses were performed using RStudio (version 1.0.143).

166 For each treatment, the whole dataset of biomarkers results was summarized in a hazard index
167 elaborated through weighted criteria which discriminate different endpoints and the magnitude of
168 effects (Sediqualsoft, Piva *et al.*, 2011). Within this quantitative model, each biomarker has a
169 "weight" based on its toxicological relevance and a "threshold" of biological significant changes
170 which consider the possibility of biphasic responses and the different responsiveness among various
171 species and tissues. Variations measured for various biomarkers are compared to their specific
172 thresholds and corrected for the weight and the statistical significance of the difference compared to
173 controls (for full details see Piva et al., 2011, Benedetti et al., 2012): the calculated Hazard Quotient

174 (HQ) does not include biomarkers with variations lower or equal to their threshold, while it averages
175 or adds the summation (Σ) respectively for those biomarkers with variations up to 2-fold or more than
176 2-fold greater than the specific threshold (Avio *et al.*, 2015; Benedetti *et al.*, 2012; Piva *et al.*, 2011;
177 Regoli *et al.*, 2014). The model finally assigns the elaborated HQ in one of five classes of hazard,
178 from Absent to Severe (Piva *et al.*, 2011). Whole calculations and assumptions have been fully given
179 elsewhere (Benedetti *et al.*, 2012; Piva *et al.*, 2011).

180

181 **3. Results**

182 Exposures to Cd caused an increase of metal concentrations in both digestive gland and gills of
183 scallops with a slightly greater, although not statistically significant accumulation in organisms
184 exposed to the metal at lower pH (Fig.1a and 1b).

185 Levels of metallothioneins were not influenced neither by Cd-exposure nor by pH-reduction, in
186 the digestive gland (Fig.2a) or gills (Fig.3a).

187 Sensitivity of antioxidant defenses showed tissue-specificities toward the investigated factors. In
188 the digestive gland, catalase (Fig.2b) was enhanced in organisms exposed to lower pH, an effect
189 which was not further modulated by concomitant presence of Cd; higher activities were observed also
190 for GST activity and Se-dependent GPx in acidic conditions (Fig.2c and 2d), but for these enzymes
191 an antagonistic effect occurred during co-exposure to Cd and high $p\text{CO}_2$ /low pH; total glutathione
192 (Fig.2g) was enhanced in all treatments involving Cd-exposure, pH reduction and their combination.
193 Variations of individual antioxidants were not reflected by any significant difference among the
194 experimental treatments of TOSC against both $\text{ROO}\cdot$ and $\text{HO}\cdot$ (Fig.2h and 2i).

195 Different interactive effects of Cd and acidification were observed on the antioxidant system of
196 gills. The activity of GST was significantly lowered in organisms exposed to the combination of
197 factors (Fig. 3c), while the inhibition of Se-dependent GPx caused by Cd alone was not observed
198 when the metal was dosed at high $p\text{CO}_2$ /low pH (fig. 3d). TOSC values toward $\text{ROO}\cdot$ were

199 significantly reduced by acidification with or without Cd (Fig. 3h), but no variations were observed
200 toward HO• (Fig.3i).

201 Lysosomal membrane stability (Fig.4a) was significantly lowered by the co-exposure to Cd and
202 acidification, no effects appeared in terms of DNA fragmentation (Fig.4b), while MN frequency (Fig.
203 4c) was strongly enhanced in all treatments with Cd and acidification, dosed alone or in combination.

204 The principal component analysis on the whole dataset of biomarkers provided a two-
205 dimensional pattern explaining almost 54.6% of the total variance (Fig.5). A clear separation was
206 observed between control and treated organisms, with a further discrimination between exposure to
207 Cd alone and those involving the acidification with or without the metal. The parameters determining
208 the separation along PC1 axis, i.e. between control pH vs. acidification-exposed organisms, were
209 antioxidants in digestive gland (CAT, GST, GPx and TOSC HO•), in gills (GST, GR and TOSC
210 ROO•), and micronuclei frequency in haemocytes; along the PC2 axis, the separation of Cd-exposure
211 was mostly determined by MTs and TGSH in both the tissues, GPxs in gills and TOSC ROO• in
212 digestive gland.

213 The elaboration of whole biomarkers data with the weighted criteria provided a synthetic hazard
214 quotient (HQ) for organisms of each experimental condition, with a level of hazard classified as
215 “Slight” for scallops exposed to Cd alone and “Moderate” for those exposed to the acidified
216 treatments; among these latter, a quantitatively higher value of HQ was summarized for scallops
217 exposed to acidification alone than in those exposed to cadmium and high $p\text{CO}_2$ /low pH (Fig.6).

218

219 **4. Discussions**

220 This study provided evidence that future projections of pH in coastal waters can modulate
221 bioaccumulation and biological effects of cadmium in the temperate scallop *F. glaber*.

222 The consequences of seawater acidification on metals bioavailability are variable: some authors
223 described increased uptake in marine organisms under reduced pH conditions (Götze *et al.*, 2014;
224 Ivanina *et al.*, 2014; Lacoue-Labarthe *et al.*, 2009, 2011; López *et al.*, 2010; Rodriguez-Romero *et*

225 *al.*, 2014; Shi *et al.*, 2016) while others did not report a similar effect (Benedetti *et al.*, 2016; Ivanina
226 *et al.*, 2016; Nardi *et al.*, 2017; Ricevuto *et al.*, 2016). In the present study, we observed a slight but
227 significant accumulation of cadmium in the digestive gland of scallops exposed to the metal, and a
228 more marked increase in the gills. The limited accumulation of cadmium in digestive gland can be
229 explained by the high basal levels of this metal in *F. glaber*, and more in general in digestive glands
230 of scallops which possess specific high molecular weight Cd-binding proteins (Bustamante *et al.*,
231 2002; Mauri *et al.*, 1990; Regoli *et al.*, 1998, 2000, 2002). When scallops were exposed under
232 hypercapnic condition, a further, slight increase of metal uptake was observed in both the tissues,
233 although this additional effect was not statistically significant. A limited influence of low pH on
234 cadmium accumulation has been reported in the Antarctic scallop *A. colbecki* and in the
235 Mediterranean mussel *M. galloprovincialis* supporting that effects of acidification on metals uptake
236 do not depend on the chemical speciation of the element, but rather reflect physiological effects of
237 CO₂ on an organism, which can not be generalized, being dependent on metal, tissue, species-specific
238 characteristics and duration of exposure (Benedetti *et al.*, 2016; Nardi *et al.*, 2017).

239 Despite cadmium is known to induce metallothioneins, levels of these proteins were not
240 modulated in *F. glaber* by metal-exposure in normocapnic or acidic conditions, neither in digestive
241 gland nor in the gills. Also this result might be related to the specific characteristics of scallops which,
242 beside the elevated tissue concentrations of cadmium and the presence of high molecular weight
243 binding proteins, also contain relatively high basal levels of MTs: these features might influence the
244 low sensitivity of metallothioneins induction pathway as previously observed in the Antarctic scallop
245 *A. colbecki* exposed to cadmium with and without acidification (Benedetti *et al.*, 2016).

246 The characterization of the antioxidant system, integrating the analyses of individual antioxidant
247 defenses with the total oxyradical scavenging capacity toward different ROS, revealed interactions
248 between cadmium and acidification. In the digestive gland, exposure to high *p*CO₂/low pH enhanced
249 the enzymatic activities of catalase, glutathione S-transferases, Se-dependent glutathione peroxidases
250 and levels of total glutathione. Catalase and Se-dependent GPx are responsible for the detoxification

251 of hydrogen peroxide, GST are also involved in metabolism of lipid hydroperoxides while GSH,
252 beside acting as direct scavenger of ROS, is a fundamental co-factor for glutathione-dependent
253 enzymes like GST and GPx (Regoli and Giuliani, 2014). These results suggest that pH reduction
254 promotes oxidative insult through the generation of peroxides, as previously hypothesized for the
255 oysters *Crassostrea virginica* and *Crassostrea gigas*, the polychaete *Sabella spallanzanii*, the scallop
256 *A. colbecki* and the Mediterranean mussel *M. galloprovincialis* (Benedetti *et al.*, 2016; Moreira *et al.*,
257 2016; Nardi *et al.*, 2017; Ricevuto *et al.*, 2016; Tomanek *et al.*, 2011). With the exception of catalase,
258 these inductive effects were typically less evident in organisms co-exposed to the metal and
259 acidification, suggesting some antagonistic interaction and/or a certain Cd-mediated impairment of
260 those antioxidant defenses (Regoli, 2012; Regoli and Giuliani, 2014). Despite the variation of these
261 antioxidants, no effects were observed for the TOSC toward ROO• and HO•, indicating a good
262 counteracting capability toward oxidative challenge in digestive gland of *F. glaber*. The observed
263 inhibitions might be compensated by the involvement of other antioxidant defenses (Regoli and
264 Giuliani, 2014), including low molecular weight scavengers or superoxide dismutase and
265 peroxiredoxines, which were reported as important mechanisms affected by acidification in the
266 mantle of *C. virginica* (Tomanek *et al.*, 2011).

267 Different effects were obtained in the gills with no induction of any antioxidant, but rather the
268 inhibition of glutathione S-transferases by co-exposure to cadmium and acidification, and of Se-
269 dependent GPx by Cd alone. Such results would indicate a lower capability to counteract oxidative
270 stress in gills compared to digestive gland of *F. glaber*. In this respect, tissue-specific effects of
271 multiple stressors were previously observed also in *A. colbecki* and *M. galloprovincialis* exposed to
272 different combinations of cadmium, temperature and acidification (Benedetti *et al.*, 2016; Nardi *et*
273 *al.*, 2017), confirming that within the complex network of oxidative interactions it is virtually
274 impossible to predict the same responses toward multiple stressors when moving to different species,
275 tissues, geographical latitudes or seasons (Camus *et al.*, 2005; de Hoop *et al.*, 2011). Despite such
276 variability, changes of antioxidant defenses still remain useful and sensitive early indicators of a

277 varied oxidative challenge, whose biological significance can be reflected by the total antioxidant
278 capacity. In our study, the marked decrease in the capability to neutralize peroxy radicals confirmed
279 the sensitivity of gills in organisms exposed to reduced pH with or without cadmium: compared to
280 digestive gland, the direct contact with seawater and the filter feeding activity can probably explain
281 the higher vulnerability of these tissues toward acidified conditions. In addition, digestive gland is a
282 detoxification/storage tissue, with the possibility to sequester cadmium in soluble or insoluble
283 compounds in Pectinidae such as *Pecten maximus* and *Chlamys varia* (Metian et al., 2007).

284 Interactive, pro-oxidant effects of acidification and cadmium exposure reduced the stability of
285 lysosomal membranes more than the single stressors, in agreement with similar effects observed in
286 haemocytes of mussels *M. edulis* and *M. galloprovincialis* (Beeseley et al., 2008; Nardi et al., 2017).
287 Concerning the onset of genotoxic damages, DNA fragmentation did not vary in terms of strand
288 breaks, while micronuclei frequency was enhanced in all treatments with comparable effects in
289 scallops exposed to cadmium, acidification or their combination. Levels of MN increased also in the
290 Antarctic scallop *A. colbecki* after exposure to various combinations of Cd and reduced pH (Benedetti
291 et al., 2016), and synergistic effects of cadmium and acidification were observed in haemocytes of
292 *M. galloprovincialis* (Nardi et al., 2017). The increased genotoxicity caused by these stressors in
293 terms of MN frequency but not as DNA strand breaks, has been hypothesized to reflect an increased
294 mitotic rate on haemocytes of such bivalves, rather than a direct effect on DNA integrity (Benedetti
295 et al., 2016; Nardi et al., 2017).

296 Principal components analysis revealed clear separation between organisms exposed to Cd alone
297 and those exposed to acidification, with or without the metal suggesting that this scallop might be
298 more sensitive to low pH than metal exposure. This hypothesis seems to be quantitatively confirmed
299 by the elaboration of the overall results through weighted criteria based on number, magnitude of
300 variations and toxicological relevance of biomarkers. The summarized hazard index was “Slight”
301 after Cd exposure, “Moderate” in organisms treated at high $p\text{CO}_2$ /low pH: among these, the
302 quantitative HQ value increased from Cd-acidified to acidified scallops. The relative high tolerance

303 of *F. glaber* to cadmium might reflect a certain cellular adaptation of this species to the elevated basal
304 content of the metal, both in terms of binding proteins and efficiency of antioxidant defenses. A
305 similar possibility has already been supposed for the Antarctic scallop *A. colbecki*, characterized by
306 Cd levels up to 5-15 folds higher than in *F. glaber* due to the presence of upwelling phenomena and
307 a natural Cd enrichment of the area: in this organism, however, the homeostatic equilibrium
308 conferring resistance to various stressors appeared overwhelmed by an additional exposure and
309 accumulation of this element, making *A. colbecki* highly sensitive to further changes of environmental
310 stressors.

311 In conclusion, this study supported the role of acidification as an environmental disturbance
312 acting both through direct mechanisms and indirect modulation of responsiveness to metal exposure:
313 however, uptake and sub-lethal effects of elements like Cd can vary in a species and tissue-specific
314 manner, highlighting the need of additional studies to elucidate the impact of multiple stressors,
315 particularly in species with elevated ecological or commercial importance.

316 **Acknowledgments:** This study was partially supported by the Italian National Program on Antarctic
317 Research (PNRA, project 2013/AZ1.14).

318

319 5. References

320 Andersen S, Grefsrud ES, Harboe T (2013). Effect of increased pCO₂ level on early shell
321 development in great scallop (*Pecten maximus* Lamarck) larvae. Biogeosciences, 10, pp. 6161-6184.

322 doi:10.5194/bg-10-6161-2013

323 Ardelan MV and Steinnes E (2010). Changes in mobility and solubility of the redox sensitive metals
324 Fe, Mn and Co at the seawater-sediment interface following CO₂ seepage. Biogeosciences, 7, pp.

325 569–583. DOI: 10.5194/bg-7-569-2010

326 Ardelan MV, Steinnes E, Lierhagen S, Linde SO (2009). Effects of experimental CO₂ leakage on
327 solubility and transport of seven trace metals in seawater and sediment. *Science of the Total*
328 *Environment*, 407, pp. 6255–6266. DOI: 10.1016/j.scitotenv.2009.09.004

329 Avio CG, Gorbi S, Milan M, Benedetti M, Fattorini D, d’Errico G, Pauletto M, Bargelloni L, Regoli
330 F (2015). Pollutants bioavailability and toxicological risk from microplastics to marine mussels.
331 *Environmental Pollution*, 198, pp. 211-222. DOI: 10.1016/j.envpol.2014.12.021

332 Beesley A, Lowe DM, Pascoe CK, Widdicombe S (2008). Effects of CO₂-induced seawater
333 acidification on the health of *Mytilus edulis*. *Climate Research*, 37, pp. 215-225. DOI:
334 10.3354/cr00765

335 Benedetti M, Ciaprini F, Piva F, Onorati F, Fattorini D, Notti A, Ausili A, Regoli F (2012). A
336 multidisciplinary weight of evidence approach for classifying polluted sediments: Integrating
337 sediment chemistry, bioavailability, biomarkers responses and bioassays. *Environment International*,
338 38, pp. 17-28. DOI: 10.1016/j.envint.2011.08.003

339 Benedetti M, Lanzoni I, Nardi A, d’Errico G, Di Carlo M, Fattorini D, Nigro M, Regoli F (2016).
340 Oxidative responsiveness to multiple stressors in the key Antarctic species, *Adamussium colbecki*:
341 Interactions between temperature, acidification and cadmium exposure. *Marine Environmental*
342 *Research*, 121, pp. 20-30. DOI: 10.1016/j.marenvres.2016.03.011

343 Bibby R, Widdicombe S, Parry H, Spicer J, Pipe R (2008). Effects of ocean acidification on the
344 immune response of the blue mussel *Mytilus edulis*. *Aquatic Biology*, 2, pp. 67-74. DOI:
345 10.3354/ab00037

346 Bustamante P, Germain P, Leclerc G, Miramand P (2002). Concentration and distribution of ²¹⁰Po
347 in the tissues of the scallop *Chlamys varia* and the mussel *Mytilus edulis* from the coasts of Charente-

348 Maritime (France). *Marine Pollution Bulletin*, 44, pp. 997-1002. DOI: 10.1016/S0025-
349 326X(02)00135-2

350 Caldeira K, Wickett M (2005). Ocean model predictions of chemistry changes from carbon dioxide
351 emissions to the atmosphere and ocean. *Journal of Geophysical Research*, 110, pp. 1–12. DOI:
352 10.1029/2004JC002671

353 Camus L, Birkely SR, Jones MB, Børseth JF, Grøsvik BE, Gulliksen B, Lønne OJ, Regoli F,
354 Depledge MH (2003). Biomarker responses and PAH uptake in *Mya truncata* following exposure to
355 oil-contaminated sediment in an Arctic fjord (Svalbard). *Science of the Total Environment* 308, pp.
356 221-234. DOI: 10.1016/S0048-9697(02)00616-2

357 Camus L, Gulliksen B, Depledge MH, Jones MB (2005). Polar bivalves are characterized by high
358 antioxidant defences. *Polar Research*, 24:1-2, pp. 111-118. DOI: 10.3402/polar.v24i1.6257

359 Cerrano C, Cardini U, Bianchelli C, Corinaldesi C, Pusceddu A, Danovaro R (2013). Red coral
360 extinction risk enhanced by ocean acidification. *Scientific Reports*, 3. DOI: 10.1038/srep01457

361 Cooley SR, Rheuban JE, Hart DR, Luu V, Glover DM, Hare JA, Doney SC (2015). An integrated
362 assessment model for helping the united states sea scallop (*Placopecten magellanicus*) fishery plan
363 ahead for ocean acidification and warming. *PLoS ONE*, 10. DOI: 10.1371/journal.pone.0124145

364 de Hoop L, Schipper AM, Leuven RSEW, Huijbregts MAJ, Olsen GH, Smit MGD, Hendriks AJ
365 (2011). Sensitivity of polar and temperate marine organisms to oil components. *Environmental*
366 *Science and Technology*, 45 (20), pp. 9017-9023. DOI: 10.1021/es202296a

367 de Orte MR, Lombardi AT, Sarmiento AM, Basallote MD, Rodríguez-Romero A, Riba I, Del Valls
368 A (2014a). Metal mobility and toxicity to microalgae associated with acidification of sediments: CO₂

369 and acid comparison. *Marine Environmental Research*, 96, pp. 136–144. DOI:
370 10.1016/j.marenvres.2013.10.003

371 de Orte MR, Sarmiento AM, DelValls TÁ, Riba I (2014b). Simulation of the potential effects of CO₂
372 leakage from carbon capture and storage activities on the mobilization and speciation of metals.
373 *Marine Pollution Bulletin*, 86, pp. 59–67. DOI: 10.1016/j.marpolbul.2014.07.042

374 Dickson AG, Sabine CL, Christian JR (2007). Guide to Best Practices for Ocean CO₂ Measurements,
375 p. 191. PICES Special Publication, 3.

376 Duckworth CG, Picariello CR, Thomason RK, Patel KS, Bielmyer-Fraser GK (2017). Responses of
377 the sea anemone, *Exaiptasia pallida*, to ocean acidification conditions and zinc or nickel exposure.
378 *Aquatic Toxicology*, 182, pp. 120-128. DOI: 10.1016/j.aquatox.2016.11.014

379 Dupont S, Havenhand J, Thorndyke W, Peck L, Thorndyke M (2008). Near-future level of CO₂-
380 driven ocean acidification radically affects larval survival and development in the brittlestar
381 *Ophiothrix fragilis*. *Marine Ecology Progress Series*, 373, pp. 285-294. DOI: 10.3354/meps07800

382 Dupont S, Ortega-Martínez O, Thorndyke M (2010). Impact of near-future ocean acidification on
383 echinoderms. *Ecotoxicology*, 19, pp. 449-462. DOI: 10.1007/s10646-010-0463-6

384 Ellis RP, Bersey J, Rundle SD, Hall-Spencer JM, Spicer JJ (2009). Subtle but significant effects of
385 CO₂ acidified seawater on embryos of the intertidal snail, *Littorina obtusata*. *Aquatic Biology*, 5, pp.
386 41-48. DOI: 10.3354/ab00118

387 Freitas R, Pires A, Moreira A, Wrona FJ, Figueira E, Soares AMVM (2016). Biochemical alterations
388 induced in *Hediste diversicolor* under seawater acidification conditions. *Marine Environmental*
389 *Research*, 117, pp. 75-84. DOI: 10.1016/j.marenvres.2016.04.003

390 Gattuso JP, Lavigne H (2009). Technical Note: Approaches and software tools to investigate the
391 impact of ocean acidification. *Biogeosciences*, 6, pp. 2121–2133. DOI: 10.5194/bg-6-2121-2009

392 Gazeau F, Quiblier C, Jansen JM, Gattuso J-P, Middelburg JJ, Heip CHR (2007). Impact of elevated
393 CO₂ on shellfish calcification. *Geophysical Research Letters*, 34. DOI: 10.1029/2006GL028554

394 Gorbi S, Regoli F (2003). Review. Total Oxyradical Scavenging Capacity as an index of
395 susceptibility to oxidative stress in marine organisms. *Comments in Toxicology* 9, pp. 303-322. DOI:
396 10.1016/S0166-445X(00)00091-6

397 Götze S, Matoo OB, Beniash E, Saborowski R, Sokolova IM (2014). Interactive effects of CO₂ and
398 trace metals on the proteasome activity and cellular stress response of marine bivalves *Crassostrea*
399 *virginica* and *Mercenaria mercenaria*. *Aquatic Toxicology*, 149, pp. 65–82. DOI:
400 10.1016/j.aquatox.2014.01.027

401 Gutowska MA, Melzner F, Langenbuch M, Bock C, Claireaux G, Pörtner HO (2010). Acid-base
402 regulatory ability of the cephalopod (*Sepia officinalis*) in response to environmental hypercapnia.
403 *Journal of comparative physiology. B, Biochemical, systemic, and environmental physiology*, 180,
404 pp. 323-335. DOI: 10.1007/s00360-009-0412-y

405 Hernroth B, Baden S, Thorndyke M, Dupont S (2011). Immune suppression of the echinoderm
406 *Asterias rubens* (L.) following long-term ocean acidification. *Aquatic Toxicology*, 103, pp. 222-224.
407 DOI: 10.1016/j.aquatox.2011.03.001

408 Hernroth B, Sköld HN, Wiklander K, Jutfelt F, Baden S (2012). Simulated climate change causes
409 immune suppression and protein damage in the crustacean *Nephrops norvegicus*. *Fish and Shellfish*
410 *Immunology*, 33, pp. 1095-1101. DOI: doi: 10.1016/j.fsi.2012.08.011

411 Hernroth B, Baden S, Tassidis H, Hörnaeus K, Guillemant J, Bergström Lind S, Bergquist J (2016)
412 Impact of ocean acidification on antimicrobial activity in gills of the blue mussel (*Mytilus edulis*).
413 Fish and Shellfish Immunology, 55, pp. 452-459. DOI: 10.1016/j.fsi.2016.04.007.

414 IPCC (2014). Climate Change 2014: Impacts, Adaptation, and Vulnerability. Contribution of
415 Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change.
416 Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 1820 pp.

417 Ivanina AV, Beniash E, Eitzkorn M, Meyers TB, Ringwood AH, Sokolova IM (2013). Short-term
418 acute hypercapnia affects cellular responses to trace metals in the hard clams *Mercenaria mercenaria*.
419 Aquatic Toxicology, 140–141, pp. 123–133. DOI: 10.1016/j.aquatox.2013.05.019

420 Ivanina AV, Hawkins C, Sokolova IM (2014). Immunomodulation by the interactive effects of
421 cadmium and hypercapnia in marine bivalves *Crassostrea virginica* and *Mercenaria mercenaria*.
422 Fish and Shellfish Immunology, 37, pp. 299–312. DOI: 10.1016/j.fsi.2014.02.016

423 Ivanina AV and Sokolova IM (2015). Interactive effects of metal pollution and ocean acidification
424 on physiology of marine organisms. Current Zoology, 6, 653–668. DOI: 10.1093/czoolo/61.4.653

425 Ivanina AV, Hawkins C, Sokolova IM (2016). Interactive effects of copper exposure and
426 environmental hypercapnia on immune functions of marine bivalves *Crassostrea virginica* and
427 *Mercenaria mercenaria*. Fish and Shellfish Immunology, 49, pp. 54-65. DOI:
428 10.1016/j.fsi.2015.12.011.

429 Jokiel PL, Rodgers KS, Kuffner IB, Andersson AJ, Cox EF, Mackenzie FT (2008). Ocean
430 acidification and calcifying reef organisms: A mesocosm investigation. Coral Reefs, 27, pp. 473-483.
431 DOI: 10.1007/s00338-008-0380-9

432 Kurihara H, Kato S, Ishimatsu A (2007). Effects of increased seawater pCO₂ on early development
433 of the oyster *Crassostrea gigas*. *Aquatic Biology*, 1, pp. 91-98.

434 Lacoue-Labarthe T, Martin S, Oberhänsli F, Teyssié J-L, Markich S, Jeffree R, Bustamante P (2009).
435 Effects of increased pCO₂ and temperature on trace element (Ag, Cd and Zn) bioaccumulation in the
436 eggs of the common cuttlefish, *Sepia officinalis*. *Biogeosciences*, 6, pp. 2561–2573. DOI:
437 10.5194/bg-6-2561-2009

438 Lacoue-Labarthe T, Réveillac E, Oberhänsli F, Teyssié JL, Jeffree R, Gattuso JP (2011). Effects of
439 ocean acidification on trace element accumulation in the early-life stages of squid *Loligo vulgaris*.
440 *Aquatic Toxicology*, 105, pp. 166–176. DOI: 10.1016/j.aquatox.2011.05.021

441 Lannig G, Eilers S, Pörtner H-O, Sokolova IM, Bock C (2010). Impact of ocean acidification on
442 energy metabolism of oyster, *Crassostrea gigas* - Changes in metabolic pathways and thermal
443 response. *Marine Drugs*, 8, pp. 2318-2339. DOI: 10.3390/md8082318

444 Le Quéré C, Raupach MR, Canadell JG, Marland G, Bopp L, Ciais P, Conway TJ, Doney SC, Feely
445 RA, Foster P, Friedlingstein P, Gurney K, Houghton RA, House JI, Huntingford C, Levy PE, Lomas
446 MR, Majkut J, Metzl N, Ometto JP, Peters GP, Prentice IC, Randerson JT, Running SW, Sarmiento
447 JL, Schuster U, Sitch S, Takahashi T, Viovy N, Van Der Werf GR, Woodward FI (2009). Trends in
448 the sources and sinks of carbon dioxide. *Nature Geoscience*, 2, 831-836. DOI: 10.1038/ngeo689

449 Lewis C, Clemow K, Holt WV (2013). Metal contamination increases the sensitivity of larvae but
450 not gametes to ocean acidification in the polychaete *Pomatoceros lamarckii* (Quatrefages). *Marine*
451 *Biology*, 160, pp. 2089–2101. DOI: 10.1007/s00227-012-2081-8

452 López IR, Kalman J, Vale C, Blasco J (2010). Influence of sediment acidification on the
453 bioaccumulation of metals in *Ruditapes philippinarum*. *Environmental Science and Pollution*
454 *Research International*, 17(9), pp. 1519–1528. DOI: 10.1007/s11356-010-0338-7

455 Marčeta T, Da Ros L, Marin MG, Codognotto VF, Bressan M (2016). Overview of the biology of
456 *Flexopecten glaber* in the North Western Adriatic Sea (Italy): A good candidate for future shellfish
457 farming aims? *Aquaculture*, 462, pp. 80-91. DOI: 10.1016/j.aquaculture.2016.04.036

458 Mauri M, Orlando E, Nigro M, Regoli F (1990). Heavy metals in the Antarctic scallop *Adamussium*
459 *colbecki*. *Marine Ecology Progress Series*, 67, pp. 27-33.

460 Mazzoldi C, Sambo A, Riginella E (2014). The Clodia database: a long time series of fishery data
461 from the Adriatic Sea. *Scientific Data*, 1, pp. 577–590. DOI: 10.1038/sdata.2014.18

462 Metian M, Warnau M, Oberhänsli F, Teyssié J-L, Bustamante P (2007). Interspecific comparison of
463 Cd bioaccumulation in European Pectinidae (*Chlamys varia* and *Pecten maximus*). *Journal of*
464 *Experimental Marine Biology and Ecology*, 353, pp. 58-67. DOI: 10.1016/j.jembe.2007.09.001

465 Miles H, Widdicombe S, Spicer JJ, Hall-Spencer J (2007). Effects of anthropogenic seawater
466 acidification on acid-base balance in the sea urchin *Psammechinus miliaris*. *Marine Pollution*
467 *Bulletin*, 54, pp. 89-96. DOI: 10.1016/j.marpolbul.2006.09.021

468 Millero FJ (2010). Carbonate constants for estuarine waters. *Marine Freshwater Research*, 61,
469 pp.139-142. DOI: 10.1071/MF09254

470 Millero FJ, Woosley R, Ditrolio B, Waters J (2009). Effect of ocean acidification on the speciation
471 of metals in seawater. *Oceanography*, 22, pp. 72–85. DOI: 10.5670/oceanog.2009.98

472 Moore MN, Allen J, McVeigh A (2006). Environmental prognostics: An integrated model
473 supporting lysosomal stress responses as predictive biomarkers of animal health status. *Marine*
474 *Environmental Research* 61, pp. 278-304. DOI: 10.1016/j.marenvres.2005.10.005

475 Moreira A, Figueira E, Soares AMVM, Freitas R (2016). The effects of arsenic and seawater
476 acidification on antioxidant and biomineralization responses in two closely related *Crassostrea*

477 species. *Science of the Total Environment*, 545-546, pp. 569-581. DOI:
478 10.1016/j.scitotenv.2015.12.029

479 Nardi A, Mincarelli LF, Benedetti M, Fattorini D, d'Errico G, Regoli F (2017). Indirect effects of
480 climate changes on cadmium bioavailability and biological effects in the Mediterranean mussel
481 *Mytilus galloprovincialis*. *Chemosphere*, 169, pp. 493-502. DOI:
482 10.1016/j.chemosphere.2016.11.093

483 Neff JM (2002). *Bioaccumulation in Marine Organisms*. Elsevier Science, Oxford, UK.

484 Nigro M, Frenzilli G, Scarcelli V, Gorbi S, Regoli F (2002). Induction of DNA strand breakage and
485 apoptosis in the eel *Anguilla Anguilla*. *Marine Environmental Research*, 54, pp. 517-520. DOI:
486 10.1016/S0141-1136(02)00178-2

487 Nikinmaa M (2013). Climate change and ocean acidification - Interactions with aquatic toxicology.
488 *Aquatic Toxicology*, 126, pp. 365-372. DOI: 10.1016/j.aquatox.2012.09.006.

489 Pan T-CF, Applebaum SL, Manahan DT (2015). Experimental ocean acidification alters the
490 allocation of metabolic energy. *Proceedings of the National Academy of Sciences of the United States*
491 *of America*, 112, pp. 4696-4701. DOI: 10.1073/pnas.1416967112

492 Pierrot D, Lewis E, Wallace DWR (2006). MS Excel Program Developed for CO2System
493 Calculations. ORNL/CDIAC-105a. Carbon Dioxide Information Analysis Center, Oak Ridge
494 National Laboratory, U.S. Department of Energy, Oak Ridge, Tennessee. DOI:
495 10.3334/CDIAC/otg.CO2SYS_XLS_CDIAC105a.

496 Pimentel MS, Faleiro F, Diniz M, Machado J, Pousão-Ferreira P, Peck MA, Pörtner HO, Rosa R
497 (2015). Oxidative stress and digestive enzyme activity of flatfish larvae in a changing ocean. *PLoS*
498 *ONE*, 10. DOI: 10.1371/journal.pone.0134082

499 Piva F, Ciaprini F, Onorati F, Benedetti M, Fattorini D, Ausili A, Regoli F (2011) Assessing sediment
500 hazard through a weight of evidence approach with bioindicator organisms: A practical model to
501 elaborate data from sediment chemistry, bioavailability, biomarkers and ecotoxicological bioassays.
502 *Chemosphere*, 83, pp. 475-485. DOI: 10.1016/j.chemosphere.2010.12.064

503 Pujolar JM, Marčeta T, Saavedra C, Bressan M, Zane L (2010). Inferring the demographic history of
504 the Adriatic Flexopecten complex. *Molecular Phylogenetics and Evolution*, 57, pp. 942-947. DOI:
505 10.1016/j.ympev.2010.08.002.

506 Regoli F (2012). Chemical pollutants and mechanisms of reactive oxygen species generation in
507 aquatic organisms. In: *Oxidative stress in aquatic ecosystems*. Abele D, Vázquez-Medina JP,
508 Zenteno-Savín T (editors). pp. 308-316.

509 Regoli F and Giuliani ME (2014). Oxidative pathways of chemical toxicity and oxidative stress
510 biomarkers in marine organisms. *Marine Environmental Research*, 93, pp. 106–117. DOI:
511 10.1016/j.marenvres.2013.07.006

512 Regoli F and Winston GW (1998). Applications of a new method for measuring the total oxyradical
513 scavenging capacity in marine invertebrates. *Marine Environmental Research*, 46, pp. 439-442. DOI:
514 doi.org/10.1016/S0141-1136(97)00119-0

515 Regoli F and Winston GW (1999). Quantification of total oxidant scavenging capacity (TOSC) of
516 antioxidants for peroxyxynitrite, peroxy radicals and hydroxyl radicals. *Toxicology and Applied
517 Pharmacology* 156, pp. 96-105. DOI: doi.org/10.1006/taap.1999.8637

518 Regoli F, Nigro M, Orlando E (1998). Lysosomal and antioxidant responses to metals in the Antarctic
519 scallop *Adamussium colbecki*. *Aquatic Toxicology*, 40, pp. 375-392. DOI: 10.1016/S0166-
520 445X(97)00059-3

521 Regoli F, Nigro M, Bompadre S, Winston GW (2000). Susceptibility to oxyradical toxicity in
522 Antarctic, Arctic, and Mediterranean scallops. *Marine Environmental Research*, 50, pp. 547.

523 Regoli F, Nigro M, Chiantore M, Winston GW (2002). Seasonal variations of susceptibility to
524 oxidative stress in *Adamussium colbecki*, a key bioindicator species for the Antarctic marine
525 environment. *Science of the Total Environment*, 289, pp. 205-211.

526 Regoli F, Nigro M, Benedetti M, Gorbi S, Pretti C, Gervasi PG, Fattorini D (2005) Interactions
527 between metabolism of trace metals and xenobiotic agonists of the aryl hydrocarbon receptor in the
528 antarctic fish *Trematomus bernacchii*: Environmental perspectives. *Environmental Toxicology and*
529 *Chemistry*, 24, pp. 1475-1482.

530 Regoli F, Pellegrini, D, Cicero AM, Nigro M, Benedetti M, Gorbi S, Fattorini D, D'Errico G, Di Carlo
531 M, Nardi A, Gaion A, Scuderi A, Giuliani S, Romanelli G, Berto D, Trabucco B, Guidi P,
532 Bernardeschi M, Scarcelli V, Frenzilli G (2014). A multidisciplinary weight of evidence approach
533 for environmental risk assessment at the Costa Concordia wreck: Integrative indices from Mussel
534 Watch. *Marine Environmental Research*, 96, pp. 92-104. DOI: 10.1016/j.marenvres.2013.09.016

535 Ricevuto E, Benedetti M, Regoli F, Spicer JI, Gambi MC (2015). Antioxidant capacity of polychaetes
536 occurring at a natural CO₂ vent system: Results of an in situ reciprocal transplant experiment. *Marine*
537 *Environmental Research*, 112, pp. 44–51. DOI: 10.1016/j.marenvres.2015.09.005

538 Ricevuto E, Lanzoni I, Fattorini D, Regoli F, Gambi MC (2016). Arsenic speciation and susceptibility
539 to oxidative stress in the fanworm *Sabella spallanzanii* (Gmelin) (Annelida, Sabellidae) under
540 naturally acidified conditions: An in situ transplant experiment in a Mediterranean CO₂ vent system.
541 *Science of The Total Environment*, 544, pp. 765–773. DOI: 10.1016/j.scitotenv.2015.11.154

542 Rodríguez-Romero A, Basallote MD, de Orte MR, DelValls TÁ, Riba I, Blasco J (2014). Simulation
543 of CO₂ leakages during injection and storage in sub-seabed geological formations: Metal mobilization
544 and biota effects. *Environment International*, 68, pp. 105–117. DOI: 10.1016/j.envint.2014.03.008

545 Rokitta SD, John U, Rost B (2012). Ocean Acidification Affects Redox-Balance and Ion-
546 Homeostasis in the Life-Cycle Stages of *Emiliana huxleyi*. *PLoS ONE*, 7. DOI:
547 10.1371/journal.pone.0052212

548 Schalkhauser B, Bock C, Stemmer K, Brey T, Pörtner H-O, Lannig G (2013). Impact of ocean
549 acidification on escape performance of the king scallop, *Pecten maximus*, from Norway. *Marine*
550 *Biology*, 160, pp. 1995-2006. DOI: 10.1007/s00227-012-2057-8

551 Shi W, Zhao X, Han Y, Che Z, Chai X, Liu G (2016). Ocean acidification increases cadmium
552 accumulation in marine bivalves: A potential threat to seafood safety. *Scientific Reports*, 6. DOI:
553 10.1038/srep20197

554 Siddiqui S, Bielmyer-Fraser GK (2015). Responses of the sea anemone, *Exaiptasia pallida*, to ocean
555 acidification conditions and copper exposure. *Aquatic Toxicology*, 167, pp. 228–239. DOI:
556 10.1016/j.aquatox.2015.08.012

557 Soriano-Santiago OS, Liñán-Cabello MA, Delgadillo-Nuño MA, Ortega-Ortiz C, Cuevas-Venegas S
558 (2013). Physiological responses to oxidative stress associated with pH variations in host tissue and
559 zooxanthellae of hermatypic coral *Pocillopora capitata*. *Marine and Freshwater Behaviour and*
560 *Physiology*, 46, pp. 275-286. DOI: 10.1080/10236244.2013.827877

561 Spicer JJ, Raffo A, Widdicombe S (2007). Influence of CO₂-related seawater acidification on
562 extracellular acid-base balance in the velvet swimming crab *Necora puber*. *Marine Biology*, 151, pp.
563 1117-1125. DOI: 10.1007/s00227-006-0551-6

564 Stumpp M, Dupont S, Thorndyke MC, Melzner F (2011). CO₂ induced seawater acidification impacts
565 sea urchin larval development II: Gene expression patterns in pluteus larvae. *Comparative*
566 *Biochemistry and Physiology - A Molecular and Integrative Physiology*, 160, pp. 320-330. DOI:
567 10.1016/j.cbpa.2011.06.023.

568 Stumpp M, Trübenbach K, Brennecke D, Hu MY, Melzner F (2012). Resource allocation and
569 extracellular acid-base status in the sea urchin *Strongylocentrotus droebachiensis* in response to CO₂
570 induced seawater acidification. *Aquatic Toxicology*, 110-111, pp. 194-207. DOI:
571 10.1016/j.aquatox.2011.12.020

572 Tomanek L, Zuzow MJ, Ivanina AV, Beniash E, Sokolova IM (2011). Proteomic response to elevated
573 PCO₂ level in eastern oysters, *Crassostrea virginica*: evidence for oxidative stress. *The Journal of*
574 *Experimental Biology*, 214, pp. 1836–1844. DOI: 10.1242/jeb.055475

575 Wallace RB, Baumann H, Grear JS, Aller R, Gobler CJ (2014). Coastal ocean acidification: The other
576 eutrophication problem. *Estuarine Coastal and Shelf Science*, 148, pp. 1-13. DOI:
577 10.1016/j.ecss.2014.05.027

578 White MM, McCorkle DC, Mullineaux LS, Cohen AL (2013). Early exposure of bay scallops
579 (*Argopecten irradians*) to high CO₂ causes a decrease in larval shell growth. *PLoS ONE*, 8. DOI:
580 10.1371/journal.pone.0061065


581 Wong PP, Losada IJ, Gattuso J-P, Hinkel J, Khattabi A, McInnes KL, Saito Y, Sallenger A (2014):
582 Coastal systems and low-lying areas. In: *Climate Change 2014: Impacts, Adaptation, and*
583 *Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working Group II to the Fifth*
584 *Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press,
585 Cambridge, United Kingdom and New York, NY, USA, pp. 361-409.

586 Table 1 - Water chemistry parameters during experimental exposure. T (temperature), S (salinity),
 587 pH_{NBS} (pH calibrated with National Bureau of Standard scale), A_T (total alkalinity), pCO_2 (partial
 588 pressure of CO_2), Ω_c and Ω_a (saturation state of respectively calcite and aragonite). Data are presented
 589 as means \pm standard deviations.

590


treatment	measured parameters				calculated parameters		
	T (°C)	S	pH (NBS scale)	A_T ($\mu\text{mol/kg}$)	pCO_2 (μatm)	Ω_c	Ω_a
CTRL	20.02 \pm 0.08	30 \pm 0.5	8.21 \pm 0.04	3464.9 \pm 138.8	407.7 \pm 20.1	8.6 \pm 0.4	5.5 \pm 0.2
Cd	20.00 \pm 0.10	30 \pm 0.5	8.19 \pm 0.04	3482.5 \pm 80.1	409.3 \pm 22.5	8.6 \pm 0.3	5.5 \pm 0.3
A	19.98 \pm 0.06	30 \pm 0.5	7.41 \pm 0.04	3431.2 \pm 179.1	3144.6 \pm 160.2	1.7 \pm 0.1	1.1 \pm 0.1
A-Cd	20.92 \pm 0.08	30 \pm 0.5	7.43 \pm 0.04	3464.9 \pm 138.8	3124.3 \pm 162.3	1.7 \pm 0.2	1.1 \pm 0.1

591




593

594 Figure 1. Cadmium concentrations in digestive gland (a) and gills (b) of exposed scallops. Data are
 595 expressed as mean values \pm standard deviations (n=5). Different letters indicate significant
 596 differences between group of means (ANOVA and Tukey HSD *post-hoc*). CTRL= Control; Cd=
 597 Cadmium; A= acidification; A-Cd= acidification + Cd.



598

599 Figure 2. Metallothioneins and antioxidant defenses in digestive gland of scallops exposed to various
600 treatments. MT: metallothioneins (a), CAT: catalase (b), GST: glutathione S-transferase (c), Se-Dep.
601 GPx: Se-dependent glutathione peroxidases (d) total GPx: sum of Se-dependent and Se-independent
602 glutathione peroxidases (e), GR: glutathione reductase (f), TGSH: total glutathione (g), TOSC ROO•:
603 total oxyradical scavenging capacity toward peroxy radical (h), TOSC HO•: total oxyradical
604 scavenging capacity toward hydroxyl radical (i). Data are given as mean values \pm standard deviations
605 (n=5). Different letters indicate significant differences between group of means (ANOVA and Tukey
606 HSD *post-hoc*). CTRL= Control; Cd= Cadmium; A= acidification; A-Cd= acidification + Cd.



607

608 Figure 3. Metallothioneins and antioxidant defenses in gills of scallops exposed to various treatments.

609 MT: metallothioneins (a), CAT: catalase (b), GST: glutathione S-transferase (c), Se-Dep. GPx: Se-

610 dependent glutathione peroxidases (d) total GPx: sum of Se-dependent and Se-independent


611 glutathione peroxidases (e), GR: glutathione reductase (f), TGSH: total glutathione (g), TOSC ROO•:

612 total oxyradical scavenging capacity toward peroxy radical (h), TOSC HO•: total oxyradical

613 scavenging capacity toward hydroxyl radical (i). Data are given as mean values \pm standard deviations


614 (n=5). Different letters indicate significant differences between group of means (ANOVA and Tukey

615 HSD *post-hoc*). CTRL= Control; Cd= Cadmium; A= acidification; A-Cd= acidification + Cd.



617




618 Figure 4. Lysosomal and genotoxic parameters in haemocytes of scallops exposed to various
 619 treatments. LMS: lysosomal membrane stability (a), DNA damage (b), MN: frequency of micronuclei
 620 (c). Data are given as mean values \pm standard deviation (n=5). Different letters indicate significant
 621 differences between group of means (ANOVA and Tukey HSD *post-hoc*). CTRL= Control; Cd=
 622 Cadmium; A= acidification; A-Cd= acidification + Cd.



623

624 Figure 5. Graphical representation of principal components analysis conducted on biological
 625 parameters analyzed in scallops tissues. CTRL= Control; Cd= Cadmium; A= acidification; A-Cd=
 626 acidification + Cd. Arrows represent the ten major variables that contribute to the separation.

627

Experimental Treatment	Hazard Quotients (HQs)	Class of Hazard	Level
Cd	9.06	SLIGHT	
A	21.07	MODERATE	
A-Cd	13.13	MODERATE	

629

630 Figure 6. Weighted classification of biomarkers data for the whole dataset of analyzed parameters in
631 each experimental condition. The quantitative hazard quotients (HQ) and the assigned class of hazard
632 are given.