

#### UNIVERSITÀ POLITECNICA DELLE MARCHE Repository ISTITUZIONALE

Slaughter performance and carcass and meat quality of Bergamasca light lambs according to slaughter age

This is the peer reviewd version of the followng article:

Original

Slaughter performance and carcass and meat quality of Bergamasca light lambs according to slaughter age / Budimir, Katarina; Trombetta, Maria Federica; Francioni, Matteo; Toderi, Marco; D'Ottavio, Paride. - In: SMALL RUMINANT RESEARCH. - ISSN 0921-4488. - ELETTRONICO. - 164:(2018), pp. 1-7. [10.1016/j.smallrumres.2018.04.006]

Availability:

This version is available at: 11566/258493 since: 2022-06-03T11:22:42Z

Publisher:

Published DOI:10.1016/j.smallrumres.2018.04.006

Terms of use:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. The use of copyrighted works requires the consent of the rights' holder (author or publisher). Works made available under a Creative Commons license or a Publisher's custom-made license can be used according to the terms and conditions contained therein. See editor's website for further information and terms and conditions. This item was downloaded from IRIS Università Politecnica delle Marche (https://iris.univpm.it). When citing, please refer to the published version.

note finali coverpage

(Article begins on next page)

1	Slaughter performance and carcass and meat quality of Bergamasca light lambs
2	according to slaughter age
3	
4	Katarina Budimir, Maria Federica Trombetta, Matteo Francioni, Marco Toderi, Paride
5	D'Ottavio*
6	
7	Dipartimento di Scienze Agrarie, Alimentari ed Ambientali, Università Politecnica delle
8	Marche, Via Brecce Bianche 10, 60131 Ancona, Italy
9	
10	
11	
12	
13	
14	
15	
16	*Corresponding author: Paride D'Ottavio
17	Dipartimento di Scienze Agrarie, Alimentari ed Ambientali
18	Università Politecnica delle Marche
19	Via Brecce Bianche 10
20	60131 Ancona, Italy
21	Tel: +39-071-2204156
22	Email: p.dottavio@univpm.it
23	

This study was designed to evaluate the effects of slaughter age (40 vs. 60 days) on slaughter 27 performance, carcass and meat quality, and fatty acid composition of intramuscular and 28 subcutaneous fat of Bergamasca lambs reared according to the traditional transhumant system 29 in central Italy. Lambs slaughtered at 60 days of age had higher carcass weight (12.44 vs. 30 10.36 kg), lower dressing percentage (47.68% vs. 52.16%), and higher proportion of non-31 carcass components and leg commercial cut (37.82% vs. 35.49%). Furthermore, after 3 and 6 32 days of storage, the meat of older lambs showed lower drip loss (3.69% vs. 6.16%; 5.73% vs. 33 9.36%, respectively). Slaughter age did not influence meat pH, cooking loss, or chemical 34 composition while older lambs had meat with higher a\* (19.43 vs. 18.91). The fatty acid 35 composition of intramuscular fat was not affected by slaughter age, except for C:13 and C:14 36 37 fatty acids, which were higher in older lambs. Subcutaneous fat of lambs slaughtered at 40 days of age showed a better fatty acid profile, as lower saturated fatty acids (52.46% vs. 38 39 55.68%) and higher mono- and polyunsaturated fatty acids (34.06% vs. 30.16%, 6.46% vs. 40 5.79%, respectively), and n-6 and n-3 polyunsaturated fatty acids. Furthermore, subcutaneous fat of lambs slaughtered at 40 days of age had better polyunsaturated/ saturated fatty acid ratio 41 (0.12 vs. 0.11) and hypocholesterolemic/ hypercholesterolemic ratio (1.42 vs. 1.03), and lower 42 atherogenic index (1.32 vs. 1.82) and thrombogenic index (1.98 vs. 2.35). For light lamb 43 production using the traditional rearing systems, slightly heavier lambs can be produced 44 without worsening chemical composition and cooking loss and fatty acid composition of the 45 longissimus lomborum muscle. However, these lambs might have lower dressing percentages 46 and a less favourable fatty acid profile of subcutaneous fat for human health. 47



#### 50 **1. Introduction**

51

Traditional lamb production in Mediterranean countries is based on light lambs that are slaughtered at early ages (i.e., at 30 to 60 days of age), so just after weaning or after a short fattening period (Juárez et al., 2009; Santos-Silva et al., 2002a). These carcasses weigh up to 13 kg and are characterised by their pale pink colour, lower amounts of fat, and good flavour (Beriain et al., 2000), compared to heavier carcasses produced in other production systems and countries (Ekiz et al., 2013; Hajji et al., 2016; Jacques et al., 2011; Lind et al., 2009; Piasentier, 2003; Priolo et al., 2002).

In recent years, the demand for lean carcasses has grown due to increased awareness of consumers for healthy meat, with a focus on the quantity and quality of fat (Font i Fournols and Guerrero, 2014). Scientific studies and nutritional guidelines recommend not only a reduction in total fat intake in the human diet, but also a focus on saturated fatty acids (SFAs) and increased consumption of polyunsaturated fatty acids (PUFAs), and especially n-3 PUFAs (Calder and Yaqoob, 2009; World Health Organisation, 2003).

The age and weight of lambs at slaughter are among the main factors that affect the 65 meat quality at both levels, in terms of the carcass and the meat. A greater weight usually 66 implies an older lamb, except when the feed is manipulated or the lamb has periods of 67 specific alimentary restrictions (Guerrero et al., 2013). Although greater slaughter age of 68 lambs results in heavier carcasses, increased adiposity and better carcass conformation 69 (D'Alessandro et al., 2013; Juárez et al., 2009), this can also result in increased intramuscular 70 fat (Abdullah and Qudsieh, 2009; della Malva et al., 2016; Pérez et al., 2002). Furthermore, 71 lambs slaughtered at a greater age can have a fatty acid (FA) composition of the meat that is 72 less favourable for human health. This can arise as a result of increased SFAs, less PUFAs 73

and n-3 PUFAs, and increased atherogenic and thrombogenic indices (Cifuni et al., 2000;
Marino et al., 2008; Santos-Silva et al., 2002b).

These aspects are of the utmost importance for the definition of strategies to enhance the production of lamb meat while also taking into consideration the market demand. In this sense, there are significant gaps of knowledge for lamb produced under quality labels, such as the Protected Geographical Indication of *'Agnello del Centro Italia'* (Lamb from Central Italy; European Union, Commission Regulation No. 475/2013). This can be produced using Bergamasca sheep, as well as some other sheep breeds.

Bergamasca sheep are an autochthonous Italian breed that originated from the 82 Lombardy region (northern Italy) and that are traditionally raised according to the 83 transhumant system (Piasentier, 2003). Nowadays, Bergamasca sheep are raised principally 84 for meat in most parts of continental Italy (Bigi and Zanon, 2008), and they are often used for 85 cross-breeding with other to improve meat yield. Male and female Bergamasca sheep can 86 reach adult weights of 105 kg and 82 kg, respectively. In the Lombardy region, Bergamasca 87 sheep are still the most important breed used to produce castrated, heavy and light lambs, as 88 the traditional products of transhumant management (Piasantier et al., 2003). 89

In the Marche region (central Italy), lamb production is mainly based on extensive 90 grazing and the transhumance is still of major importance. In summer, most of the flocks 91 graze on upland pastures. Starting from autumn, the sheep are progressively transferred to 92 lowlands, where until Spring the main forage resources are lucerne meadows, although green 93 cereals, crop residues, marginal lands, and riverbanks are sometimes also used (Caballero et 94 al., 2009; D'Ottavio and Santilocchi, 2014). Lamb production mainly occurs in the lowlands 95 for the Easter and Christmas markets, which is when lamb meat is traditionally consumed in 96 Italy (Cifuni et al., 2000). Lambs are reared on pastures with their dams until they reach the 97 optimal slaughter weight, which according to local practices starts from 40 days of age. The 98

99 lambs are not weaned so their diet is mostly based on milk (i.e., to 20 days of age), while later100 they are supplemented with concentrate and/or hay, as needed.

101 The effects of slaughter age on the quality of light lamb meat has been studied for 102 some Italian breeds, such as the Altamurana (della Malva et al., 2016; Marino et al., 2008), 103 Apulian (Cifuni et al., 2000), Leccese (D'Alessandro et al., 2013), Trimeticcio (Marino et al., 104 2008) and Italian Merino (Oriani et al., 2005) sheep. However, there is little or no such 105 information available for the lamb meat quality of the Bergamasca breed of sheep.

The aim of the present study was to evaluate the effects of slaughter age on slaughter performance, carcass traits and meat quality, including FA composition of intramuscular and subcutaneous fat, of Bergamasca light lambs reared under the traditional transhumant system of the Marche region of Italy.

110

#### 111 **2. Materials and methods**

112

All animal handling followed the recommendations of European Union Directive
2010/63/EU, which are implemented in Italian law according to Legislative Decree No.
26/2014.

116

## 117 2.1. Experimental design, diet and animal management

This study was carried out from September to October 2015 in the Marche region (central Italy) under the standard conditions for rearing and management of the transhumant sheep system that is characteristic for this region. Twenty-two male, single-born, Bergamasca lambs were included in the study. At birth, the lambs were randomly distributed into two groups that were balanced for body weight. All of these lambs stayed with their dams on grasslands dominated by alfalfa (10.3 MJ metabolisable energy kg<sup>-1</sup> dry matter [DM], 15.7% crude

protein DM, 28.5% crude fibre DM), and suckled their dams throughout the whole study 124 period. The dams grazed and had free access to alfalfa hay (11.8 MJ metabolisable energy kg 125 <sup>1</sup> DM, 15.2% crude protein DM, 22.3% crude fibre DM) and their diet was supplemented with 126 corn grain (0.5 kg head<sup>-1</sup> day<sup>-1</sup>; 16.5 MJ metabolisable energy kg<sup>-1</sup> DM, 7.9% crude protein 127 DM, 5.0% crude fibre DM). The lambs were given corn grain ad libitum in creep feeders 128 from 20 days of age, and had access to alfalfa hay. The chemical compositions of collected 129 feed samples were determined according to Martillotti et al. (1987). To calculate the average 130 daily gain (ADG), the individual lamb weights were recorded at birth and after each 20 days, 131 until slaughter. The lambs were slaughtered at two different ages: 11 lambs at an average of 132 40 days, and the other 11 lambs at an average of 60 days. 133

134

## 135 2.2. Slaughter procedure and assessment of carcass traits

136 To obtain the pre-slaughter weight, the lambs were weighed on the farm and soon after being transferred to a commercial slaughterhouse, where they were stunned and slaughtered by 137 cutting the jugular vein. After the slaughter, the non-carcass components were removed and 138 weighed (i.e., skin, head, feet, pluck [heart, lungs, liver, spleen], digestive tract). After 139 chilling at 4 °C for 24 h, the cold carcass weights were recorded, and the dressing percentages 140 were calculated. The right side of each carcass was jointed into three main commercial cuts: 141 shoulder, whole loin with flank, and leg. The weights of each commercial cut were recorded 142 and are expressed as proportions of the half carcass weight. For further analyses, the 143 longissimus lomborum muscles between the first and sixth lumbar vertebrae were obtained. 144

145

146 2.3. Meat quality parameters

The pH of the *longissimus thoracis* muscle (between the tenth and thirteenth thoracic
vertebrae) was measured 45 min and 24 h (final pH) *post-mortem* using a portable pH meter
(XS pH 110; Eutech Instruments, Singapore) equipped with a penetrating glass electrode.

The drip loss and cooking loss were determined on approximately 80 g of 3-cm-thick 150 longissimus lumborum muscle. To measure the drip loss (ASPA, 1996), the meat samples 151 were weighed and wrapped in polyethylene bags. After 24 h storage at 4 °C, the samples were 152 gently dried with paper towels, and reweighed. This procedure was carried out as two 153 replications, and was repeated for the third and the sixth days of storage. For cooking loss 154 determination (ASPA, 1996), the meat samples were initially weighed, and then placed in 155 polyethylene bags and cooked in a water bath until they reached an internal temperature of 75 156 °C. The bags with the cooked meat samples where than cooled under cold running water for 157 30 min, and then they were removed from the bags, dried with paper towels, and reweighed. 158

159 The meat colour was assessed according to the CIELAB system (CIE, 1986), as the lightness (L\*), redness (a\*) and yellowness (b\*) of the longissimus lumborum muscle was 160 recorded for the fresh meat (~24 h post-mortem) using Minolta CR 200. A D65 illuminant 161 was used at an observation angle of 10° and with an aperture of 30 mm. The instrument was 162 calibrated using white and black standard coordinates. Chroma (C\*; square root of  $[a^{*2} +$ 163  $b^{*2}$ ) and hue angle (H°; tan<sup>-1</sup> [b\*/a\*]) were also calculated. For each meat sample, three 164 colour measurements and calculations were performed for each parameter, and the means are 165 reported. After the colour measurements were performed, the meat samples were frozen (-20 166 °C), freeze-dried, ground and analysed for their chemical composition. The procedures 167 outlined by the AOAC (1999) were used to determine the DM (method ID 950.46), Kjeldahl 168 N (crude protein; method ID 981.10), fat (method ID 991.36) and ash (method ID 920.153). 169 Analyses were performed in duplicate for each sample, and the parameters and data are 170 corrected for moisture content. 171

The FA composition analysis was performed on freeze-dried samples of longissimus 174 lomborum muscle (i.e., intramuscular) fat and subcutaneous fat. The fat was extracted using a 175 Soxtec system with petroleum ether (boiling point range, 40-60 °C). The oven temperature for 176 both the pre- and post-extraction drying was within the temperature range suggested by Foss 177 for use with the Soxtec system, as  $102 \pm 3$  °C (Soxtec Avanti 2055 instruction manual). The 178 liquefied fat (~0.1 mL) was collected and dissolved in hexane (1.0 mL). The FA methyl esters 179 (FAMEs) were prepared by 'rapid' KOH-catalysed transesterification, according to method 180 181 ISO 12966-2:2011. The individual FAMEs were analysed on a gas chromatography system (HRGC MEGA 2 series; Fisons Instruments, Milan, Italy) equipped with a Rt-2560 column 182 (length, 100 m, internal diameter, 0.25 mm; df, 0.2 µm; Restek, PA, USA) and a flame-183 184 ionisation detector, with He as the carrier gas.

The identification of the FAs was based on comparisons of retention times with those of commercial standards (Supelco 37 FAMEs Mix; Supelco, Bellefonte, PA, USA). The concentrations of the individual FAMEs were expressed as percentages of the total FAMEs identified, and also grouped as follows: SFAs, monounsaturated FAs (MUFAs) and PUFAs. The PUFA/SFA and n-6/n-3 PUFA ratios were determined, along with the atherogenic and thrombogenic indices (Ulbricht and Southgate, 1991) and the hypocholesterolemic/ hypercholesterolemic ratio (h/H; Fernández et al., 2007).

192

193 2.5. Statistical analysis

The data were analysed using the JMP software version 10 (SAS Institute Inc., Cary, NC,
USA, 1989-2007). Student's t-tests were used to evaluate effects of the two slaughter ages on
all of the parameters.

### 198 **3. Results and discussion**

199

200 The lamb growth performances are given in Table 1. When grouped per slaughter age, the lambs did not differ in birth weight, weight at 20 and 40 days of age, and ADGs through 201 202 the different periods calculated. ADGs were higher than those reported by Santos-Silva et al. (2002a) for Merino Branco and crossbred Ile de France x Merino Branco light lambs, reared 203 with dams on grass and with concentrate ad libitum. The effects of slaughter age on carcass 204 traits and non-carcass components are given in Table 2. As expected, the older lambs had 205 significantly higher pre-slaughter and carcass weights (P <0.001, P <0.05, respectively). 206 According to the European carcass classification system, and based on their average weights, 207 the carcasses from both of the age categories can be classified as class C for light lambs (10.1-208 209 13.0 kg; European Union, 1994).

In agreement with previous studies (Cifuni et al., 2000; Morbidini et al., 1987), 210 211 increased slaughter age significantly decreased the dressing percentages of the light lambs (P 212 <0.001). This can be attributed to the significantly greater weight of offal of the lambs slaughtered at 60 days of age, which negatively affected their dressing percentages. In 213 contrast, Polidori et al. (2017) reported higher dressing percentages for older lambs. These 214 data might be explained on the basis of the different slaughter weights used in these two 215 studies (Polidori et al., 23.8, 37.4 kg; present study, 19.84, 25.99 kg). In the present study, the 216 younger lambs were not functional as ruminants, whereas the older lambs had started to be, 217 which will thus have reduced their dressing percentages, as also explained by Cifuni et al. 218 (2000). As for the present study, D'Alessandro et al. (2013) reported significant differences in 219 220 carcass weight between Leccese lambs slaughtered at 45 and 60 days of age, although they did not report any variations in dressing percentages. The dressing percentages of the lambs 221

slaughtered at 60 days of age were similar to those reported for heavy Bergamasca lambs
(46.3%; Piasentier et al., 2001). However, the dressing percentages of the lambs from both of
these age groups were lower than those reported for Leccese lambs slaughtered at 45 and 60
days (D'Alessandro et al., 2013) and for Apulian lambs slaughtered at 45 and 90 days of age
(Cifuni et al., 2000).

For the non-carcass components, slaughter age influenced all of the parameters investigated (Table 2). The lambs slaughtered at 60 days of age had significantly greater (P <0.001) proportions of head, skin, pluck and full digestive tract, and significantly lower (P <0.01) proportions of feet, than the lambs slaughtered at 40 days of age. All of these differences can be attributed to the older slaughter age and the increased growth of the lambs.

As expected, the older lambs had significantly greater (P <0.01) half-carcass weights 232 than the younger lambs (Table 3). Furthermore, the lambs slaughtered at 60 days of age had 233 234 significantly greater (P < 0.01) proportions of leg than the lambs slaughtered at 40 days of age, although the other commercial cuts did not differ between the slaughter ages. In a study by 235 236 Cifuni et al. (2000), first grade wholesale lamb cuts (i.e., rack, loin, leg) were not significantly 237 affected by slaughter age, while the proportions of second grade cuts (i.e., shoulder, neck, breast) decreased as age of slaughter increased. These data were attributed to the greater 238 shoulder and breast proportions of the younger animals. The proportions of leg in the present 239 240 study was higher than that reported by Skapetas et al. (2006) for lambs slaughtered at 45 and 60 days, while the proportions of shoulder were similar to those reported by Skapetas et al. 241 (2006). Similar proportions of shoulder were reported by D'Alessandro et al. (2013) for lambs 242 slaughtered in spring at a slaughter age of 45 days. However, contrary to the present data, 243 D'Alessandro et al. (2013) obtained higher proportions of leg for Leccese lambs slaughtered 244 245 at 45 days of age, compared to those slaughtered at 60 days of age. These contrary results

246 might be attributed to different methods used to dissect the carcasses into the main
247 commercial cuts.

The age at slaughter did not affect the meat pH at 45 min and 24 h *post-mortem* (Table 4). These data are in agreement with other studies where the pHs obtained did not differ between light lambs slaughtered at different ages or weights (D'Alessandro et al., 2013; della Malva et al., 2016; Juárez et al., 2009). The final pHs were within the normal ranges (Tejeda et al., 2008) and were similar to those reported for light lambs of different breeds in other studies (D'Alessandro et al., 2013; della Malva et al., 2016; Juárez et al., 2009; Mazzone et al., 2010).

255 No differences were seen between the lambs slaughtered at 40 and 60 days of age in terms of drip loss measured after 1 day of meat storage (Table 4). The drip losses were lower 256 than those reported for Appenninica light lambs (Mazzone et al., 2010). Differences in drip 257 258 loss between lambs of the two slaughter ages appeared after the third day, and continued through the sixth day of storage, whereby the meat from the younger lambs had significantly 259 higher (P <0.001) drip loss than that from the older lambs. The results might be attributed to 260 faster weakening of myofibrillar proteins by proteolytic enzymes during the storage of meat 261 of younger lambs, thereby affecting the ability of myofibrillar proteins to hold water (Lawrie 262 and Ledward, 2006). These data are in agreement with Russo et al. (2003), who reported 263 better water-holding capacity for heavier carcasses due to lower drip loss after the second day 264 of storage. 265

The cooking losses did not differ between the lambs slaughtered at 40 and 60 days of age (Table 4), and they were lower than those reported for light Appenninica lambs of similar slaughter age (Mazzone et al., 2010). On the contrary, although another method was used to determine their cooking loss, Russo et al. (2003) reported increased cooking losses with increased carcass weight. The lack of any differences in cooking losses between the lambs of the two slaughter age groups in the present study can be explained on the basis that contrary
to Russo et al. (2003), the differences in the carcass weights between the age groups of these
light lambs were lower.

274 The meat chemical compositions of lambs slaughtered at 40 and 60 days of age are given in Table 5. These data are comparable with those reported for light lambs of different 275 Italian breeds (D'Alessandro et al., 2013; della Malva et al., 2016; Mazzone et al., 2010). 276 277 Taking into consideration the intramuscular fat content, the meat analysed in the present study was of good quality, on the basis that 2% to 3% of intramuscular fat ensures the desirable 278 organoleptic properties of the meat (Wood, 1990). No significant differences were recorded 279 280 for moisture, crude protein, fat and ash of the longissimus lomborum muscle in terms of the slaughter ages. 281

These data are in agreement with other studies that have highlighted that the chemical composition of lamb meat is not affected by slaughter age (D'Alessandro et al., 2013; Marino et al., 2008) or carcass weight (Russo et al., 2003). In general, greater intramuscular fat has been reported for older lambs of different breeds (Abdullah and Qudsieh, 2009; della Malva et al., 2016; Juárez et al., 2009; Pérez et al., 2002).

In contrast to D'Alessandro et al. (2013), who observed no effects of slaughter age on 287 any of the meat colour parameters of light lambs, in the present study, some colour 288 differences were recorded (Table 5). Contrary to data reported for other breeds (Abdullah and 289 Qudsieh, 2009; della Malva et al., 2016; Juárez et al., 2009; Santos-Silva et al., 2002a; 290 Sañudo et al., 1996), in the present study, increased age and weight of the lambs did not result 291 in darker meat. Indeed, no differences were found for the lightness (L\*), yellowness (b\*) and 292 Hue and Chroma between the meat of the lambs slaughtered at 40 and 60 days of age. There 293 was significantly greater (P < 0.01) redness (a<sup>\*</sup>) for the meat of the older and heavier lambs, 294 which is in agreement with other studies (della Malva et al., 2016; Juárez et al., 2009; Sañudo 295

et al., 1996). This can be explained by the higher myoglobin content in the meat of older
lambs (Juárez et al., 2009). Furthermore, the colour differences for the meat from the lambs
slaughtered at different ages might be due to the higher consumption of milk by the younger
lambs, compared to the older ones that used proportionally more concentrate before their
slaughter (Sañudo et al., 1996).

The FA compositions of the intramuscular and subcutaneous fat are given in Tables 6 and 7, respectively. In general, the intramuscular FA composition in the present study was similar to that reported by others for light lambs with high milk content in the diet (D'Alessandro et al., 2015; Cifuni et al., 2000; Oriani et al., 2005). The total SFAs were the most abundant FAs, followed by MUFAs and PUFAs.

Slaughter age had relatively small influences on the FA composition of longissimus 306 lomborum muscle, whereby lambs slaughtered at 60 days of age had significantly greater (P 307 308 <0.05) tridecylic (C13:0) and palmitic (C16:0) SFAs than those slaughtered at 40 days of age (Table 6). The saturated palmitic FA has been shown to be hyperlipidemic (Bonamone and 309 Grundy, 1988), and therefore consumption of meat from the younger lambs might reduce the 310 311 risk of heart disease. This confirms data from other studies, where greater amounts of C16:0 were reported for the intramuscular fat of older and heavier lambs (Cifuni et al., 2000; 312 D'Alessandro et al., 2015; della Malva et al., 2016; Santos-Silva et al., 2002b). However, the 313 same studies reported better FA profile for the meat of the younger (less heavy) lambs, due to 314 an increase in the proportion of SFAs with lamb growth. This reported trend for intramuscular 315 fat was not evident in the present study, as the slaughter age did not affect any of the other 316 317 single SFAs, MUFAs or PUFAs in the lamb intramuscular fat. Consequently, the nutritional indices of the meat, which are defined here as the PUFA/SFA, n-6/n-3 and h/H ratios and the 318 atherogenic and thrombogenic indices, were similar between these Bergamasca lambs 319 slaughtered at 40 and 60 days of age. 320

For the FA composition of the subcutaneous fat, there were much more evident effects 321 322 of slaughter age (Table 7). Here, the lambs slaughtered at 40 days of age showed a better FA composition, in terms of significantly lower (P <0.01) SFAs, and significantly greater (P 323 <0.01) MUFAs and PUFAs than the lambs slaughtered at 60 days of age. Unweaned lambs 324 have a rumen that is only partially functional, and therefore their dietary unsaturated FAs are 325 326 likely to be digested and adsorbed as in non-ruminants (Sañudo et al., 1998). In contrast, the 327 older lambs start to consume a dry diet, which needs a more functional rumen to be digested. Consequently, the greater SFAs in the subcutaneous fat of the older lambs that here used 328 proportionally more concentrate before their slaughter, will be due to the extensive 329 330 hydrogenation by micro-organisms in the rumen.

In contrast to these data, Juárez et al. (2009) reported that for the FA composition of subcutaneous fat of suckling and light lambs, weight and age only influenced some of the individual SFAs, while the total SFAs, MUFAs and PUFAs, and the PUFA/SFA ratio, did not differ between the lambs of their two slaughter weights.

In the present study, the slaughter age affected the proportion of almost all of the SFAs 335 336 investigated. In particular, the subcutaneous fat from the older lambs showed significantly greater proportions of capric (C10:0; P <0.01), lauric (C12:0; P <0.05), tridecylic (P <0.01), 337 myristic (C14:0; P <0.01), pentadecanoic (C15:0; P <0.01) and palmitic (P <0.001) SFAs. It 338 has been reported that short chain SFAs C12:0, C14:0 and C16:0 can increase plasma low-339 density lipoprotein (LDL)-cholesterol (Siri-Tarino et al., 2010). Therefore, the subcutaneous 340 fat from the Bergamasca lambs slaughtered at 60 days showed less favourable levels of SFAs 341 than that of lambs slaughtered at 40 days of age. On the other hand, the subcutaneous fat of 342 the younger lambs had significantly greater heptadecanoic (C17:0; P <0.01) and stearic 343 (C18:0; P <0.001) SFAs. However, although stearic acid is a saturated FA, it will not elevate 344

plasma LDL-cholesterol (Bonamone and Grundy, 1988) as it is poorly digested and can be
easily desaturated to oleic acid.

For the MUFAs, the lambs slaughtered at 40 days of age had significantly lower (P 348 <0.01) myristoleic (C14:1) and palmitoleic (C16:1) acids, but significantly higher (P <0.01) 349 sum of oleic and cis-vaccenic acid ( $\sum$ C18:1c) in their subcutaneous fat, than the lambs 350 slaughtered at 60 days of age. Oleic acid has been reported to be hypolipidemic, which can 351 thus reduce plasma cholesterol and triglycerides (Mattson and Grundy, 1985). Therefore, 352 oleic acid can be considered as a desirable component of the diet (Baer et al., 2014).

The slaughter age also showed significantly greater (P <0.05) proportions of single PUFAs in the subcutaneous fat for the lambs slaughtered at 60 days, in terms of C18:2 trans and C20:2 n-6. In agreement with Juárez et al. (2009), the older lambs showed significantly lower linoleic acid (C18:2 n-6; P <0.001). Furthermore, the essential linolenic acid (C18:3 n-3) that has been shown to have positive health benefits was significantly lower (P <0.05) in the meat of the older lambs.

With the increased slaughter age, there were significantly decreased total PUFAs for both the n-6 (P <0.001) and n-3 (P <0.05) PUFAs. However, the ratios between these (i.e., n-6/n-3) did not differ for Bergamasca lambs between the two slaughter ages, and they both remained within the recommended value for inclusion in a human diet (i.e., <4) (Wood et al., 2003).

The lambs slaughtered at 40 days had subcutaneous fat that showed an improved PUFA/SFA ratio, as this was significantly greater (P <0.01) compared to the lambs slaughtered at 60 days of age. However, the PUFA/SFA ratios of these slaughter age groups were both low, and they did not reach the recommended value for human health (i.e., >0.4) (Wood et al., 2003). The overall changes in the FA compositions of the subcutaneous fat according to increased slaughter age showed significantly lower h/H (P <0.01), which has a negative effect on the nutritional value of the meat. The subcutaneous fat of lambs slaughtered at 40 days of age also had significantly lower, and thus more favourable, atherogenic and thrombogenic indices (P <0.01, P <0.001, respectively), compared to the lambs slaughtered at 60 days of age.

375

### **4. Conclusions**

The increase in the slaughter age for these Bergamasca lambs from 40 days to 60 days resulted in improved carcass weight, a greater proportion of leg commercial cut, and a lower drip loss after meat storage. For light lamb production using the traditional rearing systems, slightly heavier lambs can be produced without worsening chemical composition, cooking loss and fatty acid composition of the *longissimus lomborum* muscle. However, it is important to take in consideration that these heavier lambs might have lower dressing percentages and less favourable FA profile of their subcutaneous fat for human health.

For a more complete understanding of the effects of slaughter age on carcass and meat quality, additional studies are needed. These can be designed to evaluate more specifically the meat yield (i.e., from carcass or leg dissection), the effects of lamb nutrition (i.e., FA composition of feed and dams' milk), and the feed intake of the lambs.

- 390
- Abdullah, A.Y., Qudsieh, R.I., 2009. Effect of slaughter weight and aging time on the quality
  of meat from Awassi ram lambs. Meat Sci. 82, 309–316.
  doi:10.1016/j.meatsci.2009.01.027
- AOAC, 1999. Official method 950.46 moisture in meat; Official method 991.36 fat (crude) in
   meat and meat products; Official method 981.10 crude protein in meat; Official method
- 396 920.153 ash in meat. In: Cunniff. P. (Ed.), Meat and Meat Products, vol. II. 16th ed.,
- 397 Official methods of analysis of the AOAC International, Gaithersburg, MD, pp. 1–15
  398 (Chapter 39).
- ASPA, 1996. Metodiche per la determinazione delle caratteristiche qualitative della carne.
  Università di Perugia, Perugia, Italy.
- Baer, D.J., Judd, J.T., Clevidence, B.A., Tracy, R.P., 2004. Dietary fatty acids affect plasma
  markers of inflammation in healthy men fed controlled diets: a randomized crossover
  study. Am. J. Clin. Nutr. 79, 969–973.
- Beriain, M.J., Horcada, A., Purroy, A., Lizaso, G., Chasco, J., Mendizabal, J.A., 2000.
  Characteristics of Lacha and Rasa Aragonesa lambs slaughtered at three live weights. J.
  Anim. Sci. 78, 3070–3077.
- Bigi, D, Zanon, A., 2008. Atlante delle razze autoctone italiane: Bovini, Equini, Ovicaprini,
  Suini allevati in Italia. Edagricole Edizioni Agricole de il Sole 24 Ore, Milan.
- Bonamone, A., Grundy, S.M., 1988. Effect of dietary stearic acid on plasma cholesterol and
  lipoprotein levels. New Engl. J. Med. 318, 1244–1248.
- 411 Caballero R., Fernandez-Gonzalez F., Perez Badia R., Molle G., Roggero P.P., Bagella S.,
- 412 D'Ottavio P., Papanastasis V.P., Fotiadis G., Sidiropoulou A., Ipiloudis I., 2009. Grazing
- 413 systems and biodiversity in Mediterranean areas: Spain, Italy and Greece. Pastos 39, 9–

414 154.

- Calder, P.C., Yaqoob, P., 2009. Omega-3 polyunsaturated fatty acids and human health
  outcomes. Biofactors. 35, 266–272.
- 417 CIE, 1986. Colorimetry, 2nd ed. Vol. nº 15.2. Centre International de L'eclairage, Vienna.
- 418 Cifuni, G., Napolitano, F., Pacelli, C., Riviezzi, A., Girolami, A., 2000. Effect of age at
- slaughter on carcass traits, fatty acid composition and lipid oxidation of Apulian lambs.
  Small Rumin. Res. 35, 65–70. doi: 10.1016/s0921-4488(99)00060-7
- 421 D'Alessandro, A.G., Maiorano, G., Ragni, M., Casamassima, D., Marsico, G., Martemucci,
- 422 G., 2013. Effects of age and season of slaughter on meat production of light lambs:
- 423 Carcass characteristics and meat quality of Leccese breed. Small Rumin. Res. 114, 97–
- 424 104. doi:10.1016/j.smallrumres.2013.05.006
- D'Alessandro, A.G., Palazzo, M., Petrotos, K., Goulas, P., Martemucci, G., 2015. Fatty acid
  composition of light lamb meat from Leccese and Comisana dairy breeds as affected by
- slaughter age. Small Rumin. Res. 127, 36–43. doi:10.1016/j.smallrumres.2015.04.004
- D'Ottavio, P., Santilocchi, R., 2014. Mountain areas of the Marche Region (Central
  Apennine), in: Huyghe, C., De Vliegher, A., Van Gils, B., Peeters, A. (Eds.), Grasslands
  and herbivore production in Europe and effects of common policies. Éditions Quæ, pp.
  179–183.
- della Malva, A., Albenzio, M., Annicchiarico, G., Caroprese, M., Muscio, A., Santillo, A., 432 Marino, R., 2016. Relationship between slaughtering age, nutritional and organoleptic 433 properties Altamurana lamb meat. Small Rumin. Res. 39-45. 434 of 135. doi:10.1016/j.smallrumres.2015.12.020 435
- Ekiz, B., Demirel, G., Yilmaz, A., Ozcan, M., Yalcintan, H., Kocak, O., Altinel, A., 2013.
  Slaughter characteristics, carcass quality and fatty acid composition of lambs under four
  different production systems. Small Rumin. Res. 114, 26–34.

- 439 doi:10.1016/j.smallrumres.2013.05.011
- 440 European Union, 1994. European Community Standard for the Classification of Light Lambs
- 441 Carcasses, Brochure No. CM-84-94-703-ES-D. Publishing Bureau of the European
  442 Communities, L-2985 Luxembourg.
- Fernández, M., Ordóñez, J.A., Cambero, I., Santos, C., Pin, C., de la Hoz, L., 2007. Fatty acid
  compositions of selected varieties of Spanish dry ham related to their nutritional
  implications. Food Chem. 101, 107–112. doi:10.1016/j.foodchem.2006.01.006
- 446 Font i Fournols, M., Guerrero, L., 2014. Consumer preferences, behaviour and perception
  447 about meat and meat production: An overview. Meat Sci. 98, 361–371.
- Guerrero, A., Valero, M.V., Campo, M.M., Sañudo, C., 2013. Some factors that affect
  ruminant meat quality: from the farm to the fork. Review. Acta Sci. Anim. Sci. 35, 335347. doi: 10.4025/actascianimsci.v35i4.21756. Acta Sci. Anim. Sci. 35, 335–347.
- 451 Hajji, H., Joy, M., Ripoll, G., Smeti, S., Mekki, I., Gahete, F.M., Mahouachi, M., Atti, N., 2016. Meat physicochemical properties, fatty acid profile, lipid oxidation and sensory 452 characteristics from three North African lamb breeds, as influenced by concentrate or 453 pasture finishing diets. J. Food Compos. Anal. 48. 102–110. 454 doi:10.1016/j.jfca.2016.02.011 455
- ISO 12966-2:2011. Animal and vegetable fats and oils Gas chromatography of fatty acid
  methyl esters-Part 2: Preparation of methyl esters of fatty acids.
- Jacques, J., Berthiaume, R., Cinq-Mars, D., 2011. Growth performance and carcass
  characteristics of Dorset lambs fed different concentrates: Forage ratios or fresh grass.
  Small Rumin. Res. 95, 113–119. doi:10.1016/j.smallrumres.2010.10.002
- 461 JMP, Version 10. SAS Institute Inc., Cary, NC, 1989-2007.
- 462 Juárez, M.M., Horcada, A., Alcalde, M.J., Valera, M., Polvillo, O., Molina, A., 2009. Meat
- 463 and fat quality of unweaned lambs as affected by slaughter weight and breed. Meat Sci.

- 464 83, 308–313. doi:10.1016/j.meatsci.2009.05.017
- 465 Lawrie, R. A., Ledward, D. A., 2006. Lawrie's Meat Science. Seventh English, edition ed.
- 466 Cambridge England: Woodhead Publishing Limited.

- Lind, V., Berg, J., Eik, L.O., Mølmann, J., Haugland, E., Jørgensen, M., Hersleth, M., 2009.
- Meat quality of lamb: Pre-slaughter fattening on cultivated or mountain range pastures.
  Meat Sci. 83, 706–712. doi:10.1016/j.meatsci.2009.08.008
- 470 Marino, R., Albenzio, M., Annicchiarico, G., Caroprese, M., Muscio, A., Santillo, A., Sevi,
- 472 Trimeticcio lambs. Small Rumin. Res. 78, 144–151.
  473 doi:10.1016/j.smallrumres.2008.06.002

A., 2008. Influence of genotype and slaughtering age on meat from Altamurana and

- 474 Martillotti, F., Antongiovanni, M., Rizzi, L., Santi, E., Bittante, G., 1987. Metodi di analisi
  475 per la valutazione degli alimenti d'impiego zootecnico. Quaderni metodologici N. 8.
  476 IPRA, CNR, Rome.
- 477 Mattson, F.M., Grundy, S.M., 1985. Comparison of effects of dietary saturated,
  478 monounsaturated, and polyunsaturated fatty acids on plasma lipids and lipoproteins in
  479 man. J. Lipid Res. 26, 194–202.
- 480 Mazzone, G., Giammarco, M., Vignola, G., Sardi, L., Lambertini, L., 2010. Effects of the
  481 rearing season on carcass and meat quality of suckling Apennine light lambs. Meat Sci.
- 482 86, 474–478. doi:10.1016/j.meatsci.2010.05.037
- Morbidini, L., Panella, F., Polidori, P., Sarti, D.M., 1987. Observation on productivity of
  Apennine sheep. Z.N.A. 13, 147–155.
- 485 Oriani, G., Maiorano, G., Filetti, F., Di Cesare, C., Manchisi, A., Salvatori, G., 2005. Effect of
- 486 age on fatty acid composition of Italian Merino suckling lambs. Meat Sci. 71, 557–562.
  487 doi:10.1016/j.meatsci.2005.04.040
- 488 Pérez, P., Maino, M., Tomic, G., Mardones, E., Pokniak, J., 2002. Carcass characteristics and

- 489 meat quality of Suffolk Down suckling lambs. Small Rum. Res. 44, 233–240.
- 490 Piasentier, E., 2003. The effect of grazing on the quality of lamb meat. International
  491 Fachtagung Für Schafhaltung. Innsbruck.
- 492 Piasentier, E., Valusso, R., Gasperi, F., 2001. Effect of production system on the carcass and
  493 meat quality of Bergamasca lamb. In: Proceedings of the ASPA XIV Congress "Recent
  494 Progress in Animal Production Science", Firenze (Italy), June 12-15, 2001. Pp. 550-552.
- Polidori, P., Pucciarelli, S., Cammertoni, N., Polzonetti, V., Vincenzetti, S., 2017. The effects
  of slaughter age on carcass and meat quality of Fabrianese lambs. Small Rumin. Res.
  155, 12–15. doi:10.1016/j.smallrumres.2017.08.012
- Priolo, A., Micol, D., Agabriel, J., Prache, S., Dransfield, E., 2002. Effect of grass or
  concentrate feeding systems on lamb carcass and meat quality. Meat Sci. 62, 179–185.
- 500 doi:10.1016/S0309-1740(01)00244-3
- Russo, C., Prziuso, G., Verità, P., 2003. EU carcass classification system: Carcass and meat
  quality in light lambs. Meat Sci. 64, 411–416. doi:10.1016/S0309-1740(02)00209-7
- 503 Santos-Silva, J., Mendes, I.A., Bessa, R.J.B., 2002a. The effect of genotype, feeding system
- and slaughter weight on the quality of light lambs: 1. Growth, carcass composition and
  meat quality. Livest. Prod. Sci. 76, 17–25.
- Santos-Silva, J., Bessa, R.J.B., Santos-Silva, F., 2002b. Effect of genotype, feeding system
  and slaughter weight on the quality of light lambs. II. Fatty acid composition of meat.
  Livest. Prod. Sci. 77, 187–194. doi:10.1016/S0301-6226(02)00059-3
- Sañudo, C., Santolaria, M.P., Maria, G., Osorio, M., Sierra, I., 1996. Influence of carcass
  weight on instrumental and sensory lamb meat quality in intensive production systems.
  Meat Sci. 42, 195–202.
- 512 Sañudo, C., Sierra, I., Olleta, J.L., Martin, L., Campo, M.M., Santolaria, P., Wood, J.D., Nute,
- 513 G.R., 1998. Influence of weaning on carcass quality, fatty acid composition and meat

- 514 quality in intensive lamb production system. Anim. Sci. 66, 175–187
- Siri-Tarino, P.W., Sun, Q., Hu, F.B., Krauss, R.M., 2010. Saturated fatty acids and risk of
  coronary heart disease: modulation by replacement nutrients. Curr Atheroscler. Rep. 12,
  384–390.
- 518 Skapetas, B., Sinapis, E., Hatziminaoulglou, J., Karalazos, A., Katanos, J., 2006. Effect of age
- 519 at slaughter on carcass characteristics and carcass composition in lambs of mountain

520 Greek breed. Czech J. Anim. Sci. 51, 311–317.

- Tejeda, J.F., Pena, R.E., Andres, A.I., 2008. Effect of live weight and sex on physicochemical and sensorial characteristics of Merino lamb meat, Meat Sci. 80, 1061–1067.
- 523 Ulbricht, T.L.V., Southgate, D.A.T., 1991. Coronary heart disease: seven dietary factors.
  524 Lancet 338, 985–992. doi:10.1016/0140-6736(91)91846-M
- 525 Wood, J.D., Richardson, R.I., Nute, G.R., Fisher, A.V., Campo, M.M., Kasapidou, E., Sheard,
- P.R., Enser, M., 2003. Effects of fatty acids on meat quality: a review. Meat Sci. 66, 21–
  32.
- Wood, J.D., 1990. Consequences for meat quality of reducing carcass fatness. In: Wood, J.D.
  Fisher, A.V. (Eds.), Reducing Fatness in Meat Animals. Elsevier Applied Science,
  London, UK, 344–1035.
- World Health Organisation, 2003. Diet, nutrition and the prevention of chronic disease.
  Report of a joint WHO/FAO expert consultation 916.
- 533
- 534
- 535
- 536
- 537
- 538

## 540 **Table 1**

542 SD).

	Slaughter age (o	Slaughter age (days)	
	40	60	
Live Weight (kg)			
Birth	$5.37 \pm 1.05$	$5.11\pm0.83$	NS
20 days	$11.79\pm2.25$	$11.77 \pm 1.22$	NS
40 days	$19.84\pm3.07$	$19.59\pm2.35$	NS
60 days		$25.95\pm3.34$	
Average Daily Gain (g d <sup>-1</sup> )			
birth to 20 days	$318\pm73.19$	$337\pm72.57$	NS
20 to 40 days	$383\pm56.82$	$373\pm63.01$	NS
total rearing period	$351\pm60.23$	$337\pm61.93$	NS

543 NS: not significant, P > 0.05

<sup>541</sup> Growth performance of Bergamasca lambs slaughtered at 40 and 60 days of age (means  $\pm$ 

546 Carcass traits and non-carcass components of Bergamasca lambs slaughtered at 40 and 60

<sup>547</sup> days of age (means  $\pm$  SD).

	Slaughter age (day	Slaughter age (days)	
	40	60	
Pre-slaughter weight (kg)	$19.84\pm3.07$	$25.95\pm3.34$	***
Cold carcass weight (kg)	$10.36\pm1.75$	$12.44 \pm 2.14$	*
Cold dressing (%)	$52.16\pm2.50$	$47.68\pm2.65$	***
Proportion (%) on PSW			
Head	$2.63\pm0.23$	$3.92\pm0.32$	***
Skin	$9.09\pm0.60$	$11.54\pm0.85$	***
Feet	$1.01 \pm 0.11$	$0.83\pm0.10$	**
Pluck	$5.39\pm0.93$	$7.69\pm0.44$	***
Full digestive tract	$13.86\pm2.30$	$17.96 \pm 2.94$	* * *

548 PSW: pre-slaughter weight.

549 \* P < 0.05.

- 550 \*\* P < 0.01.
- 551 \*\*\* P < 0.001.

Half carcass weight and proportions of commercial cuts of Bergamasca lambs slaughtered at

	Slaughter age (	lays)	Significance
	40	60	
Half carcass weight (kg)	$4.80\pm0.85$	$6.06 \pm 1.07$	**
Proportions of commercial cuts (%)			
Shoulder	$19.81\pm0.88$	$19.53\pm1.36$	NS
Whole loin with flank	$44.69 \pm 1.71$	$42.65\pm2.84$	NS
Leg	$35.49 \pm 1.06$	$37.82 \pm 1.85$	**
NS: not significant, $P > 0.05$ .			

555 40 and 60 days of age (means  $\pm$  SD).

557 \*\* P < 0.01.

pH, drip loss and cooking loss of Bergamasca lambs slaughtered at 40 and 60 days of age 560

561 (means	$\pm$ SD).
------------	------------

	Slaughter age (days)		Significance	
	40	60		
pН				
45 min	$6.41 \pm 0.14$	$6.54\pm0.35$	NS	
24 h	$5.71\pm0.11$	$5.69\pm0.14$	NS	
Drip loss (%)				
1 day	$0.77\pm0.17$	$1.01 \pm 0.53$	NS	
3 days	$6.16 \pm 1.41$	$3.69\pm0.97$	***	
6 days	$9.36 \pm 1.45$	$5.73 \pm 1.16$	* * *	
Cooking loss (%)	$12.22 \pm 3.85$	$10.53\pm4.94$	NS	

562 NS: not significant, P > 0.05.

\*\*\* P < 0.001. 563

565 Chemical composition and colour of *longissimus lomborum* muscle of Bergamasca lambs

566	slaughtered at 40	and 60	days of age	(means $\pm$ SD).
-----	-------------------	--------	-------------	-------------------

	Slaughter age (days)	Slaughter age (days)	
	40	60	
Chemical composition (%)			
Moisture	$73.45 \pm 1.35$	$74.55 \pm 1.56$	NS
Crude Protein	$19.98\pm0.50$	$19.68\pm0.62$	NS
Fat	$2.70 \pm 1.11$	$2.15\pm0.91$	NS
Ash	$1.31\pm0.07$	$1.39\pm0.25$	NS
Colour parameters			
L*	$41.52 \pm 2.13$	$41.11 \pm 3.07$	NS
a*	$18.91 \pm 1.59$	$19.43 \pm 1.21$	*
b*	$3.3\pm0.55$	$3.43\pm0.93$	NS
Hue	$0.17\pm0.03$	$0.18\pm0.05$	NS
Chroma	$30.2 \pm 4.61$	$32.13 \pm 8.19$	NS

567 NS: not significant, P > 0.05.

568 \*\* P < 0.01.

571 Fatty acid composition (% of total fatty acids) of longissimus lomborum muscle of

572	Bergamasca	lambs slaughtered at 40 a	and 60 days of age (means $\pm$ SI	)).
-----	------------	---------------------------	------------------------------------	-----

	Slaughter age (da	Slaughter age (days) Significance	
	40	60	
C10:0	$0.64\pm0.24$	$0.75 \pm 0.18$	NS
C11:0	$0.04\pm0.01$	$0.04\pm0.01$	NS
C12:0	$1.53\pm0.57$	$1.84\pm0.50$	NS
C13:0	$0.10\pm0.03$	$0.12\pm0.02$	**
C14:0	$9.78 \pm 1.90$	$10.77\pm2.05$	NS
C15:0	$0.97\pm0.17$	$1.01\pm0.15$	NS
C16:0	$24.43 \pm 1.84$	$26.09\pm0.90$	**
C17:0	$1.20\pm0.17$	$1.12\pm0.10$	NS
C18:0	$11.98 \pm 1.72$	$11.18\pm1.79$	NS
C20:0	$0.10\pm0.04$	$0.08\pm0.03$	NS
C21:0	$0.15\pm0.07$	$0.16\pm0.03$	NS
C14:1	$0.29\pm0.11$	$0.34\pm0.11$	NS
C16:1 n-7	$1.78\pm0.40$	$1.90\pm0.32$	NS
C17:1	$0.55\pm0.14$	$0.55\pm0.10$	NS
$\sum 18.1t^{1}$	$3.31 \pm 1.59$	$3.05 \pm 1.21$	NS
$\sum 18:1c^{1}$	$30.14\pm3.68$	$28.12\pm3.29$	NS
$\sum 18:2t^1$	$1.58\pm0.26$	$1.65\pm0.27$	NS
C18:2 n-6	$4.2 \pm 1.36$	$3.56\pm0.83$	NS
C18:2 cis-9, trans-11(CLA)	$1.50\pm0.30$	$1.64\pm0.40$	NS
C18:3 n-3	$1.57\pm0.32$	$1.53\pm0.24$	NS
C20:2 n-6	$0.09\pm0.03$	$0.07\pm0.01$	NS
C20:4 n-6	$0.53\pm0.39$	$0.63\pm0.29$	NS
C20:5 n-3	$0.26 \pm 0.19$	$0.29\pm0.12$	NS
C22:5 n-3	$0.34\pm0.23$	$0.36\pm0.14$	NS
C22:6 n-3	$0.14 \pm 0.10$	$0.14\pm0.07$	NS
Other fatty acids	$2.79\pm0.26$	$2.99\pm0.19$	NS
SFA	$50.91\pm3.89$	$53.17\pm2.35$	NS
MUFA	$32.76 \pm 3.97$	$30.91 \pm 3.49$	NS
PUFA	$8.64 \pm 2.46$	$8.23 \pm 1.32$	NS
n-6 PUFA	$4.82 \pm 1.67$	$4.27 \pm 1.01$	NS
n-3 PUFA	$2.32 \pm 0.82$	$2.33\pm0.50$	NS
n-6/n-3 PUFA	$2.13 \pm 0.59$	$1.88 \pm 0.53$	NS
PUFA/SFA	$0.17 \pm 0.06$	$0.16 \pm 0.02$	NS
h/H	$1.11 \pm 0.21$	$0.95 \pm 0.15$	NS
AI	$1.66 \pm 0.40$	$1.92 \pm 0.35$	NS
TI	$1.79\pm0.31$	$1.92 \pm 0.17$	NS

<sup>573</sup>  $\sum$ C18:1t: sum of unidentified C18:1 *trans* isomers;  $\sum$ C18:1c: C18:1 $\Delta$ 9c +  $\Delta$ 11c;  $\sum$ C18:2t,

574 18:2 n-6 c,t + t,c + t,t; SFA: saturated fatty acids; MUFA: monounsaturated fatty acids;

575 PUFA: polyunsaturated fatty acids; n-6 PUFA: omega 6 fatty acids; n-3 PUFA: omega 3 fatty

- acids; PUFA/SFA: polyunsaturated/ saturated fatty acid ratio; h/H: hypocholesterolemic/ hypercholesterolemic ratio (Fernández et al., 2007); AI: atherogenic index (Ulbricht and Southgate, 1991); TI: thrombogenic index (Ulbricht and Southgate, 1991); NS: not significant, P > 0.05.; \*\* P < 0.01.
- 580

582 Fatty acid composition (% of total fatty acids) of subcutaneous adipose tissue of Bergamasca

	Slaughter age (days)		Significance	
	40	60		
C10:0	$0.63\pm0.18$	$0.86\pm0.11$	**	
C11:0	$0.03\pm0.01$	$0.04\pm0.01$	**	
C12:0	$1.09\pm0.39$	$1.50\pm0.30$	*	
C13:0	$0.06\pm0.02$	$0.08 \pm 0.02$	**	
C14:0	$7.42 \pm 1.55$	$9.28 \pm 1.24$	**	
C15:0	$0.72 \pm 0.18$	$0.90 \pm 0.09$	**	
C16:0	$19.58 \pm 2.13$	$22.99 \pm 1.51$	***	
C17:0	$1.73\pm0.17$	$1.51 \pm 0.06$	**	
C18:0	$20.94\pm2.23$	$18.23\pm1.68$	**	
C20:0	$0.17\pm0.03$	$0.17\pm0.05$	NS	
C21:0	$0.09\pm0.03$	$0.10\pm0.01$	NS	
C14:1	$0.10\pm0.02$	$0.14 \pm 0.02$	**	
C16:1 n-7	$0.80\pm0.09$	$0.94\pm0.12$	**	
C17:1	$0.42\pm0.06$	$0.39\pm0.06$	NS	
$\sum 18:1t^1$	$2.84\pm2.14$	$3.82\pm1.79$	NS	
$\overline{\Sigma}$ 18:1c <sup>1</sup>	$32.74\pm2.93$	$28.69 \pm 2.51$	**	
$\overline{\Sigma}18:2t^1$	$1.59\pm0.15$	$1.72\pm0.14$	*	
C18:2 n-6	$2.92\pm0.33$	$2.44\pm0.14$	***	
C18:2 <i>cis</i> -9, <i>trans</i> -11(CLA)	$1.71\pm0.29$	$1.71\pm0.20$	NS	
C18:3 n-3	$1.57\pm0.26$	$1.37\pm0.11$	*	
C20:2 n-6	$0.03\pm0.01$	$0.04\pm0.01$	*	
C20:4 n-6	$0.05\pm0.01$	$0.05\pm0.02$	NS	
C20:5 n-3	$0.02\pm0.01$	$0.02\pm0.01$	NS	
C22:5 n-3	$0.12\pm0.02$	$0.13\pm0.03$	NS	
C22:6 n-3	$0.04\pm0.01$	$0.03\pm0.02$	NS	
Other fatty acids	$2.59\pm0.26$	$2.83\pm0.10$	*	
SFA	$52.46 \pm 2.60$	$55.68 \pm 2.33$	**	
MUFA	$34.06\pm2.93$	$30.16\pm2.53$	**	
PUFA	$6.46\pm0.62$	$5.79\pm0.34$	**	
n-6 PUFA	$3.00\pm0.33$	$2.52\pm0.13$	***	
n-3 PUFA	$1.75\pm0.27$	$1.56\pm0.09$	*	
n-6/n-3 PUFA	$1.73\pm0.15$	$1.62\pm0.09$	NS	
PUFA/SFA	$0.12\pm0.01$	$0.11 \pm 0.01$	**	
h/H	$1.42\pm0.29$	$1.03 \pm 0.15$	**	
AI	$1.32\pm0.31$	$1.82 \pm 0.30$	**	
TI	$1.98 \pm 0.06$	$2.35 \pm 0.07$	***	

lambs slaughtered at 40 and 60 days of age (means  $\pm$  SD).

585 18:2 n-6 c,t + t,c + t,t; SFA: saturated fatty acids; MUFA: monounsaturated fatty acids;

586 PUFA: polyunsaturated fatty acids; n-6 PUFA: omega 6 fatty acids; n-3 PUFA: omega 3 fatty

<sup>584</sup>  $\sum$ C18:1t: sum of unidentified C18:1 *trans* isomers;  $\sum$ C18:1c: C18:1 $\Delta$ 9c +  $\Delta$ 11c;  $\sum$ C18:2t,

acids; PUFA/SFA: polyunsaturated/ saturated fatty acid ratio; h/H: hypocholesterolemic/ hypercholesterolemic ratio (Fernández et al., 2007); AI: atherogenic index (Ulbricht and Southgate, 1991); TI: thrombogenic index (Ulbricht and Southgate, 1991); NS: not significant, P > 0.05; \* P < 0.05; \*\* P < 0.01; \*\*\* P < 0.001.

591