



UNIVERSITÀ POLITECNICA DELLE MARCHE
Repository ISTITUZIONALE

The combinatorial invariance conjecture for parabolic Kazhdan-Lusztig polynomials of lower intervals

This is the peer reviewed version of the following article:

Original

The combinatorial invariance conjecture for parabolic Kazhdan-Lusztig polynomials of lower intervals / Marietti, Mario. - In: ADVANCES IN MATHEMATICS. - ISSN 0001-8708. - STAMPA. - 335:(2018), pp. 180-210. [10.1016/j.aim.2018.07.008]

Availability:

This version is available at: 11566/262525 since: 2022-05-21T18:23:41Z

Publisher:

Published

DOI:10.1016/j.aim.2018.07.008

Terms of use:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. The use of copyrighted works requires the consent of the rights' holder (author or publisher). Works made available under a Creative Commons license or a Publisher's custom-made license can be used according to the terms and conditions contained therein. See editor's website for further information and terms and conditions.

This item was downloaded from IRIS Università Politecnica delle Marche (<https://iris.univpm.it>). When citing, please refer to the published version.

(Article begins on next page)

THE COMBINATORIAL INVARIANCE CONJECTURE FOR PARABOLIC KAZHDAN–LUSZTIG POLYNOMIALS OF LOWER INTERVALS

MARIO MARIETTI

ABSTRACT. The aim of this work is to prove a conjecture related to the Combinatorial Invariance Conjecture of Kazhdan–Lusztig polynomials, in the parabolic setting, for lower intervals in every arbitrary Coxeter group. This result improves and generalizes, among other results, the main results of [Advances in Math. 202 (2006), 555-601], [Trans. Amer. Math. Soc. 368 (2016), no. 7, 5247–5269].

1. INTRODUCTION

Kazhdan–Lusztig polynomials play a central role in Lie theory and representation theory. They are polynomials $P_{u,v}(q)$, in one variable q , which are associated to pairs of elements u, v in a Coxeter group W . They were defined by Kazhdan and Lusztig in [19] in order to introduce the (now called) Kazhdan–Lusztig representations of the Hecke algebra of W , and soon have found applications in many other contexts.

Among others, the combinatorial aspects of Kazhdan–Lusztig polynomials have received much attention from the start, and are still a fascinating field of research. Recently, Elias and Williamson [15] proved the long-standing conjecture about the nonnegativity of the coefficients of Kazhdan–Lusztig polynomials of all Coxeter groups, thus generalizing the analogous result by Kazhdan and Lusztig on finite and affine Weyl groups appearing in [20], where $P_{u,v}(q)$ is shown to be the Poincaré polynomial of the local intersection cohomology groups of the Schubert variety associated with v at any point of the Schubert variety associated with u (in the full flag variety).

At present, from a combinatorial point of view, the most intriguing conjecture about Kazhdan–Lusztig polynomials is arguably what is usually referred to as the Combinatorial Invariance Conjecture of Kazhdan–Lusztig polynomials. It was independently formulated by Lusztig in private and by Dyer in [13].

Conjecture 1.1. *The Kazhdan–Lusztig polynomial $P_{u,v}(q)$ depends only on the isomorphism class of the interval $[u, v]$ as a poset.*

The Combinatorial Invariance Conjecture of Kazhdan–Lusztig polynomials is equivalent to the analogous conjecture on the combinatorial invariance of Kazhdan–Lusztig R -polynomials. These also are polynomials $R_{u,v}(q)$ indexed by a pair of elements u, v

2010 *Mathematics Subject Classification.* 20F55, 05E99.

Key words and phrases. Kazhdan–Lusztig polynomials, Coxeter groups, Special matchings.

This manuscript version is made available under the CC-BY-NC-ND license.

DOI: 10.1016/j.aim.2018.07.008

in W and were introduced by Kazhdan–Lusztig in the same article [19]. The Kazhdan–Lusztig R -polynomials are equivalent to the Kazhdan–Lusztig polynomials of W (in a precise sense, see Remark 2.12).

In [11], for any choice of a subset $H \subseteq S$, Deodhar introduces two modules of the Hecke algebra of W , two parabolic analogues $\{P_{u,w}^{H,x}(q)\}_{u,w \in W^H}$ of the Kazhdan–Lusztig polynomials, and two parabolic analogues $\{R_{u,w}^{H,x}(q)\}_{u,w \in W^H}$ of the Kazhdan–Lusztig R -polynomials, one for $x = q$ and one for $x = -1$. The parabolic Kazhdan–Lusztig and R -polynomials have deep algebraic and geometric significance; they are indexed by pairs of elements in the set W^H of minimal coset representatives with respect to the standard parabolic subgroup W_H , and play, in the parabolic setting, a role that is parallel to the role that the ordinary Kazhdan–Lusztig and R -polynomials play in the ordinary setting. Moreover, they generalize the ordinary Kazhdan–Lusztig and R -polynomials since these are obtained in the special trivial case when $H = \emptyset$ (for both $x = q$ and -1). As in the ordinary case, the family of parabolic Kazhdan–Lusztig polynomials is equivalent to the family of parabolic Kazhdan–Lusztig R -polynomials.

The problem of the combinatorial invariance of parabolic Kazhdan–Lusztig polynomials, which is stronger than the combinatorial invariance of the ordinary Kazhdan–Lusztig polynomials, has also attracted much attention (see, for instance, [2] and [4]). Only recently, however, the statement one gets by replacing the ordinary interval with the parabolic interval in Conjecture 1.1 has been found to be false (see [7] and [21] for counterexamples). In [21], it is proposed that the right approach to the generalization of Conjecture 1.1 to the parabolic setting could be studying to what extent the following conjecture is true.

Conjecture 1.2. *Let (W_1, S_1) and (W_2, S_2) be two Coxeter systems, $H_1 \subseteq S_1$ and $H_2 \subseteq S_2$. Let $u_1, v_1 \in W_1^{H_1}$ and $u_2, v_2 \in W_2^{H_2}$ be such that there exists a poset-isomorphism from $[u_1, v_1]$ to $[u_2, v_2]$ that restricts to a poset-isomorphism from $[u_1, v_1]^{H_1}$ to $[u_2, v_2]^{H_2}$. Then $P_{u_1, v_1}^{H_1, x}(q) = P_{u_2, v_2}^{H_2, x}(q)$ (equivalently, $R_{u_1, v_1}^{H_1, x}(q) = R_{u_2, v_2}^{H_2, x}(q)$).*

Clearly, Conjecture 1.2 reduces to Conjecture 1.1 for $H_1 = H_2 = \emptyset$.

Conjecture 1.1 and Conjecture 1.2, if true, would have interesting implications in the many contexts where ordinary and parabolic Kazhdan–Lusztig polynomials have applications. Among them, one of the most fascinating and (according to many experts in the field) surprising consequences would be in the topology of Schubert varieties of full and partial flag varieties. For the full flag variety, we refer the reader to the discussion in [3, §3]. For its generalization to the partial flag variety, the reader should have in mind the results by Kashiwara and Tanisaki [18] showing the role of the parabolic Kazhdan–Lusztig polynomials for the Schubert varieties of the partial flag variety.

In [21], Conjecture 1.2 is proved to hold true for lower intervals (that is, when u_1 and u_2 are the identity elements), in the case of doubly laced Coxeter groups (and in the case of dihedral Coxeter groups, which is much easier). The aim of this work is to prove the following more general result.

Theorem 1.3. *Conjecture 1.2 holds true for all lower intervals in every arbitrary Coxeter group.*

(Another new piece of evidence in favor of Conjecture 1.2 was recently given by Brenti in [5]).

Indeed, we prove the following slightly more general result.

Theorem 1.4. *Let (W_1, S_1) and (W_2, S_2) be two arbitrary Coxeter systems, with identity elements e_1 and e_2 , and let $H_1 \subseteq S_1$ and $H_2 \subseteq S_2$. Let $v_1 \in W_1^{H_1}$ and $v_2 \in W_2^{H_2}$ be such that there exists a poset-isomorphism ψ from $[e_1, v_1]$ to $[e_2, v_2]$ that restricts to a poset-isomorphism from $[e_1, v_1]^{H_1}$ to $[e_2, v_2]^{H_2}$. Then, for all $u, w \in [e_1, v_1]^{H_1}$, we have*

$$P_{u,w}^{H_1,x}(q) = P_{\psi(u),\psi(w)}^{H_2,x}(q) \quad \text{and} \quad R_{u,w}^{H_1,x}(q) = R_{\psi(u),\psi(w)}^{H_2,x}(q).$$

As a corollary, the parabolic Kazhdan–Lusztig polynomial $P_{u,w}^{H,x}(q)$ and R -polynomial $R_{u,w}^{H,x}(q)$ are determined by the isomorphism class of the interval $[e, w]$ and by how the parabolic interval $[e, w]^H = [e, w] \cap W^H$ embeds in $[e, w]$.

Theorem 1.4 is proved by providing an explicit method to compute the parabolic Kazhdan–Lusztig R -polynomials $R_{u,w}^{H,x}(q)$ (and so also the parabolic Kazhdan–Lusztig $P_{u,w}^{H,x}(q)$). This method is based on the concept of an H -special matching introduced in [21]: an H -special matching of w is an involution $M : [e, w] \rightarrow [e, w]$ such that

- (1) either $u \triangleleft M(u)$ or $u \triangleright M(u)$, for all $u \in [e, w]$,
- (2) if $u_1 \triangleleft u_2$ then $M(u_1) \leq M(u_2)$, for all $u_1, u_2 \in [e, w]$ such that $M(u_1) \neq u_2$,
- (3) if $u \leq w$, $u \in W^H$, and $M(u) \triangleleft u$, then $M(u) \in W^H$.

(We denote by \leq the Bruhat order and write $x \triangleleft y$ to mean that x is an immediate predecessor of y).

The set of all H -special matchings of w depends only on the isomorphism class of the interval $[e, w]$ and on how the parabolic interval $[e, w]^H$ embeds in $[e, w]$. We prove that H -special matchings may be used in place of left multiplications in the recurrence formula that computes the parabolic Kazhdan–Lusztig R -polynomials.

Theorem 1.5. *If M is an H -special matching of w , then the parabolic Kazhdan–Lusztig R -polynomial $R_{u,w}(q)$ satisfies:*

$$(1.1) \quad R_{u,w}^{H,x}(q) = \begin{cases} R_{M(u),M(w)}^{H,x}(q), & \text{if } M(u) \triangleleft u, \\ (q-1)R_{u,M(w)}^{H,x}(q) + qR_{M(u),M(w)}^{H,x}(q), & \text{if } M(u) \triangleright u \text{ and } M(u) \in W^H, \\ (q-1-x)R_{u,M(w)}^{H,x}(q), & \text{if } M(u) \triangleright u \text{ and } M(u) \notin W^H. \end{cases}$$

Theorem 1.5 directly implies Theorem 1.4. Indeed, suppose the hypotheses of Theorem 1.4 are fulfilled: thus M is an H_1 -special matching of w if and only if $M' = \psi \circ M \circ \psi^{-1}$ is an H_2 -special matching of $\psi(w)$. We choose such a matching M and apply (1.1) to both M and M' : in both computations, we fall in the same case. By iteration, we get the assertion of Theorem 1.4.

Theorems 1.3, 1.4 and 1.5 improve and generalize several results in the literature such as, for instance, the main results of [6], [12], [21], [23].

Since a special matching M is uniquely determined by its action on the dihedral intervals containing the Coxeter generator $M(e)$, special matchings of doubly laced Coxeter groups are more easily controlled than special matchings of arbitrary Coxeter

groups. Therefore, a deeper analysis on parabolic Kazhdan–Lusztig R -polynomials is needed to prove the result for arbitrary Coxeter groups. Indeed, we use several new identities among which, in particular, certain relations relating different parabolic Kazhdan–Lusztig R -polynomials indexed by elements in the same coset of dihedral standard parabolic subgroups.

The rest of the paper is devoted to the proof of Theorem 1.5.

2. NOTATION, DEFINITIONS AND PRELIMINARIES

This section reviews the background material that is needed in the rest of this work. We follow [1] and [22, Chapter 3] for undefined notation and terminology concerning, respectively, Coxeter groups and partially ordered sets.

2.1. Coxeter groups. We fix our notation on a Coxeter system (W, S) in the following list:

$m_{s,t}$	the entry of the Coxeter matrix of (W, S) in position $(s, t) \in S \times S$,
e	identity of W ,
ℓ	the length function of (W, S) ,
T	$= \{wsw^{-1} : w \in W, s \in S\}$, the set of <i>reflections</i> of W ,
$D_R(w)$	$= \{s \in S : \ell(ws) < \ell(w)\}$, the right descent set of $w \in W$,
$D_L(w)$	$= \{s \in S : \ell(sw) < \ell(w)\}$, the left descent set of $w \in W$,
W_J	the parabolic subgroup of W generated by $J \subseteq S$,
W^J	$= \{w \in W : D_R(w) \subseteq S \setminus J\}$, the set of minimal left coset representatives,
JW	$= \{w \in W : D_L(w) \subseteq S \setminus J\}$, the set of minimal right coset representatives,
\leq	Bruhat order on W (as well as usual order on \mathbb{R}),
$[u, v]$	$= \{w \in W : u \leq w \leq v\}$, the (Bruhat) interval generated by $u, v \in W$,
$w_0(J)$	the unique maximal element of $[e, w] \cap W_J$, for $J \subseteq S$,
$w_0(s, t)$	$= w_0(\{s, t\})$, for $s, t \in S$,
$[u, v]^H$	$= \{z \in W^H : u \leq z \leq v\}$, the parabolic interval generated by $u, v \in W^H$.

Given $u, v \in W$, we write $u \cdot v$ instead of simply uv when $\ell(uv) = \ell(u) + \ell(v)$ and we want to stress this additivity. On the other hand, when we write uv , $\ell(uv)$ can be either $\ell(u) + \ell(v)$ or smaller. We make use of the symbol “-” to separate letters in a word in the alphabet S when we want to stress the fact that we are considering the word rather than the element such word represents.

If $w \in W$, then a *reduced expression* for w is a word $s_1 s_2 \cdots s_q$ such that $w = s_1 s_2 \cdots s_q$ and $\ell(w) = q$. When no confusion arises, we also write that $s_1 s_2 \cdots s_q$ is a reduced expression for w .

The *Bruhat graph* of W (see [14], or, e.g., [1, §2.1] or [17, §8.6]) is the directed graph having W as vertex set and having a directed edge from u to v if and only if $u^{-1}v \in T$ and $\ell(u) < \ell(v)$. The *Bruhat order* (see, e.g., [1, §2.1] or [17, §5.9]), sometimes also

called *Bruhat-Chevalley order*, is the partial order \leq on W given by the transitive closure of the Bruhat graph of W

The following well-known characterization of Bruhat order is usually referred to as the *Subword Property* (see [1, §2.2] or [17, §5.10]), and is used repeatedly in the following sections, often without explicit mention. By a *subword* of a word $s_1s_2\cdots s_q$, we mean a word of the form $s_{i_1}s_{i_2}\cdots s_{i_k}$, where $1 \leq i_1 < \cdots < i_k \leq q$.

Theorem 2.1 (Subword Property). *Let $u, w \in W$. The following are equivalent:*

- $u \leq w$ in the Bruhat order,
- every reduced expression for w has a subword that is a reduced expression for u ,
- there exists a reduced expression for w having a subword that is a reduced expression for u .

The following results are well known (see, e.g., [10, Theorem 1.1], [1, Proposition 2.2.7] or [17, Proposition 5.9] for the first one, [1, §2.4] or [17, §1.10] for the second one, and [16, Lemma 7] for the third one).

Lemma 2.2 (Lifting Property). *Let $s \in S$ and $u, w \in W$, $u \leq w$.*

- If $s \in D_R(w)$ and $s \in D_R(u)$, then $us \leq ws$.
- If $s \notin D_R(w)$ and $s \notin D_R(u)$, then $us \leq ws$.
- If $s \in D_R(w)$ and $s \notin D_R(u)$, then $us \leq w$ and $u \leq ws$.

Symmetrically, left versions of the three statements hold.

Proposition 2.3. *Let $J \subseteq S$.*

- (i) *Every $w \in W$ has a unique factorization $w = w^J \cdot w_J$ with $w^J \in W^J$ and $w_J \in W_J$; for this factorization, $\ell(w) = \ell(w^J) + \ell(w_J)$.*
- (ii) *Every $w \in W$ has a unique factorization $w = {}_Jw \cdot {}^Jw$ with ${}_Jw \in W_J$, ${}^Jw \in {}^JW$; for this factorization, $\ell(w) = \ell({}_Jw) + \ell({}^Jw)$.*

Proposition 2.4. *Let $J \subseteq S$ and $w \in W$. The set $W_J \cap [e, w]$ has a unique maximal element $w_0(J)$, so that $W_J \cap [e, w]$ is the interval $[e, w_0(J)]$.*

Note that, by the uniqueness of the factorizations of Proposition 2.3, if $J \subseteq S$ and $w \in W$, then

$$(2.1) \quad l \in D_L({}_Jw) \iff l \in D_L(w) \cap J.$$

Furthermore, it is well known (and immediate to prove) that $v \leq w$ implies both $v^J \leq w^J$ and ${}^Jv \leq {}^Jw$.

2.2. Special matchings. Let P be a partially ordered set. An element $y \in P$ *covers* $x \in P$ if the interval $[x, y]$ coincides with $\{x, y\}$; in this case, we write $x \triangleleft y$ as well as $y \triangleright x$. The poset P is *graded* if P has a minimum and there is a function $\rho : P \rightarrow \mathbb{N}$ (the *rank function* of P) such that $\rho(\hat{0}) = 0$ and $\rho(y) = \rho(x) + 1$ for all $x, y \in P$ with $x \triangleleft y$. (This definition is slightly different from the one given in [22], but is more convenient for our purposes.) The *Hasse diagram* of P is the graph having P as vertex set and $\{\{x, y\} \in \binom{P}{2} : \text{either } x \triangleleft y \text{ or } y \triangleleft x\}$ as edge set.

A *matching* of a poset P is an involution $M : P \rightarrow P$ such that $\{v, M(v)\}$ is an edge in the Hasse diagram of P , for all $v \in V$. A matching M of P is *special* if

$$u \triangleleft v \implies M(u) \leq M(v),$$

for all $u, v \in P$ such that $M(u) \neq v$.

Now, let (W, S) be a Coxeter system and recall that W is a graded partially ordered set (under Bruhat order) having ℓ as its rank function. Given $w \in W$, we say that M is a matching of w if M is a matching of the lower Bruhat interval $[e, w]$. For $s \in D_R(w)$, we have a matching ρ_s of w defined by $\rho_s(u) = us$, for all $u \in [e, w]$. Symmetrically, for $s \in D_L(w)$, we have a matching λ_s of w defined by $\lambda_s(u) = su$, for all $u \in [e, w]$. By the Lifting Property (Lemma 2.2), such ρ_s and λ_s are special matchings of w . We call these matchings, respectively, *right* and *left multiplication matchings*.

The following two results are used several times in what follows: the first directly follows from [6, Lemma 4.3], the second is [6, Proposition 5.3]. We call an interval $[u, v]$ in a poset P *dihedral* if it is isomorphic to an interval in a Coxeter system of rank 2 ordered by Bruhat order. Moreover, given two matchings M and N , we say that M and N commute on X if the two compositions $M \circ N(x)$ and $N \circ M(x)$ are defined and equal, for all $x \in X$. We say that two matchings of w commute if they commute everywhere on $[e, w]$.

Lemma 2.5. *Let $w \in W$. Two special matchings M and N of w commute if and only if they commute on the lower dihedral intervals of $[e, w]$ containing $M(e)$ and $N(e)$.*

Lemma 2.6. *Let $J \subseteq S$, $w \in W$, and M be a special matching of w . If $M(e) \in J$, then M stabilizes $[e, w_0(J)]$.*

In particular, given two special matchings M and N of w such that $M(e) \neq N(e)$, we have that M and N commute if and only if they commute on the unique lower dihedral interval $[e, w_0(M(e), N(e))]$, and this lower dihedral interval is stabilized by both M and N .

The following definitions are taken from [21].

Definition 2.7. *A right system for $w \in W$ is a quadruple $\mathcal{R} = (J, s, t, M_{st})$ such that:*

- R1. $J \subseteq S$, $s \in J$, $t \in S \setminus J$, and M_{st} is a special matching of $w_0(s, t)$ such that $M_{st}(e) = s$ and $M_{st}(t) = ts$;
- R2. $(u^J)^{\{s, t\}} \cdot M_{st} \left((u^J)^{\{s, t\}} \cdot {}^{\{s\}}(u_J) \right) \cdot {}^{\{s\}}(u_J) \leq w$, for all $u \leq w$;
- R3. if $r \in J$ and $r \leq w^J$, then r and s commute;
- R4. (a) if $s \leq (w^J)^{\{s, t\}}$ and $t \leq (w^J)^{\{s, t\}}$, then $M_{st} = \rho_s$,
 (b) if $s \leq (w^J)^{\{s, t\}}$ and $t \not\leq (w^J)^{\{s, t\}}$, then M_{st} commutes with λ_s ,
 (c) if $s \not\leq (w^J)^{\{s, t\}}$ and $t \leq (w^J)^{\{s, t\}}$, then M_{st} commutes with λ_t ;
- R5. if $s \leq {}^{\{s\}}(w_J)$, then M_{st} commutes with ρ_s on $[e, w_0(s, t)]$.

Definition 2.8. *A left system for $w \in W$ is a quadruple $\mathcal{L} = (J, s, t, M_{st})$ such that:*

- L1. $J \subseteq S$, $s \in J$, $t \in S \setminus J$, and M_{st} is a special matching of $w_0(s, t)$ such that $M_{st}(e) = s$ and $M_{st}(t) = st$;

- L2. $(Ju)^{\{s\}} \cdot M_{st} \left((Ju)_{\{s\}} \cdot {}_{\{s,t\}}(Ju) \right) \cdot {}^{\{s,t\}}(Ju) \leq w$, for all $u \leq w$;
 L3. if $r \in J$ and $r \leq {}^Jw$, then r and s commute;
 L4. (a) if $s \leq {}^{\{s,t\}}(Jw)$ and $t \leq {}^{\{s,t\}}(Jw)$, then $M_{st} = \lambda_s$,
 (b) if $s \leq {}^{\{s,t\}}(Jw)$ and $t \not\leq {}^{\{s,t\}}(Jw)$, then M_{st} commutes with ρ_s ,
 (c) if $s \not\leq {}^{\{s,t\}}(Jw)$ and $t \leq {}^{\{s,t\}}(Jw)$, then M_{st} commutes with ρ_t ;
 L5. if $s \leq (Jw)^{\{s\}}$, then M_{st} commutes with λ_s on $[e, w_0(s, t)]$.

(As shown in [8, Lemma 4.3], Properties R5 and L5 are equivalent to the, a priori, more restrictive Properties R5 and L5 appearing in [21].)

Given a right system $\mathcal{R} = (J, s, t, M_{st})$ for w , the *matching associated* with it is the map $M_{\mathcal{R}}$ sending $u \in [e, w]$ to

$$M_{\mathcal{R}}(u) = (u^J)^{\{s,t\}} \cdot M_{st} \left((u^J)_{\{s,t\}} \cdot {}_{\{s\}}(u_J) \right) \cdot {}^{\{s\}}(u_J).$$

Symmetrically, the *matching associated* with a left system \mathcal{L} for w is the map ${}_{\mathcal{L}}M$ sending $u \in [e, w]$ to

$${}_{\mathcal{L}}M(u) = (Ju)^{\{s\}} \cdot M_{st} \left((Ju)_{\{s\}} \cdot {}_{\{s,t\}}(Ju) \right) \cdot {}^{\{s,t\}}(Ju),$$

i.e., ${}_{\mathcal{L}}M(u) = (M_{\mathcal{L}}(u^{-1}))^{-1}$, where $M_{\mathcal{L}}$ is the map on $[e, w^{-1}]$ associated to \mathcal{L} as a right system for w^{-1} .

The fact that $M_{\mathcal{R}}$ and ${}_{\mathcal{L}}M$ are actually matchings of w and the fact that the lengths add in these products are shown in [8] (respectively, in Corollary 4.10 and Proposition 4.9).

Note that $M_{\mathcal{R}}$ acts as λ_s on $[e, w_0(s, r)]$ for all $r \in J$, and as ρ_s on $[e, w_0(s, r)]$ for all $r \in S \setminus (J \cup \{t\})$; symmetrically, ${}_{\mathcal{L}}M$ acts as ρ_s on $[e, w_0(s, r)]$ for all $r \in J$, and as λ_s on $[e, w_0(s, r)]$ for all $r \in S \setminus (J \cup \{t\})$.

We comment that, if $s \in D_R(w)$, $t \in S \setminus \{s\}$, $J = \{s\}$ and $M_{st} = \rho_s$, then we obtain a right system whose associated matching is the right multiplication matching ρ_s ($M = \rho_s$ on the entire interval $[e, w]$). Symmetrically, we obtain left multiplication matchings as special cases of matchings associated with left systems. On the other hand, we may obtain matchings that are not multiplication matchings. For example, let W be the Coxeter group of type A_3 with Coxeter generators s_1, s_2 and s_3 numbered as usual (i.e. $m_{s_1, s_2} = m_{s_2, s_3} = 3$ and $m_{s_1, s_3} = 2$), and let $w = s_1 s_2 s_3 s_1 \in W$. The quadruple $\mathcal{R} = (\{s_2, s_3\}, s_2, s_1, M)$, with $M(e) = s_2$, $M(s_1) = s_1 s_2$, and $M(s_2 s_1) = s_1 s_2 s_1$, is a right system for w whose associated matching is not a multiplication matching (the reader may check that the resulting matching is the dashed special matchings in the first picture of Figure 2).

The main result of [8] is that the matchings arising from systems of w are exactly the special matchings of w . We only need one side of this characterization (see [8, Theorem 4.12]).

Theorem 2.9. *Every special matching of $w \in W$ is associated with a right or a left system of w .*

We refer the reader to [9] for a more compact characterization in terms of only one self-dual type of systems.

2.3. Kazhdan–Lusztig polynomials. Given a Coxeter system (W, S) and $H \subseteq S$, the Bruhat order induces an ordering on the set of minimal coset representatives W^H and the parabolic intervals $[u, v]^H$, for all $u, v \in W^H$.

We introduce the parabolic Kazhdan–Lusztig R -polynomials and the parabolic Kazhdan–Lusztig polynomials through the following theorems-definitions, which are due to Deodhar (see [11, §§2-3] for their proofs).

Theorem 2.10. *Let (W, S) be a Coxeter system, and $H \subseteq S$. For each $x \in \{-1, q\}$, there is a unique family of polynomials $\{R_{u,v}^{H,x}(q)\}_{u,v \in W^H} \subseteq \mathbf{Z}[q]$ such that, for all $u, v \in W^H$:*

- (1) $R_{u,v}^{H,x}(q) = 0$ if $u \not\leq v$;
- (2) $R_{u,u}^{H,x}(q) = 1$;
- (3) if $u < v$ and $s \in D_L(v)$, then

$$R_{u,v}^{H,x}(q) = \begin{cases} R_{su,sv}^{H,x}(q), & \text{if } s \in D_L(u), \\ (q-1)R_{u,sv}^{H,x}(q) + qR_{su,sv}^{H,x}(q), & \text{if } s \notin D_L(u) \text{ and } su \in W^H, \\ (q-1-x)R_{u,sv}^{H,x}(q), & \text{if } s \notin D_L(u) \text{ and } su \notin W^H. \end{cases}$$

In what follows, we often use the inductive formula of Theorem 2.10 without explicit mention.

Theorem 2.11. *Let (W, S) be a Coxeter system, and $H \subseteq S$. For each $x \in \{-1, q\}$, there is a unique family of polynomials $\{P_{u,v}^{H,x}(q)\}_{u,v \in W^H} \subseteq \mathbf{Z}[q]$, such that, for all $u, v \in W^H$:*

- (1) $P_{u,v}^{H,x}(q) = 0$ if $u \not\leq v$;
- (2) $P_{u,u}^{H,x}(q) = 1$;
- (3) $\deg(P_{u,v}^{H,x}(q)) \leq \frac{1}{2}(\ell(v) - \ell(u) - 1)$, if $u < v$;
- (4) $q^{\ell(v)-\ell(u)} P_{u,v}^{H,x}\left(\frac{1}{q}\right) = \sum_{z \in [u,v]_H} R_{u,z}^{H,x}(q) P_{z,v}^{H,x}(q)$.

The polynomials $R_{u,v}^{H,x}(q)$ and $P_{u,v}^{H,x}(q)$ are the *parabolic Kazhdan–Lusztig R -polynomials* and *parabolic Kazhdan–Lusztig polynomials* of W^H of type x .

Remark 2.12. *For a fixed $H \subset S$, the parabolic Kazhdan–Lusztig R -polynomials and the parabolic Kazhdan–Lusztig polynomials are equivalent. More precisely, given $w \in W^H$, it is possible to compute the family $\{P_{u,v}^{H,x}(q)\}_{u,v \in [e,w]^H}$ once one knows the family $\{R_{u,v}^{H,x}(q)\}_{u,v \in [e,w]^H}$, and vice versa.*

For $H = \emptyset$, $R_{u,v}^{\emptyset,-1}(q) = R_{u,v}^{\emptyset,q}(q)$ and $P_{u,v}^{\emptyset,-1}(q) = P_{u,v}^{\emptyset,q}(q)$ are the ordinary Kazhdan–Lusztig R -polynomials $R_{u,v}(q)$ and Kazhdan–Lusztig polynomials $P_{u,v}(q)$ of W .

The following result gives another relationship between the parabolic Kazhdan–Lusztig polynomials and their ordinary counterparts (see [11, Proposition 3.4, and Remark 3.8]).

Proposition 2.13. *Let (W, S) be a Coxeter system, $H \subseteq S$, and $u, v \in W^H$. We have*

$$P_{u,v}^{H,q}(q) = \sum_{w \in W_H} (-1)^{\ell(w)} P_{uw,v}(q).$$

Furthermore, if W_H is finite, then

$$P_{u,v}^{H,-1}(q) = P_{uw_0^H, vw_0^H}(q),$$

where w_0^H is the longest element of W_H .

3. PRELIMINARY RESULTS

In this section, we give some preliminary results that are needed to prove the main result of this work.

For convenience, we state the following straightforward result here for later reference.

Lemma 3.1. *The sequence $\{R_i\}_{i \geq 1} \subseteq \mathbb{Z}[q]$, defined as*

$$R_i = \begin{cases} (q-1) \left(\sum_{k=0}^{i-1} (-1)^k q^k \right), & \text{if } i \text{ is odd,} \\ (q-1)^2 \left(\sum_{k=0}^{\frac{i-2}{2}} q^{2k} \right), & \text{if } i \text{ is even,} \end{cases}$$

is the unique sequence satisfying

$$R_i = (q-1)R_{i-1} + qR_{i-2} \quad R_1 = (q-1) \quad R_2 = (q-1)^2$$

Proof. Omitted. □

We observe the following fact. Let W be a dihedral Coxeter groups, and $u, w \in W$. If $\ell(w) - \ell(u) = i$, with $i \geq 1$, then the ordinary Kazhdan–Lusztig R -polynomial $R_{u,v}(q)$ is the polynomial R_i defined in Lemma 3.1. In particular, as it is well-known, Conjecture 1.1 holds true for dihedral Coxeter groups since two intervals $[u, w]$ and $[u', w']$ in two dihedral Coxeter groups are isomorphic as posets if and only if $\ell(w) - \ell(u) = \ell(w') - \ell(u')$.

We fix an arbitrary Coxeter system (W, S) , a subset $H \subset S$, and $s, t \in S$. For notational convenience, from now on we let $\bar{s} = t$ and $\bar{t} = s$. Recall that, for every $x \in W$, the coset $W_{\{s,t\}} x = \{g_{st} x : g_{st} \in W_{\{s,t\}}\}$ is isomorphic, as a poset, to the dihedral Coxeter group $W_{\{s,t\}}$.

Proposition 3.2. *Consider an arbitrary coset $W_{\{s,t\}} \cdot x$, where (we suppose without lack of generality) $x \in {}^{\{s,t\}}W$. The intersection $(W_{\{s,t\}} \cdot x) \cap W^H$ is one of the following set:*

- (1) \emptyset ,
- (2) $\{x\}$,
- (3) $\{g_{st} \cdot x : g_{st} \in W_{\{s,t\}}, t \notin D_R(g_{st})\}$,
- (4) $\{g_{st} \cdot x : g_{st} \in W_{\{s,t\}}, s \notin D_R(g_{st})\}$,
- (5) $W_{\{s,t\}} \cdot x$.

Proof. First of all, recall that if an element w belongs to W^H , then rw belongs to W^H for all $r \in D_L(w)$. Also recall that an element not in W^H have at most one coatom in W^H (see [21, Lemma 4.1]); in particular, in the case $W_{\{s,t\}}$ is finite, the intersection $(W_{\{s,t\}} \cdot x) \cap W^H$ cannot be $(W_{\{s,t\}} \cdot x) \setminus \{w_0 \cdot x\}$, where w_0 is the longest element of $W_{\{s,t\}}$.

We prove the statement by contradiction and, by what we have just recalled, we suppose that there exist $g \in W_{\{s,t\}} \setminus \{e\}$ and $p \in \{s,t\} \setminus D_L(g)$ such that

- $g \cdot x \in W^H$,
- $p \cdot g \cdot x \notin W^H$,
- $p \cdot g$ is not the longest element of $W_{\{s,t\}}$ (if any, i.e. in the case $W_{\{s,t\}}$ is finite).

Since $p \cdot g \cdot x \notin W^H$, there exists $h \in H \cap D_R(p \cdot g \cdot x)$. Since $g \cdot x \in W^H$, we have $h \notin D_R(g \cdot x)$; by the Lifting Property (Lemma 2.2), $p \cdot g \cdot x = g \cdot x \cdot h$. Hence, both s and t belong to $D_L(p \cdot g \cdot x)$ and thus to $D_L(p \cdot g)$; by a well-known fact, this means that $p \cdot g$ is the longest element of $W_{\{s,t\}}$. \square

The following three results give formulas expressing some parabolic Kazhdan–Lusztig R -polynomials as linear combinations of other parabolic Kazhdan–Lusztig R -polynomials. (The choice of the indices could seem unnatural at this point: the reason for this choice is that, in Section 4, we apply these results in a situation where we have two missing parts w_1 and u_1 , i.e. two parts w_1 and u_1 that are both equal to e).

Lemma 3.3. *Let $w = w_2 \cdot w_3 \in W^H$ and $u = u_2 \cdot u_3 \in W^H$ with:*

- $u \leq w$,
- $w_2, u_2 \in W_{\{s,t\}}$,
- $w_3, u_3 \in {}^{\{s,t\}}W$.

If $(W_{\{s,t\}} \cdot u_3) \cap W^H = \{u_3\}$, then $u = u_3$ and

$$R_{u,w}^{H,x}(q) = (q - 1 - x)^{\ell(w_2)} R_{u_3,w_3}^{H,x}(q).$$

Proof. If $w_2 = e$, the assertions are trivial. Suppose $w_2 \neq e$ and fix $r \in \{s,t\} \cap D_L(w_2)$. By the recursive formula of Theorem 2.10 (with r as left descent of w), we have $R_{u,w}^{H,x}(q) = (q - 1 - x) R_{u,rw}^{H,x}(q)$. We get the assertion by iteration. \square

Recall that, for $r \in \{s,t\}$, we denote by \bar{r} the element in $\{s,t\} \setminus \{r\}$.

Lemma 3.4. *Let $w = w_2 \cdot w_3 \in W^H$ and $u = u_2 \cdot u_3 \in W^H$ with:*

- $u \leq w$,
- $w_2, u_2 \in W_{\{s,t\}}$,
- $w_3, u_3 \in {}^{\{s,t\}}W$,

and suppose $W_{\{s,t\}} \cdot u_3 \subseteq W^H$. Then there exists a set of polynomials $\{p_{gst}(q)\}_{gst \in W_{\{s,t\}}} \subseteq \mathbb{Z}[q]$ such that

$$\begin{aligned} R_{u,w}^{H,x}(q) &= \sum_{gst \in W_{\{s,t\}}} p_{gst}(q) R_{gst \cdot u_3, w_3}^{H,x}(q) \\ R_{u,w}(q) &= \sum_{gst \in W_{\{s,t\}}} p_{gst}(q) R_{gst \cdot u_3, w_3}(q) \end{aligned}$$

(in other words, both the parabolic and the ordinary Kazhdan–Lusztig R -polynomials indexed by u and w can be expressed as a linear combination of, respectively, the parabolic and the ordinary Kazhdan–Lusztig R -polynomials indexed by $gst \cdot u_3$ and w_3 , with $gst \in W_{\{s,t\}}$, and the two expressions have the same coefficients).

If, moreover, $|\{x \in \{s, t\} : x \leq w_3\}| \leq 1$, then $\ell(w_2) - \ell(u_2) \geq -1$ and the following statements hold.

D₋₁. If $\ell(w_2) - \ell(u_2) = -1$, then

$$R_{u,w}^{H,x}(q) = R_{pu_3,w_3}^{H,x}(q).$$

D₀. If $\ell(w_2) - \ell(u_2) = 0$, then

$$R_{u,w}^{H,x}(q) = \begin{cases} R_{u_3,w_3}^{H,x}(q) & \text{if } u_2 = w_2 \\ (q-1)R_{pu_3,w_3}^{H,x}(q) & \text{if } u_2 \neq w_2. \end{cases}$$

D₁. If $\ell(w_2) - \ell(u_2) = 1$, then

$$R_{u,w}^{H,x}(q) = \begin{cases} (q-1)R_{u_3,w_3}^{H,x}(q) + qR_{pu_3,w_3}^{H,x}(q), & \text{if } w_2 = u_2 \cdot p \\ (q-1)R_{u_3,w_3}^{H,x}(q), & \text{otherwise.} \end{cases}$$

D_i. If $\ell(w_2) - \ell(u_2) = i \geq 2$, then

$$R_{u,w}^{H,x}(q) = R_i \cdot R_{u_3,w_3}^{H,x}(q) + qR_{i-1} \cdot R_{pu_3,w_3}^{H,x}(q)$$

where the family of polynomials $\{R_j\}_{j \geq 1}$ is as in Lemma 3.1.

In the previous statements, if $|\{x \in \{s, t\} : x \leq w_3\}| \leq 1$ then $\{p\} = \{x \in \{s, t\} : x \leq w_3\}$, if $|\{x \in \{s, t\} : x \leq w_3\}| = 0$ then $R_{pu_3,w_3}^{H,x}(q) = 0$.

Proof. Let us prove the first statement. If $w_2 = e$, it is trivial. Suppose $w_2 \neq e$ and fix $r \in \{s, t\} \cap D_L(w_2)$. We apply the recursive formula of Theorem 2.10 (with r as a left descent of w) to compute both $R_{u,w}^{H,x}(q)$ and $R_{u,w}(q)$. Since $W_{\{s,t\}} \cdot u_3 \subseteq W^H$, we cannot fall into the case when the factor $(q-1-x)$ occurs, and the two computations agree. We get the assertion by iterating this argument.

Let us prove the second part of the lemma and so suppose $\{p\} \supseteq \{x \in \{s, t\} : x \leq w_3\}$.

In this proof, we use the Subword Property (Theorem 2.1), Property (2.1), and the recursive formula of Theorem 2.10 several times without explicit mention; when we apply the recursive formula of Theorem 2.10, we never fall into the case the factor $(q-1-x)$ occurs, since $W_{\{s,t\}} \cdot u_3 \subseteq W^H$.

Since $|\{x \in \{s, t\} : x \leq w_3\}| \leq 1$, the longest subword of type $s-t-s-t-\dots$ or $t-s-t-s-\dots$ of any reduced expression for w has length at most $\ell(w_2) + 1$, and hence $\ell(w_2) - \ell(u_2) \geq -1$.

Proof of D₋₁. Since $\ell(w_2) - \ell(u_2) = -1$, necessarily $u_2 = w_2 \cdot p$, as otherwise u could not be smaller than or equal to w . We have

$$R_{u,w}^{H,x}(q) = R_{w_2pu_3,w_2w_3}^{H,x}(q) = R_{pu_3,w_3}^{H,x}(q).$$

Proof of D₀. If $u_2 = w_2$, the assertion is immediate. If $u_2 \neq w_2$, there exists an element $v \in W_{\{s,t\}}$, with $\ell(v) = \ell(w_2) - 1 = \ell(u_2) - 1$, such that $u_2 = v \cdot p$ and $w_2 = l \cdot v$, where $l \in \{s, t\} \setminus D_L(v)$. We have

$$R_{u,w}^{H,x}(q) = R_{vpu_3,lvw_3}^{H,x}(q) = (q-1)R_{vpu_3,vw_3}^{H,x}(q) + qR_{lvpu_3,vw_3}^{H,x}(q) = (q-1)R_{pu_3,w_3}^{H,x}(q) + qR_{lvpu_3,vw_3}^{H,x}(q).$$

We cannot have $l \cdot v \cdot p \cdot u_3 \leq v \cdot w_3$, since all subwords of any reduced expression of $v \cdot w_3$ of type $s-t-s-t-\dots$ or $t-s-t-s-\dots$ have length at most $\ell(v) + 1$, while $\ell(l \cdot v \cdot p) = \ell(v) + 2$. Hence $R_{lvpu_3, vw_3}^{H,x}(q) = 0$, as desired.

Proof of D_1 . Since $\ell(w_2) - \ell(u_2) = 1$, we have either

- (1) $w_2 = u_2 \cdot r$, where $r \in \{s, t\} \setminus D_R(u_2)$, or
- (2) $w_2 = l \cdot u_2$, where $l \in \{s, t\} \setminus D_L(u_2)$, $u_2 \neq e$, and $l \cdot u_2$ is not the longest element of $W_{\{s,t\}}$ (if any).

In the first case, we have

$$R_{u,w}^{H,x}(q) = R_{u_2u_3, u_2rw_3}^{H,x}(q) = R_{u_3, rw_3}^{H,x}(q) = (q-1)R_{u_3, w_3}^{H,x}(q) + qR_{ru_3, w_3}^{H,x}(q).$$

If $r = p$, we get the assertion. If $r \neq p$, then $r \not\leq w_3$; thus $ru_3 \not\leq w_3$, and we get the assertion as well.

In the second case, we have

$$R_{u,w}^{H,x}(q) = R_{u_2u_3, lu_2w_3}^{H,x}(q) = (q-1)R_{u_2u_3, u_2w_3}^{H,x}(q) + qR_{lu_2u_3, u_2w_3}^{H,x}(q) = (q-1)R_{u_3, w_3}^{H,x}(q)$$

since $l \cdot u_2 \cdot u_3 \not\leq u_2 \cdot w_3$.

Proof of D_i . Suppose $\ell(w_2) - \ell(u_2) = 2$; we have either

- $w_2 = u_2 \cdot r \cdot \bar{r}$, where $r \in \{s, t\} \setminus D_R(u_2)$, or
- $w_2 = l \cdot u_2 \cdot r$, where $l \in \{s, t\} \setminus D_L(u_2)$, $r \in \{s, t\} \setminus D_R(u_2)$, $u_2 \neq e$, and $l \cdot u_2 \cdot r$ is not the longest element of $W_{\{s,t\}}$ (if any).

In the first case, we have

$$\begin{aligned} R_{u,w}^{H,x}(q) &= R_{u_2u_3, u_2r\bar{r}w_3}^{H,x}(q) = R_{u_3, r\bar{r}w_3}^{H,x}(q) = (q-1)R_{u_3, \bar{r}w_3}^{H,x}(q) + qR_{ru_3, \bar{r}w_3}^{H,x}(q) \\ &= (q-1)[(q-1)R_{u_3, w_3}^{H,x}(q) + qR_{\bar{r}u_3, w_3}^{H,x}(q)] + q[(q-1)R_{ru_3, w_3}^{H,x}(q) + qR_{\bar{r}ru_3, w_3}^{H,x}(q)] \\ &= (q-1)^2R_{u_3, w_3}^{H,x}(q) + q(q-1)[R_{\bar{r}u_3, w_3}^{H,x}(q) + R_{ru_3, w_3}^{H,x}(q)]. \end{aligned}$$

since $\bar{r}ru_3 \not\leq w_3$. Thus the assertion follows since $\{\bar{r}u_3, ru_3\} \cap \{x : x \leq w_3\} = \{pu_3\} \cap \{x : x \leq w_3\}$.

In the second case, we have

$$\begin{aligned} R_{u,w}^{H,x}(q) &= R_{u_2u_3, lu_2rw_3}^{H,x}(q) = (q-1)R_{u_2u_3, u_2rw_3}^{H,x}(q) + qR_{lu_2u_3, u_2rw_3}^{H,x}(q) \\ &= (q-1)R_{u_3, rw_3}^{H,x}(q) + qR_{lu_2u_3, u_2rw_3}^{H,x}(q) \\ &= (q-1)[(q-1)R_{u_3, w_3}^{H,x}(q) + qR_{ru_3, w_3}^{H,x}(q)] + qR_{lu_2u_3, u_2rw_3}^{H,x}(q). \end{aligned}$$

If $r = p$, then $lu_2u_3 \not\leq u_2rw_3$: thus $R_{lu_2u_3, u_2rw_3}^{H,x}(q) = 0$ and we get the assertion. If $r \neq p$, then $ru_3 \not\leq w_3$ and thus $R_{ru_3, w_3}^{H,x}(q) = 0$; on the other hand, $lu_2u_3 \leq u_2rw_3$ and $R_{lu_2u_3, u_2rw_3}^{H,x}(q) = (q-1)R_{pu_3, w_3}^{H,x}(q)$ by Statement D_0 .

Suppose $\ell(w_2) - \ell(u_2) = 3$; we have either

- $w_2 = u_2 \cdot r \cdot \bar{r} \cdot r$, where $r \in \{s, t\} \setminus D_R(u_2)$, or
- $w_2 = l \cdot u_2 \cdot r \cdot \bar{r}$, where $l \in \{s, t\} \setminus D_L(u_2)$, $r \in \{s, t\} \setminus D_R(u_2)$, $u_2 \neq e$, and $l \cdot u_2 \cdot r \cdot \bar{r}$ is not the longest element of $W_{\{s,t\}}$ (if any).

In the first case, we have

$$\begin{aligned} R_{u,w}^{H,x}(q) &= R_{u_2u_3,u_2r\bar{r}rw_3}^{H,x}(q) = R_{u_3,r\bar{r}rw_3}^{H,x}(q) = (q-1)R_{u_3,\bar{r}rw_3}^{H,x}(q) + qR_{ru_3,\bar{r}rw_3}^{H,x}(q) \\ &= (q-1)[(q-1)^2R_{u_3,w_3}^{H,x}(q) + q(q-1)R_{pu_3,w_3}^{H,x}(q)] + q(q-1)R_{u_3,w_3}^{H,x}(q) \\ &= R_3 \cdot R_{u_3,w_3}^{H,x}(q) + qR_2 \cdot R_{pu_3,w_3}^{H,x}(q), \end{aligned}$$

by the assertion (already proved) for when the difference of the length is equal to 2, and by Statement D_1 .

In the second case, we have

$$\begin{aligned} R_{u,w}^{H,x}(q) &= R_{u_2u_3,lu_2r\bar{r}w_3}^{H,x}(q) = (q-1)R_{u_2u_3,u_2r\bar{r}w_3}^{H,x}(q) + qR_{lu_2u_3,u_2r\bar{r}w_3}^{H,x}(q) \\ &= (q-1)[(q-1)^2R_{u_3,w_3}^{H,x}(q) + q(q-1)R_{pu_3,w_3}^{H,x}(q)] + q(q-1)R_{u_3,w_3}^{H,x}(q) \\ &= R_3 \cdot R_{u_3,w_3}^{H,x}(q) + qR_2 \cdot R_{pu_3,w_3}^{H,x}(q) \end{aligned}$$

by the assertion (already proved) for when the difference of the length is equal to 2, and by Statement D_1 .

Suppose $\ell(w_2) - \ell(u_2) = i$, with $i \geq 4$, and use induction on $\ell(w_2)$. The base of the induction is $u_2 = e$ and $w_2 \in \{p\bar{p}p\bar{p}, \bar{p}p\bar{p}p\}$: the assertion follows by a direct computation that we omit.

Let $\ell(w_2) > 4$ and $h \in D_L(w_2)$. If $h \in D_L(u_2)$, then $R_{u,w}^{H,x}(q) = R_{u_2u_3,hw_2w_3}^{H,x}(q) = R_{hu_2u_3,hw_2w_3}^{H,x}(q)$ and we may conclude by the induction hypothesis since $\ell(hw_2) < \ell(w_2)$, and $\ell(w_2) - \ell(u_2) = \ell(hw_2) - \ell(hu_2)$. If $h \notin D_L(u_2)$, then

$$\begin{aligned} R_{u,w}^{H,x}(q) &= R_{u_2u_3,w_2w_3}^{H,x}(q) = (q-1)R_{u_2u_3,hw_2w_3}^{H,x}(q) + qR_{hu_2u_3,hw_2w_3}^{H,x}(q) \\ &= (q-1)[R_{i-1} \cdot R_{u_3,w_3}^{H,x}(q) + qR_{i-2} \cdot R_{pu_3,w_3}^{H,x}(q)] + q[R_{i-2} \cdot R_{u_3,w_3}^{H,x}(q) + qR_{i-3} \cdot R_{pu_3,w_3}^{H,x}(q)] \\ &= [(q-1)R_{i-1} + qR_{i-2}] \cdot R_{u_3,w_3}^{H,x}(q) + q[(q-1)R_{i-2} + qR_{i-3}] \cdot R_{pu_3,w_3}^{H,x}(q) \\ &= R_i \cdot R_{u_3,w_3}^{H,x}(q) + qR_{i-1} \cdot R_{pu_3,w_3}^{H,x}(q) \end{aligned}$$

where the last equation follows by Lemma 3.1. \square

Remark 3.5. Lemma 3.3 and the first part of Lemma 3.4 hold more generally (with the same straightforward proof) if we replace $\{s, t\}$ with an arbitrary subset $J \subseteq S$.

In the proof of the following result, as well as in the proof of the main result of this work, it is essential $x \in \{q, -1\}$; indeed, we repeatedly use that x satisfies

$$(3.1) \quad (q-1)(q-1-x) + q = (q-1-x)^2.$$

Lemma 3.6. Let $w = w_2 \cdot w_3 \in W^H$ and $u = u_2 \cdot u_3 \in W^H$ with:

- $u \leq w$,
- $w_2, u_2 \in W_{\{s,t\}}$,
- $w_3, u_3 \in {}_{\{s,t\}}W$,
- $|\{x \in \{s, t\} : x \leq w_3\}| \leq 1$.

Suppose that $(W_{\{s,t\}} \cdot u_3) \cap W^H$ is a chain (see Proposition 3.2) and let $r, \bar{r} \in \{s, t\}$ be such that $r \cdot u_3 \in W^H$ and $\bar{r} \cdot u_3 \notin W^H$.

Then $\ell(w_2) - \ell(u_2) \geq -1$ and the following statements hold.

D₋₁. If $\ell(w_2) - \ell(u_2) = -1$, then

$$R_{u,w}^{H,x}(q) = R_{ru_3,w_3}^{H,x}(q).$$

D₀. If $\ell(w_2) - \ell(u_2) = 0$, then

$$R_{u,w}^{H,x}(q) = \begin{cases} R_{u_3,w_3}^{H,x}(q) & \text{if } u_2 = w_2 \\ (q-1)R_{ru_3,w_3}^{H,x}(q) & \text{if } u_2 \neq w_2 \end{cases}$$

D₁. If $\ell(w_2) - \ell(u_2) = 1$, then

$$R_{u,w}^{H,x}(q) = \begin{cases} (q-1-x)R_{u_3,w_3}^{H,x}(q) & \text{if } \bar{r} \in D_R(w_2) \\ (q-1)R_{u_3,w_3}^{H,x}(q) & \text{if } \bar{r} \notin D_R(w_2) \text{ and } u_2 \neq e \\ (q-1)R_{u_3,w_3}^{H,x}(q) + qR_{ru_3,w_3}^{H,x}(q), & \text{if } \bar{r} \notin D_R(w_2) \text{ and } u_2 = e \end{cases}$$

D₂. If $\ell(w_2) - \ell(u_2) = 2$, then

$$R_{u,w}^{H,x}(q) = \begin{cases} (q-1-x)[(q-1)R_{u_3,w_3}^{H,x}(q) + qR_{ru_3,w_3}^{H,x}(q)], & \text{if } r \in D_R(w_2) \\ (q-1)[(q-1-x)R_{u_3,w_3}^{H,x}(q) + qR_{ru_3,w_3}^{H,x}(q)], & \text{if } r \notin D_R(w_2) \end{cases}$$

D_i. If $\ell(w_2) - \ell(u_2) \geq 3$, then

$$R_{u,w}^{H,x}(q) = (q-1)(q-1-x)^{\ell(w_2)-\ell(u_2)-2}[(q-1-x)R_{u_3,w_3}^{H,x}(q) + qR_{ru_3,w_3}^{H,x}(q)]$$

Proof. In this proof, we use the Subword Property (Theorem 2.1), Property (2.1), and the recursive formula of Theorem 2.10 several times without explicit mention.

Note that we have $D_R(u_2) = \{r\}$ unless $u_2 = e$; in particular, u_2 cannot be the top element of $W_{\{s,t\}}$ (if any). If $u_2 \neq e$, we let $l \in \{s,t\}$ be such that $\{l\} = D_L(u_2)$, so that u_2 has a (unique) reduced expression starting with l and ending with r .

Since $|\{x \in \{s,t\} : x \leq w_3\}| \leq 1$, the longest subword of type $s-t-s-t-\dots$ or $t-s-t-s-\dots$ of any reduced expression for w has length at most $\ell(w_2) + 1$, and hence $\ell(w_2) - \ell(u_2) \geq -1$.

Proof of D₋₁. Since $\ell(w_2) - \ell(u_2) = -1$, necessarily $u_2 = w_2 \cdot r$, as otherwise u could not be smaller than or equal to w . We have

$$R_{u,w}^{H,x}(q) = R_{w_2ru_3,w_2w_3}^{H,x}(q) = R_{ru_3,w_3}^{H,x}(q).$$

Proof of D₀. If $u_2 = w_2$, the assertion is immediate. If $u_2 \neq w_2$, there exists an element $v \in W_{\{s,t\}}$, with $\ell(v) = \ell(w_2) - 1 = \ell(u_2) - 1$, such that $u_2 = v \cdot r$ and $w_2 = \bar{l} \cdot v$. We have

$$R_{u,w}^{H,x}(q) = R_{vru_3,\bar{l}vw_3}^{H,x}(q) = (q-1)R_{vru_3,vw_3}^{H,x}(q) + qR_{\bar{l}vru_3,vw_3}^{H,x}(q) = (q-1)R_{ru_3,w_3}^{H,x}(q) + qR_{\bar{l}vru_3,vw_3}^{H,x}(q).$$

We cannot have $\bar{l} \cdot v \cdot r \cdot u_3 \leq v \cdot w_3$, since all subwords of any reduced expression of $v \cdot w_3$ of type $s-t-s-t-\dots$ or $t-s-t-s-\dots$ have length at most $\ell(v) + 1$ while $\ell(\bar{l} \cdot v \cdot r) = \ell(v) + 2$. Hence $R_{\bar{l}vru_3,vw_3}^{H,x}(q) = 0$, as desired.

Proof of D₁. Since $\ell(w_2) - \ell(u_2) = 1$, we have either

- (1) $w_2 = u_2 \cdot \bar{r}$, or
- (2) $w_2 = \bar{l} \cdot u_2 \neq u_2 \cdot \bar{r}$ and $u_2 \neq e$, or
- (3) $w_2 = r$ and $u_2 = e$.

In the first case, we have

$$R_{u,w}^{H,x}(q) = R_{u_2u_3, u_2\bar{r}w_3}^{H,x}(q) = R_{u_3, \bar{r}w_3}^{H,x}(q) = (q-1-x)R_{u_3, w_3}^{H,x}(q).$$

In the second case, we have

$$R_{u,w}^{H,x}(q) = R_{u_2u_3, \bar{l}u_2w_3}^{H,x}(q) = (q-1)R_{u_2u_3, u_2w_3}^{H,x}(q) + qR_{\bar{l}u_2u_3, u_2w_3}^{H,x}(q) = (q-1)R_{u_3, w_3}^{H,x}(q)$$

since $\bar{l} \cdot u_2 \cdot u_3 \not\leq u_2 \cdot w_3$. In the third case, the assertion is immediate.

Proof of D₂. Since $\ell(w_2) - \ell(u_2) = 2$, we have either

- $w_2 = u_2 \cdot \bar{r} \cdot r$, or
- $w_2 = \bar{l} \cdot u_2 \cdot \bar{r} \neq u_2 \cdot \bar{r} \cdot r$ (where we set $\bar{l} = r$ if $u_2 = e$).

In the first case, we have

$$\begin{aligned} R_{u,w}^{H,x}(q) &= R_{u_2u_3, u_2\bar{r}rw_3}^{H,x}(q) = R_{u_3, \bar{r}rw_3}^{H,x}(q) = (q-1-x)R_{u_3, rw_3}^{H,x}(q) \\ &= (q-1-x)[(q-1)R_{u_3, w_3}^{H,x}(q) + qR_{ru_3, w_3}^{H,x}(q)]. \end{aligned}$$

In the second case, we have

$$\begin{aligned} R_{u,w}^{H,x}(q) &= R_{u_2u_3, \bar{l}u_2\bar{r}w_3}^{H,x}(q) = (q-1)R_{u_2u_3, u_2\bar{r}w_3}^{H,x}(q) + qR_{\bar{l}u_2u_3, u_2\bar{r}w_3}^{H,x}(q) \\ &= (q-1)(q-1-x)R_{u_3, w_3}^{H,x}(q) + q(q-1)R_{ru_3, w_3}^{H,x}(q), \end{aligned}$$

where the last equality follows from Statements D₁ and D₀.

Proof of D_i. If $\ell(w_2) - \ell(u_2) = 3$, we have either

- $w_2 = u_2 \cdot \bar{r} \cdot r \cdot \bar{r}$, or
- $w_2 = \bar{l} \cdot u_2 \cdot \bar{r} \cdot r \neq u_2 \cdot \bar{r} \cdot r \cdot \bar{r}$ (where we set $\bar{l} = r$ if $u_2 = e$).

In the first case, we have

$$\begin{aligned} R_{u,w}^{H,x}(q) &= R_{u_2u_3, u_2\bar{r}r\bar{r}w_3}^{H,x}(q) = R_{u_3, \bar{r}r\bar{r}w_3}^{H,x}(q) = (q-1-x)R_{u_3, r\bar{r}w_3}^{H,x}(q) \\ &= (q-1-x)(q-1)[(q-1-x)R_{u_3, w_3}^{H,x}(q) + qR_{ru_3, w_3}^{H,x}(q)], \end{aligned}$$

by Statement D₂. In the second case, we have

$$\begin{aligned} R_{u,w}^{H,x}(q) &= R_{u_2u_3, \bar{l}u_2\bar{r}rw_3}^{H,x}(q) = (q-1)R_{u_2u_3, u_2\bar{r}rw_3}^{H,x}(q) + qR_{\bar{l}u_2u_3, u_2\bar{r}rw_3}^{H,x}(q) \\ &= (q-1)(q-1-x)[(q-1)R_{u_3, w_3}^{H,x}(q) + qR_{ru_3, w_3}^{H,x}(q)] + q(q-1)R_{u_3, w_3}^{H,x}(q) \\ &= (q-1)[(q-1)(q-1-x) + q]R_{u_3, w_3}^{H,x}(q) + q(q-1)(q-1-x)R_{ru_3, w_3}^{H,x}(q) \\ &= (q-1)(q-1-x)^2R_{u_3, w_3}^{H,x}(q) + q(q-1)(q-1-x)R_{ru_3, w_3}^{H,x}(q) \end{aligned}$$

by Statements D₂ and D₁, and by Eq. (3.1).

If $\ell(w_2) - \ell(u_2) = 4$, we have either

- $w_2 = u_2 \cdot \bar{r} \cdot r \cdot \bar{r} \cdot r$, or
- $w_2 = \bar{l} \cdot u_2 \cdot \bar{r} \cdot r \cdot \bar{r} \neq u_2 \cdot \bar{r} \cdot r \cdot \bar{r} \cdot r$ (where we set $\bar{l} = r$ if $u_2 = e$).

In the first case, we have

$$\begin{aligned} R_{u,w}^{H,x}(q) &= R_{u_2u_3, u_2\bar{r}r\bar{r}rw_3}^{H,x}(q) = R_{u_3, \bar{r}r\bar{r}rw_3}^{H,x}(q) \\ &= (q-1-x)R_{u_3, r\bar{r}rw_3}^{H,x}(q) = (q-1)(q-1-x)^2[(q-1-x)R_{u_3, w_3}^{H,x}(q) + \\ &\quad qR_{ru_3, w_3}^{H,x}(q)]. \end{aligned}$$

In the second case, we have

$$\begin{aligned}
R_{u,w}^{H,x}(q) &= R_{u_2u_3,\bar{l}u_2\bar{r}r\bar{r}w_3}^{H,x}(q) = (q-1)R_{u_2u_3,u_2\bar{r}r\bar{r}w_3}^{H,x}(q) + qR_{\bar{l}u_2u_3,u_2\bar{r}r\bar{r}w_3}^{H,x}(q) \\
&= (q-1)^2(q-1-x)[(q-1-x)R_{u_3,w_3}^{H,x}(q) + qR_{ru_3,w_3}^{H,x}(q)] + \\
&\quad q(q-1)[(q-1-x)R_{u_3,w_3}^{H,x}(q) + qR_{ru_3,w_3}^{H,x}(q)] \\
&= (q-1)(q-1-x)[(q-1)(q-1-x) + q]R_{u_3,w_3}^{H,x}(q) + \\
&\quad q(q-1)[(q-1)(q-1-x) + q]R_{ru_3,w_3}^{H,x}(q) \\
&= (q-1)(q-1-x)^3R_{u_3,w_3}^{H,x}(q) + q(q-1)(q-1-x)^2R_{ru_3,w_3}^{H,x}(q)
\end{aligned}$$

where the last equality follows by Eq. (3.1). In both cases we have used statements that we have already proved.

Suppose $\ell(w_2) - \ell(u_2) \geq 5$ and use induction on $\ell(w_2)$. The base of the induction is $u_2 = e$ and $w_2 \in \{r\bar{r}r\bar{r}r, \bar{r}r\bar{r}r\bar{r}\}$: the assertion follows by a direct computation that we omit.

Let $\ell(w_2) > 5$: if $l \in D_L(w_2)$, then $R_{u,w}^{H,x}(q) = R_{u_2u_3,w_2w_3}^{H,x}(q) = R_{lu_2u_3,lw_2w_3}^{H,x}(q)$ and we may conclude by the induction hypothesis since $\ell(lw_2) < \ell(w_2)$ and $\ell(w_2) - \ell(u_2) = \ell(lw_2) - \ell(lu_2)$. If $l \notin D_L(w_2)$, then consider $\bar{l} \in D_L(w_2)$: we have $\bar{l} \notin D_L(u)$ since $\bar{l} \notin D_L(u_2)$, and $\bar{l}u \in W^H$ since $u_2 \neq e$. Hence, using the induction hypothesis, we have

$$\begin{aligned}
R_{u,w}^{H,x}(q) &= (q-1)R_{u_2u_3,\bar{l}w_2w_3}^{H,x}(q) + qR_{\bar{l}u_2u_3,\bar{l}w_2w_3}^{H,x}(q) \\
&= (q-1)^2(q-1-x)^{\ell(\bar{l}w_2)-\ell(u_2)-2}[(q-1-x)R_{u_3,w_3}^{H,x}(q) + qR_{ru_3,w_3}^{H,x}(q)] \\
&\quad + q(q-1)(q-1-x)^{\ell(\bar{l}w_2)-\ell(\bar{l}u_2)-2}[(q-1-x)R_{u_3,w_3}^{H,x}(q) + qR_{ru_3,w_3}^{H,x}(q)] \\
&= (q-1)(q-1-x)^{\ell(w_2)-\ell(u_2)-3}[(q-1)(q-1-x) + q]R_{u_3,w_3}^{H,x}(q) + \\
&\quad q(q-1)(q-1-x)^{\ell(w_2)-\ell(u_2)-4}[(q-1)(q-1-x) + q]R_{ru_3,w_3}^{H,x}(q) \\
&= (q-1)(q-1-x)^{\ell(w_2)-\ell(u_2)-2}[(q-1-x)R_{u_3,w_3}^{H,x}(q) + qR_{ru_3,w_3}^{H,x}(q)]
\end{aligned}$$

where the last equality follows by Eq. (3.1). \square

Remark 3.7. Lemma 3.6 implies that, under its hypotheses, $R_{u,w}^{H,x}$ is a combination of $R_{u_3,w_3}^{H,x}$ and $R_{ru_3,w_3}^{H,x}$ with coefficients in $\mathbb{Z}[q]$. Furthermore, if either

- $\ell(w_2) - \ell(u_2) \geq 3$, or
- $\ell(w_2) - \ell(u_2) = 2$ and $\{x \in \{s, t\} : x \leq w_3\} = \emptyset$,

then the coefficients of the combination depend only on $\ell(w_2) - \ell(u_2)$.

4. MAIN RESULT

In this section, we prove Theorem 1.5, whose implications were discussed in Section 1. In particular, for any arbitrary Coxeter system (W, S) , any arbitrary subset $H \subseteq S$, and any arbitrary element $w \in W^H$, we give an algorithm for computing the parabolic Kazhan–Lusztig R -polynomials $\{R_{u,w}^{H,x}(q)\}_{u \in W^H}$ once one knows the poset-isomorphism class of the interval $[e, w]$, and which elements of the interval $[e, w]$ belong to W^H .

As an immediate corollary, we have that it is possible to compute also the parabolic Kazhdan–Lusztig polynomials $\{P_{u,w}^{H,x}(q)\}_{u \in W^H}$ from the knowledge only of the poset-isomorphism class of the interval $[e, w]$ and which elements of the interval $[e, w]$ belong to W^H (see Remark 2.12).

First, we give the following general definition.

Definition 4.1. *Let P be a poset and $T \subseteq P$ be a subposet of P . A relative special matching of P with respect to T is a special matching M of P such that, if $p \in T$ and $M(p) \triangleleft p$, then $M(p) \in T$.*

Now, fix an arbitrary Coxeter system (W, S) , a subset $H \subseteq S$, and an element $w \in W^H$. An H -special matching of w is a relative special matching of $[e, w]$ with respect to $[e, w]^H$, that is a special matching M of w such that, if $u \leq w$, $u \in W^H$, and $M(u) \triangleleft u$, then $M(u) \in W^H$.

Note that the \emptyset -special matchings are exactly the special matchings and that all left multiplication matchings are H -special, for all $H \subseteq S$.

We say that an H -special matching M of w *calculates* the parabolic Kazhdan–Lusztig R -polynomials (or, simply, is *calculating*) provided

$$(4.1) \quad R_{u,w}^{H,x}(q) = \begin{cases} R_{M(u),M(w)}^{H,x}(q), & \text{if } M(u) \triangleleft u, \\ (q-1)R_{u,M(w)}^{H,x}(q) + qR_{M(u),M(w)}^{H,x}(q), & \text{if } M(u) \triangleright u \text{ and } M(u) \in W^H, \\ (q-1-x)R_{u,M(w)}^{H,x}(q), & \text{if } M(u) \triangleright u \text{ and } M(u) \notin W^H. \end{cases}$$

for all $u \in W^H$ with $u \leq w$. Clearly, all left multiplication matchings are calculating. Actually, our target is to prove that all H -special matchings are calculating.

We need the following result (see [21, Theorem 4.2]).

Theorem 4.2. *Let M be an H -special matching of w . If*

- *every H -special matching of v is calculating, for all $v \in W^H$ with $v < w$, and*
- *there exists a calculating special matching N of w commuting with M and such that $M(w) \neq N(w)$,*

then M is calculating.

We also need the following easy lemma.

Lemma 4.3. *Let $s, t \in S$, $g_{st} \in W_{\{s,t\}}$, $p \in D_R(g_{st})$, and M be a special matching of the dihedral interval $[e, g_{st}]$. If*

- *M commutes with ρ_p , and*
- *$M(x) \neq \rho_p(x)$, for all $x \in W_{\{s,t\}}$ such that $\ell(x) \neq 0, 1$ and, if $W_{\{s,t\}}$ is finite, $\ell(x) \neq m_{s,t} - 1, m_{s,t}$,*

then M is a left multiplication matching.

Proof. Without loss of generality, we suppose $M(e) = s$. We need to show $M(x) = sx$, for all $x \in [e, g_{st}]$. By contradiction, let x be minimal such that $M(x) \neq sx$.

Clearly $x \notin \{e, s\}$. By minimality, $x \triangleleft M(x)$, and $s \notin D_L(x)$ as otherwise $M(sx)$ would be x since sx would be smaller than x . Moreover, if $W_{\{s,t\}}$ is finite and g_{st} is its longest element w_0 , then $x \notin \{w_0, sw_0\}$. The element $M(x)$ cannot be xp (by hypothesis,

since at least one among x and $M(x)$ has length not in $\{0, 1, m_{s,t}-1, m_{s,t}\}$; $M(x)$ cannot be sx (by assumption); $M(x)$ cannot be tx (since $tx \triangleleft x$). The only possibility left is $M(x) = x\bar{p}$ (recall that \bar{p} is the element in $\{s, t\} \setminus \{p\}$ and notice that, if $W_{\{s,t\}}$ is finite, $x\bar{p}$ is not w_0 since otherwise w_0 would be equal to sx). Hence the element $\rho_p M \rho_p(xp)$, which is $x\bar{p}p$, would have length equal to $\ell(xp) + 3$, and so $M \rho_p(xp) \neq \rho_p M(xp)$, which contradicts the fact that M commutes with ρ_p . \square

We now recall and prove Theorem 1.5.

Theorem. *Given an arbitrary Coxeter system (W, S) and a subset $H \subset S$, let w be any element in W^H . Then all H -special matchings of w calculate the parabolic Kazhdan–Lusztig R -polynomials of W^H .*

Proof. We use induction on $\ell(w)$, the case $\ell(w) \leq 1$ being trivial. Suppose $\ell(w) > 1$.

Let M be an H -special matching of w and $u \in W^H$, with $u \leq w$. We need to show

$$R_{u,w}^{H,x}(q) = \begin{cases} R_{M(u),M(w)}^{H,x}(q), & \text{if } M(u) \triangleleft u, \\ (q-1)R_{u,M(w)}^{H,x}(q) + qR_{M(u),M(w)}^{H,x}(q), & \text{if } M(u) \triangleright u \text{ and } M(u) \in W^H, \\ (q-1-x)R_{u,M(w)}^{H,x}(q), & \text{if } M(u) \triangleright u \text{ and } M(u) \notin W^H. \end{cases}$$

We may suppose that M does not agree with a left multiplication matchings on both u and w , because otherwise the assertion is clear since left multiplication matchings are calculating.

If there exists a left multiplication matching λ of w commuting with M such that $\lambda(w) \neq M(w)$, then we can conclude by Theorem 4.2.

By Theorem 2.9, M is associated with a system (J, s, t, M_{st}) . Suppose first that (J, s, t, M_{st}) is a right system and $(w^J)^{\{s,t\}} \neq e$. Fix $l \in D_L((w^J)^{\{s,t\}})$; thus $l \in D_L(w)$ and λ_l is a special matching of w that satisfies $M(w) \neq \lambda_l(w)$ since

$$M(w) = (w^J)^{\{s,t\}} \cdot M_{st} \left((w^J)_{\{s,t\}} \cdot {}_{\{s\}}(w_J) \right) \cdot {}^{\{s\}}(w_J)$$

while

$$\lambda_l(w) = l(w^J)^{\{s,t\}} \cdot (w^J)_{\{s,t\}} \cdot {}_{\{s\}}(w_J) \cdot {}^{\{s\}}(w_J).$$

We need to show that M and λ_l commute. In order to apply Lemma 2.5, we distinguish the following cases.

(a) $l \notin \{s, t\}$.

By Property R3 of the definition of a right system, either $l \notin J$ or l commutes with s . In the first case, M acts as ρ_s on $[e, w_0(s, l)]$ and hence commutes with λ_l on $[e, w_0(s, l)]$. In the second case, M and λ_l clearly commutes on $[e, w_0(s, l)]$ since $[e, w_0(s, l)]$ is a dihedral interval with 4 elements.

(b) $l = t$.

By Property R4, M commutes with λ_t on $[e, w_0(s, t)]$.

(c) $l = s$.

We need to show that M and λ_s commute on every lower dihedral intervals $[e, w_0(s, r)]$, with $r \in S \setminus \{s\}$. For $r = t$, it follows from Property R4. For $r \neq t$, M acts on $[e, w_0(s, r)]$ as ρ_s or λ_s , and in both cases M commutes with λ_s on $[e, w_0(s, r)]$.

Now suppose that (J, s, t, M_{st}) is a left system and $(_Jw)^{\{s\}} \neq e$. Fix $l \in D_L((_Jw)^{\{s\}})$; thus $l \in D_L(w)$ and λ_l is a special matching of w that satisfies $M(w) \neq \lambda_l(w)$ since

$$M(w) = (_Jw)^{\{s\}} \cdot M_{st}\left((_Jw)_{\{s\}} \cdot {}_{\{s,t\}}(^Jw)\right) \cdot {}^{\{s,t\}}(^Jw)$$

while

$$\lambda_l(w) = l(_Jw)^{\{s\}} \cdot (_Jw)_{\{s\}} \cdot {}_{\{s,t\}}(^Jw) \cdot {}^{\{s,t\}}(^Jw).$$

In order to show that M and λ_l commute, we again apply Lemma 2.5. If $l \neq s$, then M acts as ρ_s on $[e, w_0(s, l)]$ and hence commutes with λ_l . Suppose $l = s$; we need to show that M and λ_s commute on every lower dihedral intervals $[e, w_0(s, r)]$, with $r \in S \setminus \{s\}$. If $r = t$, it follows from Property L5. If $r \neq t$, then M acts on $[e, w_0(s, r)]$ as ρ_s or λ_s , and so M commutes with λ_s .

Hence we may suppose that either

- (1) (J, s, t, M_{st}) is a right system and $(w^J)^{\{s,t\}} = e$, or
- (2) (J, s, t, M_{st}) is a left system and $(_Jw)^{\{s\}} = e$.

In the first case, we set $w_2 = (w^J)_{\{s,t\}} \cdot {}_{\{s\}}(w_J)$, $w_3 = {}^{\{s\}}(w_J)$, $u_2 = (u^J)_{\{s,t\}} \cdot {}_{\{s\}}(u_J)$, $u_3 = {}^{\{s\}}(u_J)$. In the second case, we set $w_2 = (_Jw)_{\{s\}} \cdot {}_{\{s,t\}}(^Jw)$, $w_3 = {}^{\{s,t\}}(^Jw)$, $u_2 = (_Ju)_{\{s\}} \cdot {}_{\{s,t\}}(^Ju)$, $u_3 = {}^{\{s,t\}}(^Ju)$. In both cases, we get the $W_{\{s,t\}} \times {}^{\{s,t\}}W$ -factorization of w and u :

$$w = w_2 \cdot w_3 \quad u = u_2 \cdot u_3.$$

Note:

- $u \leq w$,
- $w_2, u_2 \in W_{\{s,t\}}$,
- $w_3, u_3 \in {}^{\{s,t\}}W$,
- $u_3 \leq w_3$,
- $|\{x \in \{s, t\} : x \leq w_3\}| \leq 1$ (in the first case, this is trivial since $t \notin J$ and $w_3 \in W_J$; in the second case s and t cannot be both smaller than or equal to w_3 since otherwise $M_{st} = \lambda_s$ by Property L4 and hence $M = \lambda_s$) and, if this cardinality is 1, we let p be such that $\{p\} = \{x \in \{s, t\} : x \leq w_3\}$,
- $M(w) = M(w_2) \cdot w_3$ and $M(u) = M(u_2) \cdot u_3$,
- M acts as λ_s on $[e, w_0(s, r)]$, for all $r \in S \setminus \{t\}$ such that $r \leq w$,
- if $\{p\} = \{x \in \{s, t\} : x \leq w_3\}$, then M commutes with ρ_p on $[e, w_0(s, t)]$ (by either Property R5 or Property L4).

Recall Proposition 3.2. If $(W_{\{s,t\}} \cdot u_3) \cap W^H = \{u_3\}$, then we may conclude using Lemma 3.3.

We now suppose $W_{\{s,t\}} \cdot u_3 \subseteq W^H$ and apply Lemma 3.4. Suppose $\ell(w_2) - \ell(u_2) = i$, $i \geq 2$. By Lemma 3.4, we have

$$R_{u,w}^{H,x}(q) = R_i \cdot R_{u_3,w_3}^{H,x}(q) + qR_{i-1} \cdot R_{pu_3,w_3}^{H,x}(q).$$

If $M(u) \triangleleft u$, then $\ell(M(w_2)) - \ell(M(u_2)) = i$ and also $R_{M(u),M(w)}^{H,x}(q)$ is equal to $R_i \cdot R_{u_3,w_3}^{H,x}(q) + qR_{i-1} \cdot R_{pu_3,w_3}^{H,x}$, by Lemma 3.4.

Suppose $u \triangleleft M(u)$. If $\ell(w_2) - \ell(u_2) \geq 4$ then

$$(q-1)R_{u_2u_3, M(w_2)w_3}^{H,x}(q) + qR_{M(u_2)u_3, M(w_2)w_3}^{H,x}(q)$$

is equal to

$$\begin{aligned} &= (q-1)[R_{i-1} \cdot R_{u_3, w_3}^{H,x}(q) + qR_{i-2} \cdot R_{pu_3, w_3}^{H,x}] + q[R_{i-2} \cdot R_{u_3, w_3}^{H,x}(q) + qR_{i-3} \cdot R_{pu_3, w_3}^{H,x}] \\ &= [(q-1)R_{i-1} + qR_{i-2}] \cdot R_{u_3, w_3}^{H,x}(q) + q[(q-1)R_{i-2} + qR_{i-3}] \cdot R_{pu_3, w_3}^{H,x} \\ &= R_i \cdot R_{u_3, w_3}^{H,x}(q) + qR_{i-1} \cdot R_{pu_3, w_3}^{H,x} \end{aligned}$$

by Lemmas 3.1 and 3.4, as desired.

If $\ell(w_2) - \ell(u_2) = 3$, then Lemma 3.4 implies that

$$(q-1)R_{u_2u_3, M(w_2)w_3}^{H,x}(q) + qR_{M(u_2)u_3, M(w_2)w_3}^{H,x}(q)$$

is equal to

$$(4.2) \quad (q-1)[R_2 \cdot R_{u_3, w_3}^{H,x}(q) + qR_1 \cdot R_{pu_3, w_3}^{H,x}] + q[(q-1)R_{u_3, w_3}^{H,x}(q) + qR_{pu_3, w_3}^{H,x} \cdot \chi]$$

where $\chi = \begin{cases} 1 & \text{if } M(w_2) = M(u_2) \cdot p \\ 0 & \text{otherwise.} \end{cases}$

The term $R_{pu_3, w_3}^{H,x} \cdot \chi$ is always 0: indeed, if $p \leq w_3$, then $M(w_2) \neq M(u_2) \cdot p$ since, otherwise, $w_2 = M(M(u_2) \cdot p) = M \circ \rho_p(M(u_2)) \neq \rho_p \circ M(M(u_2)) = u_2p$ and M would not commute with ρ_p on $[e, w_0(s, t)]$. Hence the polynomial in (4.2) is always equal to

$$\begin{aligned} &= [(q-1)R_2 + q(q-1)]R_{u_3, w_3}^{H,x}(q) + q(q-1)R_1 \cdot R_{pu_3, w_3}^{H,x} \\ &= R_3 \cdot R_{u_3, w_3}^{H,x}(q) + qR_2 \cdot R_{pu_3, w_3}^{H,x}, \end{aligned}$$

as desired.

Suppose $\ell(w_2) - \ell(u_2) = 2$. If $\{x \in \{s, t\} : x \leq w_3\} = \emptyset$, then

$$(q-1)R_{u_2u_3, M(w_2)w_3}^{H,x}(q) + qR_{M(u_2)u_3, M(w_2)w_3}^{H,x}(q)$$

is equal to $(q-1)^2R_{u_3, w_3}^{H,x}(q)$ and the assertion follows. Suppose $\{x \in \{s, t\} : x \leq w_3\} = \{p\}$ and recall that, in this case, M commutes with ρ_p on $[e, w_0(s, t)]$: in order to compute

$$(4.3) \quad (q-1)R_{u_2u_3, M(w_2)w_3}^{H,x}(q) + qR_{M(u_2)u_3, M(w_2)w_3}^{H,x}(q)$$

we distinguish two cases, according to as whether p is in $D_R(M(u_2))$ or not. If $p \in D_R(M(u_2))$, then there exist $r \in \{s, t\}$ and $v \in W_{\{s, t\}}$ with $\ell(v) = \ell(u_2)$ such that $M(u_2) = v \cdot p$ and $M(w_2) = r \cdot v$. The polynomial in (4.3) is equal to

$$\begin{aligned} &= (q-1)R_{u_2u_3, rvw_3}^{H,x}(q) + qR_{vpu_3, rvw_3}^{H,x}(q) \\ &= (q-1)^2R_{u_3, w_3}^{H,x}(q) + q(q-1)R_{pu_3, w_3}^{H,x}(q) \end{aligned}$$

since Lemma 3.4 implies

- $R_{u_2u_3, rvw_3}^{H,x}(q) = (q-1)R_{u_3, w_3}^{H,x}(q)$ (notice $r \cdot v \neq u_2p$),
- $R_{vpu_3, rvw_3}^{H,x}(q) = (q-1)R_{pu_3, w_3}^{H,x}(q)$ (notice $vpu_3 \leq rvw_3$ if and only if $pu_3 \leq w_3$).

If $p \notin D_R(M(u_2))$, then $M(u_2) \cdot p = M(u_2 \cdot p) = w_2$ and $M(w_2) = u_2 \cdot p$, as otherwise $\rho_p \circ M(u_2) \neq M \circ \rho_p(u_2)$. Thus $M(u_2) \not\leq M(w_2) > M(w_2)p$ and $M(u_2) \cdot u_3 \not\leq M(w_2) \cdot w_3$, and so the polynomial in (4.3) is equal to

$$= (q-1)R_{u_2u_3, u_2pw_3}^{H,x}(q) = (q-1)[(q-1)R_{u_3, w_3}^{H,x}(q) + qR_{pu_3, w_3}^{H,x}(q)].$$

Suppose now $\ell(w_2) - \ell(u_2) = 1$. By Lemma 3.4, we have

$$R_{u,w}^{H,x}(q) = \begin{cases} (q-1)R_{u_3, w_3}^{H,x}(q) + qR_{pu_3, w_3}^{H,x}(q), & \text{if } w_2 = u_2 \cdot p \\ (q-1)R_{u_3, w_3}^{H,x}(q), & \text{otherwise.} \end{cases}$$

If $M(u_2) \triangleright u_2$, then $M(u_2) = w_2$, since M is a special matching, and

$$\begin{aligned} (q-1)R_{u_2u_3, M(w_2)w_3}^{H,x}(q) + qR_{M(u_2)u_3, M(w_2)w_3}^{H,x}(q) &= (q-1)R_{u_2u_3, u_2w_3}^{H,x}(q) + qR_{w_2u_3, u_2w_3}^{H,x}(q) \\ &= (q-1)R_{u_3, w_3}^{H,x}(q) + qR_{w_2u_3, u_2w_3}^{H,x}(q) \end{aligned}$$

where $R_{w_2u_3, u_2w_3}^{H,x}(q) = \begin{cases} R_{pu_3, w_3}^{H,x}(q), & \text{if } w_2 = u_2 \cdot p \\ 0, & \text{otherwise.} \end{cases}$

If $M(u_2) \triangleleft u_2$, then either $\{x \in \{s, t\} : x \leq w_3\} = \emptyset$, or $\{x \in \{s, t\} : x \leq w_3\} = \{p\}$ and $w_2 = u_2 \cdot p$ if and only if $M(w_2) = M(u_2) \cdot p$ since M and ρ_p commute. Hence $R_{u,w}^{H,x}(q) = R_{M(u), M(w)}^{H,x}(q)$, by Lemma 3.4.

Suppose $\ell(w_2) - \ell(u_2) = 0$. If $w_2 = u_2$ the assertion is trivial. Otherwise, $u_2 \triangleleft M(u_2) = w_0(s, t)$, $M(w_2) \triangleleft w_2$ and $M(w_2) \triangleleft u_2$. Since $u \leq w$, necessarily $p \leq w_3$, $p \in D_R(u_2)$, $u_2 = (lw_2) \cdot p$ where $l \in D_L(w_2) \setminus D_L(u_2)$, $w_0(s, t) = w_2 \cdot p = l \cdot u_2$ and, since $M \circ \rho_p(w_0(s, t)) = \rho_p \circ M(w_0(s, t))$, we have $u_2 = M(w_2) \cdot p$. Thus

$$\begin{aligned} (q-1)R_{u_2u_3, M(w_2)w_3}^{H,x}(q) + qR_{M(u_2)u_3, M(w_2)w_3}^{H,x}(q) &= (q-1)R_{M(w_2)pu_3, M(w_2)w_3}^{H,x}(q) \\ &= (q-1)R_{pu_3, w_3}^{H,x}(q) \end{aligned}$$

since $M(u_2)u_3 \not\leq M(w_2)w_3$, and we conclude by Lemma 3.4.

Suppose $\ell(w_2) - \ell(u_2) = -1$. Thus $u_2 = w_2 \cdot p$ as otherwise $u \not\leq v$, and $u_2 = w_0(s, t)$. Hence $M(u_2) \triangleleft u_2$ and we conclude by Lemma 3.4.

We are left with the case when $(W_{\{s, t\}} \cdot u_3) \cap W^H$ is a chain and we apply Lemma 3.6. Let $r, \bar{r} \in \{s, t\}$ be such that $ru_3 \in W^H$ and $\bar{r}u_3 \notin W^H$.

Suppose $\ell(w_2) - \ell(u_2) \geq 4$. By Lemma 3.6, we have

$$(4.4) \quad R_{u,w}^{H,x}(q) = (q-1)(q-1-x)^{\ell(w_2)-\ell(u_2)-2}[(q-1-x)R_{u_3, w_3}^{H,x}(q) + qR_{ru_3, w_3}^{H,x}(q)].$$

If $M(u) \triangleleft u$, then we can conclude since $\ell(M(w_2)) - \ell(M(u_2)) = \ell(w_2) - \ell(u_2)$ (see Remark 3.7).

If $u \triangleleft M(u) \notin W^H$, then

$$\begin{aligned} R_{u, M(w)}^{H,x}(q) &= (q-1)(q-1-x)^{\ell(M(w_2))-\ell(u_2)-2}[(q-1-x)R_{u_3, w_3}^{H,x}(q) + qR_{ru_3, w_3}^{H,x}(q)] \\ &= (q-1)(q-1-x)^{\ell(w_2)-\ell(u_2)-3}[(q-1-x)R_{u_3, w_3}^{H,x}(q) + qR_{ru_3, w_3}^{H,x}(q)] \end{aligned}$$

by Lemma 3.6. Hence $(q-1-x)R_{u, M(w)}^{H,x}(q)$ is equal to the right side of equation (4.4), as desired.

If $u \triangleleft M(u) \in W^H$, we separate two cases. If either $\ell(w_2) - \ell(u_2) \geq 5$, or $\ell(w_2) - \ell(u_2) = 4$ and $\{x \in \{s, t\} : x \leq w_3\} = \emptyset$, then

$$(q-1)R_{u_2 u_3, M(w_2)w_3}^{H,x}(q) + qR_{M(u_2)u_3, M(w_2)w_3}^{H,x}(q)$$

is equal to

$$\begin{aligned} &= (q-1)(q-1)(q-1-x)^{\ell(M(w_2))-\ell(u_2)-2}[(q-1-x)R_{u_3, w_3}^{H,x}(q) + qR_{ru_3, w_3}^{H,x}(q)] + \\ &\quad q(q-1)(q-1-x)^{\ell(M(w_2))-\ell(M(u_2))-2}[(q-1-x)R_{u_3, w_3}^{H,x}(q) + qR_{ru_3, w_3}^{H,x}(q)] \\ &= (q-1)(q-1-x)^{\ell(w_2)-\ell(u_2)-3}[(q-1)(q-1-x) + q]R_{u_3, w_3}^{H,x}(q) + \\ &\quad q(q-1)(q-1-x)^{\ell(w_2)-\ell(u_2)-4}[(q-1)(q-1-x) + q]R_{ru_3, w_3}^{H,x}(q) \\ &= (q-1)(q-1-x)^{\ell(w_2)-\ell(u_2)-2}[(q-1-x)R_{u_3, w_3}^{H,x}(q) + qR_{ru_3, w_3}^{H,x}(q)] \end{aligned}$$

by Lemma 3.6 and Eq. (3.1), as desired.

If $\ell(w_2) - \ell(u_2) = 4$ and $|\{x \in \{s, t\} : x \leq w_3\}| = 1$, then, by Lemma 3.6, $(q-1)R_{u_2 u_3, M(w_2)w_3}^{H,x}(q) + qR_{M(u_2)u_3, M(w_2)w_3}^{H,x}(q)$ is equal to

$$\begin{aligned} &= (q-1)(q-1)(q-1-x)[(q-1-x)R_{u_3, w_3}^{H,x}(q) + qR_{ru_3, w_3}^{H,x}(q)] + \\ &\quad q[(q-1)(q-1-x)R_{u_3, w_3}^{H,x}(q) + qAR_{ru_3, w_3}^{H,x}(q)] \\ &= (q-1)(q-1-x)[(q-1)(q-1-x) + q]R_{u_3, w_3}^{H,x}(q) + \\ &\quad q[(q-1)^2(q-1-x) + qA]R_{ru_3, w_3}^{H,x}(q) \\ &= (q-1)(q-1-x)^3R_{u_3, w_3}^{H,x}(q) + q[(q-1)^2(q-1-x) + qA]R_{ru_3, w_3}^{H,x}(q) \end{aligned}$$

where $A = \begin{cases} (q-1-x), & \text{if } r \in D_R(w_2) \\ (q-1), & \text{if } r \notin D_R(w_2) \end{cases}$ and the last equation holds by Eq. (3.1).

If $r \cdot u_3 \not\leq w_3$, we are done. Let us show that $r \cdot u_3$ cannot be smaller than or equal to w_3 by contradiction. We would have $r \leq w_3$ (so $r = p$) and M would commute with ρ_r on $[e, w_0(s, t)]$: on the other hand, $M(x) \neq xr$ for all $x \in W_{\{s, t\}}$ with $\ell(x) \neq 0, 1, m_{s, t} - 1, m_{s, t}$, since M is H -special and $M(x \cdot u_3)$ cannot be $xr \cdot u_3$. By Lemma 4.3, these two facts together would imply that M is a left multiplication matching on $[e, w_0(s, t)]$, and M would be a left multiplication matching on $[e, w]$, which is a contradiction.

Suppose $\ell(w_2) - \ell(u_2) = 3$. By Lemma 3.6

$$R_{u, w}^{H,x}(q) = (q-1)(q-1-x)[(q-1-x)R_{u_3, w_3}^{H,x}(q) + qR_{ru_3, w_3}^{H,x}(q)].$$

If $M(u) \triangleleft u$, then we conclude by Lemma 3.6.

If $u \triangleleft M(u) \notin W^H$, then

$$R_{u, M(u)}^{H,x}(q) = \begin{cases} (q-1-x)[(q-1)R_{u_3, w_3}^{H,x}(q) + qR_{ru_3, w_3}^{H,x}(q)], & \text{if } r \in D_R(w_2) \\ (q-1)[(q-1-x)R_{u_3, w_3}^{H,x}(q) + qR_{ru_3, w_3}^{H,x}(q)], & \text{if } r \notin D_R(w_2) \end{cases}$$

by Lemma 3.6. If $r \cdot u_3 \not\leq w_3$, we are done.

Let us show that $r \cdot u_3$ cannot be smaller than or equal to w_3 by contradiction.

We would have $r = p$, and M would commute with ρ_r on $[e, w_0(s, t)]$ since $r \cdot u_3 \leq w_3$ implies $r \leq w_3$: on the other hand, $M(x) \neq xr$ for all $x \in W_{\{s, t\}}$ with $\ell(x) \neq 0, 1, m_{s, t} - 1, m_{s, t}$

1, $m_{s,t}$, since M is H -special and $M(x \cdot u_3)$ cannot be $xr \cdot u_3$. By Lemma 4.3, these two facts together would imply that M is a left multiplication matching on $[e, w_0(s, t)]$ and so M would be a left multiplication matching also on $[e, w]$, which is impossible.

If $u \triangleleft M(u) \in W^H$, then

- $M(u_2) = l \cdot u_2$, where $l \in \{s, t\} \setminus D_L(u_2)$ and $l \cdot u_2$ is not the longest element in $W_{\{s,t\}}$ (if any),
- $M(u_2 \cdot \bar{r}) = l \cdot u_2 \cdot \bar{r}$ since M is H -special (as otherwise we would have $M(u_2 \cdot \bar{r} \cdot u_3) = \bar{l} \cdot l \cdot u_2 \cdot u_3 \in W^H$, with $u_2 \cdot \bar{r} \cdot u_3 \notin W^H$).

Hence the only possibility is $M(w_2) = \bar{l} \cdot l \cdot u_2$ (recall $M(w_2) \triangleleft w_2$) and $w_2 \in \{l \cdot \bar{l} \cdot l \cdot u_2, \bar{l} \cdot l \cdot u_2 \cdot \bar{r}\}$. But $w_2 = l \cdot \bar{l} \cdot l \cdot u_2$ is not allowed since M would agree with λ_l on both w and u , which is impossible. Thus $w_2 = \bar{l} \cdot l \cdot u_2 \cdot \bar{r} \neq l \cdot \bar{l} \cdot l \cdot u_2$, and

$$(q-1)R_{u_2u_3, M(w_2)w_3}^{H,x}(q) + qR_{M(u_2)u_3, M(w_2)w_3}^{H,x}(q)$$

is equal to

$$\begin{aligned} &= (q-1)R_{u_2u_3, \bar{l}lu_2w_3}^{H,x}(q) + qR_{lu_2u_3, \bar{l}lu_2w_3}^{H,x}(q) \\ &= (q-1)(q-1-x)[(q-1)R_{u_3, w_3}^{H,x}(q) + qR_{ru_3, w_3}^{H,x}(q)] + q(q-1)R_{u_3, w_3}^{H,x}(q) \\ &= (q-1)[(q-1)(q-1-x) + q]R_{u_3, w_3}^{H,x}(q) + q(q-1)(q-1-x)R_{ru_3, w_3}^{H,x}(q) \\ &= (q-1)(q-1-x)^2R_{u_3, w_3}^{H,x}(q) + q(q-1)(q-1-x)R_{ru_3, w_3}^{H,x}(q) \end{aligned}$$

by Lemma 3.6 and Eq. (3.1), as desired.

Suppose $\ell(w_2) - \ell(u_2) = 2$.

If $M(u) \triangleleft u$, then $M(u_2) = lu_2$ with $l \in D_L(u_2)$ (since M is H -special), $M(M(u_2) \cdot \bar{r}) = l \cdot M(u_2) \cdot \bar{r} = u_2 \cdot \bar{r}$, $M(w_2) = M(u_2) \cdot \bar{r} \cdot r$, and w_2 cannot be $l \cdot \bar{l} \cdot u_2$ since otherwise M and λ_l would agree on both w and u . Thus $w_2 = \bar{l} \cdot u_2 \cdot \bar{r} \neq l \cdot \bar{l} \cdot u_2$, and $w_2 = M \circ \rho_r(M(u_2) \cdot \bar{r}) \neq \rho_r \circ M(M(u_2) \cdot \bar{r})$, which implies $r \not\leq w_3$: hence $R_{u,w}^{H,x}(q) = R_{M(u), M(w)}^{H,x}(q)$ since they are both equal to

$$(q-1)(q-1-x)R_{u_3, w_3}^{H,x}(q)$$

by Lemma 3.6.

If $u \triangleleft M(u) \notin W^H$, then $M(u_2) = u_2 \cdot \bar{r}$, $w_2 \in \{l \cdot u_2 \cdot \bar{r}, \bar{l} \cdot l \cdot u_2\}$, and $M(w_2) = l \cdot u_2$, where $l \in \{s, t\} \setminus D_L(u_2)$ if $u_2 \neq e$ and $l = r$ if $u_2 = e$. We have

$$\begin{aligned} R_{u, M(w)}^{H,x}(q) &= R_{u_2u_3, lu_2w_3}^{H,x}(q) = (q-1)R_{u_2u_3, u_2w_3}^{H,x}(q) + qR_{lu_2u_3, u_2w_3}^{H,x}(q) \\ &= (q-1)R_{u_3, w_3}^{H,x}(q) + qR_{lu_2u_3, u_2w_3}^{H,x}(q) \end{aligned}$$

where the last term is 0 unless $u_2 = e$ (and so $l = r$) and $r \cdot u_3 \leq w_3$ (so $r = p$). If $\{x \in \{s, t\} : x \leq w_3\}$ is either empty or $\{\bar{r}\}$, we conclude by Lemma 3.6. If $\{x \in \{s, t\} : x \leq w_3\} = \{r\}$, then M must commute with ρ_r on $[e, w_0(s, t)]$: this implies $u_2 = e$ (since $\rho_r \circ M(u_2) = u_2 \cdot \bar{r} \cdot r$ while $\rho_r(y) \triangleleft y$ for all $y \in (W^H \cap W_{\{s,t\}}) \setminus \{e\}$), and $w_2 = \bar{r} \cdot r \cdot u_2 = \bar{r} \cdot r$ (since $\bar{r} \cdot r$ must be the element of length 2 in $W_{\{s,t\}}$ covering its matched element as otherwise $\rho_r \circ M(e)$ could not be equal to $M \circ \rho_r(e)$). The assertion then follows.

If $u \triangleleft M(u) \in W^H$, then $M(u_2) = l \cdot u_2$ with $l \in \{s, t\} \setminus D_L(u_2)$ (and $l = r$ if $u_2 = e$), and $M(w_2) = u_2 \cdot \bar{r}$. Moreover, $w_2 \neq l \cdot u_2 \cdot \bar{r}$ (as otherwise M and λ_l would agree on both w and u). Hence w_2 should be equal to $\bar{l} \cdot l \cdot u_2$ but also this is not possible since the element $w_2 \cdot u_3 = \bar{l} \cdot l \cdot u_2 \cdot u_3 \in W^H$ (which belongs to $[e, w]$ since $u_3 \leq w_3$) would be matched with $u_2 \cdot \bar{r} \cdot u_3 \notin W^H$, and this contradicts the definition of H -special.

Suppose $\ell(w_2) - \ell(u_2) = 1$.

If $M(u) \triangleleft u$, then $M(u_2) = lu_2$ with $l \in D_L(u_2)$ (since M is H -special), and $M(w_2) = (lu_2) \cdot \bar{r}$. Now $w_2 \in \{\bar{l} \cdot u_2, u_2 \cdot \bar{r}\}$, but actually both possibilities are not permitted. On one hand, w_2 cannot be $u_2 \cdot \bar{r}$, since otherwise M and λ_l would agree on both u and w . On the other hand, $w_2 \neq \bar{l} \cdot u_2$ since otherwise $\bar{l} \cdot u_2 \cdot u_3 \triangleright M(\bar{l} \cdot u_2 \cdot u_3) = M(\bar{l} \cdot u_2) \cdot u_3 = lu_2 \cdot \bar{r} \cdot u_3 \notin W^H$, with $lu_2 \cdot u_3 \in W^H$, which is impossible since M is H -special.

If $u \triangleleft M(u)$, then $M(u_2) = w_2$ since M is a special matching. The element w_2 cannot be $l \cdot u_2$, with $l \notin D_L(u)$, since otherwise M and λ_l would agree on both u and w . Thus $w_2 = u_2 \cdot \bar{r}$ and

$$R_{u,w}^{H,x}(q) = (q - 1 - x)R_{u_3,w_3}^{H,x}(q)$$

by Lemma 3.6: on the other hand, $M(u) = w_2 \cdot u_3 = u_2 \cdot \bar{r} \cdot u_3 \notin W^H$ and

$$R_{u,M(w)}^{H,x}(q) = R_{u_2u_3,u_2w_3}^{H,x}(q) = R_{u_3,w_3}^{H,x}(q),$$

and the assertion follows.

Suppose $\ell(w_2) - \ell(u_2) = 0$.

If $u_2 = w_2$, then the result is clear. Otherwise $r \leq w_3$ and M commutes with ρ_r : we have $(M(w_2), M(u_2)) = (lw_2, lu_2)$, with $l \in D_L(w_2) \setminus D_L(u_2)$, and hence M coincides with λ_l on both u and w , which is a contradiction.

Suppose $\ell(w_2) - \ell(u_2) = -1$.

Necessarily $u_2 = w_2 \cdot r = w_0(s, t)$ and Lemma 2.6 implies $M(u_2) \triangleleft u_2$. Clearly, we also have $M(w_2) \triangleleft w_2$. Hence $R_{u,w}^{H,x}(q)$ and $R_{M(u),M(w)}^{H,x}(q)$ coincide, since they are both equal to $R_{ru_3,w_3}^{H,x}(q)$ by Lemma 3.6.

The proof is completed. \square

We illustrate Theorems 1.3, 1.4, and 1.5 with an example. Let W be the Coxeter group of type A_3 with Coxeter generators s_1, s_2 and s_3 numbered as usual (i.e. $m_{s_1,s_2} = m_{s_2,s_3} = 3$ and $m_{s_1,s_3} = 2$). Let $H = \{s_2\}$, $w = s_1s_2s_3s_1 \in W^H$, and $u = s_1 \in W^H$.

Suppose that we want to compute $R_{u,w}^{H,x}(q)$ but we only know the isomorphism class of the poset $[e, w]$ and which elements of $[e, w]$ belong to W^H and which do not. In other words, we know the pieces of information that we can detect from Figure 1, where the elements represented by full (respectively, empty) bullets belong to (respectively, do not belong to) W^H .

In order to compute $R_{u,w}^{H,x}(q)$ using Theorem 1.5, we need an H -special matching M of w . There are 3 of them: we choose, for instance, the dashed H -special matching depicted in the first picture in Figure 2. Hence

$$R_{u,w}^{H,x}(q) = (q - 1 - x)R_{u,M(w)}^{H,x}(q).$$




FIGURE 1. The Hasse diagram of $[e, w]$ and how $[e, w]^H$ embeds in $[e, w]$.

Now we need an H -special matching N of $M(w)$, and we choose the dashed H -special matching depicted in the second picture in Figure 2. Hence

$$R_{u, M(w)}^{H,x}(q) = qR_{N(u), NM(w)}^{H,x}(q) + (q-1)R_{u, NM(w)}^{H,x}(q) = (q-1)R_{u, NM(w)}^{H,x}(q).$$

Finally, we need an H -special matching of $NM(w)$, and we choose the dashed H -special matching depicted in the third picture in Figure 2. Hence

$$R_{u, NM(w)}^{H,x}(q) = qR_{O(u), ONM(w)}^{H,x}(q) + (q-1)R_{u, ONM(w)}^{H,x}(q) = (q-1)R_{u, ONM(w)}^{H,x}(q).$$

Since $u = ONM(w)$ we have $R_{u, ONM(w)}^{H,x}(q) = 1$, and the computation yields

$$R_{u,w}^{H,x}(q) = (q-1-x)(q-1)^2.$$




FIGURE 2. H -special matchings.

REFERENCES

- [1] A. Björner, F. Brenti, *Combinatorics of Coxeter Groups*, Graduate Texts in Mathematics **231**, Springer-Verlag, New York, 2005.
- [2] F. Brenti, *KazhdanLusztig and R-polynomials, Young's lattice, and Dyck partitions*, Pacific J. Math., **207** (2002), 257-286.

- [3] F. Brenti, *Kazhdan-Lusztig polynomials: history, problems, and Combinatorial Invariance*, Sémin. Lothar. Combin. **49** (2003), B49b, 30pp.
- [4] F. Brenti, *Parabolic KazhdanLusztig polynomials for Hermitian symmetric pairs*, Trans. Amer. Math. Soc., **361** (2009), 1703-1729.
- [5] F. Brenti, *A twisted duality for parabolic KazhdanLusztig R-polynomials*, J. Algebra, **477** (2017), 472-482.
- [6] F. Brenti, F. Caselli, M. Marietti, *Special Matchings and Kazhdan–Lusztig polynomials*, Advances in Math. **202** (2006), 555-601.
- [7] F. Brenti, P. Mongelli, P. Sentinelli, *Parabolic KazhdanLusztig polynomials for quasi-minuscule quotients*, Adv. in Appl. Math. **78** (2016), 27-55.
- [8] F. Caselli, M. Marietti, *Special matchings in Coxeter groups*, Europ. J. Combin. **61** (2017), 151-166.
- [9] F. Caselli, M. Marietti, *A simple characterization of special matchings in lower Bruhat intervals*, Discrete Math. **341** (2018), 851-862.
- [10] V. V. Deodhar, *Some characterizations of Bruhat ordering on a Coxeter group and determination of the relative Möbius function*, Invent. Math. **39** (1977), 187-198.
- [11] V. Deodhar, *On some geometric aspects of Bruhat orderings. II. The parabolic analogue of Kazhdan–Lusztig polynomials*, J. Algebra **111** (1987), 483-506.
- [12] F. Du Cloux, *Rigidity of Schubert closures and invariance of Kazhdan-Lusztig polynomials*, Advances in Math. **180** (2003), 146-175.
- [13] M. J. Dyer, *Hecke algebras and reflections in Coxeter groups*, Ph. D. Thesis, University of Sydney, 1987.
- [14] M. Dyer, *On the Bruhat graph of a Coxeter system*, Compositio Math. **78** (1991), 185-191.
- [15] B. Elias, G. Williamson, *The Hodge theory of Soergel bimodules*, Ann. Math. **180** (2014), 1089-1136.
- [16] A. van den Hombergh, *About the automorphisms of the Bruhat-ordering in a Coxeter group*, Indag. Math. **36** (1974), 125-131.
- [17] J. E. Humphreys, *Reflection Groups and Coxeter Groups*, Cambridge Studies in Advanced Mathematics, no.29, Cambridge Univ. Press, Cambridge, 1990.
- [18] M. Kashiwara, T. Tanisaki, *Parabolic Kazhdan–Lusztig polynomials and Schubert varieties*, J. Algebra **249** (2002), 306-325.
- [19] D. Kazhdan, G. Lusztig, *Representations of Coxeter groups and Hecke algebras*, Invent. Math. **53** (1979), 165-184.
- [20] D. Kazhdan, G. Lusztig, *Schubert varieties and Poincaré duality*, Geometry of the Laplace operator (Proc. Sympos. Pure Math., Univ. Hawaii, Honolulu, Hawaii, 1979), Proc. Sympos. Pure Math., XXXVI, Amer. Math. Soc., Providence, R.I. (1980), 185-203.
- [21] M. Marietti, *Special matchings and parabolic Kazhdan–Lusztig polynomials*, Trans. Amer. Math. Soc. **368** (2016), no. 7, 5247-5269.
- [22] R. P. Stanley, *Enumerative Combinatorics*, vol.1, Wadsworth and Brooks/Cole, Monterey, CA, 1986.
- [23] A. I. Telloni, *Special matchings calculate the parabolic Kazhdan–Lusztig polynomials of the universal Coxeter groups*, J. Algebra **465** (2016), 21-40.

DIPARTIMENTO DI INGEGNERIA INDUSTRIALE E SCIENZE MATEMATICHE, UNIVERSITÀ POLITECNICA DELLE MARCHE, VIA BRECCIE BIANCHE, 60131 ANCONA, ITALY

E-mail address: m.marietti@univpm.it