



UNIVERSITÀ POLITECNICA DELLE MARCHE
Repository ISTITUZIONALE

Benthic deep-sea fungi in submarine canyons of the Mediterranean Sea

This is a pre print version of the following article:

Original

Benthic deep-sea fungi in submarine canyons of the Mediterranean Sea / Barone, G.; Rastelli, E.; Corinaldesi, C.; Tangherlini, M.; Danovaro, R.; Dell'Anno, A.. - In: PROGRESS IN OCEANOGRAPHY. - ISSN 0079-6611. - 168:(2018), pp. 57-64.

Availability:

This version is available at: 11566/266051 since: 2022-05-31T16:50:39Z

Publisher:

Published

DOI:

Terms of use:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. The use of copyrighted works requires the consent of the rights' holder (author or publisher). Works made available under a Creative Commons license or a Publisher's custom-made license can be used according to the terms and conditions contained therein. See editor's website for further information and terms and conditions.

This item was downloaded from IRIS Università Politecnica delle Marche (<https://iris.univpm.it>). When citing, please refer to the published version.

(Article begins on next page)

Manuscript Details

Manuscript number	PROOCE_2018_117
Title	Benthic deep-sea fungi in submarine canyons of the Mediterranean Sea
Article type	Full Length Article

Abstract

Fungi are ubiquitous components of microbial assemblages in aquatic ecosystems, but their quantitative relevance, ecological role and diversity in benthic deep-sea ecosystems are still largely unknown. Here, we investigated patterns and drivers of benthic fungal abundance, biomass and diversity from 200 to 1000 m depth in three submarine canyons of the Mediterranean Sea (Tricase, Crotona and Squillace canyons). The Crotona and Squillace canyons, which are close to the coast and influenced by river inputs, showed significantly higher fungal abundance, biomass and diversity (as operational taxonomic units, OTUs) compared with the Tricase canyon that was far from the coast and without nearby estuaries. Fungal biomass, ranging from 0.17 to 5.78 $\mu\text{gC g}^{-1}$, and abundance increased with increasing carbohydrate concentrations in the sediments, suggesting that deep-sea fungi have a role in the utilisation of this component of the organic matter. A total of 1742 fungal OTUs, belonging to all fungal phyla known to date, were found and Ascomycota represented the dominant phylum. However, only 36% of the reads belonged to known genera. In particular, Tricase and Crotona canyons hosted the highest proportion of unknown fungal taxa, suggesting that deep-sea sediments can harbour a high number of novel fungal lineages. Our findings also reveal that fungal assemblage composition in the investigated canyons was influenced by trophic and thermo-haline conditions, which may promote a high turnover diversity of benthic deep-sea fungal assemblages. Overall results reported here indicate that the submarine canyons of the Mediterranean Sea can represent hot-spots of abundant and highly diversified fungal assemblages and pave the way for a better understanding of the ecological role of fungi in the largest ecosystem on Earth.

Keywords	Benthic deep-sea ecosystems; fungal abundance; fungal diversity; submarine canyons; Mediterranean Sea
Manuscript category	Biological Oceanography
Corresponding Author	Antonio Dell'Anno
Corresponding Author's Institution	Università Politecnica delle Marche
Order of Authors	Giulio Barone, Eugenio Rastelli, cinzia corinaldesi, Michael Tangherlini, Roberto Danovaro, Antonio Dell'Anno
Suggested reviewers	Gaëtan Burgaud, Jan Pawlowski, Takuro Nunoura, Lucia Bongiorno
Opposed reviewers	Virginia Edgcomb

Submission Files Included in this PDF

File Name [File Type]

Barone et alii_Cover Letter.doc [Cover Letter]

Barone et alii_highlights.doc [Highlights]

Barone_et alii_main text_final.doc [Manuscript File]

To view all the submission files, including those not included in the PDF, click on the manuscript title on your EVISE Homepage, then click 'Download zip file'.

Research Data Related to this Submission

There are no linked research data sets for this submission. The following reason is given:
All data are included in the text



UNIVERSITÀ
POLITECNICA
DELLE MARCHE

Dipartimento
di Scienze
della Vita
e dell'Ambiente
DISVA

Ancona, 08.05.2018

Dear Editor,

please find enclosed the main text and figures of the manuscript entitled: "*Benthic deep-sea fungi in submarine canyons of the Mediterranean Sea*" by Giulio Barone et alii, submitted for consideration to *Progress in Oceanography* within the special issue "**Ecology and functioning of Mediterranean submarine canyons**".

This is an original manuscript not submitted or presented elsewhere in which we investigated, for the first time, the quantitative relevance and diversity of fungi in deep-sea sediments of submarine canyons of the Mediterranean Sea. In this study, we show that fungi are an important component within the benthic deep-sea food webs and that they are highly diversified. Our results also indicate that different environmental characteristics encountered in the different canyons investigated can have a major role in influencing fungal diversity and assemblage composition. We think that this work expands our knowledge on the ecology and diversity of fungi inhabiting Mediterranean submarine canyons and we hope that it might be of interest for your journal.

For any requests, please do not hesitate to contact me at the address and numbers reported here below.

Looking forward to hearing from you soon, we remain.

Best regards

On behalf of all co-authors

Antonio Dell'Anno

Prof. Antonio Dell'Anno
Department of Life and Environmental Sciences (DISVA)
Università Politecnica delle Marche, 60131, Ancona, Italy.
Phone number: +39 0712204328
E-mail: a.dellanno@univpm.it

HIGHLIGHTS

- Submarine canyons host abundant and diverse fungal communities
- Fungal abundance, biomass and diversity are driven by carbohydrate concentrations
- Deep-sea sediments can harbour a high number of novel fungal taxa
- Thermohaline and trophic conditions may promote a high turnover diversity of fungi

Benthic deep-sea fungi in submarine canyons of the Mediterranean Sea

Giulio Barone¹, Eugenio Rastelli², Cinzia Corinaldesi³, Michael Tangherlini², Roberto Danovaro^{1,2}, Antonio Dell’Anno^{1*}

¹*Department of Life and Environmental Sciences, Polytechnic University of Marche, 60131 Ancona Italy*

²*Stazione Zoologica Anton Dohrn, Villa Comunale, 80121 Naples, Italy*

³*Department of Sciences and Engineering of Materials, Environment and Urbanistics, Polytechnic University of Marche, 60131 Ancona, Italy*

*Address for correspondence: a.dellanno@univpm.it

Running title: Fungal abundance and diversity in Mediterranean canyons

Abstract

Fungi are ubiquitous components of microbial assemblages in aquatic ecosystems, but their quantitative relevance, ecological role and diversity in benthic deep-sea ecosystems are still largely unknown. Here, we investigated patterns and drivers of benthic fungal abundance, biomass and diversity from 200 to 1000 m depth in three submarine canyons of the Mediterranean Sea (Tricase, Crotone and Squillace canyons). The Crotone and Squillace canyons, which are close to the coast and influenced by river inputs, showed significantly higher fungal abundance, biomass and diversity (as operational taxonomic units, OTUs) compared with the Tricase canyon that was far from the coast and without nearby estuaries. Fungal biomass, ranging from 0.17 to 5.78 $\mu\text{gC g}^{-1}$, and abundance increased with increasing carbohydrate concentrations in the sediments, suggesting that deep-sea fungi have a role in the utilisation of this component of the organic matter. A total of 1742 fungal OTUs, belonging to all fungal phyla known to date, were found and Ascomycota represented the dominant phylum. However, only 36% of the reads belonged to known genera. In particular, Tricase and Crotone canyons hosted the highest proportion of unknown fungal taxa, suggesting that deep-sea sediments can harbour a high number of novel fungal lineages. Our findings also reveal that fungal assemblage composition in the investigated canyons was influenced by trophic and thermo-haline conditions, which may promote a high turnover diversity of benthic deep-sea fungal assemblages. Overall results reported here indicate that the submarine canyons of the Mediterranean Sea can represent hot-spots of abundant and highly diversified fungal assemblages and pave the way for a better understanding of the ecological role of fungi in the largest ecosystem on Earth.

Key Words: Benthic deep-sea ecosystems, fungal abundance, fungal diversity, submarine canyons, Mediterranean Sea

1. Introduction

Deep-sea ecosystems represent more than 65% of the world's surface and >95% of the global biosphere (Herring, 2002), and host yet undiscovered biodiversity and a significant portion of the world's genetic diversity (Danovaro et al., 2017). In benthic deep-sea ecosystems, biomass is dominated by bacteria and archaea, followed by unicellular eukaryotes and small metazoans (<0.5 mm in size, meiofauna). These organisms are essential for carbon cycling and nutrient regeneration, and thus vital for sustaining oceanic production (Dell'Anno and Danovaro, 2005; Sogin et al., 2006; Jørgensen and Boetius, 2007; Danovaro et al., 2015; Danovaro et al., 2017). Recent findings, based on culture-dependent and independent approaches, revealed that fungi are present in deep-sea environments across a variety of ecosystem types spanning from hypersaline anoxic basins (Bernhard et al., 2014; Edgcomb et al., 2017) to cold seeps (Nagahama et al., 2011; Thaler et al., 2012), from hydrothermal vents (Burgaud et al., 2009; Burgaud et al., 2010; Xu et al., 2017) to surface and subsurface sediments (Orsi et al., 2013; Pachiadaki et al., 2016). Fungi have also been reported as the dominant unicellular eukaryotic group in the marine snow in bathypelagic waters with biomass similar to that of prokaryotes (Bochdansky et al., 2017).

Theoretical estimates suggest that fungi can be the most diversified component of unicellular eukaryotes on Earth, with more than 5 million species of which only 5% have been described (Hawksworth, 1997; Blackwell, 2011). This gap applies in particular to open ocean ecosystems where a significant fraction of fungal diversity is still unknown (Jeffries et al., 2016). Recent studies suggest that a variety of environmental factors (e.g. temperature, salinity, nutrients) can influence the diversity and assemblage composition of fungi in marine ecosystems (Li et al., 2016; Tisthammer et al., 2016). However, drivers

controlling the distribution and diversity of fungi in benthic deep-sea ecosystems remain to date largely unexplored.

Fungi in terrestrial and freshwater ecosystems are among the main decomposers of organic matter and play a key role in the processing of the most refractory fraction of organic carbon (Carlile et al. 2001; Clipson et al. 2006; Hwang et al. 2006; Dighton, 2007). Since deep-sea ecosystems can contain relatively high amounts of organic carbon (Pusceddu et al., 2009), fungi might play a key role in C cycling also in these ecosystems (Hyde et al., 1998; Burgaud et al., 2009; Cathrine and Raghukumar, 2009; Jebaraj et al., 2010).

In this study, we investigated the abundance, biomass and taxonomic composition of fungal assemblages along the continental margins of the Central Mediterranean Sea. Continental margins are characterised by open slopes and submarine canyons, which are essential for C cycling and nutrient regeneration processes at a global scale (Bousquet et al., 2000; Dickens, 2003). In particular, submarine canyons can channel large amounts of organic matter photosynthetically produced from the continental shelf down to deep-sea ecosystems (Monaco et al., 1999; Sánchez-Vidal et al., 2008; Allen and Durrieu de Madron, 2009; Puig et al., 2014). For this reason, we selected three submarine canyons characterised by different environmental conditions and investigated fungal abundance, biomass and diversity at depths ranging from 200 to 1000 m. To identify the factors potentially controlling their quantitative importance and diversity in deep-sea sediments, we explored the role of environmental conditions, including the organic matter quality and quantity.

2. Materials and methods

2.1. Study area and sampling design

Sediment sampling was carried out in the Ionian Sea (Central Mediterranean Sea) during the oceanographic cruise “SAND 2016” held on board of the research vessel R/V Minerva Uno in May 2016. Sediment samples were collected within the main axis of three canyons located along the SE Italian margin at 200, 500 and 1000 m depths (Figure 1). One of the investigated canyon (hereafter defined “Tricase”) located along the Apulian margin, is far from any continental freshwater inputs. The other two investigated canyons are located along the Calabrian margin and were close to river estuaries. The Northern canyon, extending for about 30 km, is located in front of the Crotone municipality (canyon “Crotone”) and its head is close to a river mouth. The head of the canyon “Squillace” is close to the coastline in front of the Squillace municipality and is characterised by the presence of sporadic, but intense river inputs. Sediment samples were collected at each benthic site by independent multiple corer deployments. The top 1 cm of each sediment sample was used for the analysis of the quantity and biochemical composition of organic matter, fungal abundance (based on q-PCR analysis of 18S rRNA genes), biomass and diversity. At each station, temperature and salinity of bottom waters were measured using CTD casts.

2.2. Quantity and biochemical composition of organic matter

The three major biochemical classes of organic compounds (proteins, carbohydrates and lipids) in deep-sea sediments were determined according to previously described procedures (Danovaro, 2010). Protein, carbohydrate and lipid concentrations were determined spectrophotometrically and expressed as albumin, glucose and tripalmitin equivalents, respectively. All analyses were carried out in 3 replicates. Protein, carbohydrate and lipid concentrations were then converted to carbon equivalents

(conversion factors: 0.49, 0.40 and 0.75 gC g⁻¹, respectively) to determine biopolymeric C content (Dell'Anno et al., 2002).

2.3. Fungal biomass

To detect and quantify fungi in the sediment samples, fluorescence in-situ hybridisation (FISH) coupled with Calcofluor white staining (which targets chitin, cellulose and carboxylated polysaccharides) have been used following procedures previously described (Bochdansky et al., 2016). The FISH reaction was performed using the Pan-Fungal probe PF2 (5'-CTC TGG CTT CAC CCT ATT C-3') Cy-3 labelled (Kempf et al., 2000). Briefly, about 1 g of sediment was first treated using 4 ml of a mix containing EDTA, Tween 80, sodium-pyrophosphate and methanol and ultrasounds treatment to separate fungi from the sediment matrix. After centrifugation, sediment samples were washed twice with PBS buffer and then treated with increasing concentrations of ethanol (50, 80 and 96%, for 3 min each). The sediment was then suspended in 500 µl hybridisation buffer containing 0.9 M NaCl, 0.01% w/v SDS, 20 mM Tris-HCl pH 7.2, 30 %v/v formamide and 1 µM PF2 (Kempf et al., 2000), then incubated for 3 h at 46°C in the dark. Samples were then transferred in sterile tubes containing pre-warmed washing buffer (20 mM Tris-HCl pH 8.0, 0.01% w/v SDS, 5 mM EDTA, 0.112M NaCl) and incubated for 30 minutes at 48°C. After centrifugation and resuspension of the sediment samples with 0.2 µm pre-filtered water, aliquots of the slurry (n=3) were filtered on 0.2 µm polycarbonate filters (Millipore) conditions. Filters were then stained with 0.5 mM Calcofluor white and incubated in the dark for 5 min. Subsequently, slides were washed with 0.02 µm pre-filtered water and analysed under epifluorescence microscopy. The whole filter was examined, and length and width measures were taken for each fungal-like structure. Then, the average width and cumulative length were converted to a

cylinder with half-spheres at ends, and the biovolume was converted into fungal biomass, assuming 1 μm^3 of fungal biovolume equivalent to 1 pg C (Damare and Raghukumar 2008).

2.4 DNA extraction and purification for molecular analysis

The DNA was extracted and purified from the sediment samples using the PowerSoil DNA isolation kit (QIAGEN)) following the manufacturer's instruction with slight modifications to remove extracellular DNA (based on three subsequent washing steps) before DNA extraction (Danovaro, 2009; Danovaro et al., 2016).

2.5 Quantitative real-time PCR of fungal 18S rRNA gene sequences

DNA extracted from two sediment samples collected at each study site by independent multiple corer deployments was used for quantitative real-time PCR (qPCR) analysis which was performed as described in Taylor et al. (2016) with slight modifications. Briefly, fungi-specific primers FR1 5'-AIC CAT TCA ATC GGT AIT-3' and FF390 5'-CGA TAA CGA ACG AGA CCT-3' (Prevost-Boure et al., 2011) were used with the Sensi-FAST SYBR Q-PCR kit (Bioline, London, UK). The 15 μl reactions contained 8 μl Sensi-FAST master mix, 1 μl of each primer (final concentration 1 μM), 1 μl of DNA template and 5 μl nuclease-free molecular-grade water (Taylor and Cunliffe, 2016). A Bio-Rad iQ5 was used to perform qPCR. The following qPCR thermal cycles were used: 94°C for 3min, then 40 cycles of 94 °C for 10 s, annealing at 50 °C for 15 s, elongation at 72°C for 20 s and acquisition of fluorescence data at 82°C. Standard curves were generated using known concentration of *Aspergillus niger* 18S rDNA.

2.6 Fungal diversity

DNA extracted from two sediment samples collected at each study site by independent multiple corer deployments was amplified using the primer set ITS1F (5'-GGAAGTAAAAGTCGTAACAAGG-3') and ITS2 (5'-GCTGCGTTCTTCATCGATGC-3') which amplify the internal transcribed spacer-1 (ITS1) region of the fungal rRNA gene (Walters et al., 2015). Amplicons were sequenced on an Illumina MiSeq platform by LGC group (Berlin, Germany) following Earth Microbiome Project protocols (<http://www.earthmicrobiome.org/emp-standard-protocols/>). Barcodes and ITS1 primer pairs were removed before demultiplexing. Paired-end sequences were then merged with FLASH (Magoč and Salzberg, 2011). Merged sequences were quality filtered using the SEARCH tool (Edgar, 2010) to remove sequences with expected error >1.0 and analysed with the QIIME software package (Caporaso et al., 2010). Operational taxonomic units (OTUs) were assigned with a threshold of 98.5% pairwise identity as indicated by the UNITE fungal ITS database (<http://unite.ut.ee/>). Then, OTUs were classified taxonomically against the UNITE database (<http://unite.ut.ee/>, Version 7.1, November 20, 2016). To allow a proper comparison among samples, we followed the approach by Gihring et al. (2012) with sample normalisation to 2500 randomly-selected sequences (corresponding to the lowest read count obtained in our samples). Rarefaction curves highlighted that 2500 sequences used for the comparison among all samples were generally sufficient to describe the fungal diversity in the different benthic deep-sea ecosystems investigated (Figure S1).

2.7 Statistical analyses

Two-way analysis of variance (ANOVA) was performed to test for differences in organic matter content, fungal abundance, biomass and OTU richness among canyons and depths. When significant differences were encountered, post-hoc tests were also carried out.

ANOSIM analysis was performed to test for the presence of statistical differences in the trophic conditions at the seafloor between canyons. Permutational multivariate analysis of variance (PERMANOVA) was used based on Bray-Curtis similarity matrix and visualised using cluster analysis to test for differences in fungal community composition among canyons and depths. Distance-based multivariate analysis for a linear model (DistLM) forward (Anderson, 2008) was performed to identify potential factors influencing fungal abundance, biomass, OTU richness and assemblage composition. P values were obtained with 9,999 permutations of residuals under the reduced model (Anderson, 2008). Temperature, salinity and trophic resources (as protein, carbohydrate and lipid concentrations) were used as predictor variables. Distance-based redundancy analysis (dbRDA) was finally used to visualise the relationships between fungal assemblage composition of the different canyon systems and thermo-haline and trophic variables. All statistical analyses were performed using Primer 6+ software.

3. Results and discussion

The thermo-haline conditions of bottom waters of the benthic systems investigated in the present study changed across depths and canyons, with temperature values ranging from 13.77 to 15.20 °C, and salinity values ranging from 38.75 to 38.93 (Table 1). Lowest temperature and salinity values were generally observed at the greatest depth (i.e. 1000 m). Also, the analysis of organic matter quantity in the sediments revealed differences among the investigated canyons (Tables 1, TableS1), with concentrations of proteins and carbohydrates significantly higher in Crotone and Squillace canyons than in Tricase canyon ($p < 0.05$ and $p < 0.01$, for proteins and carbohydrates, respectively). The highest organic matter content in the sediments of Crotone and Squillace canyons is likely due to

their proximity to the coast and the presence of nearby river inputs which amplify the magnitude of organic matter exported from the water column and settling on the seafloor (Lopez-Fernandez et al., 2013).

The amount of organic matter in deep-sea sediments represents a significant factor influencing the abundance and distribution of benthic assemblages (Danovaro et al., 2014). Fungal abundance, expressed as number of fungal 18S rDNA copies ranged from 1.4×10^6 to 5.1×10^7 copies g⁻¹ and was significantly lower in Tricase ($0.38 \pm 0.04 \times 10^7$ copies g⁻¹) than in Crotone and Squillace canyons (2.7 ± 0.5 and $1.3 \pm 0.4 \times 10^7$ copies g⁻¹, respectively; $p < 0.01$; Figure 2a). Our results fall within previously reported ranges for deep-sea sediments of the Pacific Ocean (3.5×10^6 - 5.2×10^7 28S rDNA copies g⁻¹; Xu et al., 2014), providing the first evidence of the quantitative importance of fungi also in benthic deep-sea ecosystems of the Mediterranean Sea. In all canyons, the 18S rDNA copy number changed significantly with water depth, with highest values at the shallowest depth in Crotone and Squillace canyons and at 500 m depth in Tricase canyons.

Fungal biomass ranged from 0.17 to 5.78 $\mu\text{gC g}^{-1}$, with values significantly lower in the sediments of Tricase ($0.63 \pm 0.14 \mu\text{gC g}^{-1}$) than in Crotone and Squillace canyons (2.40 ± 0.43 and $2.73 \pm 0.49 \mu\text{gC g}^{-1}$, respectively; $p < 0.01$) (Figure 2b). The distribution of fungal biomass along the bathymetric gradients within each canyon was similar to that of 18S rDNA copy number. Data on fungal biomass are practically no existent for deep-sea surface sediments (Damare and Raghukumar, 2008). However, the fungal biomass values reported here are similar to those of other benthic components reported at equal depths in the whole Mediterranean Sea (Gambi et al., 2017) suggesting that fungi can represent a significant component of benthic biomass in deep-sea sediments.

We found a significant relationship between fungal abundance and biomass (Figure S2).

From the slope of this relationship, we estimated that 1 µg of fungal biomass could be equivalent to 7.8×10^6 fungal 18S rDNA copies. Although such relationship should be view with caution and needs to be better refined with a broader spatial scale investigation, it can provide useful information on the quantitative relevance of deep-sea fungi based on copy number determinations (Taylor and Cunliffe, 2016).

Significant positive relationships between carbohydrate concentrations and fungal abundance and biomass were found ($r=0.715$ and $r=0.893$, both $p<0.01$, for the abundance and biomass, respectively; Figure 3). Also, multivariate multiple regression analysis provided evidence that carbohydrate concentration in the sediment was the primary factor explaining the distribution of the abundance and biomass of fungi in the benthic deep-sea ecosystems investigated (Table S2). Since fungi are osmotrophic (i.e. feed by secreting enzymes into the environment to degrade organic matter externally before taking the resulting metabolites into the cell; Richards and Talbot, 2013; Richards et al., 2015), our results suggest that they could be highly specialized in the utilisation of carbohydrates which are typically characterised by a highly recalcitrant fraction, especially in benthic deep-sea ecosystems (Dell'Anno et al., 2000; Dell'Anno et al., 2013).

Our results also show that the clustering of the 1203476 fungal ITS sequences (obtained after quality check) allowed us to identify a total of 1742 fungal OTUs, belonging to all fungal phyla known to date. Ascomycota represented the dominant phylum (accounting for 68% of the total reads), followed by Basidiomycota (10%) and Chytridiomycota (4%). The dominance of such phyla has been consistently reported in other benthic deep-sea ecosystems (Zhang et al., 2016).

The number of fungal OTUs we found in the sediments of the different canyons was similar compared with that reported in other deep-sea ecosystems (Zhang et al., 2016).

The Tricase canyon displayed a significantly lower OTU number (range: 64-71 OTUs) compared to Crotone and Squillace canyons (range: 113-325 and 173-221 OTUs, respectively; $p < 0.01$; Figure 4).

In our dataset, the OTUs affiliating to currently known fungal families were represented by only 19-38% of the total reads (Figure 5). The classified fungal OTUs affiliated to 206 genera belonging to 132 families, 66 orders and 27 classes.

At all benthic sites, Pleosporales was the most represented fungal order (accounting for ca. 20% of the total reads in each sample). This group is commonly present in marine environment and can account for a relevant fraction of the fungal diversity (up to 18% of all OTUs and sequences) in benthic deep-sea ecosystems (Li et al., 2016). Moreover, members belonging to the Pleosporales order are known to be adapted to high hydrostatic pressure (Nagano and Nagahama, 2012), possibly contributing to the ecological success of such taxon in deep-sea ecosystems.

Most of the fungi that we successfully classified were affiliated to genera such as *Aspergillus*, *Penicillium*, *Epicoccum*, *Cryptococcus* and *Candida* previously encountered in other deep-sea environments (Nagahama et al., 2003; Edgcomb et al., 2011; Rédou et al., 2014). However, these genera represented overall only ca. 36% of the total reads, indicating that the majority of fungal taxa belonged to genera not represented in UNITE database (Kõljalg et al., 2013).

The majority of fungal OTUs were unclassified below the order level and overall represented up to 69% of the total sequences. The quantitative relevance of unclassified sequences in our study was much higher than that reported for coastal sediments (Picard, 2017), indicating that deep-sea ecosystems might harbour a higher richness of novel fungal lineages compared with shallow benthic ecosystems.

The composition of fungal assemblage in the sediments of the Tricase canyon was significantly different ($p < 0.01$) from that of the other canyons, which otherwise showed no significant differences (Figure 5). These results suggest that submarine canyons far from the coastline and lacking river inputs can host distinct fungal assemblages from those close to river estuaries.

The analysis of the turnover (β -)diversity highlighted that the similarity of the fungal assemblage composition among different sites was very low (Table S3 and Figure 6). Indeed, the within-canyon similarity (i.e., the similarity of fungal assemblage composition among samples collected at a different depth within the same canyon) was on average 11%, while the inter-canyon comparisons resulted in an average similarity of 7% (Table S3). Moreover, the Tricase canyon showed the highest percentage of unique OTUs (i.e., OTUs found in Tricase but not in Squillace nor Crotone canyons; Table S4). Overall, the three canyons shared only 46 out of 1742 OTUs, that cumulatively accounted for only 22% of the total sequences. Twenty-seven of these 46 shared OTUs (overall accounting for 14% of the total sequences) were not classified, while the others shared OTUs (each of them contributing for $\leq 0.45\%$ of the total sequences) included taxa belonging to *Epicoccum nigrum*, *Illyonectria robusta*, *Trichoderma bissettii*, *Cryptococcus victoriae*, *Aspergillus sydowii*, *Fusarium sp.*, *Penicillium halotolearns* and *Thermomyces lanuginosus*.

Distance-based redundancy analysis highlighted that the fungal assemblage composition in the sediments of the different canyons was related to an array of factors including organic matter content (as carbohydrates and lipid concentrations, $r = -0.624$ and $r = 0.434$, respectively) and temperature ($r = 0.980$) and salinity ($r = -0.560$; Figure 7). These results confirm that also in the deep-sea sediments investigated trophic availability and thermo-haline conditions are important drivers of fungal assemblage composition (Hanson et al.,

2008; McGuire et al., 2010; Li et al., 2016; Taylor and Cunliffe, 2016; Tisthammer et al., 2016). Our findings also suggest that changes in the thermo-haline and trophic conditions among submarine canyons may promote a high turnover diversity of benthic deep-sea fungal assemblages.

Overall results of the present study indicate that the submarine canyons of the Mediterranean Sea host abundant and highly diversified fungal assemblages most of which still unidentified and pave the way for a better understanding of the ecological role of fungi in the largest ecosystem on Earth.

Acknowledgments: This study has been conducted in the framework of the National Flag Project RITMARE (Marine Italian Research, www.ritmare.it) and supported by the EU H2020 MERCES (Marine Ecosystem Restoration in Changing European Seas) project (Grant Agreement No. 689518) and DG ENV project IDEM (Implementation of the MSFD to the Deep Mediterranean Sea; contract EU No 11.0661/2017/750680/SUB/EN V.C2).

Author Contributions: R.D., C.C., and A.D. conceived the study. G.B. participated in the oceanographic cruise for collecting sediment samples and performed laboratory analyses. G.B., E.R., M.T. and A.D. contributed to data elaboration and interpretation. G.B., E.R., and A.D. wrote the first draft of the manuscript. All authors contributed to results discussion and finalization of the manuscript.

Conflict of interest: All the other authors declare no competing financial interests.

References

- Allen, S.E., Durrieu de Madron, X., 2009. A review of the role of submarine canyons in deep-ocean exchange with the shelf. *Ocean Science* 5, 607–620. doi:10.5194/os-5-607-2009
- Anderson, D.R., 2008. Model based inference in the life sciences: A primer on evidence, *Model Based Inference in the Life Sciences: A Primer on Evidence*. Springer New York, New York, NY. doi:10.1007/978-0-387-74075-1
- Bernhard, J.M., Kormas, K., Pachiadaki, M.G., Rocke, E., Beaudoin, D.J., Morrison, C., Visscher, P.T., Cobban, A., Starczak, V.R., Edgcomb, V.P., 2014. Benthic protists and fungi of Mediterranean deep hypersaline anoxic basin redoxcline sediments. *Frontiers in Microbiology* 5, 360. doi:10.3389/fmicb.2014.00605
- Blackwell, M., 2011. The Fungi: 1, 2, 3 ... 5.1 million species? *American Journal of Botany* 98, 426–438. doi:10.3732/ajb.1000298
- Bochdansky, A.B., Clouse, M.A., Herndl, G.J., 2016. Eukaryotic microbes, principally fungi and labyrinthulomycetes, dominate biomass on bathypelagic marine snow. *The ISME Journal* 11, 362–373. doi:10.1038/ismej.2016.113
- Bousquet, P., Peylin, P., Ciais, P., Le Quere, C., Friedlingstein, P., Tans, P.P., 2000. Regional changes in carbon dioxide fluxes of land and oceans since 1980. *Science* 290, 1342–1346. doi:10.1126/science.290.5495.1342
- Burgaud, G., Arzur, D., Durand, L., Cambon-Bonavita, M.-A., Barbier, G., 2010. Marine culturable yeasts in deep-sea hydrothermal vents: species richness and association with fauna. *FEMS Microbiology Ecology* 73, 121–133. doi:10.1111/j.1574-6941.2010.00881.x

- Burgaud, G., Le Calvez, T., Arzur, D., Vandenkoornhuyse, P., Barbier, G., 2009. Diversity of culturable marine filamentous fungi from deep-sea hydrothermal vents. *Environmental Microbiology* 11, 1588–1600. doi:10.1111/j.1462-2920.2009.01886.x
- Caporaso, J.G., Kuczynski, J., Stombaugh, J., Bittinger, K., Bushman, F.D., Costello, E.K., Fierer, N., Peña, A.G., Goodrich, J.K., Gordon, J.I., Huttley, G.A., Kelley, S.T., Knights, D., Koenig, J.E., Ley, R.E., Lozupone, C.A., McDonald, D., Muegge, B.D., Pirrung, M., Reeder, J., Sevinsky, J.R., Turnbaugh, P.J., Walters, W.A., Widmann, J., Yatsunenko, T., Zaneveld, J., Knight, R., 2010. QIIME allows analysis of high-throughput community sequencing data. *Nature Methods* 7, 335–336. doi:10.1038/nmeth.f.303
- Carlile MJ, Watkinson SC, Gooday GW (2001) *The fungi*, 2nd edn. Academic Press, San Diego
- Cathrine, S.J., Raghukumar, C., 2009. Anaerobic denitrification in fungi from the coastal marine sediments off Goa, India. *Mycological Research* 113, 100–109. doi:10.1016/j.mycres.2008.08.009
- Clipson, N., Otte, M., Landy E., 2006. Biogeochemical roles of fungi in the marine and estuarine habitats. In: Gadd GM (ed) *Fungi in biogeochemical cycles*. Cambridge university press, New York, pp 436–461
- Damare, S., Raghukumar, C., 2008. Fungi and macroaggregation in deep-sea sediments. *Microb Ecol* 56, 168–177. doi:10.1007/s00248-007-9334-y
- Danovaro, R., 2009. *Methods for the Study of Deep-Sea Sediments, Their Functioning and Biodiversity*. CRC Press.
- Danovaro, R., Carugati, L., Berzano, M., Cahill, A.E., Carvalho, S., Chenuil, A., Corinaldesi, C., Cristina, S., David, R., Dell’Anno, A., Dzhenbekova, N., Garcés, E., Gasol, J.M., Goela, P., Féral, J.-P., Ferrera, I., Forster, R.M., Kurekin, A.A., Rastelli, E., Marinova, V., Miller, P.I., Moncheva, S., Newton, A., Pearman, J.K., Pitois, S.G., Reñé, A.,

- Rodríguez-Ezpeleta, N., Saggiomo, V., Simis, S.G.H., Stefanova, K., Wilson, C., Martire, Lo, M., Greco, S., Cochrane, S.K.J., Mangoni, O., Borja, A., 2016. Implementing and Innovating Marine Monitoring Approaches for Assessing Marine Environmental Status. *Frontier in Marine Science* 3, 233–25. doi:10.3389/fmars.2016.00213
- Danovaro, R., Corinaldesi, C., Dell'Anno, A., Snelgrove, P.V.R., 2017. The deep-sea under global change. *Curr. Biol.* 27, R461–R465. doi:10.1016/j.cub.2017.02.046
- Danovaro, R., Corinaldesi, C., Rastelli, E., Dell'Anno, A., 2015. Towards a better quantitative assessment of the relevance of deep-sea viruses, Bacteria and Archaea in the functioning of the ocean seafloor. *Aquatic Microbial Ecology*. 75, 81–90. doi:10.3354/ame01747
- Danovaro, R., Snelgrove, P.V.R., Tyler, P., 2014. Challenging the paradigms of deep-sea ecology. *Trends in Ecology & Evolution* 29, 465–475. doi:10.1016/j.tree.2014.06.002
- Dell'Anno, A., Fabiano, M., Mei, M.L., Danovaro, R., 2000. Enzymatically hydrolysed protein and carbohydrate pools in deep-sea sediments: estimates of the potentially bioavailable fraction and methodological considerations. *Marine Ecology Progress Series* 196, 15–23. doi:10.3354/meps196015
- Dell'Anno, A., Mei, M.L., Pusceddu, A., Danovaro, R., 2002. Assessing the trophic state and eutrophication of coastal marine systems: a new approach based on the biochemical composition of sediment organic matter. *Marine Pollution Bulletin* 44, 611–622.
- Dell'Anno, A., Danovaro, R., 2005. Ecology: Extracellular DNA plays a key role in deep-sea ecosystem functioning. *Science* 309, 2179. doi:10.1126/science.1117475
- Dell'Anno, A., Pusceddu, A., Corinaldesi, C., Canals, M., Heussner, S., Thomsen, L., Danovaro, R., 2013. Trophic state of benthic deep-sea ecosystems from two different continental margins off Iberia. *Biogeosciences*, 10, 2945–2957. doi:10.5194/bg-10-2945-2013

- Dickens, G.R., 2003. Rethinking the global carbon cycle with a large, dynamic and microbially mediated gas hydrate capacitor. *Earth and Planetary Science Letters* 213, 169–183. doi:10.1016/S0012-821X(03)00325-X
- Dighton J., 2007. Nutrient cycling by saprotrophic fungi in terrestrial habitats. In: Kubicek CP, Druzhinina IS (eds) *The Mycota IV, environmental and microbial relationships*, 2nd edn. Springer, Berlin, pp 287–300
- Edgar, R.C., 2010. Search and clustering orders of magnitude faster than BLAST. *Bioinformatics* 26, 2460–2461. doi:10.1093/bioinformatics/btq461
- Edgcomb, V.P., Beaudoin, D., Gast, R., Biddle, J.F., Teske, A., 2011. Marine subsurface eukaryotes: the fungal majority. *Environmental Microbiology* 13, 172–183. doi:10.1111/j.1462-2920.2010.02318.x
- Edgcomb, V.P., Pachiadaki, M.G., Mara, P., Kormas, K.A., Leadbetter, E.R., Bernhard, J.M., 2017. Gene expression profiling of microbial activities and interactions in sediments under haloclines of E. Mediterranean deep hypersaline anoxic basins. *The ISME Journal* 1–15. doi:10.1038/ismej.2016.58
- Fernandez-Arcaya, U., Ramirez-Llodra, E., Aguzzi, J., Allcock, A.L., Davies, J.S., Dissanayake, A., Harris, P., Howell, K., Huvenne, V.A.I., Macmillan-Lawler, M., Martín, J., Menot, L., Nizinski, M., Puig, P., Rowden, A.A., Sanchez, F., Van den Beld, I.M.J., 2017. Ecological Role of Submarine Canyons and Need for Canyon Conservation: A Review. *Frontiers in Marine Science*. 4, 69–26. doi:10.3389/fmars.2017.00005
- Gihring, T.M., Green, S.J., Schadt, C.W., 2012. Massively parallel rRNA gene sequencing exacerbates the potential for biased community diversity comparisons due to variable library sizes. *Environmental Microbiology* 14, 285–290. doi:10.1111/j.1462-2920.2011.02550.x

- Hanson, C.A., Allison, S.D., Bradford, M.A., Wallenstein, M.D., Treseder, K.K., 2008. Fungal Taxa Target Different Carbon Sources in Forest Soil. *Ecosystems* 11, 1157–1167. doi:10.1007/s10021-008-9186-4
- Hawksworth, D.L., 1997. The fascination of fungi: Exploring fungal diversity. *Mycologist* 11, 18–22. doi:10.1016/S0269-915X(97)80062-6
- Hwang, J., Druffel, E.R.M., Bauer, J.E., 2006. Incorporation of aged dissolved organic carbon (DOC) by oceanic particulate organic carbon (POC): An experimental approach using natural carbon isotopes. *Marine Chemistry* 98, 315–322. doi:10.1016/j.marchem.2005.10.008
- Hyde, K.D., Jones, E.B.G., Leañó, E., Pointing, S.B., Poonyth, A.D., Vrijmoed, L.L.P., 1998. Role of fungi in marine ecosystems. *Biodiversity and Conservation* 7, 1147–1161. doi:10.1023/A:1008823515157
- Jebaraj, C.S., Raghukumar, C., Behnke, A., Stoeck, T., 2010. Fungal diversity in oxygen-depleted regions of the Arabian Sea revealed by targeted environmental sequencing combined with cultivation. *FEMS Microbiology Ecology* 71, 399–412. doi:10.1111/j.1574-6941.2009.00804.x
- Jeffries, T.C., Curlevski, N.J., Brown, M.V., Harrison, D.P., Doblin, M.A., Petrou, K., Ralph, P.J., Seymour, J.R., 2016. Partitioning of fungal assemblages across different marine habitats. *Environmental Microbiology Reports* 8, 235–238. doi:10.1111/1758-2229.12373
- Jørgensen, B.B., Boetius, A., 2007. Feast and famine — microbial life in the deep-sea bed. *Nature Reviews Microbiology* 5, 770–781. doi:10.1038/nrmicro1745
- Kempf, V.A.J., Trebesius, K., Autenrieth, I.B., 2000. Fluorescent in situ hybridization allows rapid identification of microorganisms in blood cultures. *Journal of Clinical Microbiology* 38, 830–838.

- Kõljalg, U., Nilsson, R.H., Abarenkov, K., Tedersoo, L., Taylor, A.F.S., Bahram, M., Bates, S.T., Bruns, T.D., Bengtsson-Palme, J., Callaghan, T.M., Douglas, B., Drenkhan, T., Eberhardt, U., Dueñas, M., Grebenc, T., Griffith, G.W., Hartmann, M., Kirk, P.M., Kohout, P., Larsson, E., Lindahl, B.D., Lücking, R., Martín, M.P., Matheny, P.B., Nguyen, N.H., Niskanen, T., Oja, J., Peay, K.G., Peintner, U., Peterson, M., Põldmaa, K., Saag, L., Saar, I., Schübler, A., Scott, J.A., Senés, C., Smith, M.E., Suija, A., Taylor, D.L., Telleria, M.T., Weiss, M., Larsson, K.-H., 2013. Towards a unified paradigm for sequence-based identification of fungi. *Molecular Ecology* 22, 5271–5277. doi:10.1111/mec.12481
- Li, W., Wang, M.M., Wang, X.G., Cheng, X.L., Guo, J.J., Bian, X.M., Cai, L., 2016. Fungal communities in sediments of subtropical Chinese seas as estimated by DNA metabarcoding. *Nature Publishing Group* 1–9. doi:10.1038/srep26528
- Lopez-Fernandez, P., Calafat, A., Sanchez-Vidal, A., Canals, M., Flexas, M.M., Cateura, J., Joan B. Company, 2013. Multiple drivers of particle fluxes in the Blanes submarine canyon and southern open slope: Results of a year round experiment. *Progress in Oceanography* 118, 95–107. doi:10.1016/j.pocean.2013.07.029
- Magoč, T., Salzberg, S.L., 2011. FLASH: fast length adjustment of short reads to improve genome assemblies. *Bioinformatics* 27, 2957–2963. doi:10.1093/bioinformatics/btr507
- McGuire, K.L., Bent, E., Borneman, J., Majumder, A., Allison, S.D., Treseder, K.K., 2010. Functional diversity in resource use by fungi. *Ecology* 91, 2324–2332. doi:10.1890/09-0654.1
- Monaco, A., Durrieu de Madron, X., Radakovitch, O., Heussner, S., Carbonne, J., 1999. Origin and variability of downward biogeochemical fluxes on the Rhone continental margin (NW mediterranean). *Deep-Sea Research Part I* 46, 1483–1511. doi:10.1016/S0967-0637(99)00014-X

- 492 Nagahama, T., Hamamoto, M., Nakase, T., Takaki, Y., Horikoshi, K., 2003. *Cryptococcus*
493 *surugaensis* sp. nov., a novel yeast species from sediment collected on the deep-sea floor
494 of Suruga Bay. *International Journal of Systematic and Evolutionary Microbiology* 53,
495 2095–2098. doi:10.1099/ijs.0.02712-0
- 496 Nagahama, T., Takahashi, E., Nagano, Y., Abdel-Wahab, M.A., Miyazaki, M., 2011.
497 Molecular evidence that deep-branching fungi are major fungal components in deep-sea
498 methane cold-seep sediments. *Environmental Microbiology* 13, 2359–2370.
499 doi:10.1111/j.1462-2920.2011.02507.x
- 500 Nagano, Y., Nagahama, T., 2012. Fungal diversity in deep-sea extreme environments. *Fungal*
501 *Ecology* 5, 463–471. doi:10.1016/j.funeco.2012.01.004
- 502 Orsi, W., Biddle, J.F., Edgcomb, V.P., 2013. Deep Sequencing of Subseafloor Eukaryotic
503 rRNA Reveals Active Fungi across Marine Subsurface Provinces. *PLoS ONE* 8, e56335–
504 10. doi:10.1371/journal.pone.0056335
- 505 Pachiadaki, M.G., Rédou, V., Beaudoin, D.J., Burgaud, G., Edgcomb, V.P., 2016. Fungal and
506 Prokaryotic Activities in the Marine Subsurface Biosphere at Peru Margin and
507 Canterbury Basin Inferred from RNA-Based Analyses and Microscopy. *Frontiers in*
508 *Microbiology* 7, 364–17. doi:10.3389/fmicb.2016.00846
- 509 Puig, P., Palanques, A., Martín, J., 2014. Contemporary Sediment-Transport Processes in
510 Submarine Canyons. *Annual Review of Marine Science* 6, 53–77. doi:10.1146/annurev-
511 marine-010213-135037
- 512 Pusceddu, A., Dell’Anno, A., Fabiano M., Danovaro R., 2009. Quantity and bioavailability of
513 sediment organic matter as signatures of benthic trophic status. *Marine Ecology Progress*
514 *Series*, 375, 41–52. doi:10.3354/meps07735
- 515 Rédou, V., Ciobanu, M.-C., Pachiadaki, M.G., Edgcomb, V.P., Alain, K., Barbier, G.,
516 Burgaud, G., 2014. In-depth analyses of deep subsurface sediments using 454-

pyrosequencing reveals a reservoir of buried fungal communities at record-breaking depths. *FEMS Microbiology Ecology* 90, 908–921. doi:10.1111/1574-6941.12447

Richards, T.A., Leonard, G., Mahé, F., del Campo, J., Romac, S., Jones, M.D.M., Maguire, F., Dunthorn, M., de Vargas, C., Massana, R., Chambouvet, A., 2015. Molecular diversity and distribution of marine fungi across 130 European environmental samples. *Proc. R. Soc. B* 282, 20152243–10. doi:10.1098/rspb.2015.2243

Richards, T.A., Talbot, N.J., 2013. Horizontal gene transfer in osmotrophs: playing with public goods. *Nature Reviews Microbiology* 11, 720–727. doi:10.1038/nrmicro3108

Sánchez-Vidal, A., Pasqual, C., Kerhervé, P., Calafat, A., Heussner, S., Palanques, A., Durrieu de Madron, X., Canals, M., Puig, P., 2008. Impact of dense shelf water cascading on the transfer of organic matter to the deep western Mediterranean basin. *Geophysical Research Letters* 35, 117–125. doi:10.1029/2007GL032825

Sogin, M.L., Morrison, H.G., Huber, J.A., Welch, D.M., Huse, S.M., Neal, P.R., Arrieta, J.M., Herndl, G.J., 2006. Microbial diversity in the deep sea and the underexplored “rare biosphere.” *Proceedings of the National Academy of Sciences of the United States of America* 103, 12115–12120. doi:10.1073/pnas.0605127103

Taylor, J.D., Cunliffe, M., 2016. Multi-year assessment of coastal planktonic fungi reveals environmental drivers of diversity and abundance. *The ISME Journal* 10, 2118–2128. doi:10.1038/ismej.2016.24

Thaler, A.D., Van Dover, C.L., Vilgalys, R., 2012. Ascomycete phylotypes recovered from a Gulf of Mexico methane seep are identical to an uncultured deep-sea fungal clade from the Pacific. *Fungal Ecology* 5, 270–273. doi:10.1016/j.funeco.2011.07.002

Tisthammer, K.H., Cobian, G.M., Amend, A.S., 2016. Global biogeography of marine fungi is shaped by the environment. *Fungal Ecology* 19, 39–46. doi:10.1016/j.funeco.2015.09.003

- Walters, W., Hyde, E.R., Berg-Lyons, D., Ackermann, G., Humphrey, G., Parada, A.,
Gilbert, J.A., Jansson, J.K., Caporaso, J.G., Fuhrman, J.A., Apprill, A., Knight, R., 2015.
Improved Bacterial 16S rRNA Gene (V4 and V4-5) and Fungal Internal Transcribed
Spacer Marker Gene Primers for Microbial Community Surveys. *mSystems* 1, e00009–
15–11. doi:10.1128/mSystems.00009-15
- Xu, W., Guo, S., Pang, K.-L., Luo, Z.-H., 2017. Fungi associated with chimney and sulfide
samples from a South Mid-Atlantic Ridge hydrothermal site: Distribution, diversity and
abundance. *Deep-Sea Research Part I* 123, 48–55. doi:10.1016/j.dsr.2017.03.004
- Zhang, X.-Y., Wang, G.-H., Xu, X.-Y., Nong, X.-H., Wang, J., Amin, M., Qi, S.-H., 2016.
Exploring fungal diversity in deep-sea sediments from Okinawa Trough using high-
throughput Illumina sequencing. *Deep-Sea Research Part I* 116, 99–105.
doi:10.1016/j.dsr.2016.08.004

Captions of figures

- Figure 1.** Study area and sampling location (a). Details of benthic sites investigated within Tricase (a), Crotone (b) and Squillace (c) canyons. Bathymetry has been obtained from EMODnet (<http://portal.emodnet-bathymetry.eu>). Maps elaborated with QGIS.
- Figure 2.** Fungal abundance, expressed as 18S rDNA copy number (a), and biomass (b) in the different benthic sites of the Tricase, Crotone and Squillace canyons. Mean values and standard deviations are reported.
- Figure 3.** Relationships between carbohydrate concentrations in the sediments of the different canyons investigated and fungal abundance (a) and biomass (b)
- Figure 4.** OTU number in the different benthic sites within Tricase, Crotone and Squillace canyons. Mean values and standard deviations are reported.
- Figure 5.** Taxonomic composition (at the family level on data normalized to 2500 sequences) of the benthic fungal assemblages in the different canyons investigated. To better visualise differences among the investigated sites the output of cluster analysis is also reported.
- Figure 6.** Network visualisation based on the output of SIMPER analysis carried out on fungal community composition among the nine sites investigated. Line width is proportional to similarity values.
- Figure 7.** Output of the distance-based redundancy analysis (dbRDA) carried out on fungal community composition in the different benthic deep-sea sites in relation with thermo-haline and trophic conditions.

Table 1. Temperature, salinity and protein (PRT), carbohydrate (CHO), lipid (LIP) and biopolymeric C concentrations in the different sites of the Tricase, Crotone and Squillace canyons. Mean values and standard deviations (\pm) are reported.

Canyon	Water depth m	Temperature °C	Salinity	PRT mg g ⁻¹	CHO mg g ⁻¹	LIP mg g ⁻¹	Biopolymeric C mg g ⁻¹
Tricase	200 m	14.58±0.01	38.8±0.01	1.91±0.55	2.21±0.59	0.82±0.28	2.43±0.72
	500 m	14.23±0.05	38.75±0.01	2.42±0.73	2.37±0.29	1.22±0.4	3.05±0.78
	1000 m	13.85±0.01	38.8±0.01	0.77±0.52	2.1±0.25	1.72±0.68	2.51±0.87
Crotone	200 m	15.07±0.12	38.91±0.01	2.87±0.24	3.56±0.23	1.61±0.76	4.04±0.78
	500 m	14.4±0.03	38.88±0.01	2.09±0.48	2.44±0.23	0.48±0.19	2.36±0.47
	1000 m	13.77±0.02	38.76±0.01	2.22±0.29	2.18±0.11	0.3±0.1	2.19±0.26
Squillace	200 m	14.78±0.06	38.82±0.01	2.21±0.36	3.77±0.59	0.6±0.31	3.04±0.64
	500 m	14.64±0.05	38.92±0.01	3.5±0.78	3.08±0.22	0.28±0.05	3.16±0.5
	1000 m	13.78±0.01	38.76±0.01	2.96±0.34	2.61±0.21	0.66±0.58	2.99±0.68




Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6




Figure 7

1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980

Supplementary materials

666

667

668 **Benthic deep-sea fungi in submarine canyons of the Mediterranean Sea**

669

670 Giulio Barone, Eugenio Rastelli, Cinzia Corinaldesi, Michael Tangherlini, Roberto

671 Danovaro, Antonio Dell’Anno

672

673

674

675

676

677 Supplementary table S1-S4

678

679 Supplementary figures S1 and S2

Table S1. Reported are the outputs of the ANOSIM and SIMPER analyses carried out to test for the differences and dissimilarity in sediment organic matter contents between the different canyons investigated and the variables responsible for the estimated differences. Reported are R which represents the sample statistic (global R) and P which is the probability level. **=P <0.01; ns = not significant

	ANOSIM		Dissimilarity	Explanatory variable	SIMPER	
	R	P			Explained variance (%)	Cumulative explained variance (%)
Tricase vs. Crotone	0.153	**	22.35	Proteins	40.78	40.78
				Lipids	33.48	74.26
				Carbohydrates	25.74	100
Tricase vs. Squillace	0.449	**	27.02	Proteins	43.85	43.85
				Carbohydrates	30.74	74.59
				Lipids	25.41	100
Crotone vs. Squillace	0.12	n.s.	16.82	Proteins	n.s	n.s
				Lipids	n.s	n.s
				Carbohydrates	n.s	n.s

Table S2. Output of the multivariate multiple regression analysis carried out for testing the effects of organic matter content (proteins, carbohydrates and lipids), temperature and salinity on fungal abundance (as 18S rDNA copies) and biomass. Reported are Pseudo-F and P values (*<0.05; **<0.01; ***<0.001; ns>0.05) and the cumulative variance explained by the significant variables.

Fungal abundance (18S rDNA copies)			
Variable	Pseudo-F	P	Cumulative variance %
Carbohydrates	11.556	***	31.6
Lipids	3.814	*	41.0
Proteins	1.771	ns	-
Salinity	1.654	ns	-
Temperature	0.667	ns	-
Fungal biomass			
Carbohydrates	98.421	***	79.7
Lipids	4.249	ns	-
Proteins	2.275	ns	-
Temperature	0.82	ns	-
Salinity	2.196	ns	-

Table S3. Output of SIMPER showing the dissimilarity (turnover diversity) of fungal assemblage composition within the canyon and between the canyons investigated

Type of comparison			Turnover diversity (% Bray-Curtis dissimilarity)
within canyon	Tricase	200 m vs. 500 m	86.19
		200 m vs. 1000 m	91.97
		500 m vs. 1000 m	91.12
	Crotone	200 m vs. 500 m	89.03
		200 m vs. 1000 m	94.3
		500 m vs. 1000 m	87.88
	Squillace	200 m vs. 500 m	85.22
		200 m vs. 1000 m	88.92
		500 m vs. 1000 m	88.45
between canyons	200 m	Tricase vs. Crotone	97.01
		Tricase vs. Squillace	94.23
		Crotone vs. Squillace	88.7
	500 m	Tricase vs. Crotone	91.5
		Tricase vs. Squillace	95.42
		Crotone vs. Squillace	88.89
	1000 m	Tricase vs. Crotone	92.2
		Tricase vs. Squillace	94.52
		Crotone vs. Squillace	90.65

Table S4. Percentage of unique and shared OTUs between replicates of the same site, within the canyon and between the canyons

Type of comparison			Shared %	Unique %
between replicates of the same site	Tricase	200 m	9.4	90.6
		500 m	15.7	84.3
		1000 m	10.2	89.8
	Crotone	200 m	12.5	87.5
		500 m	14.6	85.4
		1000 m	7.6	92.4
	Squillace	200 m	12.2	87.8
		500 m	7.0	93.0
		1000 m	6.8	93.2
	Average		10.7	89.3
within canyon	Tricase	200 vs. 500 m	18.8	90.0
		200 vs. 1000 m	14.0	93.9
		500 vs. 1000 m	13.8	92.5
	Crotone	200 vs. 500 m	12.2	91.1
		200 vs. 1000 m	19.8	94.8
		500 vs. 1000 m	19.9	91.2
	Squillace	200 vs. 500 m	29.3	86.3
		200 vs. 1000 m	18.6	89.2
		500 vs 1000 m	26.9	88.0
	Average		19.2	90.8
between canyons	Tricase vs. Crotone	200 m	3.8	96.2
		500 m	7.9	92.1
		1000 m	7.6	92.4
	Tricase vs. Squillace	200 m	6.6	93.4
		500 m	5.6	94.4
		1000 m	5.9	94.1
	Crotone vs. Squillace	200 m	10.3	89.7
		500 m	10.9	89.1
		1000 m	8.1	91.9
	Average		7.4	92.6

Figure S1. Rarefaction curves calculated for each of the two independent replicates (dashed lines, 2500 sequences each) analysed in all benthic deep-sea sites of the canyons investigated.




Figure S2. Relationship between benthic fungal abundance (as 18S rDNA copies) and biomass in the sediments of the three canyons

