



UNIVERSITÀ POLITECNICA DELLE MARCHE  
Repository ISTITUZIONALE

Black Soldier Fly (*Hermetia illucens*) reared on roasted coffee by-product and *Schizochytrium* sp. as a sustainable terrestrial ingredient for aquafeeds production

This is the peer reviewed version of the following article:

*Original*

Black Soldier Fly (*Hermetia illucens*) reared on roasted coffee by-product and *Schizochytrium* sp. as a sustainable terrestrial ingredient for aquafeeds production / Zarantoniello, Matteo; Zimbelli, Andrea; Randazzo, Basilio; Compagni, Martina Delli; Truzzi, Cristina; Antonucci, Matteo; Riolo, Paola; Loreto, Nino; Osimani, Andrea; Milanović, Vesna; Giorgini, Elisabetta; Cardinaletti, Gloriana; Tulli, Francesca; Cipriani, Renato; Gioacchini, Giorgia; Olivotto, Ike. - In: AQUACULTURE. - ISSN 0044-8486. - ELETTRONICO. - 518;(2020). [10.1016/j.aquaculture.2019.734659]

*Availability:*

This version is available at: 11566/272119 since: 2024-03-27T14:27:15Z

*Publisher:*

*Published*

DOI:10.1016/j.aquaculture.2019.734659

*Terms of use:*

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. The use of copyrighted works requires the consent of the rights' holder (author or publisher). Works made available under a Creative Commons license or a Publisher's custom-made license can be used according to the terms and conditions contained therein. See editor's website for further information and terms and conditions.

This item was downloaded from IRIS Università Politecnica delle Marche (<https://iris.univpm.it>). When citing, please refer to the published version.

(Article begins on next page)



1 **Black Soldier Fly (*Hermetia illucens*) reared on roasted coffee by-product and *Schizochytrium***  
2 ***sp.* as a sustainable terrestrial ingredient for aquafeeds production**

3 Matteo Zarantoniello<sup>a</sup>, Andrea Zimbelli<sup>a</sup>, Basilio Randazzo<sup>a</sup>, Martina Delli Compagni<sup>a</sup>, Cristina  
4 Truzzi<sup>a</sup>, Matteo Antonucci<sup>a</sup>, Paola Riolo<sup>b</sup>, Nino Loreto<sup>b</sup>, Andrea Osimani<sup>b</sup>, Vesna Milanović<sup>b</sup>;  
5 Elisabetta Giorgini<sup>a</sup>, Gloriana Cardinaletti<sup>c</sup>, Francesca Tulli<sup>c</sup>, Renato Cipriani<sup>d</sup>, Giorgia Gioacchini<sup>a</sup>,  
6 Ike Olivotto<sup>\*\*</sup>.

7  
8 <sup>a</sup> *Dipartimento di Scienze della Vita e dell'Ambiente, Università Politecnica delle Marche, via*  
9 *Brecce Bianche, 60131 Ancona, Italy.*

10 <sup>b</sup> *Dipartimento di Scienze Agrarie, Alimentari e Ambientali, Università Politecnica delle Marche,*  
11 *via Brecce Bianche, 60131 Ancona, Italy.*

12 <sup>c</sup> *Department of Agricultural, Food, Environmental and Animal Science, University of Udine,*  
13 *33100, Udine, Italy.*

14 <sup>d</sup> *Società Agricola Alghitaly srl, 37066, Sommacampagna, Verona, Italy.*

15  
16 **\*\*Corresponding author:** Prof. Ike Olivotto; e-mail: [i.olivotto@univpm.it](mailto:i.olivotto@univpm.it); Dipartimento di  
17 Scienze della Vita e dell'Ambiente, Università Politecnica delle Marche, via Brecce Bianche, 60131  
18 Ancona, Italy

19  
20 **Key words:** insect meal; polyunsaturated fatty acids; circular economy; zebrafish; microbiota.

21  
22 **Abstract**

23 Promoting circularity in the aquaculture sector through the conversion of great amount of organic  
24 by-products produced on land is a valuable strategy for a further development of the aquaculture  
25 sector. In this regard, insects represent a very promising example of bio-converting organisms;  
26 their application in aquafeeds, however, still faces possible limitations because of their lack in  
27 polyunsaturated fatty acids and the presence of chitin.

28 The aim of the present study was to apply circularity to Black Soldier Fly (BSF) (*Hermetia*  
29 *illucens*) culture and to improve the insect's biomass fatty acid composition by culturing them on a  
30 land-produced by-product (coffee silverskin) enriched with a 10% *Schizochytrium sp.* The insect  
31 biomass was then used to formulate five fish diets containing 0, 25, 50, 75 and 100% of insect meal  
32 respect to fish meal, respectively. Diets were used for a feeding trial during zebrafish (*Danio rerio*)  
33 larval development (21 days) and a multidisciplinary approach including biometry, histology, gas  
34 chromatography, spectroscopy (FTIR), microbiota analyses and molecular biology was applied to

35 better understand fish responses to the new diets. Results showed that the 50% substitution of fish  
36 meal with insect meal represented the best compromise between ingredient sustainability and proper  
37 fish growth and welfare. Fish fed with higher BSF inclusions (75 and 100%) showed a severe  
38 degree of hepatic steatosis, microbiota modification, a higher lipid content (FTIR), fatty acid  
39 modification and higher expression of both stress and immune response markers.

40

## 41 **1. Introduction**

42 Aquaculture is the fastest growing food production sector worldwide, and it is estimated that by  
43 2030 62% of food-fish will come from aquaculture (FAO, 2018). The reason for this growth relies  
44 on several factors, including wild capture fisheries decline, increase in global demand for seafood  
45 products and the 9 billion people population expected on Earth by 2050, with the consequent  
46 doubling of farmed production required to meet the mid-century demand for seafood (Gerland et al.,  
47 2014; Guillen et al., 2018).

48 Because of the estimated increase in World's population, a significant rise in waste and by-products  
49 production is expected. The EC Directive No. 2008/98, which establishes the order of priority in the  
50 choice of by-products treatments (with their reuse as favoured option and their landfill disposal as  
51 last option), will play a central role for further development of a European circular economy.

52 In consideration of this, aquaculture should be more responsible, sustainable, innovative, based on  
53 the circular economy concept, and able to provide larger volumes of healthy food by using  
54 environmentally friendly ingredients while promoting fish needs and welfare (Merino et al., 2012;  
55 Tlustý and Thorsen, 2017; Stevens et al., 2018; Bohnes and Laurent, 2019).

56 For many years aquaculture has relied on the use of fish meal (FM) and fish oil (FO) as main  
57 ingredients in aquafeeds (Tacon & Metian, 2008; Shepherd and Jackson, 2013). For its further  
58 development, however, nutritious and sustainable ingredients must be identified and tested  
59 (Alhazzaa et al., 2018; Sarker et al., 2018; Vargas et al., 2018).

60 Among such ingredients, several options have already been investigated and great attention has  
61 been addressed to animal by-products (Processed Animals Proteins, PAPs), vegetable sources and  
62 microalgae (Ayadi et al., 2012; Roy and Pal, 2014; Cardinaletti et al., 2018; Xu et al., 2019), but  
63 unfortunately each of them has some downsides in the application for aquafeed formulation  
64 (Francis et al., 2001; Naylor et al., 2009; Manceron et al., 2014; Bandara, 2018; Daniel, 2018).

65 Insects represent a very promising example of bio-converting organisms (Barroso et al., 2014;  
66 Henry et al., 2015; Belghit et al., 2019).

67 Most insect species are farmed on land-produced by-products (van Huis, 2013; Čičková et al.,  
68 2015; Webster et al., 2016; Spinelli et al., 2019;) and, in addition, farming of insects is sustainable  
69 in terms of land use, water consumption and CO<sub>2</sub> production, because of their low environmental  
70 requirements (Berggren et al., 2019; Smetana et al., 2019).

71 Land organic by-products are thus efficiently bio-converted in a highly nutritious biomass which, in  
72 turn, can provide sustainable new ingredients for fish nutrition. Specifically, every year up to  
73 200.000 tons of coffee silverskin are produced by the coffee industry as by-products (Murthy and  
74 Naidu, 2012; Mussatto et al., 2011). Therefore, in a circular economy perspective, coffee silverskin  
75 may represent a suitable substrate for insect production.

76 Among several insect species, the Black Soldier Fly (*Hermetia illucens*; BSF) has a promising role  
77 for aquafeed production. From a nutritional point of view, the BSF accumulates good amounts of  
78 proteins and lipids (307.5-588.0 g kg<sup>-1</sup> and 113.0-386.0 g kg<sup>-1</sup>, respectively; Caligiani et al., 2018;  
79 Nogales-Mérida et al., 2018). However, while the essential amino acid composition is  
80 approximately similar to that of fish meal (Müller et al., 2017), the fatty acid (FA) profile is  
81 extremely different, posing some limits in the full-fat BSF meal inclusion in aquafeeds  
82 (Zarantoniello et al., 2018; Cardinaletti et al., 2019). In terms of fatty acid profile, the BSF is  
83 usually rich in saturated fatty acids (SFAs) and poor in polyunsaturated (PUFAs) ones (Barroso et  
84 al., 2014), which are extremely important for fish (Sargent et al., 1999). PUFAs deficiencies during  
85 fish farming can cause a general decrease of fish health, poor growth, low feed efficiency, anaemia

86 and high mortality (Tocher, 2010; Olivotto et al., 2011; Piccinetti et al., 2012; Dumas et al., 2018).  
87 Because of their fatty acid profile, insects are mainly used as protein source in fish nutrition and  
88 often undergo a defatting process which represents an important cost in their manufacture (Jin et al.,  
89 2012; Li et al., 2015; Wang et al., 2017). Therefore, in aquafeed production it is desirable to use  
90 full-fat insect meal. In this regard, it is known that insects are able to modulate their fatty acid  
91 composition in relation to the growth substrate (Komprda et al., 2013; Liland et al., 2017).  
92 Recently, some authors demonstrated that rearing BSF larvae on an organic substrate containing  
93 proper amounts of omega-3 fatty acids was a suitable procedure to improve the FAs profile of the  
94 final insect biomass (Barroso et al., 2017; St-Hilaire et al., 2007b).

95 New ingredients to be introduced in aquafeeds must be carefully analysed, since it is well  
96 established that different feed ingredients may have modulatory effects of on fish physiological  
97 responses and gut microbiota (Li et al., 2019; Rimoldi et al., 2019). Besides zootechnical indexes,  
98 several molecular markers involved in fish growth, stress response, lipid metabolism, appetite and  
99 immuno response (Olivotto et al., 2002; Piccinetti et al., 2015; Cardinaletti et al., 2019; Vargas-  
100 Abúndez et al., 2019) have been proposed as valid tools to precociously detect physiological  
101 responses in fish fed new diets and represent an up-to-date and important approach.

102 In addition, it is already known that insects possess natural antibiotic properties, possibly modifying  
103 fish microflora (Huyben et al., 2019; Terova et al., 2019), but knowledge concerning interactions  
104 between insect-based diets, gut microbiota, and the aforementioned markers is still fragmentary and  
105 incomplete.

106 The aim of the present study was to interconnect land and aquatic environment by: 1) recycling  
107 land-produced organic by-products (coffee silverskin) to rear BSF larvae; 2) enriching the insects'  
108 growth substrate with *Schizochytrium* sp. to improve their FAs profile; 3) producing highly  
109 nutritious full-fat BSF prepupae meal; 4) testing the biological effects of diets including graded  
110 inclusions of BSF prepupae meal in an aquatic experimental model organism, the zebrafish (*Danio*  
111 *rerio*) and 5) applying a multidisciplinary approach integrating biometric, histological, gas

112 chromatographic, molecular, microbiological and spectroscopic analyses to better understand the  
113 physiological responses of the fish. Zebrafish represents an extraordinary experimental model for  
114 aquaculture, biomedical, developmental biology, genetics, toxicology studies, due to its high  
115 reproductive rate and abundant information that has recently become available from genome  
116 sequencing (Lawrence, 2007; Reed and Jennings, 2011). Particularly, zebrafish are used to  
117 generalize how several biological processes take place in related organisms (like finfish species)  
118 and contribute to understand the mechanisms involved in fish nutrition, welfare and growth,  
119 possibly providing useful information for finfish production (Aleström et al., 2006; Dahm and  
120 Geisler, 2006; De-Santis and Jerry, 2007; Johnston et al., 2008).

121

## 122 **2. Materials and methods**

### 123 *2.1. Ethics*

124 All procedures involving animals were conducted in line with the Italian legislation on experimental  
125 animals and were approved by the Ethics Committee of the Università Politecnica delle Marche  
126 (Ancona, Italy) and the Italian Ministry of Health (626/2018-PR). Optimal rearing conditions (see  
127 further section for details) were applied throughout the study, and all efforts were made to minimize  
128 animal suffering by using an anaesthetic (MS222; Sigma Aldrich, Saint Louis, Missouri, USA).

129

### 130 *2.2. Insect feeding substrate preparation*

131 The main component of the insect feeding substrate consisted of coffee silverskin, a coffee industry  
132 by-product provided by Saccaria Caffè S.R.L. (Marina di Montemarciano, Ancona, Italy). Coffee  
133 by-product (moisture 44%) was collected and grinded in an Ariete 1769 food processor (De Longhi  
134 Appliances Srl, Italy) to a  $0.4 \pm 2$  mm particle size before the feeding substrate preparation. The  
135 insect diet was formulated including a 10% (w/w) of *Schizochytrium* sp to the coffee by-product  
136 (for details, please see Truzzi et al., in press). The freeze-dried *Schizochytrium* sp. was provided by

137 AlghItaly Società Agricola S.R.L. (Sommacampagna, VR, Italy). Insect feeding substrate was  
138 added with distilled water to reach a final moisture of ~70% (Makkar et al., 2014).

139

### 140 2.3. *Insect rearing*

141 Insects were reared in a climatic chamber at a  $27 \pm 1$  °C temperature, relative humidity of  $65 \pm 5\%$   
142 (Spranghers et al., 2017), in continuous darkness. Six days old larvae [purchased from Smart Bugs  
143 s.s. Ponzano Veneto (TV), Italy] were hand counted and divided in groups of 640 larvae per  
144 replicate (n=65) for a total of 41,600 specimens. Each replicate consisted of a plastic box  
145 (57x38x16cm) screened with fine-mesh cotton gauze and covered with a lid provided with 90  
146 ventilation holes of 0.05cm Ø (Spranghers et al., 2017). Larvae were reared at a density of 0.3/cm<sup>2</sup>  
147 (Barragan-Fonseca et al., 2018). Each larva was provided with a feeding rate of 100 mg/day (Diener  
148 et al., 2009) that was prepared and added once a week (448 g for each box). Insects were visually  
149 inspected every day and, when prepupae were identified by the change in tegument colour from  
150 white to black (May, 1961), they were collected, washed, dried and stored at -80 °C.

151

### 152 2.4. *Fish diets production*

153 Full-fat BSF prepupae were freeze dried, grinded with Retsch Centrifugal Grinding Mill ZM 1000  
154 (Retsch GmbH, Haan, Germany) and used to prepare the experimental diets. A control diet (Hi0)  
155 containing FM, wheat gluten, pea protein concentrates and FO as major ingredients, was prepared  
156 according to a commercially available standard diet for zebrafish (Zebrafeed, Sparos ltd, Portugal).  
157 The experimental diets were isonitrogenous (50%) and isolipidic (13%). Insect-based diets were  
158 prepared by including graded levels of insect meal (25%, 50%, 75%, and 100%, referred to as Hi25  
159 and Hi50, Hi75 and Hi100, respectively) in the Hi0 formulation. The conventional vegetable  
160 ingredients (pea protein concentrate and wheat gluten) used to formulate the experimental diets  
161 were maintained approximatively at constant (0.7:1 w:w) ratio in all diets. In summary, all the  
162 grounded ingredients (0.5 mm) and fish oil were thoroughly blended (Kenwood kMix KMX53



stand Mixer; Kenwood, De Longhi S.p.a., Treviso, Italy) for 20 min and then water was added to the mixture to obtain an appropriate consistency for pelleting. Pellets were obtained by using a 1 mm die meat grinder and dried at 40 °C for 48–72 h. The obtained diets were then grinded and stored in vacuum bags at –20 °C until used. Feed samples were analyzed for moisture (AOAC #950.46), crude protein, CP (AOAC #976.05), ash (AOAC #920.153) and ether extract (EE; AOAC #991.36) contents according to AOAC 2006.

The total lipid fraction of the test diets was extracted using chloroform-methanol (2:1 v:v) (Merck KGaA, Darmstadt, Germany) mixture according to Folch et al., 1957. Diet formulation and proximate composition are shown in Table 1.

TABLE 1. INGREDIENTS (g Kg<sup>-1</sup>) AND PROXIMATE COMPOSITION (g 100 g<sup>-1</sup>) OF THE EXPERIMENTAL DIETS USED IN THIS STUDY.

	Hi0 (Control)	Hi25	Hi50	Hi75	Hi100
<b><i>Ingredients (g/kg)</i></b>					
Fish meal <sup>1</sup>	470	400	250	110	-
Vegetable mix <sup>2</sup>	220	230	298	385	440
Hi meal	-	150	275	350	460
Wheat flour <sup>3</sup>	198	172	120	110	72
Fish oil	80	51	25	10	-
Soy lecithin	8	8	8	11	4
Mineral and Vitamin supplements <sup>\$</sup>	14	14	14	14	14
Binder	10	10	10	10	10
<b><i>Proximate composition (%)</i></b>					
Dry Matter	97.08 ± 0.06	95.78 ± 0.13	94.93 ± 0.05	93.63 ± 0.05	92.70 ± 0.04
Crude protein, CP	51.57 ± 0.13	50.75 ± 2.57	50.39 ± 0.28	51.23 ± 1.49	50.50 ± 3.15
Ether extract, EE	14.38 ± 0.64	13.10 ± 0.42	12.93 ± 0.38	13.24 ± 0.46	12.99 ± 0.51
NFE	21.32 ± 0.34	20.82 ± 1.00	20.64 ± 0.55	19.03 ± 0.67	18.47 ± 1.26
Ash	9.81 ± 0.25	11.11 ± 0.01	10.97 ± 0.00	10.13 ± 0.06	10.74 ± 0.13

<sup>1</sup> Raw ingredients kindly supplied by Skretting Italia; <sup>2</sup> Vegetable mix (pea protein concentrate : wheat gluten, 0.7:1 w/w) Lombarda trading srl, Casalbuttano & Uniti (CR, Italy) and Sacchetto spa (Torino, Italy); <sup>3</sup> Consorzio Agrario (Pordenone, Italy); <sup>\$</sup> Mineral and Vitamin supplement composition (% mix): CaHPO<sub>4</sub>·2H<sub>2</sub>O, 78.9; MgO,

177 2.725; KCl, 0.005; NaCl, 17.65; FeCO<sub>3</sub>, 0.335; ZnSO<sub>4</sub>.H<sub>2</sub>O, 0.197; MnSO<sub>4</sub>.H<sub>2</sub>O, 0.094; CuSO<sub>4</sub>.5H<sub>2</sub>O, 0.027;  
178 Na<sub>2</sub>SeO<sub>3</sub>, 0.067; thiamine hydrochloride (vitamin B1), 0.16; riboflavin (vitamin B2), 0.39; pyridoxine  
179 hydrochloride (vitamin B6), 0.21; cyanocobalamine (vitamin B12), 0.21; niacin (vitamin PP or B3), 2.12; calcium  
180 pantotenate, 0.63; folic acid, 0.10; biotin (vitamin H), 1.05; myo-inositol (vitamin B7), 3.15; stay C Roche  
181 (vitamin C), 4.51; tocopherol (vitamin E), 3.15; menadione (vitamin K3), 0.24; retinol (vitamin A 2500 UI kg<sup>-1</sup>  
182 diet), 0.026; cholecalciferol (vitamin D3 2400 UI kg<sup>-1</sup> diet), 0.05; choline chloride, 83.99; \* Values reported as  
183 mean of triplicate analyses; <sup>6</sup> n.d.: not determined

184

## 185 2.5. Fish

186 Zebrafish AB embryos were maintained for 48h in a Tecniplast system (Varese, Italy) in the  
187 following conditions: 28 °C temperature, pH 7.0, NO<sub>2</sub> and NH<sub>3</sub> concentrations < 0.01 mg/L, NO<sub>3</sub>  
188 concentration < 10 mg/L, and photoperiod 12L/12D, respectively (Randazzo et al., 2017). After this  
189 period, embryos were gently collected, counted under a stereomicroscope (Leica Wild M3B, Leica  
190 Microsystems, Nussloch, Germany) and randomly divided in five experimental groups (in  
191 triplicate) according to the five test diets.

192

## 193 2.6. Experimental design

194 Zebrafish larvae were maintained in fifteen 20L tanks to set up the five experimental dietary  
195 treatments; each experimental group was composed of 1500 larvae (500 larvae per tank). The water  
196 in the larval tanks had the same chemical-physical characteristics of the parent's tank and was  
197 gently replaced 10 times a day by a dripping system (Olivotto et al., 2004). The sides of each tank  
198 were covered with black panels to reduce light reflection. All tanks were siphoned 30 min after  
199 feeding (twice a day) to remove possible feed excess and dead larvae. The required larvae were  
200 sampled 20 days after fertilization (dpf), euthanized with a lethal dose of MS222 (1g/L) and  
201 properly stored for further analyses.

202

## 203 2.7. Feeding schedule

204 Starting from 5 dpf to 20 dpf, zebrafish larvae were fed as follows: Control group: larvae fed on diet  
205 0% insect meal (Hi0 diet); Group A: larvae fed on the diet including 25% BSF full-fat prepupae

meal (Hi25 diet); Group B: larvae fed on the diet including 50% of BSF full-fat prepupae meal (Hi50 diet); Group C: larvae fed on the diet including 75% BSF full-fat prepupae meal (Hi75 diet); Group D: larvae fed on the diet including 100% BSF full-fat prepupae meal (Hi100 diet). Zebrafish larvae were fed on the experimental diets (2% body weight, BW; 100-250 µm size) twice a day and, in addition, from 5 to 10 dpf, all groups were fed (one feeding in the morning) on the rotifer *Brachionus plicatilis* (5 ind/mL) according to Lawrence et al. (2012) and Piccinetti et al. (2014).

212

## 213 2.8. Biometry

Five zebrafish larvae (15 per dietary group) were randomly collected from the different tanks of each experimental group at hatching (3 dpf) and at the end of the experiment (20 dpf). The standard length was determined using a sliding calliper (Measy 2000 Typ 5921, Swiss; precision: 0.1 mm) and the wet weight using an OHAUS Explorer (OHAUS Europe GmbH, Greifensee, Switzerland) analytical balance (precision: 0.1 mg) according to Zarantonello et al. (2018). At 3 dpf, wet weight was measured on pools of five larvae in triplicate. For each experimental group, specific growth rate (SGR) was calculated as follows:  $SGR\% = [(\ln W_f - \ln W_i)/t] \times 100$ , where  $W_f$  is the final wet weight,  $W_i$ , the initial wet weight, and  $t$ , the number of days (17). Survival rate in all experimental groups was about 85%.

223

## 224 2.9. Fatty acid composition

The experimental diets and fish larvae samples were analyzed for fatty acid composition. Samples were minced and homogenized (homogenizer MZ 4110, DCG Eltronic, Monza, Italy), and larvae were also freeze-dried (Edwards EF4, Crawley, Sussex, England). Aliquots of 200 mg of each sample were added with 100 µl of Internal Standard (methyl ester of nonadecanoic acid, 99.6%, Dr. Ehrenstorfer GmbH, Germany), and extracted overnight following the method of Folch et al., 1957. Analyses were carried out on three aliquots *per* sample. All lipid extracts were evaporated under laminar flow inert gas (N<sub>2</sub>) until constant weight and re-suspended in 0.5 ml of n-eptane.

232 Fatty acid methyl esters (FAMES) were prepared according to Canonico et al., 2016 using methyl  
233 ester of nonadecanoic acid (19:0; Dr. Ehrenstorfer GmbH, Augsburg, Germany) as internal  
234 standard. FAMES were determined using an Agilent-6890 GC System (Milano, Italy) coupled to an  
235 Agilent-5973N quadrupole Mass Selective Detector (MSD) (Milano, Italy). A CPS ANALITICA  
236 CC-wax-MS (30 m × 0.25 mm ID, 0.25 µm film thickness) capillary column was used to separate  
237 FAMES. Instrumental conditions for the studied matrices were set up according to Truzzi et al.  
238 (2017, 2018). For each analysed aliquot of sample, at least three runs were performed on the  
239 GCMS.

240

#### 241 *2.10. Histology*

242 Nine zebrafish larvae per dietary group (3 larvae per tank, at 20 dpf), randomly collected from the  
243 different tanks, were fixed by immersion in Bouin's solution (Sigma-Aldrich, Milano, Italy) and  
244 then stored at 4°C for 24h. Larvae were washed three times in ethanol (70%) for ten minutes and  
245 preserved in the same ethanol solution. Larvae were then dehydrated in increasing ethanol solutions  
246 (80, 95 and 100%), washed in xylene (Bio-Optica, Milano, Italy) and embedded in paraffin (Bio-  
247 Optica). Solidified paraffin blocks were cut with a microtome (Leica RM2125 RTS, Nussloch,  
248 Germany) and 5 µm sections were stained with Mayer hematoxylin and eosin Y (Sigma-Aldrich,  
249 Milano, Italy). Sections were observed using a Zeiss Axio Imager.A2 (Oberkochen, Germany)  
250 microscope in order to study the hepatic parenchyma and intestine morphology. Images were  
251 acquired by mean of a combined color digital camera Axiocam 503 (Zeiss, Oberkochen, Germany).  
252 Moreover, to ascertain the extent of fat accumulation in liver, a quantitative analysis was performed  
253 on a significant number of histological sections from each experimental group in triplicate (n=9).  
254 No-n-evaluable areas, such as blood vessels were not considered. The percentage of fat fraction  
255 (PFF) on the total tissue areas was calculated using the ImageJ software setting a homogeneous  
256 threshold value.

257

258

## 259 2.11. FTIR analysis

260 Fifteen zebrafish larvae per dietary group (5 per tank) were randomly collected at 20 dpf. Samples  
261 were minced, homogenized (homogenizer MZ 4110, DCG Eltronic, Monza, Italy) and freeze-dried  
262 (Edwards EF4, Crawley, Sussex, England) for FTIR analysis. For each group, five aliquots of 5 mg  
263 each were analysed. FTIR analysis was performed using a Spectrum GX1 Spectrometer equipped  
264 with a U-ATR accessory and a diamond/SeZn crystal (Perkin Elmer, Waltham, Massachusetts, USA).  
265 Measurements were carried out in reflectance in the MIR region from 4000 to 800  $\text{cm}^{-1}$  (spectral  
266 resolution 4  $\text{cm}^{-1}$ ). Each spectrum was the result of 64 scans. Before each sample acquisition, a  
267 background spectrum was collected on the clean surface of the crystal. Raw IR spectra were  
268 converted in absorbance, two-points baseline linear fitted in the 4000-800  $\text{cm}^{-1}$  spectral range and  
269 vector normalized in the same interval (OPUS 7.1 software package).

270 For all experimental groups, the average absorbance spectra were calculated together with their  
271 standard deviation spectra (average absorbance spectrum  $\pm$  standard deviation spectra) and analysed  
272 to identify the most featuring IR peaks (in terms of position/wavenumbers). Then, average absorbance  
273 spectra and their standard deviation spectra were curve-fitted in the 3050-2800  $\text{cm}^{-1}$  and 1790-900  $\text{cm}^{-1}$   
274  $^1$  regions upon two-points baseline correction and vector normalization. A Gaussian algorithm was  
275 adopted, and the number and position of the underlying peaks was defined by second derivative  
276 analysis of the spectra (GRAMS/AI 9.1, Galactic Industries, Inc., Salem, NH). In the 3050-2800  $\text{cm}^{-1}$   
277 region, the following underlying peaks were identified:  $\sim 3010 \text{ cm}^{-1}$  ( $=\text{CH}$  moieties in unsaturated  
278 lipid alkyl chains);  $\sim 2959 \text{ cm}^{-1}$  and  $\sim 2872 \text{ cm}^{-1}$  ( $\text{CH}_3$  groups in lipid alkyl chains);  $\sim 2925 \text{ cm}^{-1}$  and  
279  $\sim 2854 \text{ cm}^{-1}$  ( $\text{CH}_2$  groups in lipid alkyl chains). In the 1790-900  $\text{cm}^{-1}$  region, the following underlying  
280 peaks were identified:  $\sim 1744 \text{ cm}^{-1}$  ( $\text{C}=\text{O}$  moiety in lipids and fatty acids);  $\sim 1639$  and  $\sim 1536 \text{ cm}^{-1}$   
281 (respectively Amide I and II bands of proteins);  $\sim 1457 \text{ cm}^{-1}$  (proteins side chains);  $\sim 1390 \text{ cm}^{-1}$   
282 ( $\text{COO}^-$  groups in aspartate and glutamate amino acids);  $\sim 1234 \text{ cm}^{-1}$  (collagen);  $\sim 1157 \text{ cm}^{-1}$

(glycosylated compounds);  $\sim 1080\text{ cm}^{-1}$  (phosphate groups), and  $\sim 1055\text{ cm}^{-1}$  (mucin). The integrated areas of the most relevant peaks were calculated and used to evaluate the following band area ratios: LIP/TBM (representing the overall amount of lipids), 1744/LIP (representing the amount of fatty acids compared to lipids), 2928/LIP (representing the amount of saturated fatty acids with respect to lipids), and 3010/LIP (representing the amount of unsaturated fatty acids compared to lipids); PRT/TBM (representing the overall amount of proteins), 1234/PRT (representing the amount of collagen compared to proteins), and 1055/PRT (representing the amount of mucin compared to proteins). TBM was the sum of the integrated areas of all peaks in the  $3050\text{-}2800\text{ cm}^{-1}$  and  $1790\text{-}900\text{ cm}^{-1}$  regions; LIP was the sum of the integrated areas of all peaks in the  $3050\text{-}2800\text{ cm}^{-1}$  region, while PRT was the sum of the integrated areas of the bands at  $1639$  and  $1536\text{ cm}^{-1}$ .

## 2.12. Microbiome

*RNA extraction and cDNA synthesis.* Prior to analysis, zebrafish larvae (60 larvae per dietary group) were disinfected by washing in 50 mL of ethanol (70%) on a laboratory shaker (VDRL Stirrer with thermostatic cupola, ASAL s.r.l, Milan, Italy) at 150 rpm for 1 min at room temperature, in order to analyse only the gut microbiome. These samples were thus identified as G (gut) Hi0, 25, 50, 75 and 100 for these analyses. After discharging the ethanol, the samples were rinsed in two additional washing steps in 50 mL of sterile deionized water. Subsequently, 31.5 mL of sterile peptone water (peptone, 1 g/L) was added to each sample, which was then homogenized in a Stomacher apparatus (400 Circulator, International PBI, Milan, Italy) for 3 min at 260 rpm. Then, 1.5 mL of each tenfold diluted (10<sup>-1</sup> dilution) homogenate were centrifuged at 14,000 rpm for 10 min, the supernatants were discarded, the obtained cell pellets covered with RNA later Stabilization Solution (Ambion, Foster City, CA, USA) and stored at  $-80^{\circ}\text{C}$  until use. The Quick-RNA Fungal/Bacterial Microprep kit (Zymo Research, CA, USA) was used for the extraction of total microbial RNA from the cell pellets following the manufacturer's instructions. The extracted RNAs were checked for quantity, purity and integrity as previously described by Garofalo et al. (2017). Moreover, the extracts were amplified using the PCR universal prokaryotic primers 27f and 1495r (Weisburg et al., 1991) to

309 exclude the presence of bacterial DNA contamination. Five  $\mu$ L of each RNA sample were reverse-  
310 transcribed in cDNA using the SensiFAST cDNA Synthesis Kit for RT-qPCR (Bioline, London,  
311 UK) following the manufacturer's instructions.

312

313 *16S rRNA amplicon target sequencing.* cDNA was used as template in the PCR amplifying the V3-  
314 V4 region of the 16S rRNA gene using the primers and protocols described by Klindworth et al.  
315 (2013).

316 PCR amplicons were cleaned using the Agencourt AMPure kit (Beckman Coulter, Milan, Italy) and  
317 tagged using the Nextera XT Index Kit (Illumina Inc. San Diego, CA) according to the  
318 manufacturer's instructions. After the 2<sup>nd</sup> clean-up step, amplicons were quantified using a QUBIT  
319 dsDNA Assay kit and an equimolar amount of amplicons from different samples were pooled. The  
320 library was denatured with 0.2 N NaOH, diluted to 12 pM, and combined with 20% (vol/vol)  
321 denatured 12 pM PhiX, prepared according to Illumina guidelines. The sequencing was performed  
322 with a MiSeq Illumina instrument (Illumina) with V3 chemistry and generated 250 bp paired-end  
323 reads according to the manufacturer's instructions.

324

### 325 *2.13. Molecular analyses*

326 *RNA extraction and cDNA synthesis.* Total RNA extraction from 5 zebrafish larvae from each tank  
327 (15 larvae per dietary group) was optimized using the RNeasy RT reagent (Sigma-Aldrich, R4533)  
328 according to Piccinetti et al. (2013). The total RNA extracted was eluted in 40  $\mu$ l of RNase-free  
329 water (Qiagen). The final RNA concentration was determined using a NanoPhotometer P-Class  
330 (Implen, München, Germany). RNA integrity was verified by GelRed<sup>TM</sup> staining of 28S and 18S  
331 ribosomal RNA bands on 1% agarose gel. RNA was stored at -80°C until use. Finally, 2  $\mu$ g of RNA  
332 were used for cDNA synthesis, using the High Capacity cDNA Reverse Transcription Kit (Bio-Rad,  
333 Milan, Italy) following the manufacturer's instructions.

334

335 *Real-Time PCR*. PCRs were performed with SYBER green method in an iQ5 iCycler thermal cycler  
336 (Bio-Rad Laboratories) following Vargas et al. (2018).  
337 Relative quantification of the expression of genes involved in fish growth (*igf1*, *igf2a* and *mstnb*),  
338 stress response (*nr3c1* and *hsp70.1*), long-chain polyunsaturated fatty acids biosynthesis (*elovl2*,  
339 *elovl5* and *fads2*), appetite response (*ghrl*, *npv*, *cnr1* and *lepa*), immune response (*il1b*, *il10* and  
340 *tnfa*) and enzymatic hydrolysis of chitin (*chia.2* and *chia.3*) was performed. Actin-related protein  
341 2/3 complex, subunit 1A (*arpc1a*) and ribosomal protein, large, 13 (*rpl13*) were used as internal  
342 standards in each sample in order to standardize the results by eliminating variation in mRNA and  
343 cDNA quantity and quality. No amplification products were observed in negative controls and no  
344 primer-dimer formations were observed in the control templates. Amplification products were  
345 sequenced, and homology was verified. The data obtained were analysed using the iQ5 optical  
346 system software version 2.0 (Bio-Rad) including GeneEx Macro iQ5 Conversion and genex Macro  
347 iQ5 files. Primer sequences were designed using Primer3 (210 v. 0.4.0) starting from zebrafish  
348 sequences available in ZFIN Primer sequences used were reported in Table 2.

349  
350 TABLE 2. PRIMER SEQUENCES AND THE ZEBRAFISH INFORMATION NETWORK  
351 (ZFIN) USED IN THIS STUDY

<i>Gene</i>	<i>Forward primer (5'- 3')</i>	<i>Reverse primer (5'- 3')</i>	<i>ZFIN ID</i>
<i>igf1</i>	5'-GGCAAATCTCCACGATCTCTAC-3'	5'-CGGTTTCTCTTGTCTCTCTCAG-3'	ZDB-GENE-010607-2
<i>igf2a</i>	5'-GAGTCCCATTCTCTGTTG-3'	5'-GTGGATTGGGGTTTGATGTG-3'	ZDB-GENE-991111-3
<i>mstnb</i>	5'-GGACTGGACTGCGATGAG-3'	5'-GATGGGTGTGGGGATACTTC-3'	ZDB-GENE-990415-165
<i>nr3c1</i>	5'-AGACCTTGGTCCCCTTCACT-3'	5'-CGCCTTTAATCATGGGAGAA-3'	ZDB-GENE-050522-503
<i>hsp70.1</i>	5'-TGTTCACTTCTCTGCCGTTG-3'	5'-AAAGCACTGAGGGACGCTAA-3'	ZDB-GENE-990415-91
<i>elovl2</i>	5'-CACTGGACGAAGTTGGTGAA-3'	5'-GTTGAGGACACACCACCAGA-3'	ZDB-GENE-060421-5612
<i>elovl5</i>	5'-TGGATGGGACCGAAATACAT-3'	5'-GTCTCCTCCACTGTGGGTGT-3'	ZDB-GENE-040407-2
<i>fads2</i>	5'-CATCACGCTAAACCCAACA-3'	5'-GGGAGGACCAATGAAGAAGA-3'	ZDB-GENE-011212-1
<i>ghrl</i>	5'-CAGCATGTTTCTGCTCCTGTG-3'	5'TCTTCTGCCACTCTTGGTG-3'	ZDB-GENE-070622-2
<i>npv</i>	5'-GTCTGCTTGGGGACTCTCAC-3'	5'CGGGACTCTGTTTCACCAAT-3'	ZDB-GENE-980526-438
<i>cnr1</i>	5'-AGCAAAAGGAGCAACAGGCA-3'	5'GTTGGTCTGGTACTTTCACCTTGAC-3'	ZDB-GENE-040312-3
<i>lepa</i>	5'-CTCCAGTGACGAAGGCAACTT-3'	5'GGGAAGGAGCCGGAAATGT-3'	ZDB-GENE-081001-1



<i>il1b</i>	5'-GCTGGGGATGTGGACTTC-3'	5'-GTGGATTGGGGTTTGATGTG-3'	ZDB-GENE-040702-2
<i>il10</i>	5'-ATTTGTGGAGGGCTTTCCTT-3'	5'-AGAGCTGTTGGCAGAATGGT-3'	ZDB-GENE-051111-1
<i>tnfa</i>	5'-TTGTGGTGGGGTTTGATG-3'	5'-TTGGGGCATTTTATTTGTAAAG-3'	ZDB-GENE-050317-1
<i>chia.2</i>	5'-GGTGCTCTGCCACCTTGCCTT-3'	5'-GGCATGGTTGATCATGGCGAAAGC-3'	ZDB-GENE-040426-2014
<i>chia.3</i>	5'-TCGACCCTTACCTTTGCACACACCT-3'	5'-ACACCATGATGGAGAACTGTGCCGA-3'	ZDB-GENE-040426-2891
<i>arpc1a</i>	5'-CTGAACATCTCGCCCTTCTC-3'	5'-TAGCCGATCTGCAGACACAC-3'	ZDB-GENE-040116-1
<i>rpl13</i>	5'-TCTGGAGGACTGTAAGAGGTATGC-3'	5'-AGACGCACAATCTTGAGAGCAG-3'	ZDB-GENE-031007-1

352

## 353 2.14. Statistical analysis

354 All data (except for microbiome) were analyzed by one-way ANOVA, with diet as the explanatory  
355 variable. All ANOVA tests were followed by Tukey's post-hoc tests. The statistical software  
356 package Prism5 (GraphPad Software) was used. Significance was set at  $p < 0.05$  and all results are  
357 presented as mean  $\pm$  SD. For microbiome analyses, paired-end reads were first merged using the  
358 FLASH software (Magoč and Salzberg, 2011). Joint reads were quality filtered (at Phred  $< Q20$ ) by  
359 QIIME 1.9.0 software (Caporaso et al., 2010) and the pipeline recently described (Osimani et al.,  
360 2019). Briefly, the USEARCH software version 8.1 (Edgar et al., 2011) was used for chimera  
361 filtering and clean sequences were clustered into Operational Taxonomic Units (OTUs) at 97% of  
362 similarity by UCLUST algorithms. Centroids sequences of each cluster were used for taxonomic  
363 assignment using the Greengenes 16S rRNA gene database. OTU tables were rarefied at 44412  
364 sequences. The OTU table displays the higher taxonomy resolution that was reached.

365

## 366 3. Results

### 367 3.1. Biometry

368 The increasing inclusion levels of BSF full-fat prepupae meal resulted in a statistically significant  
369 ( $p < 0.05$ ) increase in the larval specific growth rate. In particular, no significant differences ( $p > 0.05$ )  
370 were detected between Control ( $25.4 \pm 0.7\%$ ) and Hi25 group ( $25.7 \pm 1.0\%$ ), while Hi50, Hi75 and

Hi100 groups ( $27.6\pm0.5$ ,  $27.8\pm0.4$ ,  $28.4\pm0.3$  %, respectively) showed significantly ( $p<0.05$ ) higher values compared to both Control and Hi25 ones.

### 3.2. Fatty acid content and composition

*Diets.* The FAs classes percentages of the five experimental diets are presented in Figure 2a. Insect-based diets showed significantly higher ( $p<0.05$ ) percentages of SFAs ( $40.9\pm0.7$ ,  $40.0\pm2.0$ ,  $35.9\pm0.7$  and  $37.6\pm2.8$  % for Hi25, Hi50, Hi75 and Hi100 diets, respectively) compared to Control diet ( $27.8\pm1.3\%$ ). Considering mono-unsaturated fatty acids (MUFAs), all insect-based diets showed significantly lower percentages ( $p<0.05$ ) compared to Control ( $24.7\pm0.6$ ,  $19.8\pm0.3$ ,  $19.0\pm0.9$ ,  $21.5\pm0.2$  and  $20.0\pm1.0$  % for Control, Hi25, Hi50, Hi75 and Hi100, respectively). Finally, insect-based diets showed significantly ( $p<0.05$ ) lower ( $39.3\pm1.0$ ,  $41.0\pm1.0$ ,  $42.6\pm0.3$  and  $42.2\pm3.2$  % for Hi25, Hi50, Hi75 and Hi100, respectively) percentages of PUFAs compared to Control diet ( $47.4\pm1.4$  %). In addition, increasing inclusion levels of BSF full-fat prepupae meal in the diets resulted in a significant decrease ( $p<0.05$ ) of n3 percentages (from  $38.8\pm1.4\%$  for Control diet to  $11.1\pm3.1$  % for Hi100 diet) and a parallel significant ( $p<0.05$ ) increase in n6 percentages (from  $8.6\pm0.1$  for Control diet to  $31.3\pm0.9$  % for Hi100 diet; Fig. 2a). Consequently, the n6/n3 ratio showed significant differences ( $p<0.05$ ) among experimental diets, increasing from Control diet ( $0.22\pm0.01$ ) to Hi100 diet ( $2.8\pm0.2$ ) diets (Fig. 2b). Finally, considering the insect-based diets, the higher was the BSF meal dietary inclusion, the higher was the n9 content ( $10.7\pm0.2$ ,  $12.1\pm0.7$ ,  $14.6\pm0.2$  and  $15.2\pm0.7$  % for Hi25, Hi50, Hi75 and Hi100, respectively). The control diet ( $13.9\pm0.3\%$ ) showed an intermediate n9 content between Hi50 and Hi75 diets.

*Zebrafish larvae.* Figure 2c illustrates the FAs classes percentages of zebrafish larvae fed on the different diets. The FA classes of zebrafish larvae fed on the different diets was deeply influenced by the BSF meal dietary inclusion. In particular, SFAs percentage increased with the inclusion of insect meal in the diets, while both MUFAs and PUFAs generally decreased with the increasing

BSF inclusion levels in the diets (Fig. 2c). Specifically, the higher was the dietary BSF meal inclusion level, the lower was the n3 and the higher was the n6 percentage detected. Considering the n6/n3 ratio (Fig. 2d), no significant differences ( $p>0.05$ ) were detected among Control, Hi25 and Hi50 groups ( $0.5\pm0.1$ ,  $0.5\pm0.1$  and  $0.6\pm0.1$ , respectively), while Hi75 and Hi100 ( $0.8\pm0.1$  and  $1.0\pm0.1$ , respectively) showed a significantly ( $p<0.05$ ) higher value compared to the other experimental groups.

Finally, in terms of n9 percentage, only Hi25 and Hi50 groups ( $17.7\pm0.4$  and  $17.9\pm0.3$  %, respectively) showed significantly ( $p<0.05$ ) higher values compared to Control group ( $17.3\pm0.2$  %).

Table 3 shows the FA composition of total lipids of zebrafish larvae. The FA profile of larvae reared on the Control diet was characterized by high percentages of 16:0, 18:1n9, 22:6n3 (docosahexaenoic acid, DHA), and 20:5n3 (eicosapentaenoic acid, EPA) fatty acids, followed by 18:2n6, 16:1n7 and 18:0. The increasing inclusion levels of BSF meal in the diets triggered some changes in the FA composition of the larvae and the major changes are reported here.

A substantial (up to ~30-folds) increase in the lauric acid (12:0) and a significant decrease in EPA (20:5n3) percentage was detected in fish fed diets with increasing BSF meal inclusion levels. In terms of DHA, a similar, but milder trend was observed respect to EPA. Because of this the DHA/EPA ratio significantly increased with the increasing BSF meal inclusion levels in the diets (Tab. 3).

TABLE 3: FATTY ACID COMPOSITION (AS % OF TOTAL FAS) OF ZEBRAFISH LARVAE FED DIETS WHERE FM WAS REPLACED WITH 25, 50, 75 AND 100 % OF BSF MEAL (Hi25, Hi50, Hi75 AND Hi100 GROUPS).

	Zebrafish larvae				
	CTRL	Hi25	Hi50	Hi75	Hi100
10:0	$0.02\pm0.01$	$0.05\pm0.01$	$0.10\pm0.01$	$0.08\pm0.01$	$0.09\pm0.02$
12:0	$0.20\pm0.03^a$	$2.6\pm0.1^b$	$4.5\pm0.1^c$	$4.7\pm0.4^c$	$5.7\pm0.4^d$
13:0	$0.06\pm0.01$	$0.06\pm0.01$	$0.07\pm0.01$	$0.06\pm0.01$	$0.07\pm0.01$
14:0	$4.0\pm0.1^{a,b}$	$4.0\pm0.1^a$	$4.2\pm0.1^b$	$3.9\pm0.2^a$	$4.3\pm0.2^b$
15:0	$0.86\pm0.04$	$0.80\pm0.03$	$0.82\pm0.01$	$0.79\pm0.02$	$0.84\pm0.04$
16:0	$14.7\pm1.0^c$	$14.0\pm0.7^{a,b}$	$13.8\pm0.5^a$	$14.6\pm0.1^{b,c}$	$15.5\pm0.9^d$

16:1n9	1.2±0.1 <sup>a</sup>	1.3±0.1 <sup>b</sup>	1.3±0.1 <sup>c</sup>	1.5±0.1 <sup>d</sup>	1.5±0.1 <sup>d</sup>
16:1n7	8.7±0.6 <sup>c</sup>	9.0±0.3 <sup>c</sup>	8.3±0.1 <sup>b</sup>	7.6±0.4 <sup>a</sup>	7.9±0.1 <sup>a</sup>
17:0	1.2±0.1 <sup>b</sup>	1.1±0.1 <sup>a</sup>	1.1±0.1 <sup>a</sup>	1.1±0.1 <sup>a</sup>	1.2±0.1 <sup>b</sup>
18:0	6.3±0.2 <sup>c</sup>	5.5±0.1 <sup>a</sup>	5.8±0.2 <sup>b</sup>	6.3±0.4 <sup>c</sup>	6.7±0.4 <sup>d</sup>
18:1n9	14.7±0.1 <sup>a</sup>	15.3±0.3 <sup>b</sup>	15.7±0.7 <sup>c</sup>	15.1±0.3 <sup>b</sup>	15.3±0.1 <sup>b</sup>
18:1n7	5.2±0.1 <sup>d</sup>	4.9±0.2 <sup>c</sup>	4.6±0.2 <sup>b</sup>	4.3±0.1 <sup>a</sup>	4.5±0.2 <sup>b</sup>
18:2n6	9.6±0.1 <sup>a</sup>	10.6±0.2 <sup>b</sup>	11.0±0.2 <sup>c</sup>	12.6±0.3 <sup>d</sup>	12.7±0.7 <sup>d</sup>
18:3n6	0.43±0.01	0.41±0.02	0.50±0.01	0.47±0.03	0.53±0.03
18:3n3	3.7±0.1 <sup>c</sup>	3.3±0.2 <sup>b</sup>	3.5±0.2 <sup>c</sup>	2.7±0.2 <sup>a</sup>	2.8±0.3 <sup>a</sup>
20:0	0.22±0.01	0.18±0.01	0.19±0.01	0.23±0.02	0.21±0.01
20:1n9	0.90±0.05	0.75±0.01	0.63±0.02	0.48±0.02	0.39±0.02
20:2n6	0.29±0.02	0.29±0.01	0.31±0.01	0.35±0.02	0.38±0.02
20:3n6	0.46±0.04 <sup>a</sup>	0.52±0.02 <sup>b</sup>	0.70±0.04 <sup>c</sup>	0.98±0.06 <sup>d</sup>	1.1±0.1 <sup>e</sup>
20:4n6	2.3±0.2 <sup>a</sup>	2.3±0.1 <sup>a</sup>	2.4±0.1 <sup>b</sup>	3.2±0.3 <sup>c</sup>	3.5±0.1 <sup>d</sup>
20:3n3	0.21±0.03	0.19±0.01	0.21±0.01	0.17±0.01	0.17±0.02
20:5n3	10.2±0.2 <sup>e</sup>	8.7±0.7 <sup>d</sup>	7.2±0.3 <sup>c</sup>	4.6±0.1 <sup>b</sup>	3.5±0.2 <sup>a</sup>
22:1n9	0.31±0.02	0.28±0.01	0.18±0.02	0.12±0.01	0.04±0.01
22:6n3	14.0±1.3 <sup>c</sup>	13.9±1.2 <sup>c</sup>	12.9±0.8 <sup>b</sup>	14.0±1.3 <sup>c</sup>	11.2±1.0 <sup>a</sup>
24:1n9	0.18±0.02	0.09±0.01	0.08±0.01	0.05±0.01	0.03±0.01
DHA/EPA	1.4±0.1 <sup>a</sup>	1.6±0.2 <sup>b</sup>	1.8±0.1 <sup>b</sup>	3.0±0.3 <sup>c</sup>	3.2±0.2 <sup>c</sup>

Means within rows bearing different letters are significantly different ( $p < 0.05$ ). Statistical analysis was performed only for FAs > 1%. FAs with a percentage <1% were excluded from any statistical analyses because their concentrations were close to the limit of detection.

### 3.3. Histology

Histological analyses were performed in order to detect possible inflammatory events in the intestine and to evaluate lipid accumulation or steatosis in the liver. The intestine mucosa appeared unaltered and did not show any appreciable inflammatory influx in all experimental groups and fish fed on BFS-based diets showed a normal intestinal morphology, comparable to Control (Fig. 3a-j). Conversely, results evidenced a variable degree of lipid accumulation in the liver of the experimental groups (Fig. 3k-o). The most appreciable visual differences were observed in the Hi75 and Hi100

groups. Indeed, while all experimental groups presented a modest fat liver parenchima, Hi75 and Hi100 showed a severe degree of steatosis with swollen hepatocytes and abundant intracytoplasmic lipid accumulation (Fig. 3n,o). These results were confirmed by the statistical quantification of the fat percentage fraction (PPF) on liver sections that showed a significant increase in response to dietary treatments (Fig. 4). In particular, Hi50 showed a significant increase in PPF ( $48.7 \pm 2.4$ ) compared to Control ( $41.7 \pm 2.1$ ) and Hi25 ( $42.5 \pm 1.2$ ) groups. However, the highest values in PPF were detected in Hi75 and Hi100 with a significant difference ( $58.7 \pm 0.9$  and  $60.6 \pm 1.5$  respectively) compared to the other groups.

441

### 442 3.4. FTIR analysis

The average absorbance spectra of Control, Hi25, Hi50, Hi75 and Hi100 larval groups are shown in Figure 5. The analysis of IR spectra allowed to distinguish the vibrational modes of lipids, proteins, carbohydrates and phosphates (see Materials and Methods section) (Vargas et al., 2018).

In terms of biochemical composition of zebrafish larvae samples belonging to the different dietary groups, the following considerations can be drawn: (i) statistically significant higher amounts of overall lipids (LIP/TBM, Fig. 6a) were detected in Hi50, Hi75 and Hi100 zebrafish larvae compared to Control and Hi25 ones; (ii) in all zebrafish larvae fed on diets with different inclusion levels of BSF meal, a significant increase in saturated fatty acids (1744/LIP and 2925/LIP, Fig. 6b and 6c) and a significant decrease in unsaturated ones (3010/LIP, Fig. 6d) were observed; (iii) in terms of protein composition, no statistically significant changes were detected in the overall amount of proteins (PRT/TBM, Fig. 7a) among the experimental groups; (iv) a slight but statistically significant decrease of collagen (1234/PRT, Fig. 7b) was observed in all zebrafish larvae fed on diets including BFS meal, and (v) a higher amount of mucin (1055/PRT, Fig. 7c) was found in Hi50, Hi75 and Hi100 zebrafish larvae.

### 457 3.5. Microbiome

458 A total of 247,654 reads passed the filters applied through QIIME, with an average value of 49,530  
459 reads/sample, and a mean sequence length of 464 bp. The Good's coverage indicated also  
460 satisfactory coverage for all samples (>99%) (Supplementary Table 1A). Alpha-diversity indicated  
461 the highest number of OTUs after 50% of BSF inclusion if compared with control and with 25% of  
462 inclusion. Beta diversity calculation based on weighted and on unweight UniFrac distance matrix  
463 showed a clear separation of the control samples if compared with those including BSF (Fig. 8).  
464 Furthermore, 50 and 75% of BSF meal inclusion in the diets showed a similar effect on microbial  
465 composition.

466 As shown in Figure 9, the main OTUs shared among the data sets were *Cetobacterium* that reached  
467 50% of the relative abundance in sample GHi50 and about 30% in the other samples. *Vibrio* and  
468 *Mycoplasmataceae* decreased with the increasing inclusion of BSF meal in the diets;  
469 *Flavobacterium* and *Plesiomonas* were present at very low abundance in GHi100 sample, whereas  
470 in the other samples the same two genera showed a remarkable presence. Finally, among other  
471 bacteria, *Aeromonadaceae*, *Ochrobactrum* and *Tetrathiobacter* were also detected.

472

### 473 3.6. Real-time PCR results

474 Real-time PCR analyses were performed on genes involved in fish growth (*igf1*, *igf2a* and *mstnb*),  
475 stress response (*hsp70.1* and *nr3c1*), long-chain polyunsaturated fatty acids biosynthesis (*elovl2*,  
476 *elovl5* and *fads2*), appetite (*ghrl*, *npv*, *cnr1* and *lepa*), immune response (*il1b*, *il10* and *tnfa*) and  
477 enzymatic hydrolysis of chitin (*chia.2* and *chia.3*).

478 *Growth factors.* Higher mean levels in the expression of the *igf1* and *igf2a* genes were detected in  
479 all experimental groups fed on BSF-based diets compared to Control (with the exception of *igf1*  
480 gene expression in Hi25 group). In particular, the Hi75 and Hi100 groups showed significantly  
481 ( $p<0.05$ ) higher levels in comparison to the Hi25 group in terms of *igf1* gene expression, and to  
482 Control in terms of *igf2a* gene expression. Only the Hi50 and Hi100 groups showed significantly  
483 ( $p<0.05$ ) higher levels of expression of the *mstn* gene than Control (Fig. 10c).

484 *Stress response.* Considering stress markers (*nr3c1* and *hsp70.1*; Fig. 10d,e), all groups fed on BSF-  
485 based diets showed a significantly ( $p<0.05$ ) higher gene expression compared to Control (with the  
486 exception of *nr3c1* gene expression in Hi25 group). Furthermore, no significant differences  
487 ( $p>0.05$ ) were evident among the Hi50, Hi75 and Hi100 groups, while the Hi25 group showed  
488 significantly ( $p<0.05$ ) lower values compared to groups fed on diets with higher BSF meal inclusion  
489 level (with the exception of *nr3c1* gene expression in Hi50 group).

490

491 *Lipid metabolism.* The highest BSF meal inclusion (Hi100) caused the highest expression of the  
492 genes involved in long-chain polyunsaturated fatty acid elongation (*elovl2* and *elovl5*) and  
493 desaturation (*fads2*). As illustrated in Figure 10f-h, the Hi100 group presented the highest gene  
494 expression ( $p<0.05$ ) compared to all the other experimental groups. More specifically, no  
495 significant differences ( $p>0.05$ ) in the expression of the *elovl2* gene were detected between Control  
496 and Hi25 group (Fig. 10f), while both Hi50 and Hi75 had a significantly ( $p<0.05$ ) higher expression  
497 than Control. No significant differences ( $p>0.05$ ) in the expression of the *elovl5* and *fads2* genes  
498 (Fig. 10g,h) were observed among the Control, Hi25, Hi50 and Hi75 groups.

499

500 *Appetite.* As shown in Figure 10i,l, the higher was the BSF meal inclusion in the diets, the higher  
501 was the *ghrl* and *npy* gene expression. However, no significant differences ( $p>0.05$ ) were observed  
502 between Control and Hi25 in terms of *ghrl* gene expression and among Control, Hi25 and Hi50 in  
503 terms of *npy* gene expression. Figure 10m shows *cnrl* gene expression. No BSF meal dose  
504 dependency was observed in the expression of this specific gene, since only Hi25 and Hi50 groups  
505 evidenced significantly ( $p<0.05$ ) higher values than Control. , Groups fed on the highest BSF  
506 inclusion (Hi75 and Hi100) showed significantly ( $p<0.05$ ) higher gene expression of the *lepa* gene  
507 (Fig. 10n) compared to the other groups, while no significant differences ( $p<0.05$ ) were detected  
508 among Control, Hi25 and Hi50 groups.

509

510 *Immune response.* Considering genes involved in the immune response, higher BSF meal dietary  
511 inclusions resulted in a significantly ( $p<0.05$ ) higher gene expression compared to Control.  
512 Specifically, no significant differences ( $p>0.05$ ) were evident in the expression of the *il1b* gene (Fig  
513 10o) among the Control, Hi25 and Hi50 groups, while the Hi75 and Hi100 groups showed  
514 significantly ( $p<0.05$ ) higher values compared to the other experimental groups. Similarly, the  
515 Hi50, Hi75 and Hi100 groups showed a significantly ( $p<0.05$ ) higher expression of the *il10* gene  
516 (Fig 10p) compared to the Control and Hi25 groups, which did not differ significantly from each  
517 other ( $p>0.05$ ). Finally, as reported in Figure 10q, the increasing levels of inclusion of BSF meal in  
518 the diets resulted in a statistically significant ( $p<0.05$ ) dose-dependent increase in *tnfa* gene  
519 expression.

520

521 *Chitinases.* All the experimental groups fed on BSF-based diets showed an increase in chitinases  
522 gene expression (Fig. 10r,s). Specifically, no significant differences ( $p>0.05$ ) were evident in the  
523 expression of the *chia.2* and *chia.3* genes among all the experimental groups, with the exception of  
524 the Hi100 group for *chia.2* and both Hi75 and Hi100 groups for *chia.3*, which showed a  
525 significantly ( $p<0.05$ ) higher gene expression than Control.

526

#### 527 **4. Discussion**

528 The use of insects in aquafeed still faces possible limitations because of their lack in PUFAs  
529 (Barroso et al., 2017) and presence of chitin (Kroeckel et al., 2012; Xiao et al., 2018).

530 Insects are usually farmed on vegetable organic by-products, and previous studies showed that the  
531 final insect biomass had a very low PUFAs and a high SFA content (Vargas et al., 2018;  
532 Zarantoniello et al., 2018). However, it is now well established that the quantity and quality of FAs  
533 in insects can be modified by the growth substrate (Barroso et al., 2014; Spranghers et al., 2017; St-  
534 Hilaire et al., 2007a). Specifically, in the present study we demonstrated that the addition of a 10%



(w/w) of *Schyzochitrium* sp. to the growth substrate is an efficient approach to increase the PUFAs content of the final insect biomass.

This “enrichment” procedure of the insect biomass caused a progressive increase in PUFAs content in the Hi25, 50, 75 and 100 diets, respectively, highlighting that the enrichment method used is appropriate and represents, in this sense, an interesting and innovative approach on how food by-products produced on land (coffee silverskin) can be enriched and bio-converted into a valuable biomass for aquafeed production (Meneguz et al., 2018; Pinotti et al., 2019).

Compared to previous insect-based diets already tested on zebrafish (Vargas et al., 2018; Zarantoniello et al., 2019), the diets tested in this study showed good performances on fish growth, since higher BSF meal inclusion levels in the experimental diets resulted in higher larval SGR%. This result was supported by the analysed molecular markers of growth but is in contrast with previous studies reporting that BSF meal inclusion levels higher than 40% often resulted in negative effects on fish growth and welfare (Sánchez-Muros et al., 2014; Barragan-Fonseca et al., 2018; Secci et al., 2019). In particular, previous studies evidenced that such high inclusion levels affected both intestine and liver integrity and, thus, dietary nutrients absorption (Henry et al., 2015; Li et al., 2017; Cardinaletti et al., 2019).

The different diets tested in the present study also affected larval fish FA composition. However, differences among FA classes were less evident in zebrafish larvae compared to those detected in the diets. In fact, as a freshwater species, zebrafish are able to convert shorter-chain FAs in highly unsaturated ones through the elongation and desaturation pathways (Tocher, 2010). This was clearly documented in the present study by a higher *elovl2*, *elovl5* and *fads* gene expression in all the experimental groups fed on BSF meal-based diets (with emphasis on the Hi100 group).

Gas chromatographic results were also confirmed and integrated by FTIR analyses which provided data about the macromolecular composition of the analysed biological samples (Giorgini et al., 2018). Specifically, FTIR analyses showed that there were no differences in the total amount of proteins (PRT/TBM ratio) (with the exception of collagen (1234/PRT ratio) that slightly decreased)

among the experimental groups, highlighting that BSF meal is a valuable protein source for aquafeeds (Al-Qazzaz and Ismail, 2016; Lock et al., 2016; Nogales-Mérida et al., 2018).

Concerning lipids, the same analyses revealed that the increase of BSF meal percentage in the diets was associated with a general increase in total larval lipid content. Additionally, analysing specific wavelengths, FTIR confirmed the gas chromatographic results, evidencing an overall increase in saturated FA (1744/LIP and 2925/LIP ratios) and a decrease in unsaturated (3010/LIP ratio).

Molecular markers related to the appetite stimulus were fully supported by the biometric results. Fish food intake is regulated by specific regions in the brain that interpret and integrate positive (orexigenic) and negative (anorexigenic) signals derived from the hypothalamic area (neurohormones) and from the body periphery (Copeland et al., 2011; Sobrino Crespo et al., 2014).

Specifically, the orexigenic signals *ghrl* and *npv* analysed in the present study, showed a dose-dependent gene expression increase related to the BSF meal inclusion levels in the diets (the higher the BSF meal inclusion level, the higher their gene expression), while the *cnrl* gene expression was always higher in all the larval groups fed on BSF-based diets. Conversely, the results obtained from the anorexigenic signal *lepa* was not obvious. Leptin is usually involved in the inhibition of orexigenic pathways and in the stimulation of anorexigenic (Piccinetti et al., 2010; Volkoff, 2006); in the present study, however, an opposite activity was evidenced.

This anomalous pattern could be explained by the FTIR analyses that revealed an increasing total lipid content (LIP/TBM ratio) in larvae fed on diets with increasing BSF meal levels. In fact, previous studies, performed both on mammals and fish species, demonstrated a positive correlation between leptin levels and amount of adipose tissue (Chisada et al., 2014; Park and Ahima, 2015; Li et al., 2016).

Alternatively, an interconnection between leptin and proinflammatory cytokines exist (Lafrance et al., 2010). Leptin is known to enhance the production of proinflammatory cytokines, as well as proinflammatory cytokines are able to trigger leptin release (Carlton et al., 2012). In the present study, the increasing *lepa* gene expression observed in larvae fed on insect-based diets might have

587 promoted the immune-response observed in the same experimental groups. However, it should be  
588 mentioned that all insect-based diets contain chitin (Borrelli et al., 2017; Gasco et al., 2018), a  
589 molecule which still has a controversial role in aquafeeds. Indeed, some studies reported a  
590 beneficial modulatory role of chitin on fish immune system and microbiota (Ringø et al., 2012;  
591 Zhang et al., 2012; Zhou et al., 2013; Bruni et al., 2018; Henry et al., 2018), while others showed  
592 that, especially at high inclusion levels, chitin may induce inflammation of the intestinal tract and a  
593 reduction in nutrient assimilation (Kroeckel et al., 2012; Magalhães et al., 2017; Su et al., 2017;  
594 Xiao et al., 2018). In support of this conclusion, Hi50, Hi75 and Hi100 larvae showed a significant  
595 increase of mucin (1055/PRT ratio), possibly associated to a higher intestine lubrication necessary  
596 for a proper intestinal transit of these diets.

597 In the present study, no specific inflammatory events were detected through the histological  
598 analysis of the intestine samples, suggesting: 1) a possible digestion of chitin by zebrafish larvae  
599 through specific chitinases (*chia.2* and *chia.3*); 2) a possible intestinal anti-inflammatory role of  
600 lauric acid (C12). This fatty acid, which in the present study increased its dietary amount with the  
601 increasing BSF meal inclusion in the diets, has been demonstrated to have beneficial effects on fish  
602 gut's welfare by mitigating inflammatory conditions (Aleström et al., 2006; Dahm and Geisler,  
603 2006; De-Santis and Jerry, 2007; Zarantoniello et al., 2019).

604 The histological analysis of the liver showed a lipid accumulation in the hepatic parenchyma in all  
605 analysed samples, regardless of the dietary treatment. Specifically, the Hi75 and Hi100 groups  
606 showed a severe level of hepatic steatosis, a pathological condition that has previously been related  
607 to a high n-6/n-3 ratio (Di Minno et al., 2012; Leamy et al., 2013; Zarantoniello et al., 2018, 2019).  
608 This hepatic disorder is probably the cause of the higher gene expression of the stress markers  
609 (*nr3c1* and *hsp70.1*). Finally, insects are known to possess natural bioactive molecules that are  
610 known to modulate fish microbiota (Huyben et al., 2019; Terova et al., 2019) and, therefore, the  
611 bacterial community is expected to impact host metabolism and health status.

612 Although autochthonous bacterial communities were dominated by the same phyla regardless of the  
613 diet, the present study showed that bacterial populations are dynamic and can be modulated by  
614 dietary inclusion of insect meal.

615 Some bacteria were almost present in all the samples analysed, while others were mainly modulated  
616 by the insect-based diets. Among the ubiquitous species, the massive presence of *Cetobacterium* in  
617 all gut samples suggested that graded BSF inclusion levels in the diets did not influence the  
618 presence of such a fish associated microorganism. In fact, *Cetobacterium* has already been found as  
619 core genus in the gut of zebrafish with beneficial effect on fish health because of its ability to  
620 produce cobalamin (vitamin B12) (Roeselers et al., 2011; Merrifield et al., 2013; Ghanbari et al.,  
621 2015; Earley et al., 2018;). Mycoplasmataceae were also dominant in the present study, and have  
622 already been found among the dominant microorganisms in fish intestines (e.g. rainbow trout), with  
623 a possible beneficial action on host health by producing lactic acid with antibacterial effect  
624 (Rimoldi et al., 2019). Finally, Aeromonadaceae were also detected in all analysed gut samples,  
625 irrespective of the diet used. This bacterial family has already been found as naturally associated  
626 with the gut of fish or insects (Grabowski and Klein, 2017; Udayangani et al., 2017; Rimoldi et al.,  
627 2019), suggesting a contribution of BSF-based diets in the occurrence of such a bacterial family in  
628 the gut of the analysed zebrafish.

629 BSF meal inclusion in the diets modified the presence of some bacteria. Specifically, the presence  
630 of *Vibrio* was negatively influenced by the addition of increasing graded levels of BSF meal in the  
631 diets. As reported by Brugman et al., (2015), in the zebrafish intestine T lymphocytes can control  
632 the outgrowth of *Vibrio* species. As shown by the Real Time PCR analyses, the supply of BSF-  
633 based diets stimulated immune-response related genes that might be implied in *Vibrio* species  
634 control. However, further research is needed to better understand the involvement of BSF in this  
635 aspect. Finally, both *Ochrobactrum* and *Tetrathiobacter* were detected in the samples GH100. The  
636 occurrence of *Ochrobactrum* was already reported in zebrafish gut (Cantas et al., 2012) as well as in  
637 the microbiota of insects as coleoptera and lepidoptera (Grabowski and Klein, 2017), while

638 *Tetrathibacter* has already been isolated from the midgut of ticks (Li et al., 2014). These data  
639 suggest that the inclusion of BSF meal in fish diets can contribute to modify the fish microbiota  
640 with a specific possible involvement in the allochthonous microbiota modification.

641 As a general remark about gut microbiota, bacterial diversity is considered a positive indicator of  
642 gut health while a reduced diversity is often associated to dysbiosis and risk of disease in fish  
643 (Terova et al., 2019). In the present study, the PCA analyses revealed a difference between the Hi0-  
644 25 and Hi50-75-100 groups, thus suggesting a possible influence of BFS-based diets on the  
645 zebrafish gut microbiota, to be further investigated.

646

647

## 648 **5. Conclusion**

649 In order to meet aquafeed requirements, BSF biomass should be enriched in PUFAs. The present  
650 study showed that the addition of a 10% (W/W) *Schizochytrium* sp to the growth substrate is a valid  
651 method to achieve this objective. The present study evidenced that a 50% BSF meal inclusion level  
652 in the diet sustains a better fish growth and does not have any major negative effects on the fish.  
653 Higher inclusion levels affect larval liver histology and induce a general increase in lipid  
654 accumulation and stress response. This novel approach represents an interesting example of how, in  
655 the long term, a circular economy applied to the aquaculture sector may sustain animal's welfare  
656 and encourage sustainability and competitiveness.

657

## 658 **6. Acknowledgments**

659 This study was funded by, Ricerca Scientifica 2017 Cariverona, NUTRIFISH project N° 2017.0571  
660 and the authors wish to thank Prof. F. Rindi for the English revision.

661

## 662 **7. References**

663 Al-Qazzaz, M.F., Ismail, D.B., 2016. Insect meal as a source of protein in animal diet. Anim. Nutr.  
664 Feed Technol. 16, 527–547. <https://doi.org/10.5958/0974-181X.2016.00038.X>

665 Aleström, P., Holter, J.L., Nourizadeh-Lillabadi, R., 2006. Zebrafish in functional genomics and  
666 aquatic biomedicine. *Trends Biotechnol.* <https://doi.org/10.1016/j.tibtech.2005.11.004>

667 Alhazzaa, R., Nichols, P.D., Carter, C.G., 2018. Sustainable alternatives to dietary fish oil in  
668 tropical fish aquaculture. *Rev. Aquac.* <https://doi.org/10.1111/raq.12287>

669 Ayadi, F.Y., Rosentrate, K.A., Muthukumar, K., 2012. Alternative Protein Sources for Aquaculture  
670 Feeds. *J. Aquac. Feed Sci. Nutr.* 4, 1–26. <https://doi.org/10.3923/joafsnu.2012.1.26>

671 Bandara, T., 2018. Alternative feed ingredients in aquaculture: Opportunities and challenges. *J.*  
672 *Entomol. Zool. Stud.* 6, 3087–3094.

673 Barragan-Fonseca, K.B., Dicke, M., van Loon, J.J.A., 2018. Influence of larval density and dietary  
674 nutrient concentration on performance, body protein, and fat contents of black soldier fly  
675 larvae (*Hermetia illucens*). *Entomol. Exp. Appl.* <https://doi.org/10.1111/eea.12716>

676 Barroso, F.G., de Haro, C., Sánchez-Muros, M.J., Venegas, E., Martínez-Sánchez, A., Pérez-Bañón,  
677 C., 2014. The potential of various insect species for use as food for fish. *Aquaculture* 422–423,  
678 193–201. <https://doi.org/10.1016/j.aquaculture.2013.12.024>

679 Barroso, F.G., Sánchez-Muros, M.J., Segura, M., Morote, E., Torres, A., Ramos, R., Guil, J.L.,  
680 2017. Insects as food: Enrichment of larvae of *Hermetia illucens* with omega 3 fatty acids by  
681 means of dietary modifications. *J. Food Compos. Anal.*  
682 <https://doi.org/10.1016/j.jfca.2017.04.008>

683 Belghit, I., Liland, N.S., Gjesdal, P., Biancarosa, I., Menchetti, E., Li, Y., Waagbø, R., Krogdahl,  
684 Å., Lock, E.J., 2019. Black soldier fly larvae meal can replace fish meal in diets of sea-water  
685 phase Atlantic salmon (*Salmo salar*). *Aquaculture.*  
686 <https://doi.org/10.1016/j.aquaculture.2018.12.032>

687 Berggren, Å., Jansson, A., Low, M., 2019. Approaching Ecological Sustainability in the Emerging  
688 Insects-as-Food Industry. *Trends Ecol. Evol.* <https://doi.org/10.1016/j.tree.2018.11.005>

689 Bohnes, F.A., Laurent, A., 2019. LCA of aquaculture systems: methodological issues and potential  
690 improvements. *Int. J. Life Cycle Assess.* <https://doi.org/10.1007/s11367-018-1517-x>

691 Borrelli, L., Coretti, L., Dipineto, L., Bovera, F., Menna, F., Chiariotti, L., Nizza, A., Lembo, F.,  
692 Fioretti, A., 2017. Insect-based diet, a promising nutritional source, modulates gut microbiota  
693 composition and SCFAs production in laying hens. *Sci. Rep.* 7, 16269.  
694 <https://doi.org/10.1038/s41598-017-16560-6>

695 Brugman, S., Schneeberger, K., Witte, M., Klein, M.R., van den Bogert, B., Boekhorst, J.,  
696 Timmerman, H.M., Boes, M.L., Kleerebezem, M., Nieuwenhuis, E.E.S., 2015. T lymphocytes  
697 control microbial composition by regulating the abundance of *Vibrio* in the zebrafish gut. *Gut*  
698 *Microbes.* <https://doi.org/10.4161/19490976.2014.972228>

699 Bruni, L., Pastorelli, R., Viti, C., Gasco, L., Parisi, G., 2018. Characterisation of the intestinal  
700 microbial communities of rainbow trout (*Oncorhynchus mykiss*) fed with *Hermetia illucens*  
701 (black soldier fly) partially defatted larva meal as partial dietary protein source. *Aquaculture*  
702 487, 56–63. <https://doi.org/10.1016/J.AQUACULTURE.2018.01.006>

703 Caligiani, A., Marseglia, A., Leni, G., Baldassarre, S., Maistrello, L., Dossena, A., Sforza, S., 2018.  
704 Composition of black soldier fly prepupae and systematic approaches for extraction and  
705 fractionation of proteins, lipids and chitin. *Food Res. Int.*  
706 <https://doi.org/10.1016/j.foodres.2017.12.012>

707 Canonico, L., Ashoor, S., Taccari, M., Comitini, F., Antonucci, M., Truzzi, C., Scarponi, G., Ciani,

708 M., 2016. Conversion of raw glycerol to microbial lipids by new *Metschnikowia* and *Yarrowia*  
709 *lipolytica* strains. *Ann. Microbiol.* <https://doi.org/10.1007/s13213-016-1228-0>

710 Cantas, L., Sørby, J.R.T., Aleström, P., Sørum, H., 2012. Culturable Gut Microbiota Diversity in  
711 Zebrafish. *Zebrafish.* <https://doi.org/10.1089/zeb.2011.0712>

712 Caporaso, J.G., Kuczynski, J., Stombaugh, J., Bittinger, K., Bushman, F.D., Costello, E.K., Fierer,  
713 N., Peña, A.G., Goodrich, J.K., Gordon, J.I., Huttley, G.A., Kelley, S.T., Knights, D., Koenig,  
714 J.E., Ley, R.E., Lozupone, C.A., McDonald, D., Muegge, B.D., Pirrung, M., Reeder, J.,  
715 Sevinsky, J.R., Turnbaugh, P.J., Walters, W.A., Widmann, J., Yatsunenko, T., Zaneveld, J.,  
716 Knight, R., 2010. QIIME allows analysis of high-throughput community sequencing data. *Nat.*  
717 *Methods.* <https://doi.org/10.1038/nmeth.f.303>

718 Cardinaletti, G., Messina, M., Bruno, M., Tulli, F., Poli, B.M., Giorgi, G., Chini-Zittelli, G.,  
719 Tredici, M., Tibaldi, E., 2018. Effects of graded levels of a blend of *Tisochrysis lutea* and  
720 *Tetraselmis suecica* dried biomass on growth and muscle tissue composition of European sea  
721 bass (*Dicentrarchus labrax*) fed diets low in fish meal and oil. *Aquaculture* 485, 173–182.  
722 <https://doi.org/10.1016/J.AQUACULTURE.2017.11.049>

723 Cardinaletti, G., Randazzo, B., Messina, M., Zarantoniello, M., Giorgini, E., Zimbelli, A., Bruni, L.,  
724 Parisi, G., Olivotto, I., Tulli, F., 2019. Effects of Graded Dietary Inclusion Level of Full-Fat  
725 *Hermetia illucens* Prepupae Meal in Practical Diets for Rainbow Trout (*Oncorhynchus*  
726 *mykiss*). *Animals* 9, 251. <https://doi.org/10.3390/ani9050251>

727 Carlton, E.D., Demas, G.E., French, S.S., 2012. Leptin, a neuroendocrine mediator of immune  
728 responses, inflammation, and sickness behaviors. *Horm. Behav.* 62, 272–279.  
729 <https://doi.org/10.1016/j.yhbeh.2012.04.010>

730 Chisada, S., Kurokawa, T., Murashita, K., Rønnestad, I., Taniguchi, Y., Toyoda, A., Sakaki, Y.,  
731 Takeda, S., Yoshiura, Y., 2014. Leptin receptor-deficient (knockout) medaka, *Oryzias latipes*,  
732 show chronical up-regulated levels of orexigenic neuropeptides, elevated food intake and stage  
733 specific effects on growth and fat allocation. *Gen. Comp. Endocrinol.* 195, 9–20.  
734 <https://doi.org/10.1016/J.YGCEN.2013.10.008>

735 Čičková, H., Newton, G.L., Lacy, R.C., Kozánek, M., 2015. The use of fly larvae for organic waste  
736 treatment. *Waste Manag.* 35, 68–80. <https://doi.org/10.1016/J.WASMAN.2014.09.026>

737 Copeland, D.L., Duff, R.J., Liu, Q., Prokop, J., Londraville, R.L., 2011. Leptin in teleost fishes: An  
738 argument for comparative study. *Front. Physiol.* <https://doi.org/10.3389/fphys.2011.00026>

739 Dahm, R., Geisler, R., 2006. Learning from small fry: The zebrafish as a genetic model organism  
740 for aquaculture fish species. *Mar. Biotechnol.* <https://doi.org/10.1007/s10126-006-5139-0>

741 Daniel, D., 2018. A review on replacing fish meal in aqua feeds using plant protein sources. *Int. J.*  
742 *Fish. Aquat. Stud.*

743 De-Santis, C., Jerry, D.R., 2007. Candidate growth genes in finfish - Where should we be looking?  
744 *Aquaculture.* <https://doi.org/10.1016/j.aquaculture.2007.08.036>

745 Diener, S., Zurbrugg, C., Tockner, K., 2009. Conversion of organic material by black soldier fly  
746 larvae: Establishing optimal feeding rates. *Waste Manag. Res.*  
747 <https://doi.org/10.1177/0734242X09103838>

748 Dumas, A., Raggi, T., Barkhouse, J., Lewis, E., Weltzien, E., 2018. The oil fraction and partially  
749 defatted meal of black soldier fly larvae (*Hermetia illucens*) affect differently growth  
750 performance, feed efficiency, nutrient deposition, blood glucose and lipid digestibility of

751 rainbow trout (*Oncorhynchus mykiss*). Aquaculture.  
752 <https://doi.org/10.1016/j.aquaculture.2018.03.038>

753 Earley, A.M., Graves, C.L., Shiau, C.E., 2018. Critical Role for a Subset of Intestinal Macrophages  
754 in Shaping Gut Microbiota in Adult Zebrafish. Cell Rep.  
755 <https://doi.org/10.1016/j.celrep.2018.09.025>

756 Edgar, R.C., Haas, B.J., Clemente, J.C., Quince, C., Knight, R., 2011. UCHIME improves  
757 sensitivity and speed of chimera detection. Bioinformatics.  
758 <https://doi.org/10.1093/bioinformatics/btr381>

759 FAO, 2018. State of Fisheries and Aquaculture in the world. State World Fish. Aquac. 2018 - Meet.  
760 Sustain. Dev. goals. <https://doi.org/issn 10>

761 FOLCH, J., LEES, M., SLOANE STANLEY, G.H., 1957. A simple method for the isolation and  
762 purification of total lipides from animal tissues. J. Biol. Chem.

763 Francis, G., Makkar, H.P.S., Becker, K., 2001. Antinutritional factors present in plant-derived  
764 alternate fish feed ingredients and their effects in fish. Aquaculture.  
765 [https://doi.org/10.1016/S0044-8486\(01\)00526-9](https://doi.org/10.1016/S0044-8486(01)00526-9)

766 Garofalo, C., Bancalari, E., Milanović, V., Cardinali, F., Osimani, A., Sardaro, M.L.S., Bottari, B.,  
767 Bernini, V., Aquilanti, L., Clementi, F., Neviani, E., Gatti, M., 2017. Study of the bacterial  
768 diversity of foods: PCR-DGGE versus LH-PCR. Int. J. Food Microbiol.  
769 <https://doi.org/10.1016/j.ijfoodmicro.2016.11.008>

770 Gasco, L., Finke, M., van Huis, A., 2018. Can diets containing insects promote animal health? J.  
771 Insects as Food Feed 4, 1–4. <https://doi.org/10.3920/JIFF2018.x001>

772 Gerland, P., Raftery, A.E., Ševčíková, H., Li, N., Gu, D., Spoorenberg, T., Alkema, L., Fosdick,  
773 B.K., Chunn, J., Lalic, N., Bay, G., Buettner, T., Heilig, G.K., Wilmoth, J., 2014. World  
774 population stabilization unlikely this century. Science (80-. ).  
775 <https://doi.org/10.1126/science.1257469>

776 Ghanbari, M., Kneifel, W., Domig, K.J., 2015. A new view of the fish gut microbiome: Advances  
777 from next-generation sequencing. Aquaculture.  
778 <https://doi.org/10.1016/j.aquaculture.2015.06.033>

779 Giorgini, E., Randazzo, B., Gioacchini, G., Cardinaletti, G., Vaccari, L., Tibaldi, E., Olivotto, I.,  
780 2018. New insights on the macromolecular building of rainbow trout (*O. mykiss*) intestine:  
781 FTIR Imaging and histological correlative study. Aquaculture 497, 1–9.  
782 <https://doi.org/10.1016/j.aquaculture.2018.07.032>

783 Grabowski, N.T., Klein, G., 2017. Bacteria encountered in raw insect, spider, scorpion, and  
784 centipede taxa including edible species, and their significance from the food hygiene point of  
785 view. Trends Food Sci. Technol. <https://doi.org/10.1016/j.tifs.2017.01.007>

786 Guillen, J., Natale, F., Carvalho, N., Casey, J., Hofherr, J., Druon, J.N., Fiore, G., Gibin, M., Zanzi,  
787 A., Martinsohn, J.T., 2018. Global seafood consumption footprint. Ambio.  
788 <https://doi.org/10.1007/s13280-018-1060-9>

789 Henry, M., Gasco, L., Piccolo, G., Fountoulaki, E., 2015. Review on the use of insects in the diet of  
790 farmed fish: Past and future. Anim. Feed Sci. Technol.  
791 <https://doi.org/10.1016/j.anifeedsci.2015.03.001>

792 Henry, M.A., Gasco, L., Chatzifotis, S., Piccolo, G., 2018. Does dietary insect meal affect the fish  
793 immune system? The case of mealworm, *Tenebrio molitor* on European sea bass,



794 *Dicentrarchus labrax*. *Dev. Comp. Immunol.* 81, 204–209.  
795 <https://doi.org/10.1016/J.DCI.2017.12.002>

796 Huyben, D., Vidaković, A., Werner Hallgren, S., Langeland, M., 2019. High-throughput sequencing  
797 of gut microbiota in rainbow trout (*Oncorhynchus mykiss*) fed larval and pre-pupae stages of  
798 black soldier fly (*Hermetia illucens*). *Aquaculture*.  
799 <https://doi.org/10.1016/j.aquaculture.2018.10.034>

800 Jin, G., Yang, F., Hu, C., Shen, H., Zhao, Z.K., 2012. Enzyme-assisted extraction of lipids directly  
801 from the culture of the oleaginous yeast *Rhodospiridium toruloides*. *Bioresour. Technol.*  
802 <https://doi.org/10.1016/j.biortech.2012.01.152>

803 Johnston, I.A., Macqueen, D.J., Watabe, S., 2008. Molecular Biotechnology of Development and  
804 Growth in Fish Muscle, in: *Fisheries for Global Welfare and Environment*.

805 Klindworth, A., Pruesse, E., Schweer, T., Peplies, J., Quast, C., Horn, M., Glöckner, F.O., 2013.  
806 Evaluation of general 16S ribosomal RNA gene PCR primers for classical and next-generation  
807 sequencing-based diversity studies. *Nucleic Acids Res.* <https://doi.org/10.1093/nar/gks808>

808 Komprda, T., Zorníková, G., Rozíková, V., Borkovcová, M., Przywarová, A., 2013. The effect of  
809 dietary *Salvia hispanica* seed on the content of n-3 long-chain polyunsaturated fatty acids in  
810 tissues of selected animal species, including edible insects. *J. Food Compos. Anal.*  
811 <https://doi.org/10.1016/j.jfca.2013.06.010>

812 Kroeckel, S., Harjes, A.G.E., Roth, I., Katz, H., Wuertz, S., Susenbeth, A., Schulz, C., 2012. When  
813 a turbot catches a fly: Evaluation of a pre-pupae meal of the Black Soldier Fly (*Hermetia*  
814 *illucens*) as fish meal substitute - Growth performance and chitin degradation in juvenile turbot  
815 (*Psetta maxima*). *Aquaculture*. <https://doi.org/10.1016/j.aquaculture.2012.08.041>

816 Lafrance, V., Inoue, W., Kan, B., Luheshi, G.N., 2010. Leptin modulates cell morphology and  
817 cytokine release in microglia. *Brain. Behav. Immun.* 24, 358–365.  
818 <https://doi.org/10.1016/j.bbi.2009.11.003>

819 Lawrence, C., 2007. The husbandry of zebrafish (*Danio rerio*): A review. *Aquaculture* 269, 1–20.  
820 <https://doi.org/10.1016/j.aquaculture.2007.04.077>

821 Lawrence, C., Adatto, I., Best, J., James, A., Maloney, K., 2012. Generation time of zebrafish  
822 (*Danio rerio*) and medakas (*Oryzias latipes*) housed in the same aquaculture facility. *Lab*  
823 *Anim. (NY)*. <https://doi.org/10.1038/labon0612-158>

824 Li, A., Yuan, X., Liang, X.-F., Liu, L., Li, Jie, Li, B., Fang, J., Li, Jiao, He, S., Xue, M., Wang, J.,  
825 Tao, Y.-X., 2016. Adaptations of lipid metabolism and food intake in response to low and high  
826 fat diets in juvenile grass carp (*Ctenopharyngodon idellus*). *Aquaculture* 457, 43–49.  
827 <https://doi.org/10.1016/J.AQUACULTURE.2016.01.014>

828 Li, C.H., cao, J., Zhou, Y.Z., Zhang, H.S., Gong, H.Y., Zhou, J.L., 2014. The midgut bacterial flora  
829 of laboratory-reared hard ticks, *haemaphysalis longicornis*, *hyalomma asiaticum*, and  
830 *rhhipicephalus haemaphysaloides*. *J. Integr. Agric.* [https://doi.org/10.1016/S2095-3119\(13\)60517-1](https://doi.org/10.1016/S2095-3119(13)60517-1)

832 Li, S., Ji, H., Zhang, B., Zhou, J., Yu, H., 2017. Defatted black soldier fly (*Hermetia illucens*) larvae  
833 meal in diets for juvenile Jian carp (*Cyprinus carpio* var. Jian): Growth performance,  
834 antioxidant enzyme activities, digestive enzyme activities, intestine and hepatopancreas  
835 histological structure. *Aquaculture* 477, 62–70.  
836 <https://doi.org/10.1016/J.AQUACULTURE.2017.04.015>

837 Li, W., Li, Q., Zheng, L., Wang, Y., Zhang, J., Yu, Z., Zhang, Y., 2015. Potential biodiesel and  
838 biogas production from corn cob by anaerobic fermentation and black soldier fly. *Bioresour.*  
839 *Technol.* <https://doi.org/10.1016/j.biortech.2015.06.112>

840 Li, Y., Kortner, T.M., Chikwati, E.M., Munang'andu, H.M., Lock, E.J., Krogdahl, Å., 2019. Gut  
841 health and vaccination response in pre-smolt Atlantic salmon (*Salmo salar*) fed black soldier  
842 fly (*Hermetia illucens*) larvae meal. *Fish Shellfish Immunol.*  
843 <https://doi.org/10.1016/j.fsi.2018.12.057>

844 Liland, N.S., Biancarosa, I., Araujo, P., Biemans, D., Bruckner, C.G., Waagbø, R., Torstensen,  
845 B.E., Lock, E.J., 2017. Modulation of nutrient composition of black soldier fly (*Hermetia*  
846 *illucens*) larvae by feeding seaweed-enriched media. *PLoS One.*  
847 <https://doi.org/10.1371/journal.pone.0183188>

848 Lock, E.R., Arsiwalla, T., Waagbø, R., 2016. Insect larvae meal as an alternative source of nutrients  
849 in the diet of Atlantic salmon ( *Salmo salar* ) postsmolt. *Aquac. Nutr.* 22, 1202–1213.  
850 <https://doi.org/10.1111/anu.12343>

851 Magalhães, R., Sánchez-López, A., Leal, R.S., Martínez-Llorens, S., Oliva-Teles, A., Peres, H.,  
852 2017. Black soldier fly (*Hermetia illucens*) pre-pupae meal as a fish meal replacement in diets  
853 for European seabass (*Dicentrarchus labrax*). *Aquaculture* 476, 79–85.  
854 <https://doi.org/10.1016/J.AQUACULTURE.2017.04.021>

855 Magoč, T., Salzberg, S.L., 2011. FLASH: Fast length adjustment of short reads to improve genome  
856 assemblies. *Bioinformatics.* <https://doi.org/10.1093/bioinformatics/btr507>

857 Makkar, H.P.S., Tran, G., Heuzé, V., Ankers, P., 2014. State-of-the-art on use of insects as animal  
858 feed. *Anim. Feed Sci. Technol.* <https://doi.org/10.1016/j.anifeedsci.2014.07.008>

859 Manceron, S., Ben-Ari, T., Dumas, P., 2014. Feeding proteins to livestock: Global land use and  
860 food vs. feed competition . *OCL.* <https://doi.org/10.1051/ocl/2014020>

861 May, B.M., 1961. The occurrence in New Zealand and the life-history of the soldier fly *Hermetia*  
862 *illucens* (L.) (Diptera: Stratiomyidae). *New Zeal. J. Sci.*

863 Meneguz, M., Schiavone, A., Gai, F., Dama, A., Lussiana, C., Renna, M., Gasco, L., 2018. Effect of  
864 rearing substrate on growth performance, waste reduction efficiency and chemical composition  
865 of black soldier fly (*Hermetia illucens*) larvae. *J. Sci. Food Agric.*  
866 <https://doi.org/10.1002/jsfa.9127>

867 Merino, G., Barange, M., Blanchard, J.L., Harle, J., Holmes, R., Allen, I., Allison, E.H., Badjeck,  
868 M.C., Dulvy, N.K., Holt, J., Jennings, S., Mullon, C., Rodwell, L.D., 2012. Can marine  
869 fisheries and aquaculture meet fish demand from a growing human population in a changing  
870 climate? *Glob. Environ. Chang.* <https://doi.org/10.1016/j.gloenvcha.2012.03.003>

871 Merrifield, D.L., Shaw, B.J., Harper, G.M., Saoud, I.P., Davies, S.J., Handy, R.D., Henry, T.B.,  
872 2013. Ingestion of metal-nanoparticle contaminated food disrupts endogenous microbiota in  
873 zebrafish (*Danio rerio*). *Environ. Pollut.* <https://doi.org/10.1016/j.envpol.2012.11.017>

874 Müller, A., Wolf, D., Gutzeit, H.O., 2017. The black soldier fly, *Hermetia illucens* – a promising  
875 source for sustainable production of proteins, lipids and bioactive substances. *Zeitschrift für*  
876 *Naturforsch. C* 72, 351–363. <https://doi.org/10.1515/znc-2017-0030>

877 Murthy, P.S., Naidu, M.M., 2012. Resources , Conservation and Recycling Sustainable  
878 management of coffee industry by-products and value addition — A review. *Resour. Conserv.*  
879 *Recycl.* 66, 45–58. <https://doi.org/10.1016/j.resconrec.2012.06.005>

880 Mussatto, S.I., Machado, E.M.S., Martins, S., Teixeira, J.A., 2011. Production, Composition, and  
881 Application of Coffee and Its Industrial Residues. *Food Bioprocess Technol.* 4, 661–672.  
882 <https://doi.org/10.1007/s11947-011-0565-z>

883 Naylor, R.L., Hardy, R.W., Bureau, D.P., Chiu, A., Elliott, M., Farrell, A.P., Forster, I., Gatlin,  
884 D.M., Goldburg, R.J., Hua, K., Nichols, P.D., 2009. Feeding aquaculture in an era of finite  
885 resources. *Proc. Natl. Acad. Sci.* <https://doi.org/10.1073/pnas.0905235106>

886 Nogales-Mérida, S., Gobbi, P., Józefiak, D., Mazurkiewicz, J., Dudek, K., Rawski, M., Kierończyk,  
887 B., Józefiak, A., 2018. Insect meals in fish nutrition. *Rev. Aquac.* 1–24.  
888 <https://doi.org/10.1111/raq.12281>

889 Official Methods of Analysis of the Association of Official Analytical Chemists, 2006. . *J. Pharm.*  
890 *Sci.* <https://doi.org/10.1002/jps.2600600253>

891 Olivotto, I., Di Stefano, M., Rosetti, S., Cossignani, L., Pugnali, A., Giantomassi, F., Carnevali,  
892 O., 2011. Live prey enrichment, with particular emphasis on HUFAs, as limiting factor in false  
893 percula clownfish (*Amphiprion ocellaris*, Pomacentridae) larval development and  
894 metamorphosis: Molecular and biochemical implications. *Comp. Biochem. Physiol. - A Mol.*  
895 *Integr. Physiol.* <https://doi.org/10.1016/j.cbpa.2011.02.004>

896 Olivotto, I., Mosconi, G., Maradonna, F., Cardinali, M., Carnevali, O., 2002. *Diplodus sargus*  
897 interrenal-pituitary response: Chemical communication in stressed fish. *Gen. Comp.*  
898 *Endocrinol.* [https://doi.org/10.1016/S0016-6480\(02\)00024-2](https://doi.org/10.1016/S0016-6480(02)00024-2)

899 Olivotto, I., Yasumasu, S., Gioacchini, G., Maradonna, F., Cionna, C., Carnevali, O., 2004. Cloning  
900 and expression of high choriolytic enzyme, a component of the hatching enzyme system,  
901 during embryonic development of the marine ornamental fish *Chrysiptera parasema*. *Mar.*  
902 *Biol.* <https://doi.org/10.1007/s00227-004-1404-9>

903 Osimani, A., Ferrocino, I., Agnolucci, M., Cocolin, L., Giovannetti, M., Cristani, C., Palla, M.,  
904 Milanović, V., Roncolini, A., Sabbatini, R., Garofalo, C., Clementi, F., Cardinali, F.,  
905 Petruzzelli, A., Gabucci, C., Tonucci, F., Aquilanti, L., 2019. Unveiling hákarl: A study of the  
906 microbiota of the traditional Icelandic fermented fish. *Food Microbiol.*  
907 <https://doi.org/10.1016/j.fm.2019.03.027>

908 Park, H.-K., Ahima, R.S., 2015. Physiology of leptin: energy homeostasis, neuroendocrine function  
909 and metabolism. *Metabolism* 64, 24–34. <https://doi.org/10.1016/J.METABOL.2014.08.004>

910 Piccinetti, C.C., Donati, M., Radaelli, G., Caporale, G., Mosconi, G., Palermo, F., Cossignani, L.,  
911 Salvatori, R., Lopez, R.P., Olivotto, I., 2015. The effects of starving and feeding on Dover sole  
912 (*Solea solea*, Soleidae, Linnaeus, 1758) stress response and early larval development. *Aquac.*  
913 *Res.* <https://doi.org/10.1111/are.12410>

914 Piccinetti, C.C., Migliarini, B., Olivotto, I., Coletti, G., Amici, A., Carnevali, O., 2010. Appetite  
915 regulation: The central role of melatonin in *Danio rerio*. *Horm. Behav.* 58, 780–785.  
916 <https://doi.org/10.1016/J.YHBEH.2010.07.013>

917 Piccinetti, C.C., Migliarini, B., Olivotto, I., Simoniello, M.P., Giorgini, E., Carnevali, O., 2013.  
918 Melatonin and Peripheral Circuitries: Insights on Appetite and Metabolism in *Danio Rerio* .  
919 *Zebrafish.* <https://doi.org/10.1089/zeb.2012.0844>

920 Piccinetti, C.C., Montis, C., Bonini, M., Laurà, R., Guerrero, M.C., Radaelli, G., Vianello, F.,  
921 Santinelli, V., Maradonna, F., Nozzi, V., Miccoli, A., Olivotto, I., 2014. Transfer of Silica-  
922 Coated Magnetic (Fe<sub>3</sub>O<sub>4</sub>) Nanoparticles Through Food: A Molecular and Morphological  
923 Study in Zebrafish. *Zebrafish.* <https://doi.org/10.1089/zeb.2014.1037>

924 Piccinetti, C.C., Ricci, L.A., Tokle, N., Radaelli, G., Pascoli, F., Cossignani, L., Palermo, F.,  
925 Mosconi, G., Nozzi, V., Raccanello, F., Olivotto, I., 2012. Malnutrition may affect common  
926 sole (*Solea solea* L.) growth, pigmentation and stress response: Molecular, biochemical and  
927 histological implications. *Comp. Biochem. Physiol. - A Mol. Integr. Physiol.*  
928 <https://doi.org/10.1016/j.cbpa.2011.12.009>

929 Pinotti, L., Giromini, C., Ottoboni, M., Tretola, M., Marchis, D., 2019. Review: Insects and former  
930 foodstuffs for upgrading food waste biomasses/streams to feed ingredients for farm animals.  
931 *Animal* 1–11. <https://doi.org/10.1017/s1751731118003622>

932 Randazzo, B., Chemello, G., Tortarolo, I., Chiarello, G.L., Zalas, M., Santini, A., Liberatore,  
933 Margaret, Liberatore, Marco, Selli, E., Olivotto, I., 2017. A Novel Photocatalytic Purification  
934 System for Fish Culture. *Zebrafish*. <https://doi.org/10.1089/zeb.2017.1448>

935 Reed, B., Jennings, M., 2011. Guidance on the housing and care of zebrafish *Danio rerio*. *Guid.*  
936 *Hous. care zebrafish Danio rerio*.

937 Rimoldi, S., Gini, E., Iannini, F., Gasco, L., Terova, G., 2019. The Effects of Dietary Insect Meal  
938 from *Hermetia illucens* Prepupae on Autochthonous Gut Microbiota of Rainbow Trout  
939 (*Oncorhynchus mykiss*). *Animals*. <https://doi.org/10.3390/ani9040143>

940 Ringø, E., Zhou, Z., Olsen, R.E., Song, S.K., 2012. Use of chitin and krill in aquaculture - the effect  
941 on gut microbiota and the immune system: a review. *Aquac. Nutr.* 18, 117–131.  
942 <https://doi.org/10.1111/j.1365-2095.2011.00919.x>

943 Roeselers, G., Mittge, E.K., Stephens, W.Z., Parichy, D.M., Cavanaugh, C.M., Guillemin, K.,  
944 Rawls, J.F., 2011. Evidence for a core gut microbiota in the zebrafish. *ISME J.*  
945 <https://doi.org/10.1038/ismej.2011.38>

946 Roy, S. Sen, Pal, R., 2014. Microalgae in Aquaculture: A Review with Special References to  
947 Nutritional Value and Fish Dietetics. *Proc. Zool. Soc.* [https://doi.org/10.1007/s12595-013-](https://doi.org/10.1007/s12595-013-0089-9)  
948 0089-9

949 Sánchez-Muros, M.J., Barroso, F.G., Manzano-Agugliaro, F., 2014. Insect meal as renewable  
950 source of food for animal feeding: A review. *J. Clean. Prod.*  
951 <https://doi.org/10.1016/j.jclepro.2013.11.068>

952 Sargent, J., Mcevoy, L., Estevez, A., Bell, G., Bell, M., Henderson, J., Tocher, D., 1999. Lipid  
953 nutrition of marine fish during early development: current status and future directions,  
954 *Aquaculture*.

955 Sarker, P.K., Kapuscinski, A.R., Bae, A.Y., Donaldson, E., Sitek, A.J., Fitzgerald, D.S., Edelson,  
956 O.F., 2018. Towards sustainable aquafeeds: Evaluating substitution of fishmeal with lipid-  
957 extracted microalgal co-product (*nannochloropsis oculata*) in diets of juvenile nile tilapia  
958 (*oreochromis niloticus*). *PLoS One*. <https://doi.org/10.1371/journal.pone.0201315>

959 Secci, G., Mancini, S., Iaconisi, V., Gasco, L., Basto, A., Parisi, G., 2019. Can the inclusion of  
960 black soldier fly ( *Hermetia illucens* ) in diet affect the flesh quality/nutritional traits of  
961 rainbow trout ( *Oncorhynchus mykiss* ) after freezing and cooking? *Int. J. Food Sci. Nutr.* 70,  
962 161–171. <https://doi.org/10.1080/09637486.2018.1489529>

963 Shepherd, C.J., Jackson, A.J., 2013. Global fishmeal and fish-oil supply: Inputs, outputs and  
964 marketsa. *J. Fish Biol.* <https://doi.org/10.1111/jfb.12224>

965 Smetana, S., Schmitt, E., Mathys, A., 2019. Sustainable use of *Hermetia illucens* insect biomass for  
966 feed and food: Attributional and consequential life cycle assessment. *Resour. Conserv. Recycl.*

967 <https://doi.org/10.1016/j.resconrec.2019.01.042>

968 Sobrino Crespo, C., Perianes Cachero, A., Puebla Jiménez, L., Barrios, V., Arilla Ferreiro, E.,  
 969 2014. Peptides and Food Intake. *Front. Endocrinol. (Lausanne)*. 5, 58.  
 970 <https://doi.org/10.3389/fendo.2014.00058>

971 Spinelli, R., Neri, P., Pini, M., Barbi, S., Montorsi, M., Ferrari, A.M., 2019. Using black soldier  
 972 flies (*Hermetia illucens*) to bioconvert waste from the livestock production chain: A life cycle  
 973 assessment case study, in: *WIT Transactions on Ecology and the Environment*.  
 974 <https://doi.org/10.2495/WM180051>

975 Sprangers, T., Ottoboni, M., Klootwijk, C., Oryn, A., Deboosere, S., De Meulenaer, B., Michiels,  
 976 J., Eeckhout, M., De Clercq, P., De Smet, S., 2017. Nutritional composition of black soldier fly  
 977 (*Hermetia illucens*) prepupae reared on different organic waste substrates. *J. Sci. Food Agric*.  
 978 <https://doi.org/10.1002/jsfa.8081>

979 St-Hilaire, S., Cranfill, K., McGuire, M.A., Mosley, E.E., Tomberlin, J.K., Newton, L., Sealey, W.,  
 980 Sheppard, C., Irving, S., 2007a. Fish offal recycling by the black soldier fly produces a  
 981 foodstuff high in omega-3 fatty acids. *J. World Aquac. Soc.* <https://doi.org/10.1111/j.1749-7345.2007.00101.x>

982

983 St-Hilaire, S., Sheppard, C., Tomberlin, J.K., Irving, S., Newton, L., McGuire, M.A., Mosley, E.E.,  
 984 Hardy, R.W., Sealey, W., 2007b. Fly prepupae as a feedstuff for rainbow trout, *Oncorhynchus*  
 985 *mykiss*. *J. World Aquac. Soc.* <https://doi.org/10.1111/j.1749-7345.2006.00073.x>

986 Stevens, J.R., Newton, R.W., Tlustý, M., Little, D.C., 2018. The rise of aquaculture by-products:  
 987 Increasing food production, value, and sustainability through strategic utilisation. *Mar. Policy*.  
 988 <https://doi.org/10.1016/j.marpol.2017.12.027>

989 Su, J., Gong, Y., Cao, S., Lu, F., Han, D., Liu, H., Jin, J., Yang, Y., Zhu, X., Xie, S., 2017. Effects  
 990 of dietary *Tenebrio molitor* meal on the growth performance, immune response and disease  
 991 resistance of yellow catfish (*Pelteobagrus fulvidraco*). *Fish Shellfish Immunol.* 69, 59–66.  
 992 <https://doi.org/10.1016/J.FSI.2017.08.008>

993 Tacon A. G. J., Metian, M., 2008. Global overview on the use of fish meal and fish oil in  
 994 industrially compounded aquafeeds. *Aquaculture*.  
 995 <https://doi.org/10.1016/j.aquaculture.2008.08.015>

996 Terova, G., Rimoldi, S., Ascione, C., Gini, E., Ceccotti, C., Gasco, L., 2019. Rainbow trout  
 997 (*Oncorhynchus mykiss*) gut microbiota is modulated by insect meal from *Hermetia illucens*  
 998 prepupae in the diet. *Rev. Fish Biol. Fish.* <https://doi.org/10.1007/s11160-019-09558-y>

999 Tlustý, M.F., Thorsen, Ø., 2017. Claiming seafood is ‘sustainable’ risks limiting improvements.  
 1000 *Fish Fish.* <https://doi.org/10.1111/faf.12170>

1001 Tocher, D.R., 2010. Fatty acid requirements in ontogeny of marine and freshwater fish. *Aquac. Res.*  
 1002 41, 717–732. <https://doi.org/10.1111/j.1365-2109.2008.02150.x>

1003 Truzzi, C., Illuminati, S., Annibaldi, A., Antonucci, M., Scarponi, G., 2017. Quantification of fatty  
 1004 acids in the muscle of Antarctic fish *Trematomus bernacchii* by gas chromatography-mass  
 1005 spectrometry: Optimization of the analytical methodology. *Chemosphere*.  
 1006 <https://doi.org/10.1016/j.chemosphere.2016.12.140>

1007 Truzzi, C., Illuminati, S., Antonucci, M., Scarponi, G., Annibaldi, A., 2018. Heat shock influences  
 1008 the fatty acid composition of the muscle of the Antarctic fish *Trematomus bernacchii*. *Mar.*  
 1009 *Environ. Res.* <https://doi.org/10.1016/j.marenvres.2018.03.017>

- 1010 Truzzi, C., Giorgini, E., Annibaldi, A., Antonucci, M., Illuminati, S., Scarponi, G. Riolo P., Isidoro,  
1011 N., Conti C., Zarantoniello, M., Cipriani, R., Olivotto, I. (In press). **Fatty acids profile of black**  
1012 **soldier fly (*Hermetia illucens*): influence of feeding substrate based on coffee-waste silverskin**  
1013 **enriched with microalgae. Animal Feed Science and Technology, In press**
- 1014 Udayangani, R.M.C., Dananjaya, S.H.S., Nikapitiya, C., Heo, G.J., Lee, J., De Zoysa, M., 2017.  
1015 Metagenomics analysis of gut microbiota and immune modulation in zebrafish (*Danio rerio*)  
1016 fed chitosan silver nanocomposites. Fish Shellfish Immunol.  
1017 <https://doi.org/10.1016/j.fsi.2017.05.018>
- 1018 van Huis, A., 2013. Potential of Insects as Food and Feed in Assuring Food Security. Annu. Rev.  
1019 Entomol. 58, 563–583. <https://doi.org/10.1146/annurev-ento-120811-153704>
- 1020 Vargas-Abúndez, A.J., Randazzo, B., Foddai, M., Sanchini, L., Truzzi, C., Giorgini, E., Gasco, L.,  
1021 Olivotto, I., 2019. Insect meal based diets for clownfish: Biometric, histological, spectroscopic,  
1022 biochemical and molecular implications. Aquaculture.  
1023 <https://doi.org/10.1016/j.aquaculture.2018.08.018>
- 1024 Vargas, A., Randazzo, B., Riolo, P., Truzzi, C., Gioacchini, G., Giorgini, E., Loreto, N., Ruschioni,  
1025 S., Zarantoniello, M., Antonucci, M., Polverini, S., Cardinaletti, G., Sabbatini, S., Tulli, F.,  
1026 Olivotto, I., 2018. Rearing Zebrafish on Black Soldier Fly ( *Hermetia illucens* ): Biometric,  
1027 Histological, Spectroscopic, Biochemical, and Molecular Implications . Zebrafish 15, 404–  
1028 419. <https://doi.org/10.1089/zeb.2017.1559>
- 1029 Volkoff, H., 2006. The role of neuropeptide Y, orexins, cocaine and amphetamine-related  
1030 transcript, cholecystokinin, amylin and leptin in the regulation of feeding in fish. Comp.  
1031 Biochem. Physiol. Part A Mol. Integr. Physiol. 144, 325–331.  
1032 <https://doi.org/10.1016/J.CBPA.2005.10.026>
- 1033 Wang, C., Qian, L., Wang, W., Wang, T., Deng, Z., Yang, F., Xiong, J., Feng, W., 2017. Exploring  
1034 the potential of lipids from black soldier fly: New paradigm for biodiesel production (I).  
1035 Renew. Energy. <https://doi.org/10.1016/j.renene.2017.04.063>
- 1036 Webster, C.D., Rawles, S.D., Koch, J.F., Thompson, K.R., Kobayashi, Y., Gannam, A.L., Twibell,  
1037 R.G., Hyde, N.M., 2016. Bio-Ag reutilization of distiller’s dried grains with solubles (DDGS)  
1038 as a substrate for black soldier fly larvae, *Hermetia illucens*, along with poultry by-product  
1039 meal and soybean meal, as total replacement of fish meal in diets for Nile tilapia, *Oreochr.*  
1040 *Aquac. Nutr.* 22, 976–988. <https://doi.org/10.1111/anu.12316>
- 1041 Weisburg, W.G., Barns, S.M., Pelletier, D.A., Lane, D.J., 1991. 16S ribosomal DNA amplification  
1042 for phylogenetic study. J. Bacteriol.
- 1043 Xiao, X., Jin, P., Zheng, L., Cai, M., Yu, Z., Yu, J., Zhang, J., 2018. Effects of black soldier fly  
1044 (*Hermetia illucens*) larvae meal protein as a fishmeal replacement on the growth and immune  
1045 index of yellow catfish (*Pelteobagrus fulvidraco*). Aquac. Res.  
1046 <https://doi.org/10.1111/are.13611>
- 1047 Xu, W.J., Jin, J.Y., Zou, T., Han, D., Liu, H.K., Zhu, X.M., Yang, Y.X., Xie, S.Q., 2019. Growth,  
1048 feed utilization and metabolic responses of three gibel carp (*Carassius gibelio*) strains to  
1049 fishmeal and plant protein-based diets. Aquac. Nutr. <https://doi.org/10.1111/anu.12855>

- 1050 Zarantoniello, M., Bruni, L., Randazzo, B., Vargas, A., Gioacchini, G., Truzzi, C., Annibaldi, A.,  
1051 Riolo, P., Parisi, G., Cardinaletti, G., Tulli, F., Olivotto, I., 2018. Partial Dietary Inclusion of  
1052 *Hermetia illucens* (Black Soldier Fly) Full-Fat Prepupae in Zebrafish Feed: Biometric,  
1053 Histological, Biochemical, and Molecular Implications . *Zebrafish* 15, 519–532.  
1054 <https://doi.org/10.1089/zeb.2018.1596>
- 1055 Zarantoniello, M., Randazzo, B., Truzzi, C., Giorgini, E., Marcellucci, C., Vargas-Abúndez, J.A.,  
1056 Zimbelli, A., Annibaldi, A., Parisi, G., Tulli, F., Riolo, P., Olivotto, I., 2019. A six-months  
1057 study on Black Soldier Fly (*Hermetia illucens*) based diets in zebrafish. *Sci. Rep.* 9, 8598.  
1058 <https://doi.org/10.1038/s41598-019-45172-5>
- 1059 Zhang, Y., Feng, S., Chen, J., Qin, C., Lin, H., Li, W., 2012. Stimulatory effects of chitinase on  
1060 growth and immune defense of orange-spotted grouper (*Epinephelus coioides*). *Fish Shellfish*  
1061 *Immunol.* 32, 844–854. <https://doi.org/10.1016/J.FSI.2012.02.009>
- 1062 Zhou, Z., Karlsen, Ø., He, S., Olsen, R.E., Yao, B., Ringø, E., 2013. The effect of dietary chitin on  
1063 the autochthonous gut bacteria of Atlantic cod ( *Gadus morhua* L.). *Aquac. Res.* 44, 1889–  
1064 1900. <https://doi.org/10.1111/j.1365-2109.2012.03194.x>

1065

## 1066 8. Appendix

1067 TABLE 1A. OBSERVED DIVERSITY, GOOD'S COVERAGE AND NUMBER OF SEQUENCES FOR  
1068 THE 16S RRNA AMPLICONS OBTAINED FROM ZEBRAFISH GUT (G) SAMPLES.  
1069


Sample	Goods_coverage	PD_whole_tree	chao1	Observed_species	Shannon
GHi0	99.77	19.37	418.40	299	3.41
GHi25	99.88	13.26	294.02	259	4.01
GHi50	99.80	20.37	462.88	380	3.25
GHi75	99.75	21.23	470.88	388	3.49
GHi100	99.75	19.97	419.85	332	3.72

1070 Zebrafish insect-based diets were prepared by including graded levels of insect meal (25, 50, 75, and 100 %, referred to as Hi25 and Hi50, Hi75 and Hi100, respectively) in the Hi0 formulation (control diet without insect addition).  
1071  
1072  
1073

1074


## 1075 Figure Legends

1076 **Figure 1. Specific Growth Rate (% weight growth day<sup>-1</sup>) of zebrafish larvae.** Zebrafish larvae fed diets including 0,  
1077 25, 50, 75 and 100% of BSF meal (Control, Hi25, Hi50, Hi75 and Hi100). Boxplots show minimum and maximum  
1078 (whiskers), first quartile, median and third quartile (box). Different letters denote statistically significant differences  
1079 among experimental groups.




**Figure 2. Content of SFA, MUFA and PUFA (as % of total FA) and contribution of omega 3 (n3), omega 6 (n6) and omega 9 (n9) fatty acids to lipid profile. (a,b) experimental diets; (c,d) zebrafish larvae. Control, Hi25, Hi50, Hi75 and Hi100 diets were characterized by 0, 25, 50, 75 or 100% inclusion of BSF meal. Zebrafish larvae fed diets including 0, 25, 50, 75 and 100% of BSF meal (Control, Hi25, Hi50, Hi75 and Hi100). Different letters indicate statistically significant differences among experimental groups compared within the same fatty acid class ( $p < 0.05$ ). Values are presented as mean  $\pm$  SD ( $n = 12$ ).**






**Figure 3. Example of histomorphology of intestine and liver of zebrafish larvae. (a-j) intestine; (k-o) liver. Zebrafish larvae fed diets including 0, 25, 50, 75 and 100% of BSF meal (Control, Hi25, Hi50, Hi75 and Hi100). Scale bars: (a-e) 50  $\mu$ m; (f-j) 20  $\mu$ m; (k-o) 10  $\mu$ m. \* indicates lipid accumulation.**




**Figure 4. Percentage of fat fraction (PFF) in liver tissue calculated on histological sections.** in control, Hi25, Hi50, Hi75 and Hi100 groups. Values are presented as mean  $\pm$  SD (n=9). Different letters indicate statistically significant differences among the experimental groups.




**Figure 5. IR spectra of zebrafish larvae.** Average absorbance spectra of zebrafish larvae fed diets including 0, 25, 50, 75 and 100% of BSF meal (Control, Hi25, Hi50, Hi75 and Hi100). For clarity reasons, spectra are shifted along y-axis.


1104 The position (in terms of wavenumbers) of the most featuring IR peaks is reported, together with (colored boxes) the  
 1105 corresponding biochemical meaning.



1106 **Figure 6. Statistical analysis of lipid composition.** Numerical variation of IR band area ratios in zebrafish larvae fed  
 1107 diets including 0, 25, 50, 75 and 100% of BSF meal (Control, Hi25, Hi50, Hi75 and Hi100): (a) LIP/TBM (overall  
 1108 amount of lipids); (b) 1744/LIP (amount of fatty acids with respect to lipids); (c) 2925/LIP (amount of saturated fatty  
 1109 acids with respect to lipids) and (d) 3010/LIP (amount of unsaturated fatty acids with respect to lipids). Different letters  
 1110 above histograms indicate statistically significant differences among groups (p<0.05).  
 1111




**Figure 7. Statistical analysis of protein composition.** Numerical variation of IR band area ratios in Zebrafish larvae fed diets including 0, 25, 50, 75 and 100% of BSF meal (Control, Hi25, Hi50, Hi75 and Hi100): (a) PRT/TBM (overall amount of proteins); (b) 1234/PRT (amount of collagen with respect to proteins) and (c) 1055/PRT (amount of mucin with respect to proteins). Different letters above histograms indicate statistically significant differences among groups ( $p < 0.05$ ).




**Figure 8. PCA based on the OTU abundance of the zebrafish samples grouped as a function of the amount of BSF added to the diet.** The first component (PC1) accounts for the 43.14% of the variance, the second component (PC2) accounts for the 22.64 % of the variance, the third component (PC3) accounts for the 18.94% of the variance.

1124 Zebrafish insect-based diets were prepared by including graded levels of BSF meal (25, 50, 75 and 100%, referred to as  
 1125 Hi25 and Hi50, Hi75 and Hi100, respectively) in the Hi0 formulation (control diet without insect addition).



1126  
 1127  
 1128 **Figure 9.** Relative abundances as identified by MiSeq Illumina expressed as the percent ratio between the sum of reads  
 1129 of each OTUs and the total number of reads found in the zebrafish gut (G) samples.



1132 **Figure 10. Relative mRNA levels of genes analyzed in zebrafish larvae.** (a) *igf1*, (b) *igf2a*, (c) *mstnb*, (d) *nr3c1*, (e)  
 1133 *hsp70.1*, (f) *elovl2*, (g) *elovl5*, (h) *fads*, (i) *ghrl*, (l) *npy*, (m) *cnr1*, (n) *lepa*, (o) *il1b*, (p) *il10*, (q) *tnfa*, (r) *chia.2*, (s)  
 1134 *chia.3*. Different letters indicate statistically significant differences among experimental groups compared within the  
 1135 same sampling time ( $p < 0.05$ ). Values are presented as mean  $\pm$  SD ( $n = 5$ ). Zebrafish larvae fed diets including 0, 25, 50,  
 1136 75 and 100% of BSF meal (Control, Hi25, Hi50, Hi75 and Hi100).  
 1137

