



UNIVERSITÀ POLITECNICA DELLE MARCHE  
Repository ISTITUZIONALE

Citric acid bioproduction: the technological innovation change

This is the peer reviewed version of the following article:

*Original*

Citric acid bioproduction: the technological innovation change / Amato, A.; Becci, A.; Beolchini, F.. - In: CRITICAL REVIEWS IN BIOTECHNOLOGY. - ISSN 0738-8551. - 40:2(2020), pp. 199-212. [10.1080/07388551.2019.1709799]

*Availability:*

This version is available at: 11566/275004 since: 2024-03-26T14:56:10Z

*Publisher:*

*Published*

DOI:10.1080/07388551.2019.1709799

*Terms of use:*

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. The use of copyrighted works requires the consent of the rights' holder (author or publisher). Works made available under a Creative Commons license or a Publisher's custom-made license can be used according to the terms and conditions contained therein. See editor's website for further information and terms and conditions.

This item was downloaded from IRIS Università Politecnica delle Marche (<https://iris.univpm.it>). When citing, please refer to the published version.

(Article begins on next page)

**Citric acid bioproduction: ~~a patent review~~ the technological innovation  
change**

ha formattato: Inglese (Regno Unito)

Alessia Amato<sup>a\*</sup>, Alessandro Becci<sup>a</sup>, Francesca Beolchini<sup>a</sup>

<sup>a</sup> *Department of life and environmental science, Università Politecnica delle Marche, Ancona,  
Italy*

Alessia Amato: [a.amato@univpm.it](mailto:a.amato@univpm.it), ORCID: 0000-0002-0840-1584

Alessandro Becci: [a.becci@pm.unipm.it](mailto:a.becci@pm.unipm.it), ORCID: 0000-0001-5886-1557

Francesca Beolchini: [f.beolchini@univpm.it](mailto:f.beolchini@univpm.it), ORCID: 0000-0002-6211-6354

## **Citric acid bioproduction: the technological innovation change** **bioproduction, a patent review**

### **Abstract**

~~Considering its several application fields, t~~The citric acid is considered one of the most valuable weak organic acid on the market and its production by biotechnological approaches is a very interesting topic. Despite the related scientific research, the literature still lacks a state of art of the technological innovation change, necessary for the ~~identification study~~ of the inventions designed for the real scale implementation. ~~In this context, the present review took into account more than 100 worldwide patents (1929-20189), necessary for the identification of the innovative markets and the most promising fields for the economic investments. To fill this gap, the present review took into account more than 100 worldwide patents (1929-2019).~~The deepened study identified an increasing invention number, combined with the current worldwide citric acid export flows, ~~confirmed the~~with China as the leader (with an economic contribution of 75%, in 2017). ~~In order to satisfy the requests of the market which has moved towards a circular economy, the possibility to use waste substrate represents one of the main options considered in the recent patents. Furthermore, the new technology study identified the most critical aspects on which the innovation has invested: the alternative substrates, mainly scraps, and the~~The discussion highlights the sustainability improvement, achieved by the conversion ~~towards~~from a submerged technology to a solid-state fermentation (*koji* process). ~~These advancements could increase the process sustainability combining the waste exploitation with a reduction of both the consumptions and the wastewater production, in agreement with the circular economy pillars.~~ The showed results are essential for both a scientific audience and the stakeholders citric acid production, in order to have a complete and updated overview about this topic.

**Keywords:** citric acid; patent ~~review~~; technological evolution; bio-production; *Aspergillus niger*; fermentation; ~~innovative substrates~~

## 1. Introduction

The citric acid (CA) is ~~considered~~ the most valuable weak organic acid and widely used ~~weak organic acid~~ on the market for many applications [1]. ~~Initially crystallized from lemon juice, citric acid~~ It is a tricarboxylic acid with an essential role for the metabolism of aerobic organisms [2]–[2], [3]. ~~Studies about its production dates back many years when it was~~ Its production is one of the most thoroughly studied and it dates back many years, crystallized from lemon juice (18<sup>th</sup> century), thanks to the several applications, as confirmed by Curie in 1916, which describes a filamentous fungal fermentation process [3]. Currently, the ~~citric acid~~ CA biotechnological production is mainly located in China and, with amounts that have increased from 0.5 to 2 million tonnes of product for year are expected to, in the last twenty years 2020 [2], a quantity that far exceeds the production of every other organic acid made by fermentation. The uses of this agent includes ~~different several~~ fields, like including food, textile, chemical and pharmaceutical industries [3]. More in detail, about 70% of the citric acid on the market is used for food and beverage products, of which the 50% of the total for carbonated beverages [4]. This percentage is further growing for the expansion of the developing countries markets. On the other hand, pharmaceutical industry employs citric acid as a preservative for stored blood, tablets, ointments, and cosmetic preparations [5]. In the chemical sector, it is used as an antifoam agent and for the textile treatment.

Nowadays, ~~almost the entire the experimental~~ world production ~~is manufactured using of~~ citric acid CA mainly involves fungi and yeasts: *Aspergillus niger*, *A. wentii*, *A. clavatus*, *Penicillium luteum*, *P. citrinum*, *Mucor piriformis*, *Candida guilliermondii*, *Saccharomycopsis lipolytica*, *Trichoderma viride* and *Arthrobacter parafineus*. However, only *A. niger* and the closely related strain of *A. wentii* are chosen for the commercial production [4]. ~~Overall, Two main phases~~ characterize the processes include the previous fermentation, which needs high productivity and

ha formattato: Apice

yield values, an followed by the consecutive recovery and purification. The main critical parameters for the fermentation are: the high carbohydrate concentration, the maintenance of high dissolved oxygen, the constant agitation and the low pH value [5]–[9]. As concern the further recovery and purification steps, the most common approaches are On the other hand, two common processes for citric acid recovery and purification are: the lime  $\text{CaCO}_3$ /sulphuric acid  $\text{H}_2\text{SO}_4$  precipitation (with calcium hydroxide) [10]–[12], and the liquid solvent extraction (by a solvent mixture) [10]. Both options are characterized by a preliminary filtration or centrifugation for the separation of the fermentation liquor and the solid biomass. In the lime/sulphuric acid process, the fermentation liquor is treated by calcium hydroxide for the calcium citrate precipitation. The filtered product is washed for the impurity elimination and dissolved with sulphuric acid. The produced insoluble calcium sulphate is separated from the citric acid solution, which is deionised and concentrated by crystallization to form both anhydrous and monohydrated citric acid [12], [13]. The main process weakness is the production of calcium sulphate, as by product, with criticalities due to its disposal. On the other hand, the liquid solvent extraction needs a counter-current set up and it uses a mixture of tri laurylamine, n octanol and decane or undecane to extract citric acid from the fermentation broth. This step is followed by the citric acid extraction/back-extraction by water, at high temperature [13], [14]. Otherwise Alternatively, Ledakowicz et al. (2004) describe the an anion-exchange advanced method can be used for the citric acid recovery from aqueous solution using a tertiary amine resin, followed by thermal desorption [15].

In this context, the knowledge of the *A. niger* metabolism is an essential step to improve the citric acid fermentation processes and the literature has exhaustively reviewed the related mechanisms, summarized in Figure 1 [4], [15]–[18]. The production of citrate from glucose or sucrose involves many enzymatic steps occurring in two different membrane-bound cellular compartments: the cytosol and the mitochondrion. Glucose is transported into the cell and converted to the three-carbon acid, pyruvate, via the glycolytic pathway in the cytosol. The decarboxylation of one

ha formattato: Pedice

ha formattato: Pedice

ha formattato: Pedice

ha formattato: Non Evidenziato

ha formattato: Non Evidenziato

ha formattato: Non Evidenziato

Codice campo modificato

ha formattato: Non Evidenziato

ha formattato: Non Evidenziato

ha formattato: Non Evidenziato

ha formattato: Non Evidenziato

ha formattato: Non Evidenziato

Codice campo modificato

molecule of pyruvate forms acetyl-CoA by the mitochondrial pyruvate dehydrogenase complex and another is carboxylated to oxaloacetate in the cytosol by pyruvate carboxylase. The oxaloacetate must be taken into the mitochondrion (via malate) and condensed with the acetyl-CoA to form citrate. The product is transported out of the mitochondrion and finally out of the cell [10].

Considering the ~~strong interest for~~ relevance of the ~~citric acid~~ CA for the market, the recent literature has widely discussed ~~about the citric acid~~ its production by fungi, mainly *A. niger*, reporting several substrates and operative conditions [16]–[20]. ~~In this regard, m~~ Many reviews ~~have deepened the acid production this topic, often,~~ focusing on specific aspects as: ~~the fermentation variables~~ [21]–[24] and the characteristics of the solid-state fermentation [25]–[27]. Recent studies has focused on the possibility of a waste material-use as substrate [23], [28], following the circular economy principles, ~~the solid stat fermentation [23] [25] and the fermentation variables, more generally [16], [28], [29].~~ Nevertheless, the scientific literature still ~~lack~~ should be integrated by works about the state of art of the technological innovation change, represented by the registered patents, able to create an overview of the inventions designed for the application in a real scale. Currently, ~~this relevant aspect the relevance of this kind of information, represented by the registered patents,~~ was already highlighted in other application fields (e.g. waste recycling or medical applications), ~~confirming the possibility~~ to identify the most promising technologies useful for the creation of innovative markets [29]–[33].

~~In order to~~ To fill in the gap, the present work, (updated to March 2019,) analyzes 100 patents, between 1929 and 2018, showing the technological evolution of the last decades. The free access Espacenet platform was used as the main information source [34] using the keywords, for the patent search: “citric acid bioproduction”, “citric acid *Aspergillus niger*”, “citric acid *Candida*”, “citric acid *Penicillium*”, “citric acid *Aspergillus*”. This information source, created by the

European patent office, ensured a worldwide invention overview [29]. In order to simplify the review reading, the included inventions are organized following the roadmap in Figure 2.

## 2. *A. niger*, Sugar different methods to use raw material as carbon source

The main conditions with an effect on the citric acid bioproduction by *A. niger* is influenced by several suitable conditions, mainly are: low pH, high dissolved oxygen and high sugar concentration [10], [34]. This microorganism *A. niger* uses many takes advantage of different substrates for its growth (i.e. maltose, mannose, galactose, fructose, sucrose and glucose). Nevertheless, glucose and sucrose produce the best results showing both (both the greatest growth and the highest citric acid production) rate with glucose and sucrose, with a concentration of 10-14% (w/v) [35], [36]. Overall, the fungal fermentation can be carried out by three different fermentation techniques: submerged fermentation, surface fermentation (liquid surface culture) and solid-state fermentation (koji process) [27], [37], [38]. Nevertheless, the highest patent number (Figure 1A) identifies the first method as the most common, as also confirmed by the literature which reports around 80% of the world production used the by first this approach [21], [39], as also confirmed by the significant number of related patents (Figure 3).

### 2.1 Immobilization method

The fungal immobilization method was used to improve the citric acid production by *A. niger*. The fungal was embedded by calcium alginate to reduce the contact with the produced citric acid, decreasing the acid toxicity on the metabolism. This effect is due to the citrate, a strong inhibitor of the glycolytic enzymes 6-phosphofructokinase, a glycolytic enzymes [40], [41]. The patent CN102864184 describes the use of an optimal medium composed of sucrose at with a concentration of 120 g/L, at 30°C and 200 rpm [42]. A similar approach is reported in the patent CN107022541, which involves a pretreated fibrous material, as immobilization medium. An air sparging system improves the citric acid production, with a final yield of 93% (expressed as citric acid yield/glucose consumed), a citric acid concentration of 89.65 g/L and a fermentation

ha formattato: Tipo di carattere: Grassetto

rate of 1.06 g/L·h [43]. The immobilization method allows multiple advantages: the ~~increase of the citric acid~~ production efficiency increase, the ~~reduction of the~~ fermentation time reduction, the decrease of the biomass and the fermentation broth amount and a simplified separation between the ~~final~~ product and the mycelia.

## 2.2 Multiple step processes

The fungal growths using a multiple step process is an alternative technique ~~to for the production~~ increase the citric acid production. In this regard, ~~the invention CN107815421 discloses~~ describes a method where *A. niger* is cultivated ~~by in~~ three different growth phases to prevent the criticality of the spore aggregation, allowing both the highest quantity and a size uniformity of the mycelium pellets; thereafter ~~After this preliminary step,~~ the matured seem liquor is transferred to the final fermentation tank. ~~This approach solves the spore aggregation issue, permitting the highest quantity and size uniformity of the mycelium pellets.~~ Consequently, the glucose is metabolized by the fungal and the conversion rate is improved, with ~~the reduction of the~~ lower fermentation time [44]. On the other hand, the inventions CN106868061, CN102181490 describe a method which includes a first step ~~where of A. niger grows growth by CO<sub>2</sub> gas sparging with CO<sub>2</sub> sparging in a fermentation tank.~~ When the fungal biomass is grown, it is transferred to a second section where fungi continue the citric acid production ~~until decreasing the~~ reducing sugar content ~~of reducing sugar is lower than~~ up to 1-3 g/L [45], [46]. ~~Overall, this approach~~ These treatments allows both ~~higher~~ increase the conversion rate and ~~they reduce~~ lower the residual sugar, ~~than compared to the~~ traditional ~~treatments~~ approaches. The pH shifting, during the acid production, is proposed ~~An alternative, showed in patents CN104277978, CN102851330, includes the solution pH shift during the acid production.~~ During the preliminary phase, the pH is ~~maintained between 6.2 and 7.2,~~ then its rise increases the production rate [47], [48]. As ~~reported explained~~ in the invention CN102373242, the temperature ~~is another variable which modification improves could be modified during the~~ citric acid synthesis. More in detail, a 35-45°C range is selected during the rise



phase of *A. niger* respiratory quotient and growth; thereafter, the value is reduced, ~~up~~ to 30-40°C, during the second step, when the biomass is constant and the respiratory quotient decreases. ~~The~~An ~~additional -temperature~~ increase ~~s(35-45°C)-again is necessary~~ in the last step ~~(35-45°C)~~, when the sugar content is very low [49]. The fungi age change ~~could-affects~~ the process efficiency, as reported in patents CN104099253, CN104087624. *A. niger* grows in ~~the-a~~ first medium to obtain a mature seed solution; part of the resulting dispersed mycelium is transferred to a fermentation medium (5%-15% of inoculum concentration), when it reached the maximum growth rate. This second step finishes when the concentration of the reducing sugar ~~concentration~~ in the fermentation medium is lower than 0.5%. The use of the dispersed mycelium seed solution from the second step avoids the repetition of the first step with a consequent continuous production of ~~citric acid~~CA [50], [51]. A different method is described in the invention CN102443611, characterized by the addition ~~(in the same fermentation tank)~~ of a monosaccharide with 6 carbon atoms to the fermentation liquid, ~~starting~~ from 24 ~~hours~~ after the *A. niger* inoculum to 5 ~~hours~~ before the fermentation conclusion [52]. ~~The main achieved advantage is the possibility to use one fermentation tank, thanks to the continuous monosaccharide addition.~~

### 2.3 *Metal ~~utilization~~use*

~~The metal ions have an essential role in the~~ citric acidCA biosynthesis ~~The-by~~ *A. niger*. Nevertheless, the identification of the best concentration is necessary, since ~~high concentrations, at uncorrected conditions, can be translated into a low efficiency production [10], [53](Christian and Rohr 1986; Magnuson and Lasure 2004)~~ requires a variety of metal ions, at low concentrations, for the citric acid biosynthesis. Too high concentrations could be translated into a low efficiency production, nevertheless, the selection of the best conditions can significantly improve the synthesis, as described in several patents (Christian and Rohr 1986; Magnuson and Lasure 2004). ~~Therefore~~In this regard, the invention US5532148, includes the manganese-Mn (II) ~~use~~addition, with a concentration ~~between-of~~ 2.5-~~and~~ 20 ppb, at pH 1.5-3.0, achieving the highest citric acidCA

production between 4 and 7 ~~days of~~ fermentation ~~days~~. ~~The further~~ ~~Thereafter, the~~ broth is decolorized and ion-exchanged, ~~to allow the remove the color and the inorganic ions with the~~ production of an aqueous citric acid ~~CA~~ solution, with a purity around 98% [54]. ~~As an alternative to the~~ ~~Alternatively, manganese Mn, zinc Zn~~ (30-250 ppm), ~~[hexacyanoferrate ions Fe(CN)<sub>6</sub>]<sup>4-</sup>~~ (100-500 ppm) and ~~copper Cu~~ ~~proved to be useful to~~ increase the acid production, at ~~pH 1.5-2.0~~ ~~pH values between 1.5 and 2.0, as confirmed by~~ ~~in~~ patents: US5081025, US3936352, GB1392942, GB1342311 [55]–[58].

ha formattato: Pedice

ha formattato: Apice

## 2.4 ~~Other-Alternative~~ techniques

~~Additional strategies have been optimized~~ ~~Alternative methods, which include different approaches, have been studied~~ to increase the citric acid ~~CA~~ production by *A. niger*. ~~In this regard,~~ CN105586366, CN103497977 patents describe ~~the possibility of the~~ stirring speed increase during the ~~first-preliminary~~ 6-20 ~~hours~~ of incubation. This condition allows, the hypha shape changes with the formation of small and compact pellets, able to increase the oxygen exchange capacity, ~~with the consequent improvement of,~~ ~~improving~~ the acid synthesis [59], [60]. ~~On the other hand,~~ CN103695319 patent ~~explains the uses~~ a sugar solution with neutralization wastewater to replace the traditional tap water [61]. The possibility of a waste stream use, as ~~nitrogen-N~~ and ~~phosphorous-P~~ source, ~~produces a double advantage: a process combines a significant~~ cost reduction ~~and with the~~ decrease of the environmental load, ~~for the wastewater treatment.~~

~~Overall, a multi-step process represents the main option chosen for the improvement of citric acid CA production by A. niger with raw materials, as carbon sources. Around 50% of the patents in this section, referred to the last decade, confirmed an efficiency increase achieved by process suitable for the scale-up. On the other hand, the metal use is taken into account in old inventions, due to the high criticality of the element concentration which can decrease the process effectiveness [22], [62], [63].~~

ha formattato: Tipo di carattere: Corsivo

### 3. *A. niger*, food and Agriculture waste as carbon source

### 3. Food and Agriculture waste as carbon source

The possible substitution of the traditional glucose (or other sugars) by food and agriculture waste, as carbon source for the fungal metabolism, represents an interesting topic since 1920, ~~as confirmed by the related patents~~ (Figure 4.1B2). ~~In this regard,~~ The present paragraph ~~takes into account~~ focuses on the main streams considered for the technical innovation in this field.

#### 3.1 *Corns as carbon source*

Several kinds of corn have been used, as carbon source, to produce ~~citric acid~~ CA by *A. niger*, mainly: corn cobs, corn wheat, bran, soy, sorghum, corn sugar, wheat straw and rice. Usually, the substrates need a pretreatment to make the cellulose bio-available for the fungal metabolism. ~~Patents CN106119306, CN105524951 In this regard, describe a sulfuric acid~~ H<sub>2</sub>SO<sub>4</sub> washing (for 1-3.5 ~~hours~~), followed by the addition of an enzyme (e.g. cellulase or cellobiase) ~~is used as corn pretreatment, in patents CN106119306, CN105524951, to provide the hydrolysis and the conversion of cellulose into glucose. At the end of this reaction~~ At the end of this pretreatment, the fungal *A. niger* is inoculated and the fermentation starts ~~the citric acid production~~ [64], [65]. On the other hand, the inventions CN103710397 and CN107815475 ~~deseribe-report~~ a corn liquefied solution use, which needs. ~~In the first patent, the substrate is treated with the preliminary addition of beta-cyclodextrin (pH 6.0-6.8 and 15-17% (w/v) of sugar concentration between 15% and 17% w/v) and then sterilized to make it suitable for the fermentation.~~ Thereafter, the fungal is inoculated and the fermentation is carried out at 35-37°C, ensuring the oxygen supply and the necessary inorganic nitrogen ~~N~~ amount ~~of inorganic nitrogen~~ [66]. ~~The two-A two-stage fermentation is~~

ha formattato: Inglese (Regno Unito)

ha formattato: Tipo di carattere: (Predefinito) Times New Roman, 12 pt, Grassetto

ha formattato: Tipo di carattere: (Predefinito) Times New Roman, 12 pt, Grassetto, Corsivo

ha formattato: Tipo di carattere: (Predefinito) Times New Roman, 12 pt, Grassetto

Formattato: Titolo 1, Interlinea: doppia

Formattato: Rientro: Sinistro: 0.75 cm

Formattato: Rientro: Sinistro: 0.75 cm, Interlinea: doppia, Nessun elenco puntato o numerato

Formattato: Titolo 1, Rientro: Sinistro: 0.75 cm, Interlinea: doppia

ha formattato: Pedice

ha formattato: Pedice

described in the second invention. The stages fermentation process, in the second invention, is conducted in a fermentation cylinder (16-24 hours), where the fungal spores are added to sterile water and the fermentation is carried out for 16-24 hours. The first step starts after the corn liquefied addition and it carries out for 2-8 hours, at 35-39°C. In the second one a sucrose solution acts as supplementary carbon source and the citrate production continues until the sugar concentration is lower than 5 g/L. This multiply step design allows to increase both the pH and the nutrient concentration with a positive effect on the fungal growth [67]. The possibility of a cornstarch use avoids the pretreatment. More in detail, the fermentation medium described in the patent CN102864182, is prepared as following: cornstarch (20% (w/v) of cornstarch, (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> (0.2% w/v), KH<sub>2</sub>PO<sub>4</sub> (0.2% w/v), MgSO<sub>4</sub>·7H<sub>2</sub>O (0.05% w/v) and methanol-MeOH (4% w/v). The fermentation is carried out at: pH 3.0, 30°C, 200 rpm, for 7 days [68]. Alternatively, the invention RU267614 combines the addition of the nutrients to the cornstarch, with the change of both the mixing speed and the aeration rate during the fermentation period. More in detail, during the growth phase, the two parameters increase from 120 rpm and 8480 L/hmin of the first 6 hours, to 250 rpm and 321920 L/min in of the remaining 18 hours. On the other hand, during the deep fermentation period, the stirring and the aeration conditions rise from 250 rpm and 2440 L/min of the first day, to 300 rpm and 3248 L/min of the second one day to reach 400 rpm and 4056 L/min until at the process conclusion [69]. Another waste carbon source which avoids the pretreatment is Even the corn sugar does not require further process, before the fermentation. In this regard, the inventions GB738940, GB742972, CN102851328 describe the citric acid CA production by the nitrogen N source addition, in ammonia gas form, combined with further salts (i.e. KH<sub>2</sub>PO<sub>4</sub>, MgSO<sub>4</sub>, morpholine, CaCl<sub>2</sub>, MoO<sub>3</sub>, zinc Zn), adjusting the pH value at 2.6 to stimulate the acid synthesis [70]–[72]. On the other hand, the dry corn powder needs a treatment before the fermentation, as described in the invention CN101555497. A water addition produces a paste, which is mixed with amylase and heated at 92-98°C, for a first step, and at 88-90°C, for a second one, maintaining the liquefaction for 90-120 minutes 1 h. About the 90% of the resulting

solution is filtered and used for the inoculation (at 38°C), after the inorganic nitrogen-N addition ~~and the sterilization~~ [73]. Alternatively, ~~if bran is chosen as substrate to produce citric acid, A.~~ *niger* is previously grown in a glucose solution for 24-48 ~~hours~~, at 35-38°C and 150-300 rpm. ~~Thereafter,~~ it is transferred in a seed tank with the bran medium and it is cultured at 36-38°C, for 4-6 days. After the adaptation to the new substrate, it is ~~transferred to a new tank~~ mixed with a bran solution ~~to start for the fermentation step~~, as described in the patents CN103667372, CN102649971 [74], [75]. Soybean molasses or sorghum powder are chosen as organic source in the inventions CN103614421, CN102864185, without a pretreatment. In both cases the selected conditions are: 26% (w/v) of sugar content, 35°C, pH 4.5, time 4 days and ventilating to supply oxygen [76], [77]. On the other hand, the wheat straw (patent CN1884563) and the rice grain (patent CN1693470) require a preliminary enzyme addition (e.g. cellulase or alpha-amilase), at the concentration of 20 units/~~per~~ substrate gram. The further hydrolysis is conducted at the 40-45°C for 40-80 ~~hours~~, followed by a solution filtration ~~and sterilization~~, and the fungal inoculation [78], [79].

### 3.2 Fruit and lignocelluloses as carbon source

The possibility of ~~the -kiwi use as organic source a fruit use-~~ is explained in the patent CN105671093, which produces citric acid-CA by *A. niger* by ~~kiwi as organic source, using~~ a solid-state fermentation set-up. ~~The kiwi fruit offeuts residues~~ are washed and dried, at low temperature, to obtain a water content lower than 6%. Thereafter, the fruit product is ~~cut to achieve a powder where~~ pulverized and it is used for the fungal ~~is-inoculated~~ edion; ~~t~~ The fermentation starts at 27-33°C and it carries on for 4-6 days. ~~At the end of the fungal metabolism, the resulting citric acid is extracted and recovered as a crystal salt~~ [80]. Further patents use grape pomace (US4791058), apple pomace (US4767705) or other kind of fruit (GB302338). In these cases, pretreatments are not required since the glucose is available for the fungal-metabolism [81]–[83]. On the other hand, the orange peel, ~~showed in the invention KR930001261, needs is drying dried and pulverization~~ pulverized (to obtain a 15-30 mesh powder), followed by the addition of and mixed with water,

~~ammonium nitrate~~ $\text{NH}_4\text{NO}_3$  and ~~magnesium sulfate~~ $\text{MgSO}_4$  (to stimulate the fungal fermentation);.

~~After~~ a pH adjustment ~~up to at~~ 4.5, ~~and a sterilization.~~ ~~At the end of these steps,~~ the submerged fermentation starts at 10-30°C, for 3 days [84]. ~~With the same aim, A~~ pretreatment is ~~necessary~~ ~~carried out on also for~~ the lignocellulose ~~raw materials (as described by CN104805136.)~~ ~~to convert the cellulose in glucose~~ thanks to the enzymatic hydrolysis by the enzyme cellulase [85]. ~~Patent CN105506004 uses P~~ pulverized konjak ~~is used as carbon source in patent CN105506004,~~ obtaining a ~~citric acid~~ $\text{CA}$  production ~~with a final yields~~ of 150 g/L [86].

ha formattato: Pedice

ha formattato: Pedice

ha formattato: Pedice

### 3.3 Starchy materials as carbon source

~~Starchy materials~~ ~~The need a~~ preparation ~~to be used as substrate~~ ~~of the substrate is necessary when~~ ~~starchy materials are used as carbon source for A. niger metabolism, as reported in the present~~ ~~section. In this regard,~~ ~~t~~ ~~The invention CN104232699 provides suggests~~ a preliminary ~~solid fraction removal~~ ~~solid and liquid separation,~~ followed by the addition of the *A. niger*, previously cultivated in a ~~second~~ broth with untreated starchy material, for the fungal adaptation [87]. ~~As an alternative~~ ~~Alternatively,~~ the techniques ~~presented~~ in patents CN103290070, CN103146769 and CN102839203 requires an ~~water supply~~ ~~alpha~~-amylase ~~supply~~ and a high temperature to form a mixed slurry; ~~a liquefied liquid is obtained by the continuous injections liquefying technology at 70-85°C.~~ After a solid/liquid separation, the resulting solution is used for the ~~citric acid~~ production, and the ~~fungal~~ fermentation is carried out at 38-40°C. Thereafter, ~~a Nitrogen source is added the~~ ~~to the~~ medium ~~is sterilized,~~ and ~~a nitrogen source is added,~~ a ventilation is applied for ~~the first 20-36 hours,~~ feeding the culture broth in batch, for 16-22 ~~hours~~. The fermentation finishes when the residual sugar is below 1% (w/v), ~~so the medium is sterilized at 75-85°C and citric acid is recovered from the solution~~ [88]–[90]. ~~Another method, described in p~~ Patent CN101942487 ~~describes a method for~~ suitable for starchy material, ~~in general, also applied and reported by several patents for with~~ specific substrates (cheap dried sweet potatoes CN102952830, puffed dried sweet potato CN102851329, unhusked CN1415755). ~~It~~ includes the addition of ~~alpha~~-amylase enzyme

for the reduction to amino acids and the ~~implementation-increase~~ of the bio-availability, ~~avoiding-~~  
~~The four described techniques avoid the a~~ further liquid/solid separation [91]–[94]. The ~~possible~~  
 addition of a ~~nitrogen-N~~ source, ~~selected in the the~~ patent RU2007125728, ~~further~~ improves the  
 conversion rate of sugar up to ~~87.6-7-93.2%~~ [95]. ~~Alternatively, e~~Comparable results are achieved  
 in the invention RU2186850 using metals- ~~(i.e., like Zn, Fe or Cu)~~ [96].

### 3.4 Molasses as carbon source

Same roots, commonly used for the commercial sugar production, are suitable for the ~~citric acid~~CA  
 synthesis ~~by A. niger metabolism~~. Among these substrates, cassava is chosen ~~in patents~~  
~~CN103045659, CN102864183~~, without any pretreatments, ~~in patents — CN103045659,~~  
~~CN102864183, where~~carrying out the fermentation ~~is carried out at~~ 39°C, for 3-4 days and 300  
 rpm [97], [98]. Satisfying results ~~(up to 15% (w/v) of citric acid CA, oxalic acid free)~~ are obtained  
 by the molasses from beet sugar manufacturing, as described in the inventions GB951629 and  
 GB799752. In the first one, 0.3% ~~potassium ferrocyanide~~ $K_4[Fe(CN)_6]$  and 0.2% ~~potassium~~  
~~dihydrogen~~ $KH_2PO_4$  are added to the molasses, which shows a sugar amount of 6.0-7.5% (w/v)  
 and a pH ~~between 5.7- and 5.9~~. This substrate is inoculated with *A. niger* in the presence of ~~passing~~  
~~sparging~~ air and the resulting fungal pellets are added to a similar ~~sterilized~~ medium, at pH 6.8 and  
 a sugar content around 15% (w/v). The solution is agitated ~~and aerated~~ at 30°C for 40-48 hours,  
 the pH is adjusted to 3.0 by ~~hydrochloric acid~~HCl and the air supply is replaced by the oxygen  
 insufflation to stimulate the ~~citric acid~~CA production. ~~The resulting solution shows a citric acid~~  
~~content of 15% (w/v), oxalic acid free~~ [99]. In the second patent, the beet sugar molasses, with a  
 sugar amount of 10-15% (w/v), is pretreated with ~~lime~~  $CaCO_3$  ~~or an insoluble hydroxide~~ for the  
 impurity removal. Before the fungal inoculation, nutrient salts are added in the following amount  
 (w/v): 0.15-0.2% of ~~ammonium carbonate~~ $(NH_4)_2CO_3$ , 0.01-0.02% of ~~potassium acid~~  
~~phosphate~~ $KH_2PO_4$ , 0.08-0.15% of ~~hydrate magnesium sulphate~~ $MgSO_4$  and 0.0002-0.0004% of  
~~zinc~~  $Zn$ . ~~The resulting liquor contains about 10-12% of citric acid~~ [100].

ha formattato: Pedice

ha formattato: Pedice

ha formattato: Pedice

ha formattato: Pedice

ha formattato: Pedice

ha formattato: Pedice

ha formattato: Pedice

ha formattato: Pedice

ha formattato: Pedice

ha formattato: Pedice

ha formattato: Pedice

This paragraph encloses around 40% of the patents of the present review. This is due to the high price of glucose (or sucrose), as substrate from raw material, that increases the industrial production cost of the citric acid [23]. The positive cost effect achieved by the waste use can be further enhance by the use the biomass from the fermentation for both the biogas production and the fertilizers preparation [28]. Overall the main materials selected for the inventions are the corns (around 45%) and the starchy (30%), because they combine the low cost with a great availability. The patents confirm the relevance of the multiple step set-up, where the first one is necessary for the fungal adaptation.

#### 4. *A. niger*, Citric acid production by *Aspergillus niger* – mutagenic strain

Since the sixties, the researches have experimented innovative approaches which have involved the fungal genome mutations to increase the citric acid production by *A. niger* fermentation. The techniques of UV, gamma-ray-induced and chemical mutagenesis are currently accepted as routine methods. Despite of although the patent number is lower than that of the inventions where the related to fungal is used as a the wild strain, use the technique of UV, gamma ray induced and chemical mutagenesis are currently accepted as routine methods in the citric acid synthesis field [101], [102]. Considering the strong interest for this kind of approach, next paragraph shows the registered patents related to the mutation strain techniques (Figure 523).

##### 4.1 Sugar as carbon source – mutagenic strain

The patent inventions CN106755138 and US2018195052 propose the modification of the succinic semialdehyde dehydrogenase (SSD) gene of the *A. niger* genome to obtain a recombinant *Aspergillus niger* strain. The expression of the SSD gene is regulated by the low pH inducible Pgas promoter which initiates the expression of the SSD protein in *Aspergillus A. niger*. Thereby final effect is the increasing of citric acid the production of citric acid by enhancing the GABA pathway [103]. The method, described in the inventions, utilizes the *A. niger* H915-1 as a host, with an efficiency increase of 10% and a time reduction of 10 hours, at 42°C [104], [105].



Alternatively, the patent CN106635847 integrates the *A. niger* low-affinity glucose transporter LGT1 gene into the *A. niger* genome to ~~obtain-achieve~~ the *A. niger* recombinant ~~*A. niger*~~. The present ~~invention-method utilizes-involves~~ a low pH inducible promoter to initiate the expression of LGT1 protein in *A. niger*, increasing the uptake of glucose during the acidogenic phase. The result is an ~~increase-improvement~~ of the ~~citric-acid~~CA production of 6.5%, at 42°C, with a time decrease of 10 hours [106]. ~~On the other hand, T~~the invention CN108018216 ~~proposes-includes~~ the modification of the glucosyltransferase genes to increase ~~both~~ the glucose assimilation and ~~the~~ conversion rate. The new genetic engineering strain ~~permits-allows~~ a sugar concentration ~~decrease~~ ~~reduction~~ of 10% and an increase of the ~~citric-acid~~CA yield around 10% ~~(w/v)~~ [107]. The ~~*A. niger*~~ *A. niger* RCAM 02149 ~~realized-~~ from the strain VKPM F-501, ~~described-(in-patent~~ RU2013151521), is produced by a genetic mutation ~~with-by~~ chemical mutagens (ether) and UV- radiation. ~~This method allows to-achieve-about 90% of sugar conversion-to-citric-acid,~~ at 32°C and a pH ~~between-1.7-and-7.0~~ [108]. In the patents CN103952318, CN103045487 the *A. niger* FY2013 ~~and FYCA8561~~ strains is prepared ~~starting-~~ from the FY2010 strain using a low-dose compound treatment ~~at-the-following-conditions-by~~ Co<sup>60</sup>-γ-ray treatment, ~~high-temperature-(at~~ 90°C, ~~for 5 minutes),~~ adding the nitrosoguanidine (0.1 mg/ml). The resulting ~~citric-acid~~ production is about 150 g/L [109], [110]. The patent CN103194398 is designed to obtain high-yield strains, named *A. niger* TN-A09, with high tolerance levels to ~~both~~ sugar and acid concentration and high-conversion rates. Overall, the production of ~~citric-acid~~CA is 18% (w/v), with a fermentation cycle of 60 hours ~~and-an-almost-complete-conversion-~~ [109]. On the other hand, the patent WO2013082459 provides a genetic enhancement of a LaeA gene or ~~an~~ inactivation ~~ion~~ the Alg3 gene of *A. niger* to increase the ~~acid production~~ ability of the fungus ~~to-produce-more-citric-acid-~~ [111]. ~~The *A. niger* FYCA8561 strain (presented in the patent CN103045487) is produced by a genetic mutation with γ ray treated with nitrosoguanidine. The use of this new fungal produces an acid concentration of 15% (w/v) and a conversion rate of 95% [111].~~ The *A. niger* CGMCC5342 and CGMCC5343 strains, ~~included-~~ in the inventions CN102533570 and CN102399702, respectively,

are obtained by the genetic mutation of *A. niger* Co 827. The method allows a citric acid production of 15% (w/v), after 52 hours at the applied conditions: 30-40°C and a starting pH between 4.0-4 and 5.0 [112], [113]. The A portion of spore suspension is collected, after a grown of 5-6 days, in patent CN102352322, provides the *A. niger* strain mutation, using the cultured spores grown for 5-6 days, eluted with sterile water, shaken for 30-60 minutes. A minimum portion of 0.1 mL of spore suspension is collected, spread on a sterile Petri dish surface and dried at room temperature. Thereafter, the culture dish coated is placed in an ion implanter, in the presence of and mutated by nitrogen-N ions. The mutant slant is inoculated in a medium at constant temperature around 40°C, 220 rpm and cultured for 90-96 hours. The achieved citric acid production is 10% higher than that of the original strain [114]. On the other hand, the pPatent RU2428481 uses the fungal strain *A. niger* VKPM F-696 which is grown in a solution containing with sugar, in the presence of a water-soluble complex of C60 fullerene (0.5-0.7% w/v), with polyvinylchloride (0.75-1.25 mg/mL). The use of the present this technology allows an increase of the citrate synthesis activity of 30%, from the third day with an increase improvement of the citric acid product content of about around 11%, and a final conversion efficiency around of 98% [115]. The *A. niger* No CCM8210 strain (described within the invention WO9710350) is developed from a starting variant obtained by protoplasts isolation from selected *A. niger*. The protoplasts are isolated from the hyphae in a stabilized aqueous solution (0.7 M NaCl + glucose), in the presence of calcium ions,  $\text{Ca}^{2+}$ . The resulting protoplasts are filtered and washed with water. The protoplast suspensions are radiated by UV radiations, for 5 minutes. The further fusion is performed with wild strains to increase the growth and the production ability. This *A. niger* strain is subjected to a further mutation by the combination of UV radiation and chemical mutagens (i.e. 5-bromouracil, 2-aminopurine, diethylsulphate, ethylethane sulphonates and their combinations). The achieved CCM8210 strain is a high producing mutants, cultivated on sugar media, able to start fermentation at low pH value (lower than 2.8) and to avoid the production of undesirable organic acids, mainly oxalic acid. The strain has a higher conversion rate than

conventional strains, about 1.3 g/L·h of citric acidCA per liter of fermentation broth, for each hour. Overall, the variant produces up to 93 kg of citric acidCA from 100 kg of supplied sugar [116]. The *A. niger* R-3 strain (produced in the patent US4380583) is selected from the *A. niger*-119 strain using ethyleneimine, N-nitrosomethylurea and UV-radiation, in the patent US4380583. Its properties include the resistance to antagonist bacteria that may occur in the process of citric acid during the fermentation. The Up to 95-99% of the whole synthesized acid synthesized by the new strain consists of citric acidCA with a yields that reaches the 100% of sugar conversion [117].

#### 4.2 Waste as carbon source - mutagenic strain

Molasses is the principal carbon source from food and agriculture wastes, used as substrate for the metabolism of *A. niger* mutagenic strain in the citric acid production. Overall, The described patents described below show two main differences: the genome mutation kind and the final citric acid production yield. The use of *A. niger* F-718 and molasses as substrate is reported in the invention RU2125607. The process includes a first stage with a medium at pH 4.0-5.0 and a fermentation carried out at 30°C. During the first 30-48 hours of the process, a continuous feeding of both molasses and nutrient salts solution is requested. The second stage uses the microorganism suspension overflowing from the first stage, at 30°C and pH 2.0-3.0 [118]. As an alternative, *A. niger* R-1 and P-1, are involved in the patents: FR2361330, SU568677, GB1499093. Both fungi are produced by the combined activity of: UV-radiation, ethylene imine and N-nitrosomethyl urea on *A. niger* EU-119 (*A. niger* R-1) and *A. niger* FY-119 (*A. niger* P-1). The new strains increase the sugar conversion rate from 89% to 99% and the citric acidCA production yield from 94% to 97%. Furthermore, these variants show significant advantage as both a high resistance to antagonist bacteria and a low sensitive to chemical the composition of the starting molasses, allowing the decrease of the molasses low quality substrate use [119]–[121]. The use of an alternative substrate, which a mix of molasses medium with and sugar solution, at different concentrations, is proposed within in the patent RU2203322. More in detail, using the *A. niger*

VKPM-817, the obtained sugar conversions to [citric acid](#) are: 87% with a 30 g/L sugar solution, 66% with 130 g/L and 96% with 150 g/L [122]. Sorghum condensed juice, molasses and sucrose are showed, as substrate, in the invention -RU2192460 that uses *A. niger* VKPM-809 with final conversion rates ~~between of~~ 60-90% [123]. A high conversion ~~rate of~~ sugar to organic acid ~~in the range (-9-41%)~~, is ~~obtained described~~ in the patent RU95113067 by the fungal VKPM-713 strain ~~-(from the A. niger F-326 strain genetic mutation)~~ by UV, using several ~~raw~~ substrates (i.e. beet and sugar cane molasses sugar, food sugar, crude sugar, glucose and their mixtures, and fermentation conditions) [124]. Alternatively, the microorganism strains 1015, 10577, 11414, 12846, 9142, 13794, 26036, with sugar beet or date molasses as carbon source, are used in patent FR2336477. In this case, the highest acid production rates (95% of [citric acid](#)) is achieved at these conditions: substrate pretreated by [potassium ferrocyanide](#)  $K_4[Fe(CN)_6]$  (to remove ~~heavy~~ metal traces), pH 2.7, 24 ~~hours~~ [125]. A pretreatment is required in the patent GB799752, for the impurity removal from molasses, corn starch hydrolyzed or beet molasses, adding ~~lime or an insoluble hydroxide~~  $CaCO_3$ . The ~~necessary~~ nutrient salts ~~amount is are~~ added to the broth: 0.15-0.2% of [ammonium carbonate](#)  $(NH_4)_2CO_3$ , 0.01-0.02% of [potassium acid phosphate](#)  $KH_2PO_4$ , 0.08-0.15% ~~hydrated magnesium sulphate~~  $MgSO_4$  and 0.0002-0.0004% of [zinc](#)  $Zn$ , at pH 2.5. The fermentation is carried out at 24-34°C by the *A. niger* mutant strain and the resulting [citric acid](#) production is about 10% (w/v), after 8-12 days [100]. The *A. niger* R-6 and R-5 strains are used with sugar beet molasses (LV11342) and sugar cane molasses (LV11340), as substrates, respectively. The fungal fermentation is carried out with a sugar amount of 15% (w/v), for 9-12 days. The resulting sugar conversion rate reaches the 100% in the first medium and the 85% in the second one, with a the [citric acid](#) content of 95-99%, in both cases [126], [127]. Alternative substrates are used with mutagenic strain, in patents CN106367359, CN108588133 (acorn) and RU2103346 ([ethanol](#)  $EtOH$ ). In the first invention, *A. niger* AA120, obtained after mutagenic screening by the atmospheric and room temperature plasma (ARTP) technology, allows high [citric acid](#) and tannic acids production [128]. In the second one, acorns are pretreated with [ethanol](#)  $EtOH$ ,

ha formattato: Pedice

ha formattato: Pedice

ha formattato: Pedice

ha formattato: Pedice

ha formattato: Pedice

ha formattato: Pedice

ha formattato: Pedice

ha formattato: Pedice

ha formattato: Pedice

to remove phenolic substances, and liquefied before the *A. niger* CICC 2716 inoculum. The acorn pulverized use improves the [citric acid](#) production [129]. In the last patent, *A. niger* VKPM-722 synthesizes the [citric acid](#) with a conversion rate higher than 80%, using [ethanol](#) or a mix of [EtOH](#) and sucrose [130].

Considering the deepened study carried out on the traditional strains, the best operative conditions have been identified and further development could mainly focus on the innovative substrates. Therefore, the possibility of mutagenic strains use has started to gain a foothold in the market, since they allow significant [citric acid](#) synthesis improvements. Currently, the inventions are equally divided between primary and waste substrates, but the trend of patents with residue use is destined to grow in a circular economy perspective.

## 5. — Other fungi

The technological innovation literature reports the use of other fungi for the citric acid production, as an alternative to the *A. niger*. The related patents are mainly concentrated in the 70s, nevertheless, the interest has waned for the achieved low efficiencies. In this regard, the following paragraphs focus on: *Candida* sp., *Penicillium* sp. and other strain of *Aspergillus*, considered as the most representative for the technological innovation (Figure 6).

### 5.1 — *Candida* sp.

The required conditions for the citric acid production by *Candida* sp. fermentation, reported in patents DE4407441, US4178211, GB1464334, include: a carbon source, mainly glucose, with a concentration higher than 200 g/L, a nitrogen concentration between 50 and 150 mM and an atmospheric oxygen saturation in the fermenter of 20%, pH around 5.0, at 30°C [131] [133]. The glucose substitution is proposed in patent GB1418561, where the *C. guilliermondii* strain produces citric acid metabolizing cane or beet molasses after the addition of essential nutrients. The process

needs a pH value between 2.5 and 4.5 (controlled by ammonia addition), at 22–35°C, at aerated conditions obtained by sparging air. In order to maintain the high citric acid production rate, the medium is recovered and replaced with fresh solution [134]. On the other hand, the possibility to use hydrocarbon fraction is verified in the further inventions. In this regard, alpha-olefins and n-paraffin, both with 8–20 C atoms, or a mix of them, are employed for the metabolism of *C. tropicalis*, *C. lipolytica*, *C. intermedia* and *C. brumptii*, in patents US4424274 and JPS5779890. The treatments require the concentration of dissolved oxygen in a range of 10–30 ppm with a final acid concentration greater than 150 g/L [135], [136]. The invention GB1428440 uses *Candida* sp. in a two-fermentation phases; in the first step the fungal is cultured in an aerated tank containing at least one n-paraffin, for its adaptation, thereafter, it is cultured in a second fermenter where, further inorganic nitrogen is added at acid pH [137]. N-paraffin (9–20 C atoms) is employed also in patent GB1418511, for the *C. lipolytica* and *C. oleophila* metabolisms in a medium with a high fluoroacetate concentration. A specific agent (pentachlorophenol, 2,4 dinitrophenol, 4,5,6-7-tetrachloro-2-(trifluoromethyl)benzimidazole, mesoxalonitrile[p(trifluoromethoxy)–phenol]–hydrazone or dicoumarin) is added to the medium to decouple the substrate, in order to increase its bioavailability [138]. N-paraffin (10–18 C atoms, with the concentration of 3–20% v/v), sugar, glycerin, ethanol, acetic acid, butyric acid or animal or vegetable fats or oil are suitable for the *C. lipolytica* and *C. tropicalis* metabolisms, as confirmed by the inventions GB1380938, GB1297243, GB1204635. In these processes, the medium is enriched by inorganic nitrogen and other nutrient salts, at pH 1.5–3.5, under aerobic conditions, at 20–35°C [139]–[141].

## 5.2—*Penicillium* sp.

The patents US3652396, GB878151 include the fungal strains *P. adameizi* and *P. restrictum* with hydrocarbon, sucrose or cane and beet molasses, as carbon sources. In both cases, pretreatments are not necessary and inorganic nutrient (e.g. nitrogen, phosphorous and magnesium) are added. The fermentation is carried out at pH around 7, at 28°C [142], [143].

### 5.3—*Aspergillus sp.*

Patent GB581389 mentions *A. wentii*, as an alternative to *Aspergillus* strain, which metabolizes cane sugar enriched with nutrient salts (e.g. ammonium nitrate, peptone, magnesium sulphate, dipotassium hydrogen phosphate, potassium chloride, zinc sulphate, ferric chloride and calcium chloride). When the fungal has reached the maximum growth rate, the residual culture medium is recovered and replaced by a fresh fermentation liquor containing 15–20% of carbohydrates (e.g. sucrose, maltose, lactose, glucose, dextrose, levulose). The treatment is carried out at pH 3.0, maintained by calcium carbonate or an alkali metal hydroxide addition [144]. Alternatively, patent GB797390 suggests molasses as carbon source. Before the fermentation, the broth medium is treated by ion-exchange resins,  $\text{Ca}(\text{OH})_2$ , alkaline ferrocyanides or complex-forming agents, for the metal removal, mainly iron. The broth pH is maintained around 2.5 at the 32°C [145].

### 6.5. Discussion and perspectives

Figure 3 shows a whole citric acid production process which summarizes the treatments described in the present review, involving *A. niger*. As reported in the supporting materials (Tables S1), the technological innovation literature describes the further use of other fungi [131], [132], [141]–[145], [133]–[140]. Nevertheless, the related patents are mainly concentrated in the 70s and the interest has waned for the achieved low efficiencies; for this reason, these techniques have not been discussed in the present review. (reported in the Tables S1), Figure 7 shows a general citric acid production process.

The first identified variable in the scheme (Figure 3A) is the possibility of a pre-treatment, if food or agriculture waste are used as substrate for the fungal growth. This step is possibly required to remove metals and/or to make carbon source bio-available for the fungal metabolism. Currently, the industrial production mainly uses sugars from hydrolysis of plant starch. The following step includes the fermentation, irrespective of the selected conditions and the substrate kind (glucose or sucrose), the following step includes the fermentation. The average

ha formattato: Non Evidenziato

conditions which promote the fungal metabolism, with ~~a consequent~~ high sugar conversion rate (around 95%) and a high acid concentration (100-150 g/L) ~~are include~~: sugar concentration 10%-15% (w/v), pH 1.5-4.5, stirring speed 120-400 rpm, temperature 27°-35°C, air supply and nutrient salts addition (mainly 0.2% w/v of (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, ~~0.2%~~ and w/v of KH<sub>2</sub>PO<sub>4</sub> and 0.1% of MgSO<sub>4</sub>) and/or metal addition (Mn, Fe and Zn). A final phase of recovery and purification closes the treatment ~~schema~~.

As concern the technological evolution overview, the sharp increase ~~in of~~ the number of patents ~~related to the citric acid bioproduction~~, in the last 20 years, confirms the strong interest for ~~this topic~~ the acid bioproduction, especially in China. As showed in Figure [3B8](#), there is an evident change in the invention origin, from European and US to Chinese, starting from 2000. The reason can be found in a modification a of the trade routes, ~~modification~~ which currently identify China as one of the most important export-oriented country. In this regard, 824,000 tons were exported from China, principally to India (9%), Turkey (5.2%), Japan (5%), Mexico (5%) and Indonesia (3.9%), in 2015. In the same year, 1,007 tons of citric acid CA was imported to China mainly from Germany (30%), Canada (18%), Japan (17%) and Austria (12%), with a whole cost of USD 5 million [146].

Furthermore, considering the ~~further~~ increase of the citric acid CA demand, China aimed at the production process improvement with an increase of its economic exportation load of about 10%, from 2015 to 2017. In this regard, Figure [4A9a](#) attributes to China the 75% of the whole trade value connected with the citric acid CA exportation, followed by Netherland, USA and Germany (each one with a contribution around 3%). On the other hand, USA (12%), Germany (8%), Mexico (5%) and India (5%) represent the main importers in the world (Figure [49bB](#)). Whereas, the Chinese production independence is confirmed by an importation trade lower than 1% [147].

Additional information about the main critical aspects addressed in the patents can be deduced from the trends in Figure [3B8](#). In this regard, the current technological innovation, has focused on



the identification of innovative substrates, mainly agriculture waste, in agreement with the scientific literature which tends to fulfill the circular economy rudiments [148].

## 6. Conclusions

The present review proves the necessity of a technological innovation study to understand the real state-of-art in a specific field. A patented invention is designed for the implementation on a real scale and it should be commercially available. Furthermore, for the grant of a patent it is necessary to prove the actual novelty level, ensuring the innovation level of the review. Compared to the traditional literature, the invention overview helps with the market previsions thanks to the additional information related to the origin and the state of development [27]. This focus on the patents about the ~~citric acid~~ production, over a long period of time, highlighted the growing interest for this topic and a technology progress, in agreement with the market demand. In this regard, a relevant increase of the waste substrate proves the development of the circular economy model. Further studies, with a similar approach, could deep the aspect of the ~~Furthermore, there is a change in the~~ fermentation design change. The scientific literature confirms as the research ~~which has~~ moved towards the conversion of the most consolidated submerged process to the innovative solid-state fermentation (*koji* process) [4], [149]–[151]. This simple and eco-sustainable technique combines the possibility of an agro-industrial waste use with a low energy demand and a minimum wastewater ~~production~~ production [27]. ~~The choice of a limited in-depth of this topic is due to the~~ low number of the related patents, probably connected with the current scale-up difficulties. This technique cannot be implemented in the bioreactors for submerged process and many criticalities are still due to the lack of standardized processes. ~~n-~~

**Declarations of interest:** The authors report no declarations of interest.

## References

[1] Luo H, Cheng X, Liu G, et al. Citric acid production using a biological electrodialysis with bipolar membrane. J Memb Sci. 2017;523:122–128.

ha formattato: Tipo di carattere: (Predefinito) Times New Roman, 12 pt, Grassetto

Formattato: Interlinea: doppia

ha formattato: Tipo di carattere: (Predefinito) Times New Roman

ha formattato: Tipo di carattere: Non Grassetto

- [2] [Ciriminna R, Meneguzzo F, Delisi R, et al. Citric acid: Emerging applications of key biotechnology industrial product. Chem Cent J. 2017;11\(1\):1–9.](#)
- [3] [Sun X, Lu H, Wang J. Recovery of citric acid from fermented liquid by bipolar membrane electrodialysis. J Clean Prod. 2017;143:250–256.](#)
- [4] [Kuforiji OO, Kuboye AO, Odunfa SA. Orange and pineapple wastes as potential substrates for citric acid production. Int J Plant Biol. 2010;1\(1\):19–21.](#)
- [5] [Dhillon GS, Brar SK, Verma M, et al. Utilization of different agro-industrial wastes for sustainable bioproduction of citric acid by \*Aspergillus niger\*. Biochem Eng J. 2011;54\(2\):3–92.](#)
- [6] [Zhang A, Roehr M. Effects of varied phosphorus concentrations on citric acid fermentation by \*Aspergillus niger\*. Acta Biotechnol. 2002;22\(3–4\):383–390.](#)
- [7] [Zhang A, Roehr M. Citric acid fermentation and heavy metal ions - II. The action of elevated manganese ion concentrations. Acta Biotechnol. 2002;22\(3–4\):375–382.](#)
- [8] [Ferreira P, Lopes M, Mota M, et al. Oxygen transfer rate and pH as major operating parameters of citric acid production from glycerol by \*Yarrowia lipolytica\* W29 and CBS 2073. Chem Pap. 2016;70\(7\):869–876.](#)
- [9] [Liu X, Xu J, Xia J, et al. Improved production of citric acid by \*Yarrowia lipolytica\* using oleic acid as the oxygen-vector and co-substrate. Eng Life Sci. 2016;16\(5\):424–431.](#)
- [10] [Magnuson JK, Lasure LL. Organic Acid Production by Filamentous Fungi. Boston \(MA\): Springer; 2004.](#)
- [11] [Dhillon GS, Brar SK, Verma M, et al. Recent advances in citric acid bio-production and recovery. Food Bioprocess Technol. 2011;4\(4\):505–529.](#)
- [12] [Pazouki M, Panda T. Recovery of citric acid – a review. Bioprocess Eng. 1998;19\(6\):435–439.](#)
- [13] [Wennersten R. The extraction of citric acid from fermentation broth using a solution of a tertiary amine. J Chem Technol Biotechnol. 2008;33\(2\):85–94.](#)
- [14] [Dhillon GS, Brar SK, Verma M, et al. Recent advances in citric acid bio-production and recovery. Food Bioprocess Technol. 2011;4\(4\):505–529.](#)
- [15] [Ledakowicz S, Jamroz T, Sencio B, et al. Equilibrium and dynamic investigations of organic acids adsorption onto ion-exchange resins. Bioprocess Biosyst Eng. 2004;26\(3\):185–190.](#)
- [16] [Yu D, Shi Y, Wang Q, et al. Application of methanol and sweet potato vine hydrolysate as enhancers of citric acid production by \*Aspergillus niger\*. Bioresour Bioprocess. 2017;4\(1\):35.](#)
- [17] [Wang B, Chen J, Li H, et al. Pellet-dispersion strategy to simplify the seed cultivation of \*Aspergillus niger\* and optimize citric acid production. Bioprocess Biosyst Eng. 2017;40\(1\):45–53.](#)
- [18] [Auta HS, Abidoye KT, Tahir H, et al. Citric acid production by \*Aspergillus niger\* cultivated on \*Parkia biglobosa\* fruit pulp. Int Sch Res Not. 2014;2014:1–8.](#)
- [19] [Maharani V, Reeta D, Sundaramanickam A, et al. Original research article isolation and characterization of citric acid producing \*Aspergillus niger\* from spoiled coconut. Int J Curr Microbiol App Sci. 2014;3\(3\):700–705.](#)

- [20] Liu X, Lv J, Zhang T, et al. Citric acid production from hydrolysate of pretreated straw cellulose by *Yarrowia lipolytica* SWJ-1b using batch and fed-batch cultivation. *Prep Biochem Biotechnol*. 2015;45(8):825–835.
- [21] Show PL, Oladele KO, Siew QY, et al. Overview of citric acid production from *Aspergillus niger*. *Front Life Sci*. 2015;8(3):271–283.
- [22] Soccol CR, Vandenberghe LPS, Rodrigues C. New perspectives for citric acid production and application. *Food Technol Biotechnol*. 2006;44(2):141–149.
- [23] Hu W, Jian Li W, Quan Yang H, et al. Current strategies and future prospects for enhancing microbial production of citric acid. *Appl Microbiol Biotechnol*. 2019;103(1):201–209.
- [24] Tong Z, Zheng X, Tong Y, et al. Systems metabolic engineering for citric acid production by *Aspergillus niger* in the post-genomic era. *Microb Cell Fact*. 2019;18(1):1–15.
- [25] Soccol CR, Vandenberghe LP. Overview of applied solid-state fermentation in Brazil. *Biochem Eng J*. 2003;13(2–3):205–218.
- [26] Gowthaman MK, Krishna C, Moo-Young M. Fungal solid state fermentation - an overview. *App Mycology Biotech*. 2001;1:305–352.
- [27] Letti LAJ, Soccol CR, Karp SG, et al. Recent developments and innovations in solid state fermentation. *Biotechnol Res Innov*. 2017;1(1):52–71.
- [28] Sawant O. Fungal citric acid production using waste materials: a mini-review. *J Microbiol Biotechnol Food Sci*. 2018;8(2):821–828.
- [29] Amato A, Beolchini F. End of life liquid crystal displays recycling: A patent review. *J Environ Manage*. 2018;225:1–9.
- [30] Garcia R, Calantone R. A critical look at technological innovation typology and innovativeness: a literature review. *J Prod Innov Manag*. 2002;19(2):110–132.
- [31] Rocchetti L, Amato A, Beolchini F. Printed circuit board recycling: A patent review. *J Clean Prod*. 2018;178:814–832.
- [32] Zhang QW, Ye ZD, Shi L. c-Met kinase inhibitors: an update patent review (2014-2017). *Expert Opin Ther Pat*. 2019;29(1):25–41.
- [33] Li X, Wang C, Jiang H, et al. A patent review of arginine methyltransferase inhibitors (2010–2018). *Expert Opin Ther Pat*. 2019;29(2):97–114.
- [34] Worldwide.espacenet.com [internet]. Available from: <https://worldwide.espacenet.com/>
- [35] Murad AEH, Khalaf SAD. Citric acid production from whey with sugars and additives by *Aspergillus niger*. *African J Biotechnol*. 2003;2(10):356–359.
- [36] Xu D, Madrid C, Rohr M, et al. The influence of type and concentration of the carbon source on production of citric acid by *Aspergillus niger* Ding-Bang. *Appl Microbiol Biotechnol*. 1989;30(6):553–558.
- [37] Hang YD, Woodams EE. Solid state fermentation of apple pomace for citric acid production. *J Appl Microbiol Biotechnol*. 1986;2(2):283–287.

- [38] Vandenberghe LPS, Soccol CR, Prado FC, et al. Comparison of citric acid production by solid-state fermentation in flask, column, tray, and drum bioreactors. *Appl Biochem Biotechnol*. 2004;118(1–3):293–304.
- [39] Vandenberghe LPS, Soccol CR, Pandey A, et al. Microbial production of citric acid. *Brazilian Arch Biol Technol*. 1999;42(3):263–276.
- [40] Arts E, Kubicek CP, Rohr M. Regulation of phosphofructokinase from *Aspergillus niger*: Effect of fructose 2,6-bisphosphate on the action of citrate, ammonium ions and AMP. *Microbiology*. 1987;133(5):1195–1199.
- [41] Ates S, Dingil N, Bayraktar E, et al. Enhancement of citric acid production by immobilized and freely suspended *Aspergillus niger* using silicone oil. *Process Biochem*. 2002;38(3):433–436.
- [42] Jin Y, inventor. Taicang Maotong Huajian Co LTD, assignee. Method for producing citric acid by utilizing immobilized *Aspergillus niger*. China patent CN 102,864,184. 2013 Jan 09.
- [43] Ying H, Yu B, Chen Y, et al, inventor. Univ Nanjing Tech, assignee. Immobilization method of *Aspergillus niger*. China patent CN 107,022,541. 2017 Aug 08.
- [44] Shi G, Liu L; Chen J, et al, inventor. Jiangsu Guoxin Xielian Energy Co LTD, Univ Jiangnan, assignee. Method for seed culture of *Aspergillus niger* and preparing citric acid. China patent CN 107,815,421. 2018 Mar 20.
- [45] Chen B, inventor. Qingdao Hua'nian Shengyuan Fruit Ind Co LTD, assignee. Preparation method of citric acid. China patent CN 106,868,061. 2017 Jun 20.
- [46] Yong Z, Zongmei L, Jixue Z, et al, inventor. Anhui BBKA Biochemical Co LTD, assignee. Method for producing citric acid. China patent CN 102,181,490. 2011 Sep 14.
- [47] Li B, Zhou Y, Lu Z, et al, inventor. Anhui BBKA Biochemical Co LTD, assignee. *Aspergillus niger* seed liquid preparation method and citric acid fermentation preparation method. China patent CN 104,277,978. 2015 Jan 14.
- [48] Xia L, Lu Z, Shao Y, et al, inventor. Anhui BBKA Biochemical Co LTD, assignee. Method for preparing citric acid through fermentation. China patent CN 102,851,330. 2013 Jan 02.
- [49] Yongsheng Z, Siliang Z, Yong Z, et al, inventor. Anhui BBKA Biochemical Co LTD, Univ East China Science & Tech, assignee. Citric acid preparation method. China patent CN 102,373,242. 2012 Mar 14.
- [50] Shi G, Wang B, Zhang J, et al, inventor. Univ Jiangnan, Yixing Union Biochemical Co LTD, assignee. Citric acid *Aspergillus niger* seed continuous culture method based on mycelium pellet dispersion technology. China patent CN 104,099,253. 2014 Oct 15.
- [51] Shi G, Wang B, Zhang J, et al, inventor. Univ Jiangnan, Yixing Union Biochemical Co LTD, assignee. Method for producing citric acid by continuous fermentation of *Aspergillus niger*. China patent CN 104,087,624. 2014 Oct 08.
- [52] Yongsheng Z, Yong Z, Zongmei L, et al, inventor. Anhui BBKA Biochemical Co LTD, assignee. Production method of citric acid. China patent CN 102,443,611. 2012 May 09.

- [53] Schreiferl G, Kubicek CP, Röhr M. Inhibition of citric acid accumulation by manganese ions in *Aspergillus niger* mutants with reduced citrate control of phosphofructokinase. *J Bacteriol.* 1986;165(3):1019–1022.
- [54] Datta R, Bergemann EP, inventor. NTEC INC, assignee. Process for producing of citric acid and monovalent citrate salts. United States patent US 5,532,148. 1996 Jul 02.
- [55] Kirkovits A, Edlauer H, inventor. Jungbunzlauer AG, assignee. Process for the fermentative production of citric acid from carbohydrates. United States patent US 5,081,025. 1992 Jan 14.
- [56] Kabil AJ, inventor. Jungbunzlauer Spiritus, assignee. Process for the production of citric acid. United States patent US 3,936,352. 1976 Feb 03.
- [57] Jungbunzlauer Spiritus UND CH, assignee. Process for the production of citric acid by submerged fermentation. Great Britain patent GB 1,392,942. 1975 May 07.
- [58] Jungbunzlauer Spiritus, assignee. Process for the production of citric acid by submerged fermentation. Great Britain patent GB 1,342,311. 1974 Jan 03.
- [59] Shi G, Chen J, Wang B, et al, inventor. Univ Jiangnan, Jiangsu Guoxin Union Energy Co LTD, assignee. Method for improving fermentation performance of citric acid on basis of mycelium structure control. China patent CN 105,586,366. 2016 May 18.
- [60] Luo H, Xiong J, Lu Z, inventor. Anhui BBKA Biochemical Co LTD, assignee. Method for preparing citric acid by fermentation. China patent CN 103,497,977. 2014 Jan 08.
- [61] Li R, Shang H, Yang W, et al, inventor. Anhui BBKA Fermentation Technology Engineering Res Co LTD, assignee. Bacterial strain for producing citric acid and method for preparing citric acid by fermenting same. China patent CN 103,695,319. 2014 Apr 02.
- [62] Anastassiadis S, Morgunov I, Kamzolova S, et al. Citric acid production patent review. *Recent Pat Biotechnol.* 2008;2(2):107–123.
- [63] Zhang M, Roehr A. Citric acid fermentation and heavy metal ions. *Acta Biotechnol.* 2002;22(3–4):363–373.
- [64] Yin L, inventor. Liuzhou Mingpin Tech Co LTD, assignee. Method for preparing citric acid from corncobs. China patent CN 106,119,306. 2016 Nov 16.
- [65] Shen X, Shen D, inventor. Shen Dechao, Zibo Huichuang Biotechnology LTD Company, assignee. Method for preparing fermentation liquor of citric acid by utilizing extrudate obtained after low-temperature extrusion and enzymolysis. China patent CN 105,524,951. 2016 Apr 27.
- [66] Li J, Li S, Ma Q, et al, inventor. Weifang Ensign Industry Co LTD, assignee. Technique for preparing citric acid by fermenting beta-cyclodextrin mother solution. China patent CN 103,710,397. 2014 Apr 09.
- [67] Shi G, Liu L, Chen J, et al, inventor. Jiangsu Guoxin Xielian Energy Co LTD; Univ Jiangnan, assignee. Method for producing citric acid fermentation liquor by two-stage fermentation. China patent CN 107,815,475. 2018 Mar 20.
- [68] Jin Y, inventor. Taicang Maotong Huajian Co LTD, assignee. Method for preparing citric acid by fermenting corn starch by using *Aspergillus niger*. China patent CN 102,864,182. 2013 Jan 09.

- [69] Sharova NY, Vybornova AA, Printseva TV, et al, inventor. FED Gosudarstvennoe Byudzhethoe Nauchnoe Uchrezhdenie FED Nauchnyj Tsentr Pishchevykh Sistem IM V M, assignee. Method for producing invertase and citric acid. Russia patent RU 2,676,144. 2018 Dec 27.
- [70] Miles LAB, assignee. Improvements in or relating to the production of citric acid. Great Britain GB 738,940. 1955 Oct 19.
- [71] Miles LAB, assignee. Improvements in or relating to the production of citric acid. Great Britain GB 742,972. 1956 Jan 14.
- [72] Jin Y, inventor. Taicang Maotong Huajian Co LTD, assignee. Method for preparing citric acid through fermenting corn sugar solution by immobilized *Aspergillus niger*. China patent CN 102,851,328. 2013 Jan 02.
- [73] Yongsheng Z, Huiping Z, Sixiang L, et al, inventor. Anhui BBKA Biochemical Co LTD, assignee. Preparation method of citric acid fermentation solution. China patent CN 101,555,497. 2009 Oct 14.
- [74] Sun W, Han H, Wei N, et al, inventor. Tianjin Res Inst Of Ind Microbiology, assignee. Method for preparing bran starter for citric acid by liquid inoculation. China patent CN 103,667,372. 2014 Mar 26.
- [75] Hu L, Yun M, Yingang S, et al, inventor. Cofco Biochemical Anhui Co LTD, assignee. Method for cultivating *Aspergillus niger* mouldy bran and method for preparing citric acid through fermentation. China patent CN 102,649,971. 2012 Aug 29.
- [76] Cheng Q, Meng L, Yi Y, et al, inventor. Univ Guangxi Sci & Technology, assignee. Method for producing citric acid by fermenting soybean molasses. China patent CN 103,614,421. 2014 Mar 05.
- [77] Li R, inventor. Anhui BBKA Fermentation Technology Engineering Res Co LTD, assignee. Method for producing citric acid by fermenting sorghum powder. China patent CN 102,864,185. 2013 Jan 09.
- [78] Chen HZ, inventor. INST Process ENG CAS, assignee. Method for fermenting and producing citric acid using steam-explosion straw as raw material. China patent CN 1,884,563. 2006 Dec 27.
- [79] Guo BZ, inventor. Guo Bing, assignee. New method of producing citric acid using paddy fermentation. China patent CN 1,693,470. 2005 Nov 09.
- [80] Tan R, Shen J, inventor. Wuchuan County Jinfeng Kiwi Crop Farmer Professional Coop, assignee. Method for producing citric acid from kiwi fruit offcut. China patent CN 105,671,093. 2016 Jun 15.
- [81] Hang YD, inventor. Cornell Res Foundation INC, assignee. Grape pomace as substrate for microbial production of citric acid. United States patent US 4,791,058. 1988 Dec 13.
- [82] Hang YD, inventor. Cornell Res Foundation INC, assignee. Apple pomace as substrate for microbial production of citric acid. United States patent US 4,767,705. 1988 Aug 30.
- [83] Montan UND Industrialwerke, assignee. A process for the production of citric acid by fermentation. Great Britain patent GB 302,338. 1929 Oct 17.

- [84] Kang SK, Choe JG, Kwon IB, inventor. Lotte Confectionery Co LTD, assignee. Process for preparing citric acid by tangerin peelings. Korea patent KR 930,001,261. 1993 Mar 27.
- [85] Bao J, Meng J, Zhang J, inventor. Univ East China Science & Tech, assignee. Method for producing citric acid by using lignocellulose raw material. China patent CN 104,805,136. 2015 Jul 29.
- [86] Tang H, Yang W, Qin Q, et al, inventor. Anhui BBFA Fermentation Tech Eng Res Co LTD, assignee. Method for producing citric acid by fermenting konjak powder residues. China patent CN 105,506,004. 2016 Apr 20.
- [87] Gao Z, Zhou G, Xiong J, et al, inventor. Anhui BBFA Biochemical Co LTD, assignee. Method for fermentation production of citric acid. China patent CN 104,232,699. 2014 Dec 24.
- [88] Liu C, Liu F, inventor. Nantong Kaisai Biochemical Engineering Equipment Co LTD, assignee. Method for producing citric acid through continuous batch feeding fermentation. China patent CN 103,290,070. 2013 Sep 11.
- [89] Luo H, Lu Z, Yang R, et al, inventor. Anhui BBFA Biochemical Co LTD, assignee. Method for preparing citric acid by fermentation. China patent CN 103,146,769. 2013 Jun 12.
- [90] Zhou Y, Man Y, Zhang X, et al, inventor. Cofco Biochemical Anhui Co LTD, assignee. Enzymolysis method of starch raw material and method for preparing citric acid by fermenting. China patent CN 102,839,203. 2012 Dec 26.
- [91] Guiyang S, Liang Z, Xiaodong J, et al, inventor. Univ Jiangnan, Yixing Union Biochemical Co LTD, assignee. Method for fermenting and preparing citric acid by adding saccharifying enzyme. China patent CN 101,942,487. 2011 Jan 12.
- [92] Jin Y, inventor. Taicang Maotong Chemcial & Building Materials Co LTD, assignee. Method for preparing citric acid through fermenting cheap dried sweet potatoes by utilizing *Aspergillus niger*. China patent CN 102,952,830. 2013 Mar 06.
- [93] Jin Y, inventor. Taicang Maotong Huajian Co LTD, assignee. Method for preparing citric acid through fermenting puffed dried sweet potato raw material by *Aspergillus niger*. China patent CN 102,851,329. 2013 Jan 02.
- [94] Guo B, Zhu J, inventor. Guo Bing, assignee. Method for producing citric acid by fermenting paddy rice. China patent CN 1,415,755. 2003 May 07.
- [95] Sharova NJ, Aleksandrovna PT, Vladimirovna VT, et al, inventor. State Institution All-Russian Research Institute of Food Flavors, Acids and Dyes of the Russian Academy of Agricultural Sciences, assignee. Method For Preparation Of Citric Acid,  $\alpha$ -Amilase And Glucoamylase. Russia patent RU 2,007,125,728. 2009 Jan 20.
- [96] Sharova NJ, Mushnikova LN, Pozdnjakova TA, et al, inventor. G Uchrezhdenie Vserossijskij N; Orov Kislot I Krasitelej Raskh; I Skij I Pishchevykh Aromatiza, assignee. Method of citric acid producing. Russia patent RU 2,186,850. 2002 Oct 10.
- [97] Li R, Shang H, Yang W, et al, inventor. Anhui BBFA Fermentation Technology Engineering Res Co LTD, assignee. Method for detoxification in production of citric acid by utilizing cassava fermentation. China patent CN 103,045,659. 2013 Apr 17.

- [98] Jin Y, inventor. Taicang Maotong Huajian Co LTD, assignee. Method for preparing citric acid by fermenting cassava raw material residue-removed clear solution with *Aspergillus niger*. China patent CN 102,864,183. 2013 Jan 09.
- [99] Svenska Sockerfabriks AB, assignee. A method of producing citric acid. Great Britain GB 951,629. 1964 Mar 11.
- [100] Miles LAB, assignee. Citric acid production. Great Britain GB 799,752. 1958 Aug 13.
- [101] Gupta S, Sharma CB. Citric acid fermentation by the mutant strain of the *Aspergillus niger* resistant to manganese ions inhibition. *Biotechnol Lett*. 1995;17(3):269–274.
- [102] Khurshid IH, Ashraf A, Rajoka Q. Mutation of *Aspergillus niger* for hyperproduction of citric acid from black strap molasses. *World J Microbiol Biotechnol*. 2001;17(1):35–37.
- [103] Kumar S, Puneekar NS. The metabolism of 4-aminobutyrate (GABA) in fungi. *Mycol Res*. 1997;101(4):403–409.
- [104] Jin S, Hu Z, Peng Y, et al, inventor. Jiangsu Guoxin Union Energy Co LTD, assignee. Method of increasing yield of citric acid produced by *Aspergillus niger* fermentation. China patent CN 106,755,138. 2017 May 31.
- [105] Liu L, Chen J, Du G, et al, inventor. Univ Jiangnan, Jiangsu Guoxin Union Energy Co LTD, assignee. Method for increasing citric acid production by *Aspergillus niger* fermentation. United States patent US 2,018,195,052. 2018 Jul 12.
- [106] Sun F, Hu Z, Jiang X, et al, inventor. Jiangsu Guoxin Union Energy Co LTD, assignee. Recombinant *Aspergillus niger* capable of improving yield of citric acid and preparation method of recombinant *Aspergillus niger*. China patent CN 106,635,847. 2017 May 10.
- [107] Li M, Lu F, Wei R, inventor. Univ Tianjin Science & Tech, assignee. Method for increasing sugar utilization rate and yield of citric acid in citric acid fermentation and application. China patent CN 108,018,216. 2018 May 11.
- [108] Vybornova TV, Sharova N, inventor. State Scientific Institution All-Russian Research Institute of Food Flavors, Acids and Dyes of the Russian Academy of Agricultural Sciences, assignee. Strain *Aspergillus niger* - Producer of citric acid. Russia patent RU 2,013,151,521. 2015 May 27.
- [109] Li R, Shang H, Xu B, inventor. Anhui Bbca Fermentation Technology Engineering Res Co LTD, assignee. Citric acid high-yielding *Aspergillus niger* FY2013 and application thereof. China patent CN 103,952,318. 2014 Jul 30.
- [110] Li R, Mu X, Li W, et al, inventor. Anhui BBKA Fermentation Technology Engineering Res Co LTD, assignee. Bacterial strain for producing citric acid and method for fermenting and producing citric acid through fermentation of bacterial strain. China patent CN 103,045,487. 2013 Apr 17.
- [111] Dai Z, Baker SE, inventor. Battelle Memorial Institute, assignee. Enhanced citric acid production in *Aspergillus*. United States patent WO 2,013,082,459. 2013 Jun 06.
- [112] Xiu C, Zongmei L, Hua Z, et al, inventor. Anhui BBKA Biochemical Co LTD, assignee. *Aspergillus niger*, application of *Aspergillus niger* and method for preparing citric acid by fermentation. China patent CN 102,533,570. 2012 Jul 04.



- [113] [Junhua Z, Yong Z, Hua Z, et al, inventor. Anhui BBKA Biochemical Co LTD, assignee. \*Aspergillus niger\* and application thereof as well as citric acid preparation method through fermentation. China patent CN 102,399,702. 2012 Apr 04.](#)
- [114] [He H, Shuang L, Jing X, et al, inventor. Univ Nanjing, assignee. High-temperature-resistant citric acid producing strain. China patent CN 102,352,322. 2012 Feb 15.](#)
- [115] [Nikiforova TJA, Komov VP, Vybornova TJV, et al, inventor. G Uchrezhdenie Vrnii Pishchevykh Aromatizatorov Kislot I Krasitelej Rossijskoj Akademii Sel Skokhozj, assignee. Citric acid production method. Russia patent RU 2,428,481. 2011 Sep 10.](#)
- [116] [Minarik M, Skvarenina D, Michalik P, et al, inventor. Likospol SRO, Minarik Martin, Skvarenina Dusan, Michalik Peter, Sitkey Vladimir, Visacky Viliam, assignee. Process, apparatus and microorganism strain for the manufacture of citric acid. Slovakia patent WO 9,710,350. 1997 Mar 20.](#)
- [117] [Karklin RY, Rumba AA, Azanda VK, inventor. Karklin Roman Y, Rumba Alma A, Azanda Via K, assignee. Method of preparing seeding material for production of citric acid. United States patent US 4,380,583. 1983 Apr 19.](#)
- [118] [Avchieva PB, Berezhnoj JD, inventor. Nerubio Investments BV, assignee. Continuous method of producing citric acid and its salts. Russia patent RU 2,125,607. 1999 Jan 27.](#)
- [119] [Rumba AA, Karklin RY, Azanda VK, et al, inventor. Ex Z Biokhim, assignee. Citric acid prepn. by fermentation - using \*Aspergillus niger\* R-1 strain on nutrient medium contg. molasses. France patent FR 2,361,330. 1978 Mar 10.](#)
- [120] [Rumba AA, Karklinsh RY, Azanda VK, et al, inventor. Ex Z Biokhim Preparatov An Lat, assignee. \*Aspergillus niger\* P-1 strain as producer of citric acid. Switzerland patent SU 568,677. 1977 Aug 15.](#)
- [121] [Ex Z Biokhim Preparatov I Mikr, assignee. Method of preparing citric acid. Great Britain patent GB 1,499,093. 1978 Jan 25.](#)
- [122] [Shcherbakova EJ, Nikiforova TA, Vova EBL, inventor. Orov Kislot I Krasitelej Raskh, I Skij I Pishchevykh Aromatiza, G Uchrezhdenie Vserossijskij N, assignee. Diploid strain of \*Aspergillus niger\* as producer of citric acid. Russia patent RU 2,203,322. 2003 Apr 27.](#)
- [123] [Krasikova NV, Nikiforova TA, Fin KVM, inventor. G Uchrezhdenie Vserossijskij N, Orov Kislot I Krasitelej Raskh, I Skij I Pishchevykh Aromatiza, assignee. Strain of fungus \*Aspergillus niger\* as producer of citric acid. Russia patent RU 2,192,460. 2002 Nov 10.](#)
- [124] [Ermakova VP, Golubtsova VM, Sakson EE, et al, inventor. Sankt Peterburgskogo ZD A Limonnoj Kisloty TOO, assignee. Strain of fungus \*Aspergillus niger\* F-713 - a producer of citric acid. Russia patent RU 95,113,067. 1997 Jun 20.](#)
- [125] [Fazeli Abbas, assignee. Citric acid prodn by fermentation of sugar beet or date molasses - using \*Aspergillus niger\* of \*Mucor piriformis\* sp. organism. France patent FR 2,336,477. 1977 Jul 22.](#)
- [126] [Karklins R, Skrastina I, inventor. Romans Karklins, Ingemara Skrastina, assignee. A method for preparing of citric acid producent \*Aspergillus niger\* ation of combustible materials especially of industrial and household waste. Latvia patent LV 11,342. 1996 Jun 20.](#)

- [127] Karklins R, Skrastina I, inventor. Romans Karklins, Ingemara Skrastina, assignee. Microscopical fungus *Aspergillus niger* R-5 as citric acid producent. Latvia patent LV 11,340. 1996 Jun 20.
- [128] Zhang N, Jiang J, Wei M, et al, inventor. Inst Chem Ind Forest Prod CAF, assignee. *Aspergillus niger* and application thereof to preparing citric acid from fermented acorns. China patent CN 106,367,359. 2017 Feb 01.
- [129] Zhang B, Zhu L, Li L, et al, inventor. Univ Shaanxi Normal, assignee. Method for preparing citric acid through liquid state fermentation of acorns. China patent CN 108,588,133. 2018 Sep 28.
- [130] Vinarov AJ, Sidorenko TE, Smetanina SE, et al, inventor. GNI Skij I, Biosinteza Belkovykh Veshchest, assignee. Fungus strain *Aspergillus niger* - a producer of citric acid. Russia patent RU 2,103,346. 1998 Jan 27.
- [131] Anastasiadis S, Aivasidis A, Wandrey C, inventor. Forschungszentrum Juelich GMBH, assignee. Fermentation process for the continuous production of citric acid. Germany patent DE 4,407,441. 1994 Sep 15.
- [132] Leavitt RI, inventor. Ethyl CORP, assignee. Process for producing citric acid. United States patent US 4,178,211. 1979 Dec 11.
- [133] Rech Et Dactivites Petrolieres, assignee. Process for the production of citric and isocitric acids by a culture of a diploid *Candida lipolytica* yeast. Great Britain patent GB 1,464,334. 1977 Feb 09.
- [134] Pfizer LTD, assignee. Citric acid production. Great Britain patent GB 1,418,561. 1975 Dec 24.
- [135] Matsumoto T, Ichikawa Y, Nagata T, inventor. Showa Oil, assignee. Process for producing citric acid from hydrocarbons by fermentation. United States patent US 4,424,274. 1984 Jan 03.
- [136] Fujimaki A, Osada T, Matsumoto T, inventor. Showa Oil, assignee. Method for suppressing production of isocitric acid in citric acid fermentation of petroleum hydrocarbon. Japan patent JPS 5,779,890. 1982 May 19.
- [137] Inst Francais Du Petrole, Rech Et Dactivites Petroliere, assignee. Fermentative production of citric acid. Great Britain patent GB 1,428,440. 1976 Mar 17.
- [138] Benckiser GMBH Joh A, assignee. Process for the production of citric acid and salts thereof. Great Britain patent GB 1,418,511. 1975 Dec 24.
- [139] Hitachi Chemical Co LTD, assignee. Process for producing citric acid by fermentation. Great Britain patent GB 1,380,938. 1975 Jan 15.
- [140] Werkwijze voor het langs microbiologische weg bereiden van citroenzuur. Great Britain patent GB 1,297,243. 1972 Nov 22.
- [141] Okumura S, Tsugawa R, Kamijo H, et al, inventor. Ajinomoto KK, assignee. A fermentation process for the production of citric acid. Great Britain patent GB 1,204,635. 1970 Sep 09.
- [142] Tanaka K, Kimura K, inventor. Kyowa Hakko Kogyo KK, assignee. Process for preparing citric acid by fermentation. United States patent US 3,652,396. 1972 Mar 28.
- [143] Kinoshita S, Tanaka K, Akita S, inventor. Kyowa Hakko Kogyo KK, assignee. Process for the production of citric acid by fermentation. Great Britain patent GB 878,151. 1961 Sep 27.

- [144] Merck & Co INC, assignee. Improvements in or relating to the production of citric acid by fermentation. Great Britain patent GB 581,389. 1946 Oct 10.
- [145] Usines De Melie, assignee. Improvements in or relating to a process for producing citric acid by aerobic fermentation of solutions containing molasses. Great Britain patent GB 797,390. 1958 Jul 02.
- [146] China Citric Acid Industry Went Cold [Internet]. China: Jiangsu Guoxin Union Energy CO LTD 2016 Aug 28. Available from: <http://www.jsqx.net/gxxl/Pages/China-Citric-Acid-Industry-Went-Cold.aspx>
- [147] D. of E. and S. A. D. United Nations International Trade Statistics Database [Internet]. ALL COMMODITIES | Imports and Exports | 2017. Available from: [https://trandeconomy.com/data/commodity\\_h2?commodity=291814&trade\\_flow=Export,Import&time\\_p](https://trandeconomy.com/data/commodity_h2?commodity=291814&trade_flow=Export,Import&time_period=2017)  
eriod=2017.
- [148] Singh G, Kaur S, Verma M. Utilization of different agro-industrial wastes for sustainable bioproduction of citric acid by *Aspergillus niger*. *Biochem Eng J.* 2011;54(2):83–92.
- [149] Karthikeyan A, Sivakumar N. Bioresource Technology Citric acid production by Koji fermentation using banana peel as a novel substrate. *Bioresour Technol.* 2010;101(14):5552–5556.
- [150] Alam Z, Bari N, Muyibi SA, et al. Solid state bioconversion of oil palm empty fruit bunches for production of citric acid by wild strains of *Aspergillus niger*. *Food Biotechnol.* 2010;24(1):19–36.
- [151] Kumar D, Jain VK, Shanker G, et al. Citric acid production by solid state fermentation using sugarcane bagasse. *Process Biochem.* 2003;38(12):1731–1738.

## References

- [1] H. Luo *et al.*, “Citric acid production using a biological electrodialysis with bipolar membrane,” *J. Memb. Sci.*, vol. 523, no. September 2016, pp. 122–128, 2017.
- [2] R. Ciriminna, F. Meneguzzo, R. Delisi, and M. Pagliaro, “Citric acid: Emerging applications of key biotechnology industrial product,” *Chem. Cent. J.*, vol. 11, no. 1, pp. 1–9, 2017.
- [3] X. Sun, H. Lu, and J. Wang, “Recovery of citric acid from fermented liquid by bipolar membrane electrodialysis,” *J. Clean. Prod.*, vol. 143, pp. 250–256, 2017.
- [4] O. O. Kuforiji, A. O. Kuboye, and S. A. Odunfa, “Orange and pineapple wastes as potential substrates for citric acid production,” *Int. J. Plant Biol.*, vol. 1, no. 1, pp. 19–21, 2010.
- [5] G. S. Dhillon, S. K. Brar, M. Verma, and R. D. Tyagi, “Utilization of different agro-industrial wastes for sustainable bioproduction of citric acid by *Aspergillus niger*,” *Biochem. Eng. J.*, vol. 54, no. 2, pp. 83–92, 2011.
- [6] A. Zhang and M. Roehr, “Effects of Varied Phosphorus Concentrations on Citric Acid Fermentation by *Aspergillus niger*,” *Acta Biotechnol.*, vol. 22, no. 3–4, pp. 383–390, Jul. 2002.
- [7] A. Zhang and M. Roehr, “Citric Acid Fermentation and Heavy Metal Ions—II. The Action of Elevated Manganese Ion Concentrations,” *Acta Biotechnol.*, vol. 22, no. 3–4, pp. 375–382, Jul. 2002.
- [8] P. Ferreira, M. Lopes, M. Mota, and I. Belo, “Oxygen transfer rate and pH as major operating parameters of citric acid production from glycerol by *Yarrowia lipolytica* W29

Codice campo modificato

- and CBS 2073," *Chem. Pap.*, vol. 70, no. 7, pp. 869–876, 2016.
- [9] X. Liu, J. Xu, J. Xia, J. Lv, Z. Wu, and Y. Deng, "Improved production of citric acid by *Yarrowia lipolytica* using oleic acid as the oxygen vector and co-substrate," *Eng. Life Sci.*, vol. 16, no. 5, pp. 424–431, 2016.
- [10] J. K. Magnuson and L. L. Lasure, "Organic Acid Production by Filamentous Fungi," in *Advances in fungal biotechnology for industry, agriculture, and medicine*, J. S. Tkacz and L. Lange, Eds. Boston: Springer, 2004, pp. 307–340.
- [11] G. S. Dhillon, S. K. Brar, M. Verma, and R. D. Tyagi, "Recent Advances in Citric Acid Bio-production and Recovery," *Food Bioprocess Technol.*, vol. 4, no. 4, pp. 505–529, 2011.
- [12] M. Pazouki and T. Panda, "Recovery of citric acid—a review," *Bioprocess Eng.*, vol. 19, no. 6, pp. 435–439, Dec. 1998.
- [13] R. Wennersten, "The extraction of citric acid from fermentation broth using a solution of a tertiary amine," *J. Chem. Technol. Biotechnol. Biotechnol.*, vol. 33, no. 2, pp. 85–94, Apr. 2008.
- [14] G. S. Dhillon, S. K. Brar, M. Verma, and R. D. Tyagi, "Recent Advances in Citric Acid Bio-production and Recovery," *Food Bioprocess Technol.*, vol. 4, no. 4, pp. 505–529, 2011.
- [15] S. Ledakowicz, T. Jamroz, B. Sencio, and P. Gluszez, "Equilibrium and dynamic investigations of organic acids adsorption onto ion-exchange resins," *Bioprocess Biosyst. Eng.*, vol. 26, no. 3, pp. 185–190, Apr. 2004.
- [16] D. Yu, Y. Shi, Q. Wang, X. Zhang, and Y. Zhao, "Application of methanol and sweet potato vine hydrolysate as enhancers of citric acid production by *Aspergillus niger*," *Bioresour. Bioprocess.*, vol. 4, no. 1, p. 35, 2017.
- [17] B. Wang, J. Chen, H. Li, F. Sun, Y. Li, and G. Shi, "Pellet dispersion strategy to simplify the seed cultivation of *Aspergillus niger* and optimize citric acid production," *Bioprocess Biosyst. Eng.*, vol. 40, no. 1, pp. 45–53, 2017.
- [18] H. S. Auta, K. T. Abidoye, H. Tahir, A. D. Ibrahim, and S. A. Aransiola, "Citric Acid Production by *Aspergillus niger* Cultivated on *Parkia biglobosa* Fruit Pulp," *Int. Sch. Res. Not.*, vol. 2014, pp. 1–8, Nov. 2014.
- [19] V. Maharani, D. Reeta, A. Sundaramanickam, S. Vijayalakshmi, and T. Balasubramanian, "Original Research Article Isolation and characterization of citric acid producing *Aspergillus niger* from spoiled coconut," vol. 3, no. 3, pp. 700–705, 2014.
- [20] X. Liu, J. Lv, T. Zhang, and Y. Deng, "Citric Acid Production from Hydrolysate of Pretreated Straw Cellulose by *Yarrowia lipolytica* SWJ-1b Using Batch and Fed-Batch Cultivation," *Prep. Biochem. Biotechnol.*, vol. 45, no. 8, pp. 825–835, Nov. 2015.
- [21] P. L. Show, K. O. Oladele, Q. Y. Siew, F. A. Aziz Zakry, J. C. W. Lan, and T. C. Ling, "Overview of citric acid production from *Aspergillus niger*," *Front. Life Sci.*, vol. 8, no. 3, pp. 271–283, 2015.
- [22] C. R. Soccol, L. P. S. Vandenberghe, and C. Rodrigues, "New Perspectives for Citric Acid Production and Application," *Food Technol. Biotechnol.*, vol. 44, no. 2, pp. 141–149, 2006.
- [23] W. Hu, W. jian Li, H. quan Yang, and J. hong Chen, "Current strategies and future prospects for enhancing microbial production of citric acid," *Appl. Microbiol. Biotechnol.*, vol. 103, no. 1, pp. 201–209, 2019.
- [24] Z. Tong, X. Zheng, Y. Tong, Y. C. Shi, and J. Sun, "Systems metabolic engineering for citric acid production by *Aspergillus niger* in the post-genomic era," *Microb. Cell Fact.*, vol. 18, no. 1, pp. 1–15, 2019.
- [25] C. R. Soccol and L. P. S. Vandenberghe, "Overview of applied solid-state fermentation in Brazil," *Biochem. Eng. J.*, vol. 13, no. 2–3, pp. 205–218, Mar. 2003.
- [26] M. K. Gowthaman, C. Krishna, and M. Moo-Young, "Fungal solid state fermentation—

- an overview," in *Applied Mycology and Biotechnology*, vol. 1, 2001, pp. 305–352.
- [27] L. A. J. Letti, C. R. Soccol, S. G. Karp, E. S. F. da Costa, L. P. de S. Vandenberghe, and A. L. Woiciechowski, "Recent developments and innovations in solid-state fermentation," *Biotechnol. Res. Innov.*, vol. 1, no. 1, pp. 52–71, 2017.
- [28] O. Sawant, "Fungal Citric Acid Production Using Waste Materials: a Mini-Review," *J. Microbiol. Biotechnol. Food Sci.*, vol. 8, no. 2, pp. 821–828, 2018.
- [29] A. Amato and F. Beolchini, "End of life liquid crystal displays recycling: A patent review," *J. Environ. Manage.*, vol. 225, no. July, pp. 1–9, Nov. 2018.
- [30] R. Garcia and R. Calantone, "A critical look at technological innovation typology and innovativeness: a literature review," *J. Prod. Innov. Manage.*, vol. 19, no. 2, pp. 110–132, 2002.
- [31] L. Rocchetti, A. Amato, and F. Beolchini, "Printed circuit board recycling: A patent review," *J. Clean. Prod.*, vol. 178, pp. 814–832, 2018.
- [32] Q. W. Zhang, Z. D. Ye, and L. Shi, "e-Met kinase inhibitors: an update patent review (2014–2017)," *Expert Opin. Ther. Pat.*, vol. 29, no. 1, pp. 25–41, Jan. 2019.
- [33] X. Li, C. Wang, H. Jiang, and C. Luo, "A patent review of arginine methyltransferase inhibitors (2010–2018)," *Expert Opin. Ther. Pat.*, vol. 29, no. 2, pp. 97–114, Feb. 2019.
- [34] E. P. Office, "Espacenet."
- [35] A. E. H. Murad and S. A. D. Khalaf, "Citric acid production from whey with sugars and additives by *Aspergillus niger*," *African J. Biotechnol.*, vol. 2, no. 10, pp. 356–359, Oct. 2003.
- [36] D. Xu, C. Madrid, M. Rohr, and C. Kubicek, "The influence of type and concentration of the carbon source on production of citric acid by *Aspergillus niger* Ding-Bang," *Appl. Microbiol. Biotechnol.*, vol. 30, no. 6, pp. 553–558, Jun. 1989.
- [37] Y. D. Hang and E. E. Woodams, "Solid-state fermentation of apple pomace for citric acid production," *MIRCEN J. Appl. Microbiol. Biotechnol.*, vol. 2, no. 2, pp. 283–287, 1986.
- [38] L. P. S. Vandenberghe, C. R. Soccol, F. C. Prado, and A. Pandey, "Comparison of Citric Acid Production by Solid-State Fermentation in Flask, Column, Tray, and Drum Bioreactors," *Appl. Biochem. Biotechnol.*, vol. 118, no. 1–3, pp. 293–304, 2004.
- [39] L. P. S. Vandenberghe, C. R. Soccol, A. Pandey, and J. M. Lebeault, "Microbial Production of Citric Acid," *Brazilian Arch. Biol. Technol.*, vol. 42, no. 3, pp. 263–276, 1999.
- [40] E. Arts, C. P. Kubicek, and M. Rohr, "Regulation of Phosphofructokinase from *Aspergillus Niger*: Effect of Fructose 2,6-Bisphosphate on the Action of Citrate, Ammonium Ions and AMP," *Microbiology*, vol. 133, no. 5, pp. 1195–1199, May 1987.
- [41] S. Ates, N. Dingil, E. Bayraktar, and U. Mehmetoglu, "Enhancement of citric acid production by immobilized and freely suspended *Aspergillus niger* using silicone oil," *Process Biochem.*, vol. 38, no. 3, pp. 433–436, 2002.
- [42] Y. Jin, "Method for producing citric acid by utilizing immobilized *aspergillus niger*," CN102864184, 2013.
- [43] H. Ying, B. Yu, Y. Chen, X. Zhang, W. Sun, and N. Zhao, "Immobilization method of *Aspergillus niger*," CN107022541, 2017.
- [44] G. Shi *et al.*, "Method for seed culture of *Aspergillus niger* and preparing citric acid," CN107815421, 2018.
- [45] B. Chen, "Preparation method of citric acid," CN106868061, 2017.
- [46] Z. Yong, L. Zongmei, Z. Jixue, and F. Zhifei, "Method for producing citric acid," CN102181490, 2011.
- [47] B. Li, Y. Zhou, Z. Lu, and R. Yang, "Aspergillus niger seed liquid preparation method and citric acid fermentation preparation method," CN104277978, 2015.
- [48] L. Xia, Z. Lu, Y. Shao, R. Yang, and H. Zhong, "Method for preparing citric acid through fermentation," CN102851330, 2013.

- [49] Z. Yongsheng, Z. Siliang, Z. Yong, F. Zhifei, and Y. Ruwen, "Citric acid preparation method," CN102373242, 2012.
- [50] G. Shi *et al.*, "Citric acid aspergillus niger seed continuous culture method based on mycelium pellet dispersion technology," CN104099253, 2014.
- [51] G. Shi *et al.*, "Method for producing citric acid by continuous fermentation of *Aspergillus niger*," CN104087624, 2014.
- [52] Z. Yongsheng, Z. Yong, L. Zongmei, Z. Jixue, and Y. Zhichen, "Production method of citric acid," CN102443611, 2012.
- [53] P. Christian and M. A. X. Rohr, "Accumulation by," vol. 165, no. 3, pp. 1019–1022, 1986.
- [54] R. Datta and E. P. Bergemann, "Process for producing of citric acid and monovalent citrate salts," US5532148, 1996.
- [55] A. Kirkovits and H. Edlauer, "PROCESS FOR THE FERMENTATIVE PRODUCTION OF CITRIC ACID FROM CARBOHYDRATES," US5081025, 1992.
- [56] A. J. Kabil, "Process for the production of citric acid," US3936352, 1976.
- [57] J. S. U. CH, "PROCESS FOR THE PRODUCTION OF CITRIC ACID BY SUBMERGED FERMENTATION," GB1392942, 1975.
- [58] J. SPIRITUS, "PROCESS FOR THE PRODUCTION OF CITRIC ACID BY SUBMERGED FERMENTATION," GB1342311, 1974.
- [59] G. Shi *et al.*, "Method for improving fermentation performance of citric acid on basis of mycelium structure control," CN105586366, 2016.
- [60] H. Luo, J. Xiong, Z. Lu, M. Wang, and X. Wu, "Method for preparing citric acid by fermentation," CN103497977, 2014.
- [61] R. Li *et al.*, "Bacterial strain for producing citric acid and method for preparing citric acid by fermenting same," CN103695319, 2014.
- [62] S. Anastassiadis, I. Morgunov, S. Kamzolova, and T. Finogenova, "Citric Acid Production Patent Review," *Recent Pat. Biotechnol.*, vol. 2, no. 2, pp. 107–123, 2008.
- [63] M. ZHANG, A., ROEHR\*, "Citric Acid Fermentation and Heavy Metal Ions," *Acta Biotechnol.*, vol. 22, no. 3–4, pp. 363–373, 2002.
- [64] L. Yin, "Method for preparing citric acid from corneobs," CN106119306, 2016.
- [65] X. Shen and D. Shen, "Method for preparing fermentation liquor of citric acid by utilizing extrudate obtained after low temperature extrusion and enzymolysis," CN105524951, 2016.
- [66] J. Li, S. Li, Q. Ma, M. Yu, and J. Chen, "Technique for preparing citric acid by fermenting beta cyclodextrin mother solution," CN103710397, 2014.
- [67] G. Shi *et al.*, "Method for producing citric acid fermentation liquor by two-stage fermentation," CN107815475, 2018.
- [68] Y. Jin, "Method for preparing citric acid by fermenting corn starch by using aspergillus niger," CN102864182, 2013.
- [69] N. Y. Sharova, A. A. Vybornova, Tatyana Vladimirovna Printseva, and A. R. Yushkauskajte, "Method for producing invertase and citric acid," RU2676144, 2018.
- [70] M. LAB, "Improvements in or relating to the production of citric acid," GB738940, 1955.
- [71] "Improvements in or relating to the production of citric acid," GB742972, 1956.
- [72] Y. Jin, "Method for preparing citric acid through fermenting corn sugar solution by immobilized *Aspergillus niger*," CN102851328, 2013.
- [73] Z. Yongsheng, Z. Huiping, L. Sixiang, and G. Cui, "Preparation method of citric acid fermentation solution," CN101555497, 2009.
- [74] W. Sun, H. Han, N. Wei, Q. Zhao, and Y. Zhang, "Method for preparing bran starter for citric acid by liquid inoculation," CN103667372, 2014.
- [75] L. Hu, M. Yun, S. Yingang, Z. Hua, and M. Jing, "Method for cultivating aspergillus niger mouldy bran and method for preparing citric acid through fermentation,"

- CN102649971, 2012.
- [76] Q. Cheng, L. Meng, Y. Yi, T. Sun, and J. Huang, "Method for producing citric acid by fermenting soybean molasses," CN103614421, 2014.
  - [77] R. Li, "Method for producing citric acid by fermenting sorghum powder," CN102864185, 2013.
  - [78] H. Z. Chen, "Method for fermenting and producing citric acid using steam explosion straw as raw material," CN1884563, 2006.
  - [79] B. Z. Guo, "New method of producing citric acid using paddy fermentation," CN1693470, 2005.
  - [80] R. Tan and J. Shen, "Method for producing citric acid from kiwi fruit offcut," CN105671093, 2016.
  - [81] Y. D. Hang, "Grape pomace as substrate for microbial production of citric acid," US4791058, 1988.
  - [82] Y. D. Hang, "Apple pomace as substrate for microbial production of citric acid," US4767705, 1988.
  - [83] M. U. INDUSTRIALWERKE, "A process for the production of citric acid by fermentation," GB302338, 1929.
  - [84] S. K. Kang, J. Y. Choe, and I. B. Kwon, "PROCESS FOR PREPARING CITRIC ACID BY TANGERIN PEELINGS," KR930001261, 1993.
  - [85] J. Bao, J. Meng, and J. Zhang, "Method for producing citric acid by using lignocellulose raw material," CN104805136, 2015.
  - [86] H. Tang, W. Yang, Q. Qin, J. Zhang, B. Xu, and X. Zhang, "Method for producing citric acid by fermenting konjak powder residues," CN105506004, 2016.
  - [87] Z. Gao, G. Zhou, J. Xiong, and X. Lu, "Method for fermentation production of citric acid," CN104232699, 2014.
  - [88] C. Liu and F. Liu, "Method for producing citric acid through continuous batch feeding fermentation," CN103290070, 2013.
  - [89] H. Luo, Z. Lu, R. Yang, and X. Lu, "Method for preparing citric acid by fermentation," CN103146769, 2013.
  - [90] Y. Zhou, Y. Man, X. Zhang, H. Zhang, and J. Zhu, "Enzymolysis method of starch raw material and method for preparing citric acid by fermenting," CN102839203, 2012.
  - [91] S. Guiyang *et al.*, "Method for fermenting and preparing citric acid by adding saccharifying enzyme," CN101942487, 2011.
  - [92] Y. Jin, "Method for preparing citric acid through fermenting cheap dried sweet potatoes by utilizing aspergillus niger," CN102952830, 2013.
  - [93] Y. Jin, "Method for preparing citric acid through fermenting puffed dried sweet potato raw material by Aspergillus niger," CN102851329, 2013.
  - [94] B. Guo and J. Zhu, "Method for producing citric acid by fermenting paddy rice," CN1415755, 2003.
  - [95] H. IO. Hapona *et al.*, "METHOD FOR PREPARATION OF CITRIC ACID,  $\alpha$ -AMYLASE AND GLUCOAMYLASE," RU2007125728, 2009.
  - [96] N. J. Sharova, L. N. Mushnikova, T. A. Pozdnjakova, and T. A. Nikiforova, "METHOD OF CITRIC ACID PRODUCING," RU2186850, 2002.
  - [97] R. Li, H. Shang, W. Yang, Y. Deng, and H. Tang, "Method for detoxification in production of citric acid by utilizing cassava fermentation," CN103045659, 2013.
  - [98] Y. Jin, "Method for preparing citric acid by fermenting cassava raw material residue-removed clear solution with aspergillus niger," CN102864183, 2013.
  - [99] S. S. AB, "A method of producing citric acid," GB951629, 1964.
  - [100] M. LAB, "Citric acid production," GB799752, 1958.
  - [101] S. Gupta and C. B. Sharma, "Citric acid fermentation by the mutant strain of the Aspergillus niger resistant to manganese ions inhibition," *Biotechnol. Lett.*, vol. 17, no.



- 3, pp. 269–274, Mar. 1995.
- [102] Ikram-ul-Haq, Khurshid, Ali, Ashraf, Qadeer, and Rajoka, “Mutation of *Aspergillus niger* for hyperproduction of citric acid from black strap molasses,” *World J. Microbiol. Biotechnol.*, vol. 17, no. 1, pp. 35–37, 2001.
- [103] S. Kumar and N. S. Punekar, “The metabolism of 4 aminobutyrate (GABA) in fungi,” *Mycol. Res.*, vol. 101, no. 4, pp. 403–409, Apr. 1997.
- [104] S. Jin *et al.*, “Method of increasing yield of citric acid produced by *aspergillus niger* fermentation,” CN106755138, 2017.
- [105] L. Liu *et al.*, “Method for increasing citric acid production by *Aspergillus niger* fermentation,” US2018195052, 2018.
- [106] F. Sun *et al.*, “Recombinant *aspergillus niger* capable of improving yield of citric acid and preparation method of recombinant *aspergillus niger*,” CN106635847, 2017.
- [107] M. Li, F. Lu, and R. Wei, “Method for increasing sugar utilization rate and yield of citric acid in citric acid fermentation and application,” CN108018216, 2018.
- [108] T. B. Buttophona and H. IO. Ilapona, “STRAIN *Aspergillus niger* PRODUCER OF CITRIC ACID,” RU2013151521, 2015.
- [109] R. Li, H. Shang, and B. Xu, “Citric acid high-yielding *Aspergillus niger* FY2013 and application thereof,” CN103952318, 2014.
- [110] R. Li, X. Mu, W. Li, and Z. Chang, “Bacterial strain for producing citric acid and method for fermenting and producing citric acid through fermentation of bacterial strain,” CN103045487, 2013.
- [111] Z. Dai and S. E. Baker, “ENHANCED CITRIC ACID PRODUCTION IN *ASPERGILLUS*,” WO2013082459, 2013.
- [112] C. Xiu, L. Zongmei, Z. Hua, W. Xiaoyan, and X. Lihong, “*Aspergillus niger*, application of *Aspergillus niger* and method for preparing citric acid by fermentation,” CN102533570, 2012.
- [113] Z. Junhua, Z. Yong, Z. Hua, L. Zongmei, and F. Zhifei, “*Aspergillus niger* and application thereof as well as citric acid preparation method through fermentation,” CN102399702, 2012.
- [114] H. He, L. Shuang, Z. Jing, X. Qing, and M. Yun, “High-temperature-resistant citric acid producing strain,” CN102352322, 2012.
- [115] T. J. A. Nikiforova, V. P. Komov, T. J. V. Vybornova, L. B. Piotrovskij, M. A. E. Dumpis, and E. V. Litasova, “CITRIC ACID PRODUCTION METHOD,” RU2428481, 2011.
- [116] M. Minarik, D. Skvarenina, P. Michalik, V. Sitkey, and V. Visacky, “PROCESS, APPARATUS AND MICROORGANISM STRAIN FOR THE MANUFACTURE OF CITRIC ACID,” WO9710350, 1997.
- [117] R. Y. Karklin, A. A. Rumba, and V. K. Azanda, “Method of preparing seeding material for production of citric acid,” US4380583, 1983.
- [118] P. B. Avchieva and J. D. Berezhnoj, “CONTINUOUS METHOD OF PRODUCING CITRIC ACID AND ITS SALTS,” RU2125607, 1999.
- [119] A. A. Rumba, R. Y. Karklin, V. K. Azanda, V. Y. Berzinsh, A. A. Lurin, and V. F. Fedoseev, “Citric acid prepn. by fermentation—using *Aspergillus niger* R-1 strain on nutrient medium contg. molasses,” FR2361330, 1978.
- [120] A. A. Rumba, R. Y. Karklinsh, V. K. Azanda, V. Y. Berzinya, A. A. Lurinsh, and V. F. Fedoseev, “*ASPERGILLUS NIGER* P-1 STRAIN AS PRODUCER OF CITRIC ACID,” SU568677, 1977.
- [121] E. Z. B. P. I. MIKR, “METHOD OF PREPARING CITRIC ACID,” GB1499093, 1978.
- [122] E. J. Shecherbakova, T. A. Nikiforova, and E. B. L. Vova, “DIPLOID STRAIN OF *ASPERGILLUS NIGER* AS PRODUCER OF CITRIC ACID,” RU2203322, 2003.
- [123] N. V. Krasikova, Nikiforova T. A., and K. V. M. Fin, “STRAIN OF FUNGUS



- ASPERGILLUS NIGER AS PRODUCER OF CITRIC ACID," RU2192460, 2002.
- [124] V. P. Ermakova, V. M. Golubtsova, E. E. Sakson, V. V. Ajukov, L. A. Sergeeva, and A. A. Veselova, "STRAIN OF FUNGUS ASPERGILLUS NIGER F-713—A PRODUCER OF CITRIC ACID," RU95113067, 1997.
- [125] F. ABBAS, "Citric acid prodn. by fermentation of sugar beet or date molasses—using *Aspergillus niger* of *Mucor piriformis* sp. organism," FR2336477, 1977.
- [126] R. Karklins and I. Skrastina, "A METHOD FOR PREPARING OF CITRIC ACID PRODUCENT ASPERGILLUS NIGERATION OF COMBUSTIBLE MATERIALS ESPECIALLY OF INDUSTRIAL AND HOUSEHOLD WASTE," LV11342, 1996.
- [127] R. Karklins and I. Skrastina, "MICROSCOPICAL FUNGUS ASPERGILLUS NIGER R-5 AS CITRIC ACID PRODUCENT," LV11340, 1996.
- [128] N. Zhang, J. Jiang, M. Wei, J. Yang, and J. Zhao, "Aspergillus niger and application thereof to preparing citric acid from fermented acorns," CN106367359, 2017.
- [129] B. Zhang, L. Zhu, L. Li, and Y. Pei, "Method for preparing citric acid through liquid-state fermentation of acorns," CN108588133, 2018.
- [130] A. J. Vinarov, T. E. Sidorenko, S. E. Smetanina, and V. N. Jashina, "FUNGUS STRAIN ASPERGILLUS NIGER—A PRODUCER OF CITRIC ACID," RU2103346, 1998.
- [131] S. Anastassiadis, A. Aivasidis, and C. Wandrey, "Fermentation process for the continuous production of citric acid," DE4407441, 1994.
- [132] R. I. Leavitt, "Process for producing citric acid," US4178211, 1979.
- [133] R. E. D. PETROLIERES, "PROCESS FOR THE PRODUCTION OF CITRIC AND ISOCITRIC ACIDS BY A CULTURE OF A DIPLOID CANDIDA LIPOLYTICA YEAST," GB1464334, 1977.
- [134] P. LTD, "CITRIC ACID PRODUCTION," GB1418561, 1975.
- [135] T. Matsumoto, Y. Ichikawa, and T. Nagata, "Process for producing citric acid from hydrocarbons by fermentation," US4424274, 1984.
- [136] A. Fujimaki, T. Osada, and T. Matsumoto, "METHOD FOR SUPPRESSING PRODUCTION OF ISOCITRIC ACID IN CITRIC ACID FERMENTATION OF PETROLEUM HYDROCARBON," JPS5779890, 1982.
- [137] I. F. DU PETROLE and R. E. D. PETROLIERE, "FERMENTATIVE PRODUCTION OF CITRIC ACID," GB1428440, 1976.
- [138] B. G. J. A., "PROCESS FOR THE PRODUCTION OF CITRIC ACID AND SALTS THEREOF," GB1418511, 1975.
- [139] H. C. C. LTD, "PROCESS FOR PRODUCING CITRIC ACID BY FERMENTATION," GB1380938, 1975.
- [140] "WERKWIJZE VOOR HET LANGS MICROBIOLOGISCHE WEG BEREIDEN VAN CITROENZUUR,," GB1297243, 1972.
- [141] S. Okumura, R. Tsugawa, H. Kamijo, A. Kamimura, and K. Komagata, "A FERMENTATION PROCESS FOR THE PRODUCTION OF CITRIC ACID," GB1204635, 1970.
- [142] K. Tanaka and K. Kimura, "PROCESS FOR PREPARING CITRIC ACID BY FERMENTATION," US3652396, 1972.
- [143] S. Kinoshita, K. Tanaka, and S. Akita, "Process for the production of citric acid by fermentation," GB878151, 1961.
- [144] MERCK & CO INC, "Improvements in or relating to the production of citric acid by fermentation," GB581389, 1946.
- [145] USINES DE MELIE, "Improvements in or relating to a process for producing citric acid by aerobic fermentation of solutions containing molasses," GB797390, 1958.
- [146] "Jiangsu Guoxin Union Energy CO LTD," 2016.
- [147] D. of E. and S. A. D. United Nations International Trade Statistics Database, "ALL COMMODITIES | Imports and Exports | 2017." [Online]. Available:

[https://trendeconomy.com/data/commodity\\_h2?commodity=291814&trade\\_flow=Export,Import&time\\_period=2017](https://trendeconomy.com/data/commodity_h2?commodity=291814&trade_flow=Export,Import&time_period=2017). [Accessed: 18 Mar 2019].

- [148] G. Singh, S. Kaur, and M. Verma, "Utilization of different agro-industrial wastes for sustainable bioproduction of citric acid by *Aspergillus niger*," *Biochem. Eng. J.*, vol. 54, no. 2, pp. 83–92, 2011.
- [149] A. Karthikeyan and N. Sivakumar, "Bioresource Technology Citric acid production by Koji fermentation using banana peel as a novel substrate," *Bioresour. Technol.*, vol. 101, no. 14, pp. 5552–5556, 2010.
- [150] Z. Alam, N. Bari, S. A. Muyibi, P. Jamal, and Abdullah Al Mamun, "Solid State Bioconversion of Oil Palm Empty Fruit Bunches for Production of Citric Acid by Wild Strains of *Aspergillus Niger*," *Food Biotechnol.*, vol. 24, no. 1, pp. 19–36, Mar. 2010.
- [151] D. Kumar, V. K. Jain, G. Shanker, and A. Srivastava, "Citric acid production by solid state fermentation using sugarcane bagasse," *Process Biochem.*, vol. 38, no. 12, pp. 1731–1738, Jul. 2003.