



UNIVERSITÀ POLITECNICA DELLE MARCHE
Repository ISTITUZIONALE

Towards a marine strategy for the deep Mediterranean Sea: Analysis of current ecological status

This is the peer reviewed version of the following article:

Original

Towards a marine strategy for the deep Mediterranean Sea: Analysis of current ecological status / Danovaro, R.; Fanelli, E.; Canals, M.; Ciuffardi, T.; Fabri, M. -C.; Taviani, M.; Argyrou, M.; Azzurro, E.; Bianchelli, S.; Cantafaro, A.; Carugati, L.; Corinaldesi, C.; de Haan, W. P.; Dell'Anno, A.; Evans, J.; Foglini, F.; Galil, B.; Gianni, M.; Goren, M.; Greco, S.; Grimalt, J.; Guell-Bujons, Q.; Jadaud, A.; Knittweis, L.; Lopez, J. L.; Sanchez-Vidal, A.; Schembri, P. J.; Snelgrove, P.; Vaz, S.; Angeletti, L.; Barsanti, M.; Borg, J. A.; Bosso, M.; Brind'Amour, A.; Castellán, G.; Conte, F.; Delbono, I.; Galgani, F.; Morgana, G.; Prato, S.; Schirone, A.; Soldevila, E. - In: MARINE POLICY, ISSN 0308-597X, 112 (2020). [10.1016/j.marpol.2019.103781]
This version is available at: 11566/275931 since: 2024-03-25 10:20:19Z

Publisher:

Published

DOI:10.1016/j.marpol.2019.103781

Terms of use:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. The use of copyrighted works requires the consent of the rights' holder (author or publisher). Works made available under a Creative Commons license or a Publisher's custom-made license can be used according to the terms and conditions contained therein. See editor's website for further information and terms and conditions.

This item was downloaded from IRIS Università Politecnica delle Marche (<https://iris.univpm.it>). When citing, please refer to the published version.

(Article begins on next page)

Marine Policy

February 2020, Volume 112 Pages 103781 (18p.)

<https://doi.org/10.1016/j.marpol.2019.103781>

<https://archimer.ifremer.fr/doc/00609/72110/>

Towards a marine strategy for the deep Mediterranean Sea: Analysis of current ecological status

Danovaro R. ^{1,2,*}, Fanelli E. ¹, Canals M. ³, Ciuffardi T. ⁴, Fabri Marie-Claire ⁵, Taviani M. ^{2,6,7}, Argyrou M. ⁸, Azzurro E. ^{2,9}, Bianchelli S. ¹, Cantafaro A. ¹⁰, Carugati L. ¹, Corinaldesi C. ¹¹, De Haan W.P. ³, Dell'anno A. ¹, Evans J. ¹⁰, Foglini F. ⁶, Galil B. ¹², Gianni M. ¹³, Goren M. ¹², Greco S. ², Grimalt J. ¹⁴, Güell-Bujons Q. ³, Jadaud Angelique ¹⁵, Knittweis L. ¹⁰, Lopez J.L. ¹⁴, Sanchez-Vidal A. ³, Schembri P.J. ¹⁰, Snelgrove P. ¹⁶, Vaz Sandrine ¹⁵, Angeletti L. ¹⁷, Barsanti M. ¹⁸, Borg J.A. ¹⁹, Bosso M. ¹⁸, Brind'Amour Anik ²⁰, Castellan G. ¹⁷, Conte F. ¹⁸, Delbono I. ¹⁸, Galgani Francois ²⁰, Morgana G. ¹⁸, Prato S. ¹⁸, Schirone A. ¹⁸, Soldevila E. ²¹

¹ Department of Life and Environmental Sciences, Polytechnic University of Marche, 60131, Ancona, Italy

² Stazione Zoologica Anton Dohrn Naples, 80122, Naples, Italy

³ CRG Marine Geosciences, Department of Earth and Ocean Dynamics, Faculty of Earth Sciences, University of Barcelona, 08028, Barcelona, Spain

⁴ Italian National Agency for New Technologies, Energy and Sustainable Economic Development (ENEA), Department for Sustainability, S. Teresa Marine Environment Research Centre, 19100, La Spezia, Italy

⁵ Institut Français de Recherche pour l'exploitation de la Mer (Ifremer), Département Océanographie et Dynamique des Ecosystèmes, 83500, La Seyne sur Mer, France

⁶ Institute of Marine Sciences (ISMAR), CNR, 40129, Bologna, Italy

⁷ Biology Department, Woods Hole Oceanographic Institution, MA, 02543, USA

⁸ Department of Fisheries and Marine Research (DFMR), 1416, Nicosia, Cyprus

⁹ Institute for Environmental Protection and Research (ISPRA) STS Livorno, 57122, Italy

¹⁰ Department of Biology, University of Malta, Msida, MSD2080, Malta

¹¹ Department of Sciences and Engineering of the Materials, Environment and Urban Planning, Polytechnic University of Marche, Italy

¹² The Steinhardt Museum of Natural History, Israel National Center for Biodiversity Studies, Tel Aviv University, Tel Aviv, 69978, Israel

¹³ Deep Sea Conservation Coalition, Postbus, 59681, Amsterdam, Netherlands

¹⁴ Department of Environmental Chemistry, Institute of Environmental Assessment and Water Research (CSIC), 08034, Barcelona, Spain

¹⁵ UMR Marbec, Ifremer, IRD, Université de Montpellier, CNRS, 34203, Sète Cedex, France

¹⁶ Departments of Ocean Sciences and Biology, Memorial University of Newfoundland, St. John's, NL A1C 5S7, Canada

¹⁷ Institute of Marine Sciences (ISMAR), CNR, 40129, Bologna, Italy

¹⁸ Italian National Agency for New Technologies, Energy and Sustainable Economic Development (ENEA), Department for Sustainability, S. Teresa Marine Environment Research Centre, 19100, La Spezia, Italy

¹⁹ Department of Biology, University of Malta, Msida, MSD2080, Malta

²⁰ Institut Français de Recherche pour l'exploitation de la Mer (Ifremer), Département Océanographie et Dynamique des Ecosystèmes, 83500, La Seyne sur Mer, France

²¹ CRG Marine Geosciences, Department of Earth and Ocean Dynamics, Faculty of Earth Sciences, University of Barcelona, 08028, Barcelona, Spain

* Corresponding author : R. Danovaro, email address : r.danovaro@univpm.it

Abstract :

The Marine Strategy Framework Directive (MSFD), introduced in June 2008, was adopted to achieve a Good Environmental Status (GES) in the EU's marine waters and to protect resources of socio-economic interest. The MSFD extends to the marine area over which a Member State exercises jurisdictional rights in accordance with the United Nations Convention on the Law of the Sea (UNCLOS), including the deep-sea waters, seafloor and sub-seafloor of the Exclusive Economic Zones (EEZ). However, currently the MSFD focuses on coastal habitats and the shallow-water seafloor to the detriment of the deeper habitats. Despite the huge dimension of the deep sea (below 200 m of depth) covering more than 65% of the Earth's surface and including >95% of the global biosphere, the relevance of the dark portion of the seas and oceans is still almost completely neglected. Given the important bi-directional links between shallow and deep ecosystems, there is a clear need for extending the implementation of the MSFD into the deep sea, to define a sound ecosystem-based approach for the management and protection of deep-sea ecosystems and attain GES. We assembled data on drivers, anthropogenic pressures and impacts concerning the MSFD descriptors pertaining to the Mediterranean deep sea. We list deep-sea monitoring activities and the main sources providing benchmark conditions, and discuss knowledge and geographic coverage gaps. MSFD descriptors apply to the deep sea as to coastal waters, and ought to be monitored contemporaneously. We provide recommendations for guidelines for future deep-sea monitoring in the Mediterranean Sea.

Highlights

► MSFD fails to cover the huge dimension of deep-sea environments and important bi-directional link with shallow ones. ► Extending MSFD to the deep sea and defining an ecosystem-based approach for its management and protection is urgently needed. ► Data on drivers, anthropogenic pressures and impacts regarding the MSFD descriptors for deep-sea Mediterranean were reviewed. ► Deep-sea monitoring activities were discussed and knowledge and geographic coverage gaps evidenced. ► Recommendations for guidelines for future deep-sea monitoring were provided.

Keywords : Marine strategy framework directive, Deep-sea ecosystems, Mediterranean basin

60
61
62 **6 List of acronyms and abbreviations**

- 63 7 ABNJ: Areas Beyond National Jurisdiction
64 8 ACCOBAMS: Agreement on the Conservation of Cetaceans in the Black Sea Mediterranean Sea and
65 9 Contiguous Atlantic Area
66 10 AUV: Autonomous Underwater Vehicle
67 11 CBD: Convention on Biological Diversity
68 12 CFP: Common Fisheries Policy
69 13 CWC: Cold-Water Corals
70 14 DCF: Data Collection Framework
71 15 DSF: Deep-Sea Fisheries
72 16 DSWC: Dense Shelf Water Cascading
73 17 EEZ: Exclusive Economic Zone
74 18 EFH: Essential Fish Habitats
75 19 EIA: Environmental Impact Assessment
76 20 EMT: Eastern Mediterranean Transient
77 21 EwE: Ecopath with Ecosim
78 22 FAO: Food and Agricultural Organization
79 23 FRA: Fishery Restricted Areas
80 24 GFCM: General Fisheries Commission for the Mediterranean
81 25 GSA: Geographical Sub Area
82 26 ICCAT: International Commission for the Conservation of Atlantic Tunas
83 27 LIW: Levantine Intermediate Water
84 28 MBES: Multibeam Echosounder
85 29 MEDIAS: Mediterranean International Acoustic Survey
86 30 MEDITS: Mediterranean International Trawl Survey
87 31 MS: Member States
88 32 MSFD: Marine Strategy Framework Directive
89 33 MSY: Maximum Sustainable Yield
90 34 NIS: Non-Indigenous Species
91 35 OMZ: Oxygen Minimum Zones
92 36 PAH: Polycyclic Aromatic Hydrocarbons
93 37 POM: Particulate Organic Matter
94 38 POP: Persistent Organic Pollutants
95 39 ROV: Remotely Operated Vehicle
96 40 SAC: Scientific Advisory Committee
97 41 SCA: Stomach Content Analysis
98 42 SSB: Spawning Stock Biomass
99 43 SIA: Stable Isotope Analysis
100 44 SSS: Side-Scan Sonar
101 45 STECF: Scientific, Technical and Economic Committee for Fisheries
102 46 VME: Vulnerable Marine Ecosystem
103 47 UNCLOS: United Nations Convention on the Law of the Sea
104 48 UNGA: United Nations General Assembly
105 49

71 1. Introduction

72 1.1 The Mediterranean Sea

73 The Mediterranean is a semi-enclosed basin between the European and African coasts with
74 the narrow and shallow Strait of Gibraltar connecting its waters and life to the Atlantic Ocean. The
75 Suez Canal creates a man-made connection to the Red Sea, recently doubled, which allows the
76 penetration of tropical Indo-pacific species (Longhurst, 2017). Concomitantly, the Mediterranean
77 Basin is experiencing major climatic-related changes that are strongly influencing its oceanography
78 in terms of water mass characteristics (e.g., temperature, salinity, dissolved oxygen), currents,
79 nutrients and relative sea levels. The interplay of these factors has, since historical times, strongly
80 influenced the diversity and colonization of the Mediterranean Sea (Taviani, 2002; Danovaro et al.,
81 2010).

82 Unique hydrology characterizes the present-day Mediterranean, including microtidal regime,
83 oligotrophy, high salinity (37.5-39.5 psu), homoeothermic temperatures from 300–500 m to the
84 bottom, with values at the seafloor ranging from ca 13-13.5 °C in the western basin to 13.5-15.5 °C
85 in the eastern basin, and the almost complete lack of thermal boundaries (Emig and Geistdoerfer,
86 2005). These features uniquely make the Mediterranean one of the “warmest” deep-sea basins of
87 the world. Strong energy gradients also characterize the Mediterranean, with primary production
88 and food supply to the deep decreasing from the western to the eastern region of the basin and
89 from shallower to deeper waters (Danovaro et al. 1999).

90 These historical, topographic and environmental characteristics complicate deep-sea
91 biodiversity patterns of the Mediterranean Sea but raise intriguing questions. Numerous studies
92 document that the Mediterranean Sea, although modest in size (0.82% and 0.32% of the global
93 ocean surface and volume, respectively; Bianchi and Morri, 2000), is a biodiversity hot spot with
94 overall ca 17000 species, which represent 7.5% of the species richness of the oceans (Danovaro
95 and Pusceddu, 2007; Coll et al., 2010; Lejeusne et al., 2010). However, although data on the
96 species richness of its deeper habitats are incomplete (two thirds of the deep species – excluding
97 prokaryotes – have not been censused yet; Ramirez-Llodra et al., 2009; Coll et al., 2010, Danovaro
98 et al., 2010), it appears that the biodiversity of the deep Mediterranean basin is lower than that of
99 other oceans (Danovaro et al. 2010).

100 The biodiversity of the deep Mediterranean Sea depends largely from the heterogeneity of
101 habitats, which include submarine canyons and seamounts, continental rise deposits, mud
102 volcanoes and extreme environments such as hydrothermal vents, cold seeps and deep-

hypersaline anoxic basins (Olu-Le Roy et al., 2004; Danovaro et al., 2010; Taviani, 2011, 2014; Fernandez-Arcaya et al., 2017). Even seemingly “featureless” soft bottom habitats host unique and vulnerable species and habitats (e.g. sponge fields, gorgonian and pennatulacean meadows) (Danovaro et al., 2010).

The Mediterranean basin is threatened by multiple stressors associated with the rapid expansion of coastal populations, urbanization, changes in agricultural, industrial and shipping patterns, overfishing and exploration and extraction of offshore minerals and hydrocarbons, which exert increasing pressures through habitat destruction, chemical pollution, and dumping of waste and litter (EEA, 1999; Danovaro et al., 1993). In concert with climate change, these stressors may act synergistically to affect the dynamics, and potentially the resilience, of fragile deep ecosystems (WWF/IUCN, 2004; UNEP/MAP, 2012). Direct stressors and processes also occur on the adjacent shelf and in the epi-mesopelagic zones, including Dense Shelf Water Cascading (DSWC) events down the continental slope, open-sea convection and severe coastal storms, which may transport sediments and organic matter to the continental slope and beyond, influencing deep-sea biodiversity and ecosystem functioning (Canals et al., 2006; Ulses et al., 2008; Sanchez-Vidal et al., 2012; Durrieu de Madron et al., 2013; Taviani et al., 2016). In particular, bottom-contacting fisheries, specifically bottom-trawling and longlines, represent the most significant anthropogenic threats to deep-water biota, severely impacting sensitive habitats and species such as cold-water corals (CWCs) and/or sponge gardens (Rogers, 1999; Koslow et al., 2000). Additional evidence attributes a significant proportion of deep-sea litter to the fishing industry (Bo et al., 2014; Tubau et al., 2015; D'Onghia et al., 2017; Mecho et al. 2017), along with land- and ship-based sources (Ryan et al. 2009).

Given the increasing pressures on deep-sea habitats, scientists and managers are becoming conscious of the need to develop standardised tools and harmonized observation systems for long-term biological monitoring, in order to enable the collection of scientifically-validated data and a better understanding of the consequences of the present and future anthropogenic impacts (Danovaro et al., 2017, Aguzzi et al., 2019; Danovaro et al., 2019).

1.2 Implementing the Marine Strategy Framework Directive in the deep Mediterranean Sea

The Marine Strategy Framework Directive (MSFD 2008/56/EC) represents the EU's Integrated Maritime Policy tool to achieve Good Environmental Status (GES) of marine waters, with an initial target for 2020. The MSFD applies to the area of marine waters over which a

135 Member State exercises jurisdictional rights in accordance with the UNCLOS (see Figure 1, for the
136 definition of territorial waters and EEZs in the Mediterranean). These include also deep-sea
137 waters, seabed and sub-seafloor. At present, MSFD implementation focuses mostly on coastal
138 habitats or those impacted by commercial fisheries (Raicevich et al., 2017). However, over long-
139 time scales, global nutrient and carbon cycles depend on a functioning deep sea (e.g. Snelgrove et
140 al., 2018). Moreover, the life-cycle stages of some coastal species use offshore environments, thus
141 achieving GES for marine ecosystems associated with continental shelves, must link to the
142 achievement of GES for deep Mediterranean environments and Areas Beyond National Jurisdiction
143 (ABNJ or “High Seas”). Otherwise, the MSFD will largely disregard the precautionary principle and
144 undermine an ecosystem-based approach to marine management.

145 An effective governance and management of the Mediterranean Sea requires consideration
146 of the complexity of these environmental issues, and meaningful international cooperation (de
147 Vivero and Rodriguez Mateos, 2015). Given the transboundary nature of most of the deep waters,
148 their inclusion in MSFD complicates the requirement for each Member State to apply the Directive
149 to areas within its national jurisdiction. This emphasizes the need for Member States (MS) to
150 cooperate in order to ensure coordinated and harmonized development of marine strategies at
151 the scale of region/sub-region in the Mediterranean Basin, where EU MS and developing countries
152 co-exist.

153 MSFD implementation currently suffers from a lack of standardized and consistent
154 methodology for deep waters. To address this gap, we identify approaches, variables, and
155 methodologies to enable MSFD implementation in the deep Mediterranean Sea. This synthesis
156 summarises available information on MSFD descriptors for the deep Mediterranean Sea, with
157 respect to the criteria listed in the European Commission Decision (COMM/DEC/2017/848), and
158 anthropogenic pressures, uses and human activities affecting the marine environment (Table 2 of
159 Annex III of COMM/DEC/2017/848).

161 2. State of the knowledge of MSFD descriptors in the deep Mediterranean

162 2.1 Descriptor 1: Biological diversity

163 Descriptor 1 (D1) states that *“The quality and occurrence of habitats and the distribution
164 and abundance of species are in line with prevailing physiographic, geographic and climatic
165 conditions”* (MSFD, 2008/56/EC, Annex I, summarised in Table 1). The species groups specified in
166 Part II of the Annex to COMM/DEC/2017/848, these include birds, mammals, reptiles, fish and

167 cephalopods, some of which are present, diverse and abundant in the deep sea, such as fishes and
168 cephalopods, in addition to deep diving and feeding cetaceans. Deep-sea organisms play an
169 important role in marine food webs, either as predators or as important prey of a large set of high
170 trophic level predators, including other fishes and cephalopods and marine mammals (Fanelli et
171 al., 2012, 2013; Quetglas et al., 2013).

172 Data on these components from the deep Mediterranean Sea are largely included in the
173 MEDITS database, which also represent the only extensive time series available for the deep
174 Mediterranean (Bertrand et al., 2002), [although with the limitation that it is a destructive sampling](#)
175 [method with discrete sampling time, which exclude the possibility to detect any displacements of](#)
176 [demersal species \(Aguzzi et al., 2009; 2013\)](#). MEDITS is funded as part of the EU Data Collection
177 multi-annual sampling program (DC-MAP), which limits the sampling frequency to yearly surveys
178 confined to the northern part of the Mediterranean Basin. MEDITS mostly targets demersal fish
179 (including deep-sea sharks), but includes also commercial invertebrates and other macro and
180 mega-invertebrates (as by-catch species). MEDITS provides detailed information on their
181 abundance and biomass, including the population structure of several species (including length
182 frequency distributions by sex and maturity stages for different target species), which allow us to
183 obtain information on the size spectra, maturity ogives, sex ratios and mortality rates. This
184 information contributes to both the census of shallow and deep marine biodiversity and stock
185 assessments carried out by the GFCM and the STECF of the European Commission (see
186 Vasilakopoulos et al., 2014; [Cardinale et al., 2017](#)) [see Descriptor 3 below]. In the case of meso-
187 and bathypelagic species (MEDIAS Handbook, 2015; i.e. Galil, 2004; Papiol et al., 2013; Fanelli et
188 al., 2013, 2015; Valls et al., 2014), species of non-commercial interest, hard bottom habitats
189 between 200 and 800 m depth, and in general all habitats below 800 m depth, only scattered
190 information without temporal datasets exist.

191 Another important gap concerns the smaller size biota, such as meiofauna, which are a key
192 component in the deep-sea ecosystems and are driven by water depth, regional setting and
193 geomorphological characteristics of the deep Mediterranean habitats (Bianchelli and Danovaro,
194 2019). Meiofauna are highly diversified (possibly hyper-diverse), and play a fundamental
195 ecological role in the biogeochemical cycles and in food webs and are sensitive to environmental
196 and anthropogenic pressures ([Pusceddu et al., 2014](#)). Since this component, which increases its
197 [ecological](#) relevance, in terms of abundance and functional role, with increasing water depths
198 (Danovaro et al., 2015), has been recently suggested for inclusion in the D1 for the

199 implementation of the MSFD (Semprucci et al., 2014; Bianchelli et al., 2016a; 2018). It is even
200 more evident that it should be taken into consideration in the implementation of the MSFD in the
201 deep sea.

202 The MSFD deep-sea habitats included in the COMM/DEC/2017/848 are: a) upper bathyal
203 rocks and biogenic reefs, b) upper bathyal sediments, c) lower bathyal rock and biogenic reef, d)
204 lower bathyal sediments, and e) abyssal seafloor. These include other benthic habitats such as:
205 canyons (which may include rocky and sedimentary substrates), rocky bottoms with coral banks
206 (including Cold-water corals) or large bivalves, different types of sedimentary bottoms in bathyal
207 or abyssal plains (muds, sands or coarser sediment), chemosynthetic ecosystems (hydrothermal
208 vents and mud volcanoes), and seamounts.

209 Previous studies on the deep Mediterranean Sea reported a west-east decreasing gradient
210 of food availability (Danovaro et al., 1999; Danovaro et al., 2008), which explains the presence of a
211 significant decreasing gradient in the abundance and biomass of most deep-sea benthic
212 components along that gradient, from meiofauna to megafauna (Sardá et al., 2004; Bianchelli and
213 Danovaro, 2019; Fanelli et al., 2018). The CWCs apparently follow the same gradient (Taviani et al.,
214 2017; Chimienti et al., 2019), and the presence of Levantine Intermediate Water (LIW) likely
215 strongly influences their distribution (Freiwald et al., 2009; Taviani et al., 2016, 2017).

216 Trawl surveys provide most of the available information on deep-sea habitats and their
217 characteristics (see Table 1, Annex III, MSFD), but only for soft bottom habitats including those
218 dominated by *Isidella elongata* and *Funiculina quadrangularis* (Lauria et al., 2017; Vasilis et al.,
219 2019). Oceanographic cruises using ROVs offer the possibility to conduct non-destructive image
220 and sample collections able to contribute significantly to the study of deep-sea habitats. Most ROV
221 surveys to date have focused on CWC habitats and coral gardens, and provide important
222 information on the composition, abundance, and biomass of the communities within these
223 habitats (Taviani et al., 2005, 2011, 2015, 2019; Schembri et al., 2007; Fabri et al., 2014, 2017; Bo
224 et al., 2015; Evans et al., 2016; Fanelli et al., 2017; Chimienti et al., 2019; Moccia et al., 2019).
225 Most available information focuses on deep-sea canyons (Migeon et al., 2012), seamounts (Wurtz
226 and Rovere, 2015) and mud volcanoes (Masclé et al., 2014), with major data gaps for deep-sea
227 pelagic habitats, notwithstanding there is an increasing information on deep-water zooplankton
228 and micronekton (Koppelman et al., 2009; Fanelli et al. 2011, 2014; Cartes et al., 2013; Denda
229 and Christiansen, 2014; Danovaro et al., 2017; Conese et al., 2019). Descriptor 1 is directly linked
230 to D2, D3, D4 and D6 (habitats), and monitoring efforts in the deep sea can therefore gather

contextual information on all these descriptors. The ecosystem criteria listed in COMM/DEC/2017/848, which link Descriptors 1 and 4, consider trophic guilds. These are highly relevant to the deep sea and can therefore be immediately described, as the already available data would provide the required background information.

2.2 Descriptor 2: Non-indigenous species introduced by human activities

The number of recorded [Non-Indigenous Species](#) (NIS) in the Mediterranean Sea greatly exceeds that in other European seas (Galil et al., 2014; Zenetos, 2019). Their establishment alters biotic assemblages and ecosystem functions (Galil, 2007; Katsanevakis et al., 2007; Fanelli et al., 2015; Galil et al., 2016, 2017; Goren et al., 2016; [Azzurro et al., 2019](#)). The Suez Canal is an important pathway for Red Sea species, which indeed represent 2/3 of the NIS in the Mediterranean Sea (Galil et al., 2017). In the past, it was assumed that NIS could establish only in shallow waters, however, the deep sea is not immune to species invasions. NIS have been rarely documented in the deep sea, a notable exception is the red king crab *Paralithodes camtschaticus* in the Barents Sea (Jørgensen and Nilssen, 2011). Yet, a growing number of Erythraean species were reported from the deeper part of the continental shelf, beyond the shelf break and in the upper slope (Özcan et al., 2008; Corsini-Foka et al., 2010; Innocenti et al., 2017; Özgür Özbek et al., 2017). For example, the lethally poisonous silver-cheeked toadfish, *Lagocephalus sceleratus*, has been collected from 350-400 m depth off Spain (Izguendo-Munoz and Izguendo-Gomez, 2014). The invasive lionfish, *Pterois miles*, that was initially present only in the upper shelf has been recently recorded at depths down to 110-150 m (Yağlioğlu and Ayas, 2016; Jimenez et al., 2019). In the southern Levantine Sea, three carnivorous species of Erythraean origin have been observed at 200 m depth: the crocodile toothfish *Champsodon nudivittis*, the burrowing goby *Trypauchen vagina* and the red-eye round herring *Etrumeus golanii* (Galil et al. 2018). The presence of deeper dwelling populations suggests that thermal niche assessments based only on a species' native range may underestimate their ability to tolerate lower temperatures (Parravicini et al. 2015). Wider thermal tolerance of some Erythraean species may facilitate their bathymetric and geographic expansion to depths where unique, diverse, and fragile mesophotic 'animal forests' occur. The lately observed "descent" of NIS from the upper to lower continental shelf may be an indication of temperature-dependent range expansion at increasing water depths, and appears to be accelerating. Therefore, even if abundances of NIS at levels of true invasions have not been reported yet in the deep Mediterranean, these vulnerable environments should be monitored also for D2, as they could be future targets of NIS invasions.

2.3 Descriptor 3: Populations of commercially exploited fish and shellfish

Descriptor 3 determines that Member States should maintain commercially exploited stocks of fish and shellfish in a healthy state. This descriptor implies sustainable exploitation that does not exceed the Maximum Sustainable Yield (MSY), i.e. the maximum yield catch that can be taken annually without reducing the fish stock productivity. Heavy fishing pressures, such as overexploitation or overfishing, produce negative environmental and socio-economic impacts, ranging from loss of significant potential yield of targeted stocks to severe stock depletion and fisheries collapse (Gascuel et al., 2016). Overfishing can also reduce fish stocks dramatically to the point where they lose internal genetic diversity and, with it, their capacity to adapt to environmental change (Pinsky and Palumbi, 2014; Allendorf et al., 2014). Fish communities may also change, such as altered size structures, when fisheries target or discard particular-sized individuals of a species, may potentially affect predator and prey dynamics (Fanelli et al., 2010), i.e. Descriptor 4 addresses the question of trophic relationships and marine food webs. The MSFD builds on existing EU legislation such as the Common Fishery Policy (CFP), and the criteria describing stock status follow internationally acknowledged best practices. The SAC of the GFCM, the STECF of the European Commission and the ICCAT (for highly migratory species, such as tunas or swordfishes, which account for more than 10 % of the value of the total catches in the Mediterranean) collectively monitor exploitation of fisheries resources in the Mediterranean marine sub-regions. The FAO and GFCM oversee collection of fisheries monitoring data in the Mediterranean Sea within GSAs (Geographical Sub Areas management divisions, according to resolution GFCM/33/2009/2, www.gfcm.org, for the correspondence between GSA numbers and their names Fig. 2), often assessing stocks over one or several GSAs. However, the MSFD sub-regions do not match with the GSAs. Furthermore, when we focus our attention on depths >200 m, the distinction between shallow and deep-water species is often irrelevant because distribution, exploitation and assessment of many stocks often cover wide depth ranges.

Descriptor 3 stipulates the need for fishery-induced mortality, yielding (but not exceeding) MSY (D3C1), and that populations of all commercially exploited species should remain within safe biological limits (D3C2), with a population age and size distribution (D3C3) indicative of a healthy stock. Fulfilling D3 criteria for deep-sea stocks requires: (1) sustainable exploitation consistent with high long-term yields, (2) maintaining full reproductive capacity in order to maintain stock biomass, and (3) maintaining or increasing the proportion of older and larger fish/shellfish, an

295 indicator of a healthy stock. Achieving GES also for a deep-sea stock requires fulfilling all three of
296 these attributes and, for the reasons highlighted above, D3 indicators require trans-national
297 cooperation at the level of each MFSD sub-region.

298 In the Mediterranean Sea, the enforcement of the CFP and, more recently, of the MSFD,
299 continues to fall far short of achieving its objectives for exploited living marine resources (e.g.,
300 Colloca et al., 2013; Vasilakopoulos et al., 2014). Notwithstanding the enforcement of the EU-Data
301 Collection Regulation (EU, 2000) in the early 2000s by all EU Member States, and the rapid
302 increase in the number of assessed stocks by the GFCM and the STECF, industries continue to
303 exploit Mediterranean Sea marine resources above MSY levels, with few signs of population
304 recovery (Vasilakopoulos et al., 2014; Cardinale et al., 2017; Colloca et al., 2017).

305 Management practices of DSF and VMEs in the Mediterranean were reviewed in 2016 and
306 2017 (FAO, 2016 and GFCM, 2017). UNGA Resolutions 51/2006 and the FAO International
307 Guidelines for the Management of Deep-sea Fisheries in the High Seas (FAO, 2009), identify DSF in
308 the Mediterranean Sea, as those: a) using bottom-contact gears or b) using deep-pelagic trawls
309 and c) targeting species associated with the sea floor between 300/400 m and 1000 m depth (FAO,
310 2016 and GFCM, 2017). This grouping considers shallower fisheries that extend below 400 m.
311 These reports identify deep-water red shrimps (*Aristaeomorpha foliacea* and *Aristeus antennatus*)
312 as the primary DSF targets in the Mediterranean deep-sea habitats, which are harvested mostly at
313 300/400–800 m depths, along with European hake (*Merluccius merluccius*), Norway lobster
314 (*Nephrops norvegicus*) and deep-water rose shrimp (*Parapenaeus longirostris*). In addition, gillnet
315 fisheries and demersal long-liners target both *M. merluccius* and the blackspot seabream (*Pagellus*
316 *bogaraveo*). Also in Spain, deep-sea fisheries target *Plesionika* shrimps below 300 m depth (see
317 also IDEM 2018a). GFCM banned the use of towed dredges and trawl nets at depths beyond 1,000
318 m in 2005 (Recommendation GFCM/29/2005/1), protecting over ca. 1,700,000 km² of
319 Mediterranean Sea seafloor habitats (about 59% of the GFCM area of application) (FAO, 2018).

320 To date, the GFCM has established a number of Fishery Restricted Areas (FRAs) to protect
321 Essential Fish Habitats (EFHs) or/and VMEs from excessive fishing mortality or the significant
322 adverse impact of fishing activities through bottom-contact fishing gears, respectively. The FRAs
323 encompass a total marine area of ca. 22,500 km² (FAO, 2018). Four of the FRAs were declared
324 within a multiannual management plan for deep-sea fisheries, in order to protect EFHs for
325 spawners of several species that are heavily exploited, to maintain habitat of the continental slope
326 (canyons and submarine canyons), and to preserve all the species of the area (commercially

exploited or not), i.e., one in the Gulf of Lion in France (Le Corre and Farrugio, 2011), one in the Jabuka-Pomo Pit (Elahi et al., 2018) and two FRAs south of Sicily. The existence of management plans, however, does not necessarily imply regular completion of accurate stock evaluation. Moreover, the existence of the FRA does not necessarily imply banning bottom trawling and may simply represent an effort management tool (to prevent further effort increase as seen in freezing fishing effort in the Gulf of Lion FRA).

Member States shall establish a list of commercially exploited species to which the criteria apply in each assessment area through regional or sub-regional cooperation, and update that list for each six-year evaluation period, taking into account Council Regulations (EU) 1251/2016, 1380/2013, 1343/2011, and 1967/2006, in accordance with article 43 (3) of the Treaty on the Functioning of the European Union, article 9 of Regulation (EU) No 1380/2013 and article 19 of Regulation (EC) No 1967/2006.

The MSFD criteria available for coastal environments cannot often be directly utilised for deep-sea species. This is because stock assessments are limited and not sufficiently monitored, hampering the assessment of stock exploitation at MSY (Criterion D3C1). The available information decreases eastwards and southwards. In addition, a gap of knowledge is also present in terms of time-series coverage of Spawning Stock Biomass (SSB) (Criterion D3C2) trend data hampering the possibility to define appropriate reference points.

The third criterion, i.e. "Healthy age and size structure" (criterion D3C3), assumes that a stock with sufficient large, and therefore old, fish corresponds to a healthy stock, thus reflecting good status. The larger and older fish stocks indicate healthier conditions, but this criterion has not been developed because GES lacks accepted thresholds (European Environment Agency, 2018).

This gap suggests a need to identify and test suitable indicators, metrics, and thresholds for populations and age size distributions for each deep-sea stock (ICES, 2016, 2017). In conclusion, our analysis points out that there is a potential to inform D3 criteria, but more data need to be collected in the future to propose sound stocks analyses and reference conditions, likely though the extension of the EU Data Collection Multiannual Programme (DC-MAP, EU Regulation 2016/1701) to include more deep-sea species.

2.4 Descriptor 4: Marine food webs

Descriptor 4 addresses the functional aspects of marine food webs, especially the rates of energy transfer within the system and levels of productivity in key components. In the context of the MSFD, this descriptor reaches GES when “All elements of the marine food webs, to the extent that they are known, occur at normal abundance and diversity and population levels capable of ensuring the long-term abundance of the species and the retention of their full reproductive capacity” (MSFD, 2008/56/EC, Annex I).

Deep-sea food webs critically depend on input of organic carbon from the photic zone (Thomsen et al., 2017). The microbial loop and viral infections can play an important role in the functioning of deep-sea food webs both in the water column and in sediments, and in the control of the biogeochemical cycles (Danovaro et al., 2008). At the same time, deep-sea meiofauna represents a potential basic linkage in energy transfer from the benthic detritus and microbes to the macro- megafauna and demersal fishes (Van Oevelen et al., 2011; Gambi et al., 2014).

D4 is one of the most controversial MSFD Descriptors in terms of protocols, criteria, and thresholds (ICES, Report 2015). D4 is generally investigated along with Descriptors D1 and D6 or D3. Studies of food web properties typically utilize two different approaches: a) Stomach Contents' Analyses (SCA) and b) Stable Isotope Analyses (SIA), and fatty acid trophic markers. Modelling techniques, in contrast, can provide insights regarding the potential structure of food webs (Rombouts et al., 2013). SIA identified three trophic levels among deep-sea supra-benthos (Fanelli et al., 2009) and four levels within macrozoobenthos (Iken et al., 2001; Fanelli et al., 2011a) and macrozooplankton/micronekton (Fanelli et al., 2011b). Fishes, decapods, and cephalopods dominate higher trophic levels in deep-sea demersal communities. Despite the availability of a large dataset for the deep Mediterranean Sea (MEDITS, 2002), this dataset includes few commercial species from bathyal depths: the European hake *Merluccius merluccius* and the greater forkbeard *Phycis blennoides*, some sharks (mostly *Etmopterus spinax* and *Galeus melastomus*), decapods (the red shrimps *Aristaeus antennatus* and *Aristaeomorpha foliacea*, the rose shrimp *Parapenaeus longirostris* and the Norway lobster *Nephrops norvegicus*). We lack sufficient data on other dominant deep-sea species, such as macrourids, or key predators such as deep-sea sharks (e.g., *Centroscymus coelolepis*; Stefanescu et al., 1994; Massutí et al., 2004; Anastosopoulou et al., 2016).

Most studies on trophic functional groups have focused on macro- and megafauna (i.e. fish, decapods, cephalopods and echinoderms), whereas few studies are available on other biotic compartments such as meiofauna or mesozooplankton (Danovaro et al., 2010). The northwestern

Mediterranean portion is better studied (Fanelli et al., 2009, 2011a, b, 2013, 2015, 2016; Papiol et al., 2013; Cresson et al., 2014), whereas the Ionian and Aegean seas have been much less investigated (Carlier et al., 2009; Koppelman et al., 2009; Tecchio et al., 2013; Cartes et al., 2014; Naumann et al., 2015). On the other hand, these types of studies rarely consider some key species/taxa, such as mesopelagic fishes or megazooplankton (Fanelli et al., 2014; Valls et al., 2014).

COMM/DEC/848/2017 sets criteria and methodological standards for monitoring and assessment of GES within the theme "Ecosystems". For example, selection criteria require that at least one of the three trophic guilds monitored should focus on primary producers. This criterion is the major drawback for this descriptor given that, aside from the few, very localized ecosystems that depend on chemosynthesis (hydrothermal vents, cold seeps, or wood and whale falls; Luna et al., 2012; Molari et al., 2013), the vast majority of the deep sea lacks primary production. The data gap between experimental and functional data adds further complication.

Stable isotope analysis may comply with the primary criterion D4C1 (diversity of trophic guilds) and the secondary criterion D4C3 (distribution of individuals across the trophic guild). Moreover, in combination with abundance, biomass, and other biological data available from MEDITS data, it may offer inputs into ecosystem models that could generate useful outputs, such as identification of unrecognized keystone species, a gap not presently considered. Italy addresses D4 under its fishery monitoring program (i.e. D3) and specifically with three subprograms aimed at: i) defining, testing, and applying ecosystem indicators through models (essentially EwE, <http://www.ecopath.org>); ii) identifying functional groups through the application of stable isotope analysis of monitored species within the DCF; and iii) integrating analysis of commercial species with those for benthos, zooplankton and Particulate Organic Matter (POM) samples every three years. Spain and France recently introduced SIA and SCA of species collected during MEDITS or MEDIAS surveys for use in D4.

In conclusion, the analysis of D4 can be realistically initiated for the deep Mediterranean, using the available technologies, protocols and monitoring programs and adapting the criteria, by neglecting the relevance of primary production, which could be replaced by the analysis of the inputs of primary organic matter and/or by starting from primary consumers and/or including the chemosynthetic primary production.

2.5 Descriptor 5: Human-induced eutrophication

422 Eutrophication refers to “a process driven by enrichment of water by nutrients, especially
423 compounds of nitrogen and/or phosphorus, leading to increased primary production and biomass
424 of algae, changes in the balance of organisms, and water quality degradation” (Ferreira et al.,
425 2010). According to the MSFD, GES is achieved with respect to eutrophication when “Human-
426 induced eutrophication is minimised, especially adverse effects thereof, such as losses in
427 biodiversity, ecosystem degradation, harmful algae blooms and oxygen deficiency in bottom
428 waters” (MSFD, 2008/56/EC, Annex I).

429 In coastal environments, eutrophication means the increase of primary production generally
430 monitored through the chlorophyll-*a* (Chl *a*) content and/or macroalgal biomass (Ferreira et al.,
431 2007; Xiao et al., 2007; Borja et al., 2008; Bricker et al., 2008; Nixon, 2009). The effects of
432 eutrophication are the lowering of oxygen concentrations, losses of submerged aquatic
433 vegetation, and mass mortalities especially at the sediment water interface (Claussen et al., 2009;
434 Ferreira et al., 2011). The global occurrence and expansion of hypoxic/anoxic events at bathyal
435 depths worldwide point to the need for better understanding and monitoring of the effects of
436 such phenomena on deep-sea benthic communities (Doya et al., 2015; Breitburg et al., 2018).
437 Recent models foresee an oxygen decline from 1 to 7% in the next 100 years (Keeling et al., 2010)
438 with an increase in the extension of Oxygen Minimum Zones (OMZs) (Stramma et al., 2008).
439 Farther, OMZs, with their naturally occurring low pH and oxygen, offer some hints as to the
440 structure of deep-sea ecosystems affected by eutrophication (Levin, 2003; Moffit et al., 2015).

441 Deep-sea ecosystems have been historically considered as a food-poor environment, and
442 this is typically true for the deep Mediterranean Sea, especially in its eastern basin, but some areas
443 may experience symptoms of eutrophication and oxygen depletion (Danovaro et al., 2014). For
444 instance, it has been reported that massive phytodetritus exports from highly productive coastal
445 waters to the deep-sea floor (Billet et al., 1983). Excessive C inputs in combination with the high
446 bottom temperatures can cause episodic oxygen depletion in the deep sea (Ferreira et al., 2011;
447 Danovaro et al., 2014). Recent studies highlighted that deep-sea trophic status can be also
448 affected by climate change, as the Western basin is expected to become more oligotrophic and
449 the Eastern basin more eutrophic (Piroddi et al., 2017). In addition, predicted increasing surface
450 temperatures may affect water mass stratification and the formation of cold oxygenated deep
451 water, modifying global ocean circulation and the dissolved oxygen availability in deep-water
452 masses (Ramirez-Llodra et al., 2011). Local scale eutrophication could affect deep-sea sediments

453 facing highly productive areas of the Mediterranean Sea, such as the Gulf of Lions, the northern
454 Aegean Sea and the Ionian Sea receiving inputs from the Adriatic Sea.

455 However, the MSFD, in relation to the qualitative Descriptor 5, calls for an assessment of
456 nutrients and organic matter inputs (Annex III of Directive 2017/845) and the use of the following
457 criteria (Directive 2017/848): i) nutrient concentrations in the water column, ii) chlorophyll a in the
458 water column, iii) harmful algal blooms, iv) photic limit, v) dissolved oxygen at the bottom of the
459 water column, vi) opportunistic macroalgae, vii) macrophyte communities and viii) macrofaunal
460 communities. These criteria can only be partially applied to the deep sea. Firstly, primary
461 producers (e.g., macrophyte, macroalgae, harmful algal bloom) must be excluded, and the
462 assessment of trophic status using variables measured in the water column can lead to misleading
463 classifications (Dell'Anno et al., 2002, see also Fabri et al., 2018). Considering that oxygen
464 depletion is one of the main causes of benthic faunal mortality, it is important to measure
465 physical-chemical parameters and indicators also in the sediments (Mercado et al., 2015). In
466 addition, the concentration of organic matter accumulated in surface sediments can provide a
467 good indication of the eutrophication process occurring on the seafloor (Dell'Anno et al., 2002;
468 Pusceddu et al., 2009). The current conceptual framework suggests the need to introduce new
469 criteria and indicators, related to benthic ecosystems and, particularly, to the deep sea.

470 A group of core indicators is already utilised to monitor eutrophication in open waters,
471 including: i) nutrients (nitrate, ammonium, phosphate), ii) dissolved oxygen and iii) phytoplankton
472 (chlorophyll a, dominance). Zooplankton biomass is considered a potential, though not fully
473 mature indicator (UNEP(DEPI)/MED, 2007 and references therein) because of incomplete
474 knowledge of its relationship to eutrophication. The monitoring of benthic ecosystems should
475 include i) quantity and quality of sedimentary organic matter, and ii) biodiversity and taxonomic
476 composition of benthic invertebrates (Dell'Anno et al., 2002; Pusceddu et al., 2009, 2011, 2014;
477 Bianchelli et al., 2016a). Among indicators recently proposed to assess benthic trophic status of
478 marine ecosystems, the quantity and biochemical composition of sedimentary organic matter has
479 received the widest application, both in coastal and deep-sea ecosystems (Pusceddu et al., 2009;
480 see also Fabri et al., 2018). The concentrations of biopolymeric C (defined as the sum of C deriving
481 from proteins, carbohydrates and lipids) and its algal fraction have been used to assess impacts of
482 humans on benthic trophic status in different oceanic and coastal regions and varying water
483 depths, within the Mediterranean basin (Dell'Anno et al., 2002; Pusceddu et al., 2009, 2011, 2014;
484 Bianchelli et al., 2016b). Changes in the quantity and quality of organic matter in the deep sea

lead to responses from all benthic components, from prokaryotes to foraminifera, and from meiofauna to macrofauna (Danovaro et al., 1999). Also the functional traits of macrofauna have been widely used as indicators of alteration and for measure the health status of marine benthic ecosystems (Borja et al., 2008). Further, meiofauna could be considered a good indicator as it is highly sensitive to environmental changes, and particularly to organic enrichment due to eutrophication (Pusceddu et al., 2011). For these reasons, meiofauna have been recently proposed for the monitoring of eutrophication effects and for assessing the environmental quality of both coastal and deep-sea ecosystems (Bianchelli et al., 2016a; Pusceddu et al., 2016).

2.6 Descriptor 6: Sea floor integrity

Descriptor 6 requires that seafloor integrity is *“at a level that ensures that the structure and functions of the ecosystems are safeguarded and benthic ecosystems, in particular, are not adversely affected”* (Commission Decision 2010/477/EU). This involves addressing *“physical damage, having regard to substrate characteristics”*, and the *“condition of benthic community”*, the latter being directly related to Descriptor 1. The relevant pressures identified in the Commission Decision (EU) 2017/848 generically refer to “physical loss (due to permanent change of seabed substrate or morphology and to extraction of seabed substrate)” and to “physical disturbance to seabed (temporary or reversible)”. Four primary criteria address these points, three of which (D6C1 to D6C3) are specific for Descriptor 6, while two are also relevant for Descriptor 1 (D6C4 and D6C5).

The deep Mediterranean seafloor experiences two dominant physical disturbances associated with human activities: i) bottom-contact fisheries and ii) oil and gas activities (Boschen et al., 2013; D’Onghia et al., 2017; Lauria et al., 2016; Holler et al., 2017). Fisheries using bottom-contacting gear lead to direct alteration of seafloor morphology at large, medium and small scales (Puig et al., 2012; Martin et al., 2014). Bottom trawling is a key driver for large-scale seascape change as it smoothens the natural topography (Puig et al., 2012). Direct and indirect biological effects of bottom trawling have been demonstrated in terms of biogeochemical changes (e.g. less total amino acid concentration in sediments) and faunal desertification (Pusceddu et al., 2014). The Mediterranean Sea shows the highest fisheries footprint per unit landings in Europe (Eigaard et al., 2017), with peak intensities in the Tyrrhenian and the Adriatic Sea. In the Catalan margin, trawling impact is major down to 800 m depth (Puig et al., 2012). Sediment resuspension from fishing grounds can propagate to wider and deeper areas eventually leading to suffocation and

517 burial of VME (Martin et al., 2008). Downslope moving gravity-driven resuspension flows enhance
518 sedimentation rates far beyond fishing grounds, such as in canyon axes. Other types of fishing gear
519 such as bottom touching longlines and gillnets could also have a significant adverse impact on
520 vulnerable benthic communities and organisms such as black corals, gorgonians, scleractinians and
521 many other habitat-forming species (GFCM, 2017), because of breaking while pulling, ghost fishing
522 or entanglement.

523 Activities undertaken by the offshore oil and gas industry may cause physical loss of the
524 natural deep seabed. Physical (and chemical) impacts on the seafloor and subseafloor range from
525 the installation of drilling rigs, wellheads and other structures on the seabed to the accumulation
526 of litter including lost or abandoned equipment, consumables and other materials. Today's deep-
527 water (>200 m) oil and gas production in the Mediterranean Sea, or advanced prospects for it,
528 takes place essentially offshore Egypt, Israel, Lebanon, Syria and Cyprus (The Petroleum Economist
529 Ltd, 2013; Galil and Herut, 2011). The environmental approach for the hydrocarbon industry in the
530 Mediterranean Sea is developed in the Offshore Protocol of the Barcelona Convention, adopted in
531 October 1994, which obliges countries to perform comprehensive EIAs after entering into force in
532 December 2012. The EU adopted the Directive on Safety of Offshore Oil and Gas Prospection,
533 Exploration and Production Activities in July 2013, which provides a blueprint of the best
534 international practice also for non-EU countries in the Eastern Mediterranean that are new to the
535 energy industry (Livnat, 2014). Further disturbance occurs in case of cable deployment, for not the
536 cables and pipelines per se, rather for the impact of the anchoring of the supply vessel during the
537 deployment of the cable.

538 Dumping of industrial waste in the deep Mediterranean Sea is a matter of concern for
539 habitat integrity. Submarine canyons with heads close to the coast are favoured sites for direct
540 deep-sea disposal (Ramirez-Llodra et al., 2015). Two aluminium-processing plants have discharged
541 red mud waste in the deep Mediterranean Sea: one in France (Cassidaigne Canyon, Gulf of Lion)
542 (Dauvin, 2010; Fontanier et al., 2012, 2014; see also Fabri et al., 2018) and one in Greece (Gulf of
543 Corinth, Antikyra Bay) (Varnavas et al., 1986; Varnavas and Archilleopoulos, 1995; Poulos et al.,
544 1996). Since 1988, Coal Fly-Ash (CFA) from the Hadera power plant, in Israel, has been dumped
545 into a 16 km² disposal site some 70 km offshore, at a water depth of 1400 m, where a 0.5-1.0 cm
546 thick ash layer has been noticed (Kress et al., 1996, 1998) together with severe impoverishment of
547 benthic fauna. Israel allowed also long-term disposal of dredged sediments and industrial waste

1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121

548 (1,900,000 m³) polluted with Hg, Cd, Pb, tributyltin and organotins, and PCBs at a site 1300 m deep
549 (Herut et al., 2010).

550 Different proved tools is currently available to assess seafloor integrity. High-resolution
551 maps of benthic substrata and habitats are increasingly required both to underpin environmental
552 and socioeconomic impact assessments and to help in developing effective management
553 measures (Kenny et al., 2003; Brown et al., 2011; Stephens and Diesing, 2014; Holler et al., 2017;
554 Fabri et al., 2018). Multibeam Echo-Sounders (MBES) and side scan sonars (SSS), map seabed areas
555 with 100% spatial coverage at a resolution finer than 1 m², depending on the depth of data
556 collection and on distance-to-bottom of the sensors (Kenny et al., 2003). Ground-truthing
557 methods, [such as the use of remotely operated vehicles \(ROVs\) and autonomous underwater](#)
558 [vehicles \(AUVs\) \(Fabri et al., 2014; Lastras et al., 2016\)](#), are widely available and could be applied
559 according to the size and the nature of the area of interest (Kenny et al., 2003; Brown and Blondel,
560 2009; Brown et al., 2011; Holler et al., 2017). Habitat suitability models try to predict the
561 distribution of some habitats such as CWCs (Lo Iacono et al., 2012; Bargain et al., 2017, 2018; Fabri
562 et al., 2017; Angeletti et al., 2019; Lo Iacono et al., 2018; Giusti et al., 2014, 2017; Lauria et al.,
563 2017). However, because such models often include a large degree of uncertainty, decisions based
564 entirely on modelling approaches may involve significant risk.

565 The revision of the MSFD ([through the COMM. DEC. 2017/848/EU](#)) emphasised that
566 “Physical loss shall be understood as a permanent change to the seabed which has lasted or is
567 expected to last for a period of two reporting cycles (12 years) or more”, but for this to be
568 implemented, a very long time perspective is needed. All impacts described in this section have
569 immediate effects (and sometimes also delayed effects) on seafloor communities, which in most
570 cases could represent either a tipping point (e.g. large-scale seascape change) or require long time
571 before any significant recovery could take place. A time-span of 12 years is possibly too short and
572 it is urgent to proceed with a sound extensive evaluation of the current status of the deep benthic
573 habitats in the Mediterranean Sea before human impact severely modifies or erases them from
574 the face of our planet.

575
576 **Descriptor 7: Permanent alteration of hydrographical conditions**

577 Descriptor 7 is geared towards addressing the problem of the permanent alteration of
578 hydrographical conditions. These conditions are often affected by the presence of coastal
579 infrastructure and other man-made activities (ports, artificial reefs, etc.). However, in most cases

1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180

these structures impact coastal areas and only rarely can reach higher depths. Conversely, global climate change, combined with episodic climate-driven events, can alter the “hydrographical conditions” also at depths. In recent decades, deep Mediterranean waters have experienced drastic changes resulting in an alteration of the stratification associated to temperature increases and salinity shifts (Schroeder et al., 2009, 2016). At the end of the 1980s, climate change, changing hydrographic properties, surface circulation, and deep-water convection caused a ‘regime shift’ on a global scale (Reid et al., 2016) and throughout the Mediterranean basin (Conversi et al., 2010). During that period, the main site of deep-water formation shifted from the southern Adriatic to the Aegean sub-basins. This “Eastern Mediterranean Transient” (EMT; 1987-1994) event resulted in increased oxygen consumption (Roether and Well, 2001; Klein et al., 2003) and, in the eastern Ionian, in a nutricline shoaling by about 150 m (Klein et al., 1999). Other rapid hydrological changes have also occurred in the Western Mediterranean Sea. The “Western Mediterranean Transient” (WMT; 2004/05 and 2005/06) was characterised by the formation of warmer and denser new deep waters over the continental shelf as a result of cooling and evaporation of the surface layer and downslope cascading (Canals et al., 2006; Palanques et al., 2006; Schroeder et al., 2009). The high volumes of newly formed deep waters generated during intense cascading and convection events dramatically altered the hydrological structure of the basin, completely de-stratifying the water column and transferring massive heat and salt to the deep layers (Canals et al., 2006; Schroeder et al., 2009; Martin et al., 2010). The 2004/05 event was the first of a series of similar events in the last decade that greatly altered the structure of the intermediate and, especially, the deep layers of the Western Basin (Durrieu de Madron et al., 2013). Cascading events transport huge amounts of nutrients and organic matter, to bathyal depths (Canals et al., 2006; Sanchez-Vidal et al., 2008; Danovaro et al., 1999; Company et al., 2008). Hydrographic preconditioning (heat and salt content and structure of the water column before the onset of convection), and atmospheric forcing (heat, freshwater and buoyancy fluxes) triggered deep-water formation (Fabri et al., 2018). Moreover, progressive increase in heat and salt content in the intermediate layer, advected from east to west, favoured new dense water formation in the North-Western Mediterranean basin. Multiple heat and saline anomalies characterised the Mediterranean Sea from 1950 to 2000 (Rixen et al., 2005; Kress et al., 2014) and although these alterations cannot be considered permanent, all of these changes have long-term effects.

The multidisciplinary Mediterranean Targeted Projects MTP-I and MTP-II/MATER (1993-2000; Monaco and Peruzzi, 2002), the MEDAR/MEDATLAS database (Fichaut et al., 2003), the

SeaDataNet (i.e., <https://www.seadatanet.org/>) and EMODnet (<http://www.emodnet-physics.eu>) infrastructures created datasets on temperature, salinity oxygen, silicate, nitrates and phosphates, but in some cases with insufficient coverage in the eastern region (including Tunisia, Libya, Croatia and Turkey; Simoncelli et al., 2015).

Since the 2000's, national and international programmes (e.g. EU-PERSEUS and MedSeA, IT-VECTOR, FR-MERMEX, E-RADMED) produced hydrological data on the whole Mediterranean Basin. Regular cruises, mooring lines, and deployment of new instruments and infrastructure (Argo floats, gliders) now support intensive collection of *in situ* observations in the North-Western Mediterranean Sea. Argo floats, autonomous profiling floats, drift at a given depth for a given time period. After drifting for a set time, they sink to 2000 m and profile temperature and salinity during the upcast. These data proven useful in describing deep-water formation (Smith et al., 2008). Moreover, in the last decade, glider technology mainly in the North-Western Mediterranean Sea has enabled repeated cross-basin transects at depths from the surface to 1000 m. In addition, numerical models implemented at regional or local scales use these data to elucidate water mass formation and spreading, and basin-scale hydrological dynamics (Bonaldo et al., 2015; Estournel et al., 2005).

Long-term monitoring of basic hydrological parameters (temperature and salinity), collected as time series with appropriate temporal resolution (i.e. sampling intervals that resolve all relevant timescales) represent a science priority in the context of climate change study for key locations in the Mediterranean Sea (e.g. straits and channels, zones of dense water formation and spreading, deep basins) (Schroeder et al., 2013; Aguzzi et al., 2019). The HYDROCHANGES network aims to address this need by deploying moorings fitted with Conductivity, Temperature, Depth sensors (CTDs) at key locations for monitoring temperature and salinity (Schroeder et al., 2013). The FixO3 (Fixed point Open Ocean Observatory network, <http://earthvo.fixo3.eu/>) programmes and the EMSO (European Multidisciplinary Seafloor and water column Observatories, <http://emso.eu/>) EU Infrastructure provide additional near-bottom data from fixed-point observations. For example, DYFAMED and LION deep-water stations, and ANTARES neutrino telescope in the North-Western Mediterranean or the KM3NET in the Ionian Sea (Tamburini et al., 2013; Aguzzi et al., 2018), continuously monitor sets of specific parameters. Long time series provided by these mooring stations have contributed pivotal findings on the deep dynamics of the Mediterranean Sea in last years. In order to partially interpolate data within homogeneous habitats, scientists have generated gridded products through objective analysis of available observations (such as

numerical models with data assimilation delivered by Copernicus downstream services (<http://marine.copernicus.eu/services-portfolio/access-to-products/>).

2.8 Descriptor 8: Concentrations of contaminants giving rise to pollution effects

The input of xenobiotic substances represent one of the major threats for ocean health (Halpern et al., 2008). Hydrophobic pollutants, such as organo-halogenates and polycyclic aromatic hydrocarbons (PAH), enter the marine environment through effluent discharges, atmospheric deposition, runoff and other means (Iwata et al., 1994). Once in the water column, the adsorption onto particulate matter transfers these compounds from the surface to the deep waters and sediments (Buesseler, 1998). Particle settling is also favoured by biological processes, and by lateral transport from continental shelves (Heussner et al., 2006; Martin et al., 2006; Zuñiga et al., 2009). DSWC is a massive mechanism of pollutant transfer to the open deep ocean (Canals et al., 2006). Higher fluxes of organohalogen pollutants and Polycyclic Aromatic Hydrocarbons (PAH) occur during these cascading events (Salvadó et al., 2017). PAH settling fluxes in the north-western Mediterranean Sea vary widely (Lipiatou et al., 1993; Raoux et al., 1999), with highest concentrations of contaminants in the Alboran Sea (Dachs et al., 1996), and much lower values in Sardinia and in the Southern Ionian Sea (Bouloubassi et al., 2006; Tsapakis et al., 2006). However, these values greatly exceed atmospheric deposition of PAH in central sites of the Western Mediterranean Sea, thus highlighting the role of river discharge (Heussner et al., 2006; Bonnin et al., 2008; Palanques et al., 2008). Qualitative differences are also observed in relation to these transfer processes. Sediments of coastal areas, continental shelves and slopes have higher proportions of petrogenic PAHs whereas the deep basin of the north-western Mediterranean Sea is characterized by high amounts of pyrogenic PAHs.

Organochlorine compounds such as Polychlorobiphenils (PCBs) and chlorinated pesticides, characterised a group of Persistent Organic Pollutants (POPs) of worldwide concern due to their toxic effects (Harmon, 2015). Notwithstanding the discontinued use of these compounds in most world areas, thanks to relevant national regulations and international agreements such as the Stockholm Convention, their extensive occurrence is still observed. Their high lipophilicity, hydrophobicity, chemical stability and resistance to biological degradation have led to their accumulation in biological tissues and biomagnification through the food chain.

Radioactive compounds in the Mediterranean Sea are derived from the fallout of nuclear weapon testing and the Chernobyl accident. In sediments, concentrations of ^{137}Cs and $^{239+240}\text{Pu}$

have been measured in various parts of the Mediterranean Sea, including deep basins. The concentrations are generally higher in coastal ecosystems because land-based sources can exceed atmospheric inputs (Durrieu de Madron et al., 2011; Garcia-Orellana et al., 2009). Concentrations in biota are presently undistinguishable from those in areas without point sources. Hence, the relevance of these contaminants lies in their usefulness as process tracers, more than on their impact on the environment. Studies on radionuclides in marine organisms also underscore that the radionuclide levels are constantly decreasing due to modifications of the inputs. Very little work has been done to examine the trophic transfers of man-made radionuclides (Harmelin-Vivien et al., 2012). Finally, neglected impacts that can be very important in several areas of world are military activities. Information on their impacts on the environment are relatively scarce and are often studied after several years from their production and without any baseline available (Lawrence et al., 2015; Danovaro et al., 2019).

Atmospheric inputs constitute one of the major sources of Trace Elements (TE) to the deep Mediterranean Sea (Migeon et al., 2012; Guerzoni et al., 1999), where TE concentrations in waters are typically higher than in other areas of the world ocean. In addition, Cd, Cu and Ni (as well as Cr) are dominated by lateral advection and vertical mixing rather than by biogeochemical cycling (Morley et al., 1997). The hydrologic regime of the Mediterranean Sea tends to transfer the pollutants and nutrients to the Atlantic by bottom water flow transport. Our knowledge on the concentrations, fluxes, and behaviour of trace elements, radionuclides and organic substances in the deep waters and sediment and their toxicological impacts on habitats and organisms is scarce (Durrieu de Madron et al., 2011). Pollutants with hydrophobic properties, e.g. PCBs and mercury, accumulate in biota and thus in the food web. In case of chronic pollution events, the concentration of contaminants should be analysed in sediments collected by sediment cores, which enable the reconstruction of temporal trends. Sample collection of water at different depths and analysis of the dissolved and particulate matter would also be important. Abundance of populations and estimates of the extent of habitats adversely affected by chronic pollution should be assessed concurrently.

Nonetheless, information on pollutants in the deep sea is almost completely lacking, and this represent the main gap in the application of the criteria needed to determine the D8.

2.9 Descriptor 9: Contaminants in fish and other seafood for human consumption

Descriptor 9 focuses on the accumulation of toxic, persistent and liable substances in wild deep-sea organisms used for human consumption (i.e., mostly teleost and decapod crustaceans) and the contaminants considered by D9 are only part of those of interest for the D8 (cfr. Regulation EC 1881/2006 and its amendments EC 2006, 2008). Each Member State may ignore specific contaminants and/or include additional ones (EC 2017) (Fliedner et al., 2018). In any case the monitoring of the contaminants accumulated in the deep-sea biota should at least consider the following compounds for which regulatory levels have been set: i) heavy metals (lead, cadmium and mercury); ii) PAHs; iii) dioxins (including dioxin-like PCBs). In addition, the following contaminants of relevance should be monitored: i) non-dioxins like PCBs; ii) phthalates; iii) organochlorine pesticides; iv) organotin compounds; v) brominated flame retardants; vi) polyfluorinated compounds. Also, artificial radionuclides should be monitored in case of nuclear accidents or any other radioactive emergencies that could lead to or has led to significant radioactive contamination of food.

Contaminants in fish and other seafood might derive from numerous anthropogenic sources described for the D8. Chemical contamination in fish and seafood results from a complex process that balances inputs of contaminants, mostly through diet, and their excretion (Solé et al., 2001; Trudel and Rasmussen, 2001; Cresson et al., 2014).

Investigating contamination levels in fish and seafood requires understanding which contaminants exceed regulatory limits, how much they alter food webs, and what metabolic processes are involved in detoxification. The presence of xenobiotics in the deep Mediterranean organisms has been repeatedly documented (Galil et al., 1995; Storelli et al., 2009) with deep-sea Mediterranean fishes tending to exhibit higher levels of metal accumulation than those of populations inhabiting other areas such as the Atlantic Ocean (Damiano et al., 2011). Red-shrimps, *Aristeus antennatus* and *Aristaeomorpha foliacea* may be useful indicator species of levels of deep-sea contamination (e.g., see data on *A. antennatus*, Koenig et al., 2012). Contaminants in fish muscle and liver have been investigated in the most abundant deep-sea megafaunal species, e.g. *Alepocephalus rostratus*, *Coelorinchus mediterraneus*, *Coelorhynchus caelorhincus*, *Trachyrincus trachyrincus* and *Nezumia sclerorhynchus*, *Chimaera monstrosa*, *Lophius budegassa*, *Lepidion lepidion* (Koenig et al., 2013c), revealing mercury concentration exceeding $0.5 \mu\text{g g}^{-1}$ muscle wet weight in all species but one. This represents the threshold value indicated by the European Commission as acceptable for human consumption. High mercury concentration is a distinct feature of the Mediterranean Sea (the so-called “Mediterranean mercury anomaly”; Cossa and

Coquery, 2005; Cossa et al., 2012). The Hg concentration is even higher in some deep-sea fish species (Koenig et al., 2013c; Cresson et al., 2014; Chauvelon et al., 2018). Most deep-sea species are long-lived and slow-growing, which favours the bioaccumulation of pollutants (Drazen and Haedrich, 2012; Koenig et al., 2013a, b, c). Since some deep-sea species are of commercial interest, the high contamination level poses serious risks for human health (Rotllant et al., 2006; Carbery et al., 2018). Despite this, few studies have investigated the concentrations of contaminants in the deep Mediterranean fauna fished for human consumption (Storelli et al., 2004, 2007; Koenig et al., 2013c; Cresson et al., 2014). In this regard, the Gulf of Lion is the best-investigated area, whereas the Levantine Basin (with the exception of Israel waters; Galil et al., 1995) are the least studied. A comparison of the concentrations of xenobiotics in fish collected in the NW Mediterranean Sea, in 1996 and 2009 (Koenig et al., 2013a; Porte et al., 2000; Solé et al., 2001) indicate that their contamination did not change over time.

The application of the criteria of Descriptor 9 should consider the concentration, the thresholds, and the contamination sources. Regulators should also consider the species of interest for human diets and their ability to bioaccumulated pollutants. GES would be achieved if all contaminants occur at levels below those established for safe human consumption.

2.10 Descriptor 10: Marine litter

Two primary and two secondary criteria are associated to Descriptor 10: i) the composition, amount and spatial distribution of litter (D10C1) and of micro-litter (D10C2) “on the coastline, in the surface layer of the water column, and on the seabed, are at levels that do not cause harm to the coastal and marine environment” (primary) and ii) the amount of litter ingested by marine animals, which should not reach a level that adversely affect the health of the species (D10C3) and the number of individuals which are adversely affected due to litter, such as by entanglement, other types of injury or mortality, or health effects (D10C4) (secondary). Each sub-region should assess the outcomes for all criteria and as well as threshold values.

Marine litter represents a threat for the health of the deep Mediterranean Sea due to its limited exchange with other basins, dense population, touristic and industrialized coastlines, and heavy maritime traffic (UNEP, 2015). The sources of marine litter to the deep-sea floor of the Mediterranean Sea are either from land (river discharge, storm drains, sewage treatment plants and industrialized areas) or marine (fishing activities, commercial and recreational shipping, aquaculture, direct dumping), and include plastics (accounting for >70% of the total), glass, metal,

clinker, cardboard and fabrics (Galgani et al., 2000; Ramirez-Llodra et al., 2013; Fabri et al., 2014; Pham et al., 2014; Tubau et al., 2015; UNEP, 2015; [Mecho et al., 2017](#)). The quantity and composition of marine litter differs among regions and changes with depth, probably as a result of a complex set of interactions between hydrodynamics, geomorphology, and anthropogenic sources (Pham et al., 2014; Tubau et al., 2015; UNEP, 2015). The abundance of marine litter items in the deep Mediterranean Sea varies from 500 items km⁻² on the continental slopes off Malta and Cyprus (Mifsud et al., 2013; Ioakeimidis et al., 2014), the Tyrrhenian Sea (Angiolillo et al., 2015), or the Adriatic Sea (Galgani et al., 2000), to more than 2,000 items km⁻² in the Antalya Bay in the Eastern Mediterranean or in the submarine canyons of the Gulf of Lion and of the Catalan Sea (Galgani et al., 2000; Tubau et al., 2015). Astonishingly high litter abundance of up to 1.3 million of items km⁻² were reported at 300-600 m depth in the Messina Strait canyons (Central Mediterranean Sea) (Pierdomenico et al., 2019). Litter abundance found in submarine canyons and depths greater than 500 m typically exceeds that at shallower depths, suggesting that submarine canyons can act as primary conduits of litter from the coast to the deep sea (Galgani et al., 2000; Tubau et al., 2015). Superposition of highly efficient source-to-sink sedimentary transport (with flash-flood generated hyperpycnal flows) and strong urbanization of the coastal area promote the occurrence of large litter hotspots in the deep sea (Pierdomenico et al., 2019). In addition to large marine debris, concern has grown about microplastics (i.e., <1-5 mm in diameter; Desforges et al., 2014), which can directly enter the ocean also through cosmetic abrasives (i.e. microbeads), preproduction plastic pellets, or textile fibres known as primary plastics. Additionally, combined mechanical, biological, photic and thermal actions can break down larger plastic objects into numerous small fragments, which are defined as secondary microplastics. Depending on the density of the polymer, microplastics may sink and behave as very fine-grained sediments (for example polyester; Woodall et al., 2014), or they may float and subsequently sink following colonization by organisms, adsorption to phytoplankton, and/or aggregation with organic debris. Fibres appear to dominate the microplastics reported in Mediterranean deep-sea sediments (van Cauwenberghe et al., 2013; Woodall et al., 2014; Sanchez-Vidal et al., 2018). Recent studies indicate that the Mediterranean deep-sea floor might act as a long-term sink for microplastics where the abundances of such particles can exceed those in surface waters (Sanchez-Vidal et al., 2018). Marine litter in the Mediterranean deep sea may significantly affect different ecological compartments and, consequently, human health, with potentially severe economic impacts. Biotic effects of large and small items include entanglement, ingestion, colonization and rafting (Gregory,

2009; Murray and Cowie, 2011; Anastasopoulou et al., 2012; Ramirez-Llodra et al., 2013; Bo et al., 2014; Pham et al., 2014; Angiolillo et al., 2015; Tubau et al., 2015). However, information on the actual effects of (micro)plastics on deep-sea organisms and trophic webs is still limited (Taylor et al., 2016).

Marine litter in deep sea produces economic impacts primarily on the fishery sector, damaging vessels and fishing equipment due to entanglement of catch, loss of target species through ghost fishing, or reduced reproductive capacity of benthic organisms consuming microplastics (Newman et al., 2015). Furthermore, marine litter may contain pollutants (hazardous plastic additives, POPs) that exert toxic and endocrine disruptive effects on marine organisms that ingest plastics (Oehlmann et al., 2009).

2.11 Descriptor 11: Introduction of energy including underwater noise

Descriptor 11 deals with introduction of energy into the marine environment. Underwater noise can be pulsed or continuous. MSFD currently focuses on two criteria: anthropogenic pulsed (D11C1) and continuous low-frequency (D11C2) sounds in water. D11C1 addresses the space-time distribution of pulsed noise sources, whereas D11C2 addresses levels of continuous noise, using in situ measurements and models. Pulsed noise may cause direct acute effects such as hearing loss, tissue damage, and death of individuals of sensitive species such as cetaceans. Whereas continuous or chronic noise exposure mainly causes stress and behavioural alterations, with negative effects on deep-sea organisms (Nowacek et al., 2015). The proposed strategy on noise monitoring recommends several adaptations in the case of the deep Mediterranean. Particularly, both indicators are closely related to the acoustic biology of deep-diving marine mammal species, such as sperm whale and Cuvier's beaked whale. Pulsed noise can be monitored by setting up a register of anthropogenic activities, reporting on date, location, proportion of days within a given period and over a given geographical scale in which activities generating pulsed sounds occur. This could be done through the deployment of hydrophones (e.g., permanent or semi-permanent PAM) on new infrastructures (or implementation of the existing infrastructures) and their subsequent future development into larger geographic network: Rountree et al., 2019).

A variety of phenomena generates noise in the ocean, either from physical processes, such as wind-generated waves, earthquakes, precipitation, or from biological phenomena such as whale songs, dolphin clicks, and fish vocalizations (Montgomery and Radford, 2017); not all reach the deep sea. Fish produce sounds for their navigation, habitat selection and mating, as well as to

835 communicate (Simpson et al., 2005). Marine mammals use sound as a primary tool for underwater
836 communication (Wartzok and Ketten, 1999), mating and social interaction (Edds-Walton, 1997),
837 and for tracking the prey (Au, 1993).

838 Anthropogenic noise can reach the deep sea, through commercial shipping, oil and gas
839 exploration, fishing, and scientific research; all of these sources currently contribute to the general
840 noise budget of the ocean (Montgomery and Radford, 2017). The impact of noise on marine is
841 being increasingly investigated (Wenz, 1962; Hildebrand, 2009). Noise sources are divided into
842 three frequency bands: low (10 to 500 Hz), medium (500 Hz to 25 kHz) and high (>25 kHz).
843 Anthropogenic sources dominate the low-frequency band, and include commercial shipping and
844 seismic emissions for hydrocarbon exploration. Minimal attenuation of low-frequency sound
845 allows long-distance propagation. Sea-surface agitation (breaking waves, spray, bubble formation
846 and collapse, and rainfall) and various sonars (e.g. military and multibeam seabed mapping), as
847 well as small vessels produce most medium frequency sound. Greater attenuation limits
848 propagation of noise in the mid-frequency band over long distances, and only local or regional
849 (10s of km distant) sound sources contribute to this ambient noise field. At high frequencies,
850 extreme acoustic attenuation confines all noise sources to the area close to the receiver.

851 Oil industry operations have traditionally focused on shallow, continental shelf waters, but
852 exploration is moving in deeper waters (>500-1000 m). Expansion of oil exploration into deeper
853 water has increased the potential for long-range propagation of seismic reflection signals. Indeed,
854 sound in deep waters can propagate greater distances than in shallow-water ecosystems, by
855 moving through the deep sound channel (Hildebrand, 2009). Seismic surveys currently target all
856 regional seas in the south-eastern Mediterranean, apart from the Aegean Sea (Maglio et al., 2016).
857 This expansion stresses the transboundary aspect of seismic surveys and calls for international
858 cooperation.

859 Anthropogenic noise can cause physical and biological damage, such as behavioural changes
860 and stress, especially in marine mammals, sea turtles and fish (Popper et al., 2014; Peng et al.,
861 2015). The occurrence of low frequency noise in the deeper part of the basins is particularly
862 important for deep-diving marine mammals, such as toothed whales, because the ambient noise
863 they use as a background for echolocation decreases rapidly with depth (Foote et al., 2004; André
864 et al., 2011; Azzellino et al., 2011). [Indeed, small odontocetes produce high frequency sounds](#)
865 [\(ranging from 70 kHz to more than 150 kHz\), while sperm whales, *Physeter macrocephalus*, during](#)
866 [diving, make sound with frequencies ranging to more than 30 kHz which are detectable within 10-](#)

15 km. The fin whale *Balaenoptera physalus*, the only mysticete constantly present in the Mediterranean Sea, emits mostly infrasonic signals (20-40 Hz), which are emitted in long sequences and can be detected at large distances.

Research on sea turtles in the South-Eastern Mediterranean region revealed that they can detect low frequency sounds that overlap with seismic airgun frequencies, these are high intensity, low-frequency impulsive noise at regular intervals, mostly between 10 and 300 Hz (Carroll et al., 2017; Nelms et al., 2016). Airguns can stress the sea turtles, *Caretta caretta* (DeRuiter et al., 2012). The impacts of anthropogenic noise on sharks and rays are poorly studied, with most research to date focusing outside the Mediterranean region (Weilgart, 2017). Fish sensitivity to certain frequencies varies among species (Carroll et al., 2017). Recent studies demonstrate negative effects of seismic survey airgun operations even in zooplankton (McCauley et al., 2017).

In situ acoustic measurements can document continuous low-frequency sound, gathering field data on ambient noise in a given location. Understanding the large-scale influence of artificial noise on marine organisms and ecosystems represents the main gap to the application of the D11 on the deep sea. Deep-sea observatories offer new opportunities to assess the presence and effects of noise in on deep-sea life (Aguzzi et al., 2019). Deep-sea cabled observatories (i.e. NEMO-SN1 in the Western Ionian Sea, Caruso et al. 2015, Favali et al., 2013; ANTARES in the Ligurian Sea, André et al., 2017; and PYLOS in the South Ionian Sea, <http://www.fixo3.eu/observatory/pylos/>) are equipped with hydrophones for passive acoustic monitoring. Besides these measurements, and especially for monitoring continuous low-frequency sound in deep sea, modelling approaches (both for single sources or distributed sources of noise, from the most advanced Dynamic Ambient Noise Prediction System elaborated by the U.S. for modelling multiple sources, to the Acoustic Integration Model used for modelling the effects of noise on cetaceans; NRC, 2003) may reduce the time required to establish trends and patterns.

3. Future implementation

In order to effectively implement the deep-sea MSFD, we need to identify the criteria to achieve or maintain GES in open waters and deep-sea bottoms, including “spatial protection measures, contributing to coherent and representative networks of marine protected areas, adequately covering the diversity of the constituent ecosystems, such as special areas of conservation pursuant to the Habitats Directive, special protection areas pursuant to the Birds

899 *Directive, and marine protected areas as agreed by the Community or Member States concerned in*
900 *the framework of international or regional agreements to which they are parties"* (MSFD,
2008/56/EC, Article 13).

902 The MSFD takes an overarching and integrated approach by focusing on achieving GES and
903 targets, and we therefore recommend exploring and assessing synergies between the different
904 treaties, directives, and conventions (e.g. see Descriptor 6 section) so that, wherever possible, the
905 programme of measures and proposed MSFD monitoring simultaneously address the
906 requirements of other legislations.

907 Our analysis indicates that the 11 Descriptors promulgated by the MSFD (MSFD,
2008/56/EC) can be adapted and applied to the deep sea. Several Descriptors (D1, D2, D3, D6, D8,
D10) can be readily implemented, others (D4, D9 and D11) require additional data in order to set
up benchmark and threshold values, while two (D5, D7) require changes in the assumptions
and/or modification in the concept of "permanent".

912 Priority ecological variables, spatial distribution, extent of pressures and impacts ought to
be identified and standardized in order to establish targets and indicators addressing the distinct
conditions in the deep sea. The expertise, tools and resources required for deep-sea monitoring
are not universally available to all Mediterranean Member States (MS), nor to countries in the
southern and easternmost Mediterranean. These limitations may be overcome by initiating deep
sea MSFD-focused monitoring in already data-rich locations (presumably off MS), and pioneering
joint-effort monitoring in collaboration with non-MS, to enhance awareness, capacity building,
and gain much needed data on scantily studied regions. Given the costs entailed by scientific and
technical expertise, tools and infrastructure required for deep-sea research and monitoring, we
advocate for EU-level financial support for MS/non-MS collaboration spanning joint fieldwork,
training (early career research fellowships, workshops) and *public awareness* communication.

923 Since the millennium, increased awareness of the vulnerability of deep-sea ecosystems has
924 changed attitudes concerning their protection and conservation (Ramirez-Llodra et al, 2011). Yet,
for effective regulatory measures, legislators and managers require scientific evidence, which
follows from basic scientific research and monitoring. Ecosystem-based management of the
Mediterranean deep sea pressingly requires comprehensive analysis of available data, new data
from yet unexplored regions, and impact assessment studies. Mounting evidence points to the
vulnerability of the deep biota to anthropogenic disturbance that may result in biodiversity loss,

930 urgent implementation of the MSFD in the Mediterranean deep sea will go a long way towards
931 conserving its unique biodiversity and habitats.

932

933 **Acknowledgements**

934 This study has been supported by the DG ENV project IDEM (Implementation of the MSFD
935 to the Deep Mediterranean Sea; contract EU No 11.0661/2017/750680/SUB/EN V.C2). MC and AS-
936 V from University of Barcelona acknowledge support from the Spanish government through Red
937 BAMAR (ref.: CGL2016-81854-REDT), a network on marine litter, and RTD projects NUREIEV (ref.
938 CTM2013-44598-R) and NUREIEVA (ref. CTM2016-75953-C2-1-R) on far-field and near-field
939 impacts of the Portman Bay, SE Spain, coastal submarine mine tailings disposal site. Generalitat de
940 Catalunya autonomous government funding to CRG Marine Geosciences (ref. 2017 SGR 315)
941 within its support scheme to excellence research groups is equally acknowledged.

943 **References**

- 944 [Aguzzi J., Bahamon N., 2009. Modeled day-night biases in decapod assessment by bottom trawling](#)
945 [survey. Fisheries Research 100: 274-280.](#)
- 946 [Aguzzi J., Company J.B., Bahamon N., Flexas M.M., Tecchio S., Fernandez-Arcaya U., García J.A.,](#)
947 [Mechó A., Koenig S., Canals M. 2013. Seasonal bathymetric migrations of deep-sea fishes](#)
948 [and decapod crustaceans in the NW Mediterranean Sea. Progress in Oceanography 118:](#)
949 [210-221.](#)
- 950 [Aguzzi, J., Fanelli, E., Ciuffardi, T., Doya, C., Kawato, M., Miyazaki, M., Furushima, Y., Costa, C.,](#)
951 [Fujiwhara, Y., 2018. Faunal activity rhythms influencing early community succession of an](#)
952 [implanted whale carcass offshore Sagami Bay, Japan. Scientific Report, 8, 11163.](#)
- 953 [Aguzzi, J., Chatzievangelou, D., Marini, S., Fanelli, E., Danovaro, R., Flögel, S., Lebris, N., Juanes, F.,](#)
954 [De Leo, F., Del Rio, J., Thomsen, L.S., Costa, C., Riccobene, G., Tamburini, C., Lefevre, D.,](#)
955 [Gojak, C., Poulain, P.M., Favali, P., Griffa, A., Purser, A., Cline, D., Edgington, D., Navarro J.,](#)
956 [Stefanni, S., Company, J.B., 2019. New high-tech interactive and flexible networks for the](#)
957 [future monitoring of deep-sea ecosystems. Env. Sci. Technol., 53\(12\), 6616-6631.](#)
- 958 [Allendorf, F.W., Berry, O., & Ryman, N., 2014. So long to genetic diversity, and thanks for all the](#)
959 [fish. Molecular ecology, 23\(1\), 23-25.](#)

- 960 Andradi-Brown, D. A., 2019. Invasive lionfish (*Pterois volitans* and *P. miles*): distribution, impact,
961 and management. In: Loya Y., Puglise K., Bridge T. (eds) Mesophotic Coral Ecosystems. Coral
962 Reefs of the World, vol 12. Springer, Cham, pp. 931-941.
- 963 Anastasopoulou, GA., Mytilineou, C., Smith, C., Papadopoulou, K., 2012. Plastic debris ingested by
964 deep-water fish of the Ionian Sea (Eastern Mediterranean). Deep Sea Research I, 74, 11-13.
- 965 Anastasopoulou, A., Biandolino, F., Chatzisprou, A., Hemida, F., Guijarro, B., Kousteni, V.,
966 Mytilineou, Ch., Pattoura P., Prato, E., 2016. New Fisheries-related data from the
967 Mediterranean Sea (November, 2016). Mediterranean Marine Science, 17, 822-827.
- 968 André, M., Caballé, A., Van Der Schaar, M., Solsona, A., Houégnigan, L., Zaugg, S., and ANTARES
969 consortium, 2017. Sperm whale long-range echolocation sounds revealed by ANTARES, a
970 deep-sea neutrino telescope. Scientific Reports, 7, 45517.
- 971 André, M., Solé, M., Lenoir, M., Durfort, M., Quero, C., Mas, A., Lombarte, A., van der Schaar, M.,
972 Lopez-Bejar, M., Morell, M., et al., 2011. Low-frequency sounds induce acoustic trauma in
973 cephalopods, Frontiers in Ecology and the Environment, 9, 489-493.
- 974 Angeletti, L., Bargain, A., Campiani, E., Foglini, F., Grande, V., Leidi, Mercorella, A., Prampolini, M.,
975 Taviani, M., 2019. Cold-Water Coral Habitat Mapping in the Mediterranean Sea:
976 Methodologies and Perspectives. In: Mediterranean Cold-Water Corals: Past, Present and
977 Future, C. Orejas, C. Jiménez (eds.), Coral Reefs of the World 9, Springer International
978 Publishing AG, part of Springer Nature 2019. doi.org/10.1007/978-3-319-91608-8_16
- 979 Angiolillo, M., Lorenzo, B., Farcomeni, A., Bo, M., Bavestrello, G., Santangelo, G., Cau, A.,
980 Mastascusa, V., Sacco, F., Canese, S., 2015. Distribution and assessment of marine debris in
981 the deep Tyrrhenian Sea (NW Mediterranean Sea, Italy). Marine Pollution Bulletin, 92, 149-
982 159.
- 983 Au, W.W.L., 1993. The sonar of dolphins. Springer-Verlag, New York.
- 984 Azzellino, A., Lanfredi, C., D'amico, A., Pavan, G., Podestà, M., Haun, J., 2011. Risk mapping for
985 sensitive species to underwater anthropogenic sound emissions: model development and
986 validation in two Mediterranean areas. Marine Pollution Bulletin, 63, 56-70.
- 987 Azzurro, E., Sbragaglia, V., Cerri, J., Bariche, M., Bolognini, L. Ben Souissi, J., Busoni, G., Coco, S.,
988 Chryssanthi, A., Fanelli, E., Ghanem, R., Garrabou, J., Gianni, F., Grati, F., Kolitari, J., Letterio,
989 G., Lipej, L., Mazzoldi, C., Milone, N., Pannacciulli, F., Pešić, A., Samuel-Rhoads, Y., Saponari,
990 L., Tomanic, J., Topçu, N.E., Vargiu, G., Moschella, P., 2019. Climate change, biological

1889
1890
1891 991 [invasions, and the shifting distribution of Mediterranean fishes: A large-scale survey based](#)
1892 [on local ecological knowledge. Global Change Biology, 25, 2779– 2792.](#)
1893 992
1894
1895 993 Bargain, A., Marchese, F., Savini, A., Taviani, M., Fabri, M.-C., 2017. Santa Maria di Leuca Province
1896 994 (Mediterranean Sea): Identification of suitable mounds for cold-water coral settlement using
1897
1898 995 geomorphometric proxies and Maxent methods, *Frontiers in Marine Science* 4, 338, doi:
1899 10.3389/fmars.2017.00338.
1900 996
1901 997 Bargain, A., Foglini, F., Pairaud, I., Bonaldo, D., Carniel, S., Angeletti, L., Taviani, M., Rochette, S.,
1902
1903 998 Fabri, M.-C., 2018. Predictive habitat modeling in two Mediterranean canyons including
1904
1905 999 hydrodynamics variables. *Progress in Oceanography*, 169, 151-168.
1906
1907 1000 Bertrand, J., De Sola, L., Papaconstantinou, C., Relini, G., Souplet, A., 2002. The general
1908 1001 specifications of the MEDITS surveys. *Sci. Mar.* 66, 9–17.
1909
1910 1002 Béthoux, J.P., Gentili, B., 1996. The Mediterranean Sea, coastal and deep-sea signatures of climatic
1911
1912 1003 and environmental changes. *Journal of Marine Systems*, 7, 383–394.
1913
1914 1004 Bianchelli, S., Danovaro, R., 2019. Meiofaunal biodiversity in submarine canyons of the
1915 1005 Mediterranean Sea: A meta-analysis. *Progress in Oceanography*, 170, 69-80.
1916
1917 1006 Bianchelli, S., Pusceddu, A., Buschi, E., Danovaro, R., 2016a. Trophic status and meiofauna
1918 1007 biodiversity in the Northern Adriatic Sea: Insights for the assessment of good environmental
1919
1920 1008 status. *Marine Environmental Research*, 113, 18–30.
1921
1922 1009 Bianchelli, S., Buschi, E., Danovaro, R., Pusceddu, A., 2016b. Biodiversity loss and turnover in
1923
1924 1010 alternative states in the Mediterranean Sea: a case study on meiofauna. *Scientific Reports*, 6,
1925 1011 34544.
1926
1927 1012 Bianchi, N., Morri, C., 2000. Marine biodiversity of the Mediterranean Sea: Situation, problems and
1928
1929 1013 prospects for future research. *Marine Pollution Bulletin*, 40, 367–376.
1930
1931 1014 Bo, M., Bava, S., Canese, S., Angiolillo, M., Cattaneo-Vietti, R., Bavestrello, G., 2014. Fishing impact
1932 1015 on deep Mediterranean rocky habitats as revealed by ROV investigation. *Biological*
1933
1934 1016 *Conservation*, 171, 167-176.
1935
1936 1017 Bo, M., Bavestrello, G., Angiolillo, M., Calcagnile, L., Canese, S., Cannas, R., Cau, A., D'Elia, M.,
1937 1018 D'Oriano, F., Follesa, M.C. and Quarta, G., 2015. Persistence of pristine deep-sea coral
1938
1939 1019 gardens in the Mediterranean Sea (SW Sardinia). *PloS One*, 10(3), p.e0119393.
1940
1941 1020 Bonaldo, D., Benetazzo, A., Bergamasco, A., Campiani, E., Foglini, F., Sclavo, M., Trincardi, F.,
1942 1021 Carniel, S., 2015. Interactions among Adriatic continental margin morphology, deep
1943
1944 1022 circulation and bedform patterns, *Marine Geology*, 375, 82-98.
1945
1946
1947

1948
1949
1950
1951 1023 Bonnin, J., Heussner, S., Calafat, A., Fabres, J., Palanques, A., Durrieu de Madron, X., Canals, M.,
1952 1024 Puig, P., Avril, J., Delsaut, N. (2008) Comparison of horizontal and downward particle fluxes
1953
1954 1025 across canyons of the Gulf of Lions (NW Mediterranean): Meteorological and
1955
1956 1026 hydrodynamical forcing. *Continental Shelf Research*, 28, 1957-1970.

1957 1027 Borja, A., Bricker, S.B., Dauer, D.M., Demetriades, N.T., Ferreira, J.G., Forbes, A.T., Hutchings, P.,
1958
1959 1028 Jia, X.P., Kenchington, R., Marques, J.C., et al., 2008. Overview of integrative tools and
1960
1961 1029 methods in assessing ecological integrity in estuarine and coastal systems worldwide,
1962 1030 *Marine Pollution Bulletin* 56, 9, 1519-1537.

1963
1964 1031 Boschen, R.E., Rowden, A.A., Clark, M.R., Gardner, J.P.A., 2013. Mining of deep-sea seafloor
1965
1966 1032 massive sulfides: A review of the deposits, their benthic communities, impacts from mining,
1967 1033 regulatory frameworks and management strategies, *Ocean & Coastal Management* 84,
1968
1969 1034 Supplement C, 54-67.

1970
1971 1035 Bouloubassi, I., Mejanelle, L., Pete, R., Fillaux, J., Lorre, A., Point, V., 2006. PAH transport by sinking
1972 1036 particles in the open Mediterranean Sea: A 1-year sediment trap study. *Marine Pollution*
1973
1974 1037 *Bullettin*, 52, 560-571.

1975
1976 1038 Breitburg, D., Levin, L.A., Oschlies, A., Grégoire, M., Chavez, F.P., Conley, D.J., Garçon, V., Gilbert,
1977 1039 D., Gutiérrez, D., Isensee, K., Jacinto, G.S., 2018. Declining oxygen in the global ocean and
1978
1979 1040 coastal waters. *Science*, 359(6371).

1980
1981 1041 Bricker, S.B., Longstaff, B., Dennison, W., Jones, A., Boicourt, K., Wicks, C., Woerner, J., 2008.
1982 1042 Effects of nutrient enrichment in the nation's estuaries: A decade of change, *Harmful Algae*
1983
1984 1043 8, 1, 21-32.

1985
1986 1044 Brown, C.J., Blondel, P., 2009. Developments in the application of multibeam sonar backscatter for
1987 1045 seafloor habitat mapping, *Applied Acoustics*, 70, 1242-1247.

1988
1989 1046 Brown, C.J., Smith, S.J., Lawton, P., Anderson, J.T., 2011. Benthic habitat mapping: A review of
1990
1991 1047 progress towards improved understanding of the spatial ecology of the seafloor using
1992
1993 1048 acoustic techniques, *Estuarine Coastal and Shelf Science*, 92, 502-520.

1994 1049 Canals, M., Puig, P., Durrieu de Madron, X., Heussner, S., Palanques, A., Fabres, J., 2006. Flushing
1995
1996 1050 submarine canyons. *Nature*, 444, 354-357.

1997
1998 1051 Carbery, M., O'Connor, W., Palanisami, T., 2018. Trophic transfer of microplastics and mixed
1999
2000 1052 contaminants in the marine food web and implications for human health. *Environment*
2001 1053 *International*, 15, 400-409.

2002
2003
2004
2005
2006

2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065

1054 Cardinale, M., Osio G.C., Scarcella, G., 2017. Mediterranean Sea: a failure of the European fisheries
1055 management system. *Frontiers in Marine Science*, 4, p.72,
1056 <https://doi.org/10.3389/fmars.2017.00072>.

1057 Carlier, A., Le Guilloux, E., Olu, K., Sarrazin, J., Mastrototaro, F., Taviani, M., Clavier, J., 2009.
1058 Trophic relationships in a deep Mediterranean cold-water coral bank (Santa Maria di Leuca,
1059 Ionian Sea), *Marine Ecology-Progress Series* 397, 125-137.

1060 Carroll, A.G., Przeslawski, R., Duncan, A., Gunning, M., Bruce, B., 2017. A critical review of the
1061 potential impacts of marine seismic surveys on fish & invertebrates, *Marine Pollution*
1062 *Bulletin* 114, 9-24.

1063 Cartes, J.E., Lolocono, C., Mamouridis, V., López-Pérez, C., Rodríguez, P., 2013. Geomorphological,
1064 trophic and human influences on the bamboo coral *Isidella elongata* assemblages in the
1065 deep Mediterranean: To what extent does *Isidella* form habitat for fish and invertebrates?
1066 *Deep Sea Research Part I*, 76, 52-65.

1067 Cartes, J.E., Fanelli, E., Kaporis, K., Bayhan, Y.K., Ligas, A., López-Pérez, C., Murenu, M., Papiol, V.,
1068 Rumolo, P., Scarcella, G., 2014. Spatial variability in the trophic ecology and biology of the
1069 deep-sea shrimp *Aristaeomorpha foliacea* in the Mediterranean Sea, *Deep Sea Research Part*
1070 *I*, 87, Supplement C, 1-13.

1071 Caruso, F., Sciacca, V., Bellia, G., De Domenico, E., Larosa, G., Papale, E., Pellegrino, C., Pulvirenti,
1072 S., Riccobene, G., Simeone, F., and KM3NET Consortium, 2015. Size Distribution of Sperm
1073 Whales Acoustically Identified during Long Term Deep-Sea Monitoring in the Ionian Sea, *Plos*
1074 *One* 10, 12.

1075 Chimienti, G., Bo, M., Taviani, M., Mastrototaro, F., 2019. Occurrence and biogeography of cold
1076 water coral communities in Mediterranean hard and soft bottoms. In: *Mediterranean Cold-*
1077 *Water Corals: Past, Present and Future*, C. Orejas, C. Jiménez (eds.), *Coral Reefs of the World*
1078 9, Springer International Publishing AG, part of Springer Nature 2019.

1079 Chouvelon, T., Cresson, P., Bouchouca, M., Brach-Papa, C., Bustamante, P., Crochet, S., Marco-
1080 Miralles, F., Thomas, B., Knoery, J., 2018. Oligotrophy as a major driver of mercury
1081 bioaccumulation in medium-to high-trophic level consumers: A marine ecosystem-
1082 comparative study, *Environmental Pollution* 233, Supplement C, 844-854.

1083 Claussen, U., Zevenboom, W., Brockmann, U., Topcu, D., Bot, P., 2009. Assessment of the
1084 eutrophication status of transitional, coastal and marine waters within OSPAR, *Hydrobiologia*
1085 629, 1, 49-58, doi: 10.1007/s10750-009-9763-3.

2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124

1086 Coll, M., Piroddi, C., Kaschner, K., Ben Rais Lasram, F., Steenbeek, J., et al., 2010. The biodiversity
1087 of the Mediterranean Sea: Status, patterns and threats. PloS ONE. 5 (8): e11842.
1088 doi:10.1371/journal.pone.0011842.

1089 Colloca, F., Cardinale, M., Maynou, F., Giannoulaki, M., Scarcella, G., Jenko, K., Bellido, J.M.,
1090 Fiorentino, F. (2013). Rebuilding Mediterranean fisheries: a new paradigm for ecological
1091 sustainability. Fish and Fisheries, 14, 89–109. DOI: 10.1111/j.1467-2979.2011.00453.x

1092 Company, J.B., Puig, P., Sarda, F., Palanques, A., Latasa, M., Scharek, R., 2008. Climate Influence on
1093 deep sea populations, Plos One 3, 1, e1431, doi: 10.1371/journal.pone.0001431.

1094 Conese, I., Fanelli, E., Miserocchi, S., Langone, L., 2019. Food web structure and trophodynamics of
1095 deep-sea plankton from the Bari Canyon and adjacent slope (Southern Adriatic, central
1096 Mediterranean Sea). *Progress in Oceanography* 175, 92–104

1097 Conversi, A., Fonda-Umani, S., Peluso, T., Molinero, J.C., Santojanni, A., Edwards, M., 2010. The
1098 Mediterranean Sea Regime Shift at the End of the 1980s, and Intriguing Parallelisms with
1099 Other European Basins, Plos One 5, 5, doi: 10.1371/journal.pone.0010633.

1100 Coro, G., Vilas, L. G., Magliozzi, C., Ellenbroek, A., Scarponi, P., & Pagano, P. (2018). Forecasting the
1101 ongoing invasion of *Lagocephalus sceleratus* in the Mediterranean Sea. *Ecological Modelling*,
1102 371, 37-49.

1103 Corsini-Foka M., Pancucci-Papadopoulou A., Kondilatos G., Kalogirou, S., 2010. *Gonioinfradens*
1104 *paucidentatus* (A. Milne Edwards, 1861) (Crustacea, Decapoda, Portunidae): a new alien crab
1105 in the Mediterranean Sea. *Mediterranean Marine Science*, 11, 331–340

1106 Cossa, D., Coquery, M., 2005. The Mediterranean mercury anomaly, a geochemical or a biological
1107 issue. In: Saliot, A. (ed.) *The Mediterranean Sea - The Handbook of Environmental Chemistry*,
1108 n° 5, Part K, pp. 177-208, Springer-Verlag, Berlin.

1109 Cossa, D., Harmelin-Vivien, M., Mellon-Duval, C., Loizeau, V., Averty, B., Crochet, S., Chou, L.,
1110 Cadiou, J.F., 2012. Influences of bioavailability, trophic position, and growth on
1111 Methylmercury in Hakes (*Merluccius merluccius*) from Northwestern Mediterranean and
1112 Northeastern Atlantic. *Environmental Science & Technology*, 46, 4885-4893.

1113 Cresson, P., Fabri, M.C., Bouchoucha, M., Brach Papa, C., Chavanon, F., Jadaud, A., Knoery, J.,
1114 Miralles, F., Cossa, D., 2014. Mercury in organisms from the Northwestern Mediterranean
1115 slope: Importance of food sources. *Science of the Total Environment*, 497-498, 229-23,.

1116 D'Onghia, G., Calculli, C., Capezzuto, F., Carlucci, R., Carluccio, A., Grehan, A., Indennidate, A.,
1117 Maiorano, P., Mastrototaro, F., Pollice, A., Russo T., 2017. Anthropogenic impact in the Santa

2125
2126
2127 1118 Maria di Leuca cold-water coral province (Mediterranean Sea): Observations and
2128 conservation straits. Deep Sea Research, Part II, Topical Studies in Oceanography, 145, 87-
2129 101.
2130
2131 1120
2132 1121 Dachs, J., Bayona, J.M., Fowler, S.W., Miquel, J.-C., Albaigés, J., 1996. Vertical fluxes of polycyclic
2133 aromatic hydrocarbons and organochlorine compounds in the western Alboran Sea
2134 (southwestern Mediterranean). Marine Chemistry, 52, 75-86.
2135
2136 1123
2137 1124 Damiano, S., Papetti, P., Menesatti, P., 2011. Accumulation of heavy metals to assess the health
2138 status of swordfish in a comparative analysis of Mediterranean and Atlantic areas, Marine
2139 Pollution Bulletin 62, 1920-1925.
2140
2141 1126
2142 1127 Danovaro, R., Pusceddu, A. 2007. Ecomanagement of biodiversity and ecosystem functioning in
2143 the Mediterranean Sea: concerns and strategies. Chemistry and Ecology, 23, 347-360.
2144
2145 1128
2146 1129 Danovaro, R., Dinet, A., Duineveld, G., Tselepidis, A., 1999. Benthic response to particulate fluxes
2147 in different trophic environments: a comparison between the Gulf of Lions-Catalan Sea
2148 (Western Mediterranean) and the Cretan Sea (Eastern-Mediterranean). Progress in
2149 Oceanography, 44, 287-312.
2150
2151 1132
2152 1133 Danovaro, R., 2003. Pollution threats in the Mediterranean Sea: an overview. Chemistry and
2153 Ecology 19 (1), 15-32.
2154
2155 1134
2156 1135 Danovaro, R., Gambi, C., Dell'Anno, A., Corinaldesi, C., Fraschetti, S., Vanreusel, A., Vincx, M. and
2157 Gooday, A.J., 2008. Exponential decline of deep-sea ecosystem functioning linked to benthic
2158 biodiversity loss. Current Biology, 18, 1-8.
2159
2160 1137
2161 1138 Danovaro, R., Corinaldesi, C., D'Onghia, G., Galil, B., Gambi, C., Gooday, A.J., Lampadariou, N.,
2162 Luna, G.M., Morigi, C., Olu, K., Polymenakou, P., 2010. Deep-sea biodiversity in the
2163 Mediterranean Sea: The known, the unknown, and the unknowable. PloS One, 5(8), p.e
2164 11832.
2165
2166 1141
2167 1142 Danovaro, R., Snelgrove, P.V.R., Tyler, P., 2014. Challenging the paradigms of deep-sea ecology.
2168 Trends in Ecology & Evolution 29, 8, 465-475.
2169
2170 1143
2171 1144 Danovaro, R., Corinaldesi, C., Rastelli, E., Dell'Anno, A., 2015. Towards a better quantitative
2172 assessment of the relevance of deep-sea viruses, Bacteria and Archaea in the functioning
2173 of the ocean seafloor. Aquat. Microb. Ecol. 75: 81-90. <https://doi.org/10.3354/ame01747>
2174
2175 1146
2176 1147 Danovaro, R., Carugati, L., Boldrin, A., Calafat, A., Canals, M., Fabres, J., Finlay, K., Heussner, S.,
2177 Miserocchi, S., Sanchez-Vidal, A., 2017. Deep-water zooplankton in the Mediterranean Sea:

2184
2185
2186
2187 1149 results from a continuous, synchronous sampling over different regions using sediment
2188 1150 traps. Deep Sea Research, Part I, 103, 103-114.
2189
2190 1151 Danovaro, R., Fanelli, E., Aguzzi, J., Billett, D., Carugati, L., Corinaldesi, C., Dell'Anno, A., Gjerde, K.,
2191 1152 Jamieson, A.J., Kark, S., McClain, C., Levin, L., Levin, N., Ramirez-Llodra, E., Ruhl, H., Smith,
2192 1153 C.R., Snelgrove, P.V.R., Thomsen, L., Van Dover, C., Yasuhara, M., 2019. Ecological variables
2193 1154 for developing a global deep-ocean monitoring and conservation strategy. Nature Ecol. Evol.,
2194
2195 1155 in press.
2196
2197 1156 Dauvin, J.C., 2010. Towards an impact assessment of bauxite red mud waste on the knowledge of
2198
2199 the structure and functions of bathyal ecosystems: The example of the Cassidaigne canyon
2200 1157 (north-western Mediterranean Sea), Marine Pollution Bulletin 60, 197-206.
2201
2202 1158 de Vivero J.L.S., Rodriguez Mateos J C, 2015. Marine Governance in the Mediterranean Sea. In
2203 1159 Gilek M., Kern K. (eds.), Governing Europe's Marine Environment: Europeanization of
2204
2205 1160 Regional Seas or Regionalization of EU policies? 2015. Published by Ashgate Publishing.
2206
2207 1161 Dell'Anno, A., Mei, M.L., Pusceddu, A., Danovaro, R., 2002. Assessing the trophic state and
2208 1162 eutrophication of coastal marine systems: a new approach based on the biochemical
2209
2210 1163 composition of sediment organic matter. Marine Pollution Bulletin 44, 611-622.
2211
2212 1164 Denda, A., Christiansen, B., 2014. Zooplankton distribution patterns at two seamounts in the
2213 1165 subtropical and tropical NE Atlantic. Marine ecology, 35(2), 159-179.
2214
2215 1166 DeRuiter, S.L., Larbi Doukara, K., 2012. Loggerhead turtles dive in response to airgun sound
2216
2217 1167 exposure, Endangered Species Research 16, 55-63.
2218
2219 1168 Desforges J.P.W., Galbraith M., Dangerfield N., Ross P.S., 2014. Widespread distribution of
2220 1169 microplastics in subsurface seawater in the NE Pacific Ocean Marine Pollution Bulletin, 79:
2221
2222 1170 94-99.
2223
2224 1171 Doya, C., Aguzzi, J., Chatzievangelou, D., Costa, C., Company, J.B., Tunnicliffe, V., 2015. The
2225 1172 seasonal use of small-scale space by benthic species in a transiently hypoxic area. Journal of
2226
2227 1173 Marine Systems, 154, 280-290.
2228
2229 1174 Drazen, J.C., Haedrich, R.L., 2012. A continuum of life histories in deep-sea demersal fishes, Deep-
2230 1175 Sea Research, Part I , 61, 34-42.
2231
2232 1176 Durrieu de Madron, X., Houpert, L., Puig, P., Sanchez-Vidal, A., Testor, P., Bosse, A., Estournel,
2233 1177 C., Somot, S., Bourrin, F., Bouin, M. N., Beauverger, M., Beguery, L., Calafat, A., Canals,
2234
2235 1178 M., Cassou, C., Coppola, L., Dausse, D., D'Ortenzio, F.; Font, J., Heussner, S., Kunesch,
2236
2237 1179 S., Lefevre, D., Le Goff, H., Martín, J., Mortier, L., Palanques, A., Raimbault, P., 2013.
2238
2239 1180

2243
2244
2245 1181 Interaction of dense shelf water cascading and open-sea convection in the northwestern
2246 Mediterranean during winter 2012. *Geophysical Research Letters*, American Geophysical
2247 1182 Union, 2013, 40, 1379-1385.
2248
2249 1183
2250 1184 Durrieu de Madron, X., Guieu, C., R. Sempéré, P. Conan, D. Cossa, F. D'Ortenzio, C. Estournel, F.
2251 Gazeau, C. Rabouille, L. Stemmann, S. Bonnet, F. Diaz, P. Koubbi, O. Radakovitch, M. Babin,
2252 1185 M. Baklouti, C. Bancon-Montigny, S. Belviso, N. Bensoussan, B. Bonsang, I. Bouloubassi, C.
2253 Brunet, J.-F. Cadiou, F. Carlotti, M. Chami, S. Charmasson, B. Charrière, J. Dachs, D. Doxaran,
2254 1186 J.-C. Dutay, F. Elbaz-Poulichet, M. Eléaume, F. Eyrolles, C. Fernandez, S. Fowler, P. Francour,
2255 J.C. Gaertner, R. Galzin, S. Gasparini, J.-F. Ghiglione, J.-L. Gonzalez, C. Goyet, L. Guidi, K.
2256 1187 Guizien, L.-E. Heimbürger, S.H.M. Jacquet, W.H. Jeffrey, F. Joux, P. Le Hir, K. Leblanc, D.
2257 1188 Lefèvre, C. Lejeusne, R. Lemé, M.-D. Loÿe-Pilot, M. Mallet, L. Méjanelle, F. Mélin, C. Mellon,
2258 B. Mérigot, P.-L. Merle, C. Migon, W.L. Miller, L. Mortier, B. Mostajir, L. Mousseau, T.
2259 1189 Moutin, J. Para, T. Pérez, A. Petrenko, J.-C. Poggiale, L. Prieur, M. Pujo-Pay, Pulido-Villena, P.
2260 Raimbault, A.P. Rees, C. Ridame, J.-F. Rontani, D. Ruiz Pino, M.A. Sicre, V. Taillandier, C.
2261 1190 Tamburini, T. Tanaka, I. Taupier-Letage, M. Tedetti, P. Testor, H. Thébault, B. Thouvenin, F.
2262 1191 Touratier, J. Tronczynski, C. Ulses, F. Van Wambeke, V. Vantrepotte, S. Vaz, R. Verney, 2011.
2263 Marine ecosystems' responses to climatic and anthropogenic forcings in the Mediterranean,
2264 1192 *Progress in Oceanography*, 91, 97-166.
2265
2266 1193
2267 1194 EC 2006. No 1881/2006 of 19 December 2006 setting maximum levels for certain contaminants in
2268 foodstuffs. *Official Journal of the European Union*, L364/5-24.
2269 1195
2270 EC 2008. DIRECTIVE 2008/105/EC OF THE EUROPEAN PARLIAMENT AND OF THE COUNCIL of 16
2271 1196 December 2008 on environmental quality standards in the field of water policy, amending
2272 and subsequently repealing Council Directives 82/176/EEC, 83/513/EEC, 84/156/EEC,
2273 1197 84/491/EEC, 86/280/EEC and amending Directive 2000/60/EC of the European Parliament
2274 1198 and of the Council. *Official Journal of the European Union*, L348/84-97.
2275
2276 1199 EC 2017. COMMISSION DECISION (EU) 2017/848 of 17 May 2017 laying down criteria and
2277 methodological standards on good environmental status of marine waters and specifications
2278 1200 and standardised methods for monitoring and assessment, and repealing Decision
2279 1201 2010/477/EU. *Official Journal of the European Union*, L125/43-74.
2280
2281 1202
2282
2283 1203
2284 1204
2285 1205
2286 1206
2287
2288 1207
2289
2290 1208
2291
2292 1209
2293
2294 1210 Edds-Walton, P.L., 1997. Acoustic communication signals of mysticete whales, *Bioacoustics*, 8, 47-
2295 60.
2296 1211
2297
2298
2299
2300
2301

2302
2303
2304
2305 1212 EEA, 1999. State and pressures of the marine and coastal Mediterranean environment.
2306 1213 Environmental Issue series, No5, European Commission, 137 pp.
2307
2308 1214 Eigaard, O.R., Bastardie, F., Hintzen, N.T., Buhl-Mortensen, L., Buhl-Mortensen, P., Catarino, R.,
2309 1215 Dinesen, G.E., Egekvist, J., Fock, H.O., Geitner, K., Gerritsen, H.D., González, M.M., Jonsson,
2310 1216 P., Kavadas, S., Laffargue, P., Lundy, M., Gonzalez-Mirelis, G., Nielsen, J.R., Papadopoulou, N.,
2311 1217 E. Posen, P.E., Pulcinella, J., Russo, T., Sala, A., Silva, C., Smith, C.J., Vanelslander, B.,
2312 1218 Rijnsdorp, A.D., 2017. The footprint of bottom trawling in European waters: distribution,
2313 1219 intensity, and seabed integrity, Ices Journal of Marine Science 74, 847-865.
2314
2315 1220 Elahi, R., Ferretti, F., Bastari, A., Cerrano, C., Colloca F., Kowalik, J., Ruckelshaus, M., Struck, A.,
2316 1221 Micheli, F., 2018. Leveraging vessel traffic data and a temporary fishing closure to inform
2317 1222 marine management. *Front. Ecol. Environ.*, 16(8), 1-7.
2318
2319 1223 Emig, C.C., Geistdoerfer, P., 2004. The Mediterranean deep-sea fauna: historical evolution,
2320 1224 bathymetric variations and geographical changes. *Carnets de Geologie, Madrid*, 4(A01), 10 p.
2321 1225 doi: 10.4267/2042/3230
2322
2323 1226 Estournel, C., Zervakis, V., Marsaleix, P., Papadopoulos, A., Auclair, F., Perivoliotis, L., Tragou, E.,
2324 1227 2005. Dense water formation and cascading in the Gulf of Thermaikos (North Aegean), from
2325 1228 observations and modelling. *Cont. Shelf Res.* 25, 2366-2386.
2326
2327 1229 Evans, J., Aguilar, R., Alvarez, H., Borg, J.A., Garcia, S., Knittweis, L., Schembri, P.J., 2016. Recent
2328 1230 evidence that the deep sea around Malta is a biodiversity hotspot. 41st Congress of the
2329 1231 Mediterranean Science Commission (CIESM), Kiel, Germany, 12 - 16 September 2016
2330
2331 1232 Fabri, M.C., Bargain, A., Pairaud, I., Pedel, L., Taupier-Letage, I., 2017. Cold-water coral ecosystems
2332 1233 in Cassidaigne Canyon: An assessment of their environmental living conditions, *Deep Sea*
2333 1234 *Research, Part II*, 137, 436-453.
2334
2335 1235 Fabri, M.C., Pedel, L., Beuck, L., Galgani, F., Hebbeln, D., Freiwald, A., 2014. Megafauna of
2336 1236 vulnerable marine ecosystems in French Mediterranean submarine canyons: Spatial
2337 1237 distribution and anthropogenic impacts, *Deep Sea Research, Part I*, 104, 184-207.
2338
2339 1238 Fabri, M.C., Vinha, B., Allais, A.G., Bouhier, M.-E., Dugornay, O., Gaillot, A., & Arnaubec, A. (under
2340 1239 revision). Evaluating ecological status of cold-water coral habitats using non-invasive
2341 1240 methods, an example from Cassidaigne canyon, northwestern Mediterranean Sea. *Progress*
2342 1241 *in Oceanography in press*
2343
2344 1242 Fabri, M.C., Brind'Amour, A., Jadaud, A., Galgani, F., Vaz, S., Taviani, M., Scarcella, G., Canals, M.,
2345 1243 Sanchez, A., Grimalt, J., Galil, B., Goren, M., Schembri, P., Evans, J., Knittweis, Leyla,
2346
2347
2348
2349
2350
2351
2352
2353
2354
2355
2356
2357
2358
2359
2360

2361
2362
2363
2364
2365
2366
2367
2368
2369
2370
2371
2372
2373
2374
2375
2376
2377
2378
2379
2380
2381
2382
2383
2384
2385
2386
2387
2388
2389
2390
2391
2392
2393
2394
2395
2396
2397
2398
2399
2400
2401
2402
2403
2404
2405
2406
2407
2408
2409
2410
2411
2412
2413
2414
2415
2416
2417
2418
2419

Cantafaro, A.-L., Fanelli, E., Carugati, L., Danovaro, R., 2018. Review of literature on the implementation of the MSFD to the deep Mediterranean Sea.IDEM project, Deliverable 1.1. 228 p. www.msfd-idem.eu. doi.org/10.13155/53809

Fanelli, E., Cartes, J.E., Rumolo, P., Sprovieri, M., 2009. Food-web structure and trophodynamics of mesopelagic-suprabenthic bathyal macrofauna of the Algerian Basin based on stable isotopes of carbon and nitrogen, Deep Sea Research, Part I, 56, 1504-1520.

Fanelli, E., Badalamenti, F., D'Anna, G., Pipitone, P., Romano C., 2010. Trophodynamic effects of trawling on the feeding ecology of Pandora, *Pagellus erythrinus* off the northern Sicily coast (Mediterranean Sea). Marine and Freshwater Research 61, 408–417.

Fanelli, E., Papiol, V., Cartes, J.E., Rumolo, P., Brunet, C., Sprovieri, M., 2011a. Food web structure of the megabenthic, invertebrate epifauna on the Catalan slope (NW Mediterranean): evidence from $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ analysis. Deep Sea Research, Part I, 58, 98-109.

Fanelli, E., Cartes, J.E., Papiol, V., 2011b. Food web structure of deep-sea macrozooplankton and micronekton off the Catalan slope: Insight from stable isotopes, Journal of Marine Systems, 87, 79-89.

Fanelli, E., Cartes, J.E. Papiol, V., 2012. Assemblage structure and trophic ecology of deep-sea demersal cephalopods in the Balearic basin (NW Mediterranean). Marine and Freshwater Research, 63, 264-274.

Fanelli, E., Cartes, J. E., Papiol, V., López-Pérez, C., 2013. Environmental drivers of megafaunal assemblage composition and biomass distribution over mainland and insular slopes of the Balearic Basin (Western Mediterranean). Deep Sea Research, Part I, 78, 79-94.

Fanelli, E., Papiol, V., Cartes, J.E., Rodriguez-Romeu, O., 2014. Trophic ecology of *Lampanyctus crocodilus* on north-west Mediterranean Sea slopes in relation to reproductive cycle and environmental variables, Journal of Fish Biology 84, 1654-1688.

Fanelli, E., Azzurro, E., Bariche, M., Cartes, J.E., Maynou, F., 2015. Depicting the novel Eastern Mediterranean food web: a stable isotopes study following Lessepsian fish invasion, Biological Invasions 17, 2163-2178.

Fanelli, E., Cartes, J.E., Papiol, V., López-Pérez, C., Carrasson, M., 2016. Long-term decline in the trophic level of megafauna in the deep Mediterranean Sea: a stable isotopes approach, Climate Research, 67, 191-207.

2420
2421
2422
2423 1274 Fanelli, E., Delbono, I., Ivaldi, R., Pratellesi, M., Cocito, S., Peirano, A., 2017. Cold water coral
2424 1275 *Madrepora oculata* in the eastern Ligurian Sea (NW Mediterranean): historical banks and
2425 recent findings. Aquatic Conservation: Marine Freshwater Ecosystems, 27, 965-975.
2426 1276
2427 1277 Fanelli E., Bianchelli S., Danovaro R., 2018. Deep-sea mobile megafauna of Mediterranean
2428 submarine canyons and open slopes: analysis of spatial and bathymetric gradients. Progress
2429 1278 in Oceanography, 168, 23-24.
2430
2431 1279
2432 1280 FAO, 2009. International Guidelines for the Management of Deep-sea Fisheries in the High Seas.
2433 FAO, Rome, Italy, 73pp.
2434 1281
2435 1282 FAO, 2016. Report of the FAO Workshop on Deep-sea Fisheries and Vulnerable Marine Ecosystems
2436 of the Mediterranean, Rome, Italy, 18–20 July 2016. FAO Fisheries and Aquaculture Report
2437 1283 No. 1183, Rome, Italy.
2438
2439 1284
2440 1285 FAO, 2018. The State of Mediterranean and Black Sea Fisheries. General Fisheries Commission for
2441 the Mediterranean. Rome, 172pp
2442 1286
2443 1287 Favali, P., Chierici, F., Marinaro, G., Giovanetti, G., Azzarone, A., Beranzoli, L., and KMNET
2444 consortium, 2013. NEMO-SN1 abyssal cabled observatory in the Western Ionian Sea. IEEE
2445 Journal of Oceanic Engineering, 38, 358-374.
2446 1288
2447 1289
2448 1290 Fernandez-Arcaya, U., Ramirez-Llodra E., Aguzzi J., Allcock A.L., Davies J.S., Dissanayake A., Harris
2449 P., Howell K., Huvenne, V.A.I., Ismail K., Macmillan-Lawler M., Martín J., Menot L., Nizinski
2450 M., Puig P., Rowden A., Sanchez F., Steward H.A., Van den Beld I. 2017. Ecological role of
2451 1291 submarine canyons and need for canyon conservation: a review. Front. Mar. Sci. 31.
2452 1292
2453 1293
2454 1294 Ferreira J.G., Andersen J.H., Borja A., Bricker S.B., Camp J., Cardoso da Silva M., Garcés E.,
2455 Heiskanen A.S., Humborg C., Ignatiades L., Lancelot C., Menesguen A., Tett P., Hoepffner N.,
2456 1295 Claussen U., 2011. Overview of eutrophication indicators to assess environmental status
2457 within the European Marine Strategy Framework Directive. Estuarine, Coastal and Shelf
2458 1296 Science, 93, 117-131.
2459 1297
2460 1298
2461 1299 Ferreira, J.G., Andersen, J.H., Borja, A., Bricker, S.B., Camp, J., Cardoso da Silva, M., Garcés, E.,
2462 Heiskanen, A.S., Humborg, C., Ignatiades, L., Lancelot, C., Menesguen, A., Tett, P., Hoepffner,
2463 1300 N., Claussen, U., 2010. Marine Strategy Framework Directive Task Group 5 Report
2464 Eutrophication. EUR 24338
2465 1301
2466 1302
2467 1303 Ferreira, J.G., Vale, C., Soares, C.V., Salas, F., Stacey, P.E., Bricker, S.B., Silva, M.C., Marques, J.C.,
2468 2007. Monitoring of coastal and transitional waters under the EU water framework directive,
2469 Environmental Monitoring and Assessment, 135, 195-216.
2470 1304
2471 1305
2472
2473
2474
2475
2476
2477
2478

2479
2480
2481 1306 Fichaut, M., Garcia, M.J., Giorgetti, A., Iona, A., Kuznetsov, A., Rixen, M., Group, M., 2003.
2482
2483 1307 MEDAR/MEDATLAS 2002: A Mediterranean and Black Sea database for operational
2484
2485 1308 oceanography. In: Dahlin, H., Flemming, N.C., Nittis, K., Petersson, S.E. (eds.), Elsevier
2486
2487 1309 Oceanography Series, n° 69, pp. 645-648, Elsevier, doi: [https://doi.org/10.1016/S0422-](https://doi.org/10.1016/S0422-9894(03)80107-1)
2488 1310 [9894\(03\)80107-1](https://doi.org/10.1016/S0422-9894(03)80107-1). 0422-9894
2489
2490 1311 Fliedner, A., Rüdell, H., Knopf, B., Lohmann, N., Paulus, M., Jud, M., Pirntke, U., Koschorreck, J.
2491
2492 1312 (2018). Assessment of seafood contamination under the marine strategy framework
2493 1313 directive: contributions of the German environmental specimen bank. Environmental
2494
2495 1314 Science and Pollution Research, 25(27), 26939-26956.
2496
2497 1315 Fontanier, C., Fabri, M.C., Buscail, R., Biscara, L., Koho, K., Reichart, G.J., Cossa, D., Galaup, S.,
2498 1316 Chabaud, G., Pigot, L. (2012) Deep-sea foraminifera from the Cassidaigne Canyon (NW
2499
2500 1317 Mediterranean): Assessing the environmental impact of bauxite red mud disposal. Marine
2501
2502 1318 Pollution Bulletin, 64, 1895-1910. doi : 10.1016/j.marpolbul.2012.06.016
2503 1319 Fontanier, C., Biscara, L., Mamo, B., Delord, E., 2014. Deep-sea benthic foraminifera in an area
2504
2505 1320 around the Cassidaigne Canyon (NW Mediterranean) affected by bauxite discharges, Marine
2506
2507 1321 Biodiversity 44, 4, doi: 10.1007/s12526-014-0281-9.
2508 1322 Foote, A.D., Osborne, R.W., Hoelzel, A.R., 2004. Environment - Whale-call response to masking
2509
2510 1323 boat noise, Nature 428, 6986, 910-910, doi: 10.1038/428910a.
2511
2512 1324 Freiwald, A., Beuck, L., Rüggeberg, A., Taviani, M., Hebbeln, D., R/V Meteor M70-1 Participants,
2513 1325 2009. The white coral community in the central Mediterranean Sea revealed by ROV surveys.
2514
2515 1326 Oceanography, 22(1), 58-74.
2516
2517 1327 Galgani, F., Leaute, J.P., Moguedet, P., Souplet, A., Verin, Y., Carpentier, A., Goraguer, H., Latrouite,
2518 1328 D., Andral, B., Cadiou, Y., Mahe, J.C., Poulard, J.C., Nerisson, P., 2000. Litter on the sea floor
2519
2520 1329 along European coasts. Marine Pollution Bulletin 40 (6), 516-527.
2521
2522 1330 Galil, B. S. (2004). The limit of the sea: the bathyal fauna of the Levantine Sea. Scientia Marina,
2523
2524 1331 68(S3), 63-72.
2525 1332 Galil, B.S., 2007. Loss or gain? Invasive aliens and biodiversity in the Mediterranean Sea, Marine
2526
2527 1333 Pollution Bulletin 55, 7-9, 314-322, doi: 10.1016/j.marpolbul.2006.11.008.
2528
2529 1334 Galil, B.S., Danovaro, R., Rothman, S.B.S., Gevili, R. and Goren, M., 2019. Invasive biota in the
2530 1335 deep-sea Mediterranean: an emerging issue in marine conservation and management.
2531
2532 1336 Biological invasions, 21(2), 281-288.
2533
2534 1337 Galil, B., and Herut, B. (2011). Marine environmental issues of deep-sea oil and gas exploration

2538
2539
2540
2541
2542
2543
2544
2545
2546
2547
2548
2549
2550
2551
2552
2553
2554
2555
2556
2557
2558
2559
2560
2561
2562
2563
2564
2565
2566
2567
2568
2569
2570
2571
2572
2573
2574
2575
2576
2577
2578
2579
2580
2581
2582
2583
2584
2585
2586
2587
2588
2589
2590
2591
2592
2593
2594
2595
2596

and exploitation activities off the coast of Israel. IOLR Report H15/2011, 24 p.

Galil, B.S., Golik, A., Türkay, M., 1995. Litter at the bottom of the sea: A sea bed survey in the Eastern Mediterranean, *Marine Pollution Bulletin* 30, 1, 22-24, doi: [https://doi.org/10.1016/0025-326X\(94\)00103-G](https://doi.org/10.1016/0025-326X(94)00103-G).

Galil, B.S., Marchini, A., Occhipinti-Ambrogi, A., 2016. East is east and West is west? Management of marine bioinvasions in the Mediterranean Sea, *Estuarine, Coastal and Shelf Science*, doi: <http://dx.doi.org/10.1016/j.ecss.2015.12.021>

Galil, B.S., Marchini, A., Occhipinti-Ambrogi, A., Minchin, D., Narscius, A., Ojaveer, H., Olenin, S., 2014. International arrivals: widespread bioinvasions in European Seas, *Ethology Ecology & Evolution* 26, 2-3, 152-171, doi: 10.1080/03949370.2014.897651.

Galil, B., Marchini, A., Occhipinti-Ambrogi, A., Ojaveer, H., 2017. The enlargement of the Suez Canal - Erythraean introductions and management challenges, *Management of Biological Invasions* 8, 2, 141-152, doi: 10.3391/mbi.2017.8.2.02.

Galil, B.S., Danovaro, R., Rothman, S.B.S., Gevili, R., Goren, M., 2018. Invasive biota in the deep-sea Mediterranean: an emerging issue in marine conservation and management. *Biological Invasions*, 21(2): 281-288

Gambi, C., Pusceddu, A., Benedetti-Cecchi, L., & Danovaro, R. (2014). Species richness, species turnover and functional diversity in nematodes of the deep Mediterranean Sea: searching for drivers at different spatial scales. *Global ecology and biogeography*, 23(1), 24-39.

Gascuel, D., Coll, M., Fox, C., Guénette, S., Guitton, J., Kenny, A., Knittweis, L., Nielsen, J. R., Piet, G., Raid, T., Travers-Trolet, M., and Shephard, S., 2016. Fishing impact and environmental status in European seas: a diagnosis from stock assessments and ecosystem indicators, *Fish and Fisheries*, 17, 31-55.

GFCM, 2017. Report of the first meeting of the Working Group on Vulnerable Marine Ecosystems (WGVME). Malaga, Spain, 3-5 April 2017

Giusti, M., Innocenti, C., & Canese, S. (2014). Predicting suitable habitat for the gold coral *Savalia savaglia* (Bertoloni, 1819) (Cnidaria, Zoantharia) in the South Tyrrhenian Sea. *Continental Shelf Research*, 81, 19-28.10.1016/j.csr.2014.03.011

Giusti, M., Bo, M., Angiolillo, M., Cannas, R., Cau, A., Follesa, M.C., Canese, S. (2017). Habitat preference of *Viminella flagellum* (Alcyonacea: Ellisellidae) in relation to bathymetric variables in southeastern Sardinian waters. *Continental Shelf Research*, 138, 41-50.10.1016/j.csr.2017.03.004

2597
2598
2599 1370 Goren, M., Galil, B.S., Diamant, A., Stern, N., Levitt-Barmatz, Y., 2016. Invading up the food web?
2600
2601 1371 Invasive fish in the southeastern Mediterranean Sea, *Marine Biology*, 163, 8, doi:
2602
2603 1372 10.1007/s00227-016-2950-7.
2604
2605 1373 Gregory, M., 2009. Environmental implications of plastic debris in marine settings entanglement,
2606 1374 ingestion, smothering, hangers-on, hitch-hiking and alien invasions. *Philos Trans R Soc Lond*
2607
2608 1375 *B Biol Sci.*, 364 (1526), 2013–2025.
2609
2610 1376 Gress, E., Andradi-Brown, D.A., Woodall, L., Schofield, P.J., Stanley, K., Rogers, A.D., 2017. Lionfish
2611 1377 (*Pterois* spp.) invade the upper-bathyal zone in the western Atlantic, *Peerj* 5, doi:
2612
2613 1378 10.7717/peerj.3683.
2614
2615 1379 Guerzoni, S., Chester, R., Dulac, F., Herut, B., Loye-Pilot, M.D., Measures, C., Migon, C., Molinaroli,
2616 1380 E., Moulin, C., Rossini, P., Saydam, C., Soudine, A., Ziveri, P., 1999. The role of atmospheric
2617
2618 1381 deposition in the biogeochemistry of the Mediterranean Sea, *Progress in Oceanography* 44,
2619
2620 1382 1-3, 147-190.
2621
2622 1383 Keeling, R.F., Kortzinger, A., Gruber, N., 2010. Ocean Deoxygenation in a Warming World, *Annual*
2623 1384 *Review of Marine Science* 2, 199-229, doi: 10.1146/annurev.marine.010908.163855.
2624
2625 1385 Halpern BS, Walbridge S, Selkoe KA, Kappel CV, Micheli F, D'Agrosa C, Bruno JF, Casey KS, Ebert C,
2626 1386 Fox HE, Fujita R, Heinemann D, Lenihan HS, Madin EMP, Perry MT, Selig ER, Spalding M,
2627
2628 1387 Steneck R, Watson R (2008). A global map of human impact on marine ecosystems, *Science*
2629
2630 1388 319, 5865, 948-952.
2631
2632 1389 Harmelin-Vivien M., Bodiguel X., Charmasson S., Loizeau V., Mellon-Duval C., Tronczynski J., Cossa
2633 1390 D., 2012. Differential biomagnification of PCB, PBDE, Hg and Radiocesium in the food web of
2634
2635 1391 the European hake from the NW Mediterranean, *Marine Pollution Bulletin* 64, 5, 974-983.
2636
2637 1392 Harmon, S.M., 2015. [The Toxicity of Persistent Organic Pollutants to Aquatic Organisms](#), in
2638 1393 ["Persistent Organic Pollutants \(POPs\): Analytical Techniques, Environmental Fate and](#)
2639
2640 1394 [Biological Effects"](#) by Eddy Y. Zeng (Editor), [Comprehensive Analytical Chemistry](#), 67, 587-
2641
2642 1395 [613](#).
2643
2644 1396 Herut, B., Galil, B., Shefer, E., 2010. Monitoring Alpha – disposal site of dredged material in the
2645 1397 Mediterranean Sea. Results of monitoring July-September 2009. *Israel Oceanographic &*
2646
2647 1398 *Limnological Research Reports* (in Hebrew) 9, 47
2648
2649 1399 Heussner, S., Durrieu de Madron, X., Calafat, A., Canals, M., Carbonne, J., Delsaut, N., Saragoni, G.,
2650 1400 2006. Spatial and temporal variability of downward particle fluxes on a continental slope:
2651
2652
2653
2654
2655

2656
 2657
 2658 1401 Lessons from an 8-yr experiment in the Gulf of Lions (NW Mediterranean), Marine Geology,
 2659 234, 63-92.
 2660 1402
 2661
 2662 1403 Hildebrand, J.A., 2009. Anthropogenic and natural sources of ambient noise in the ocean, Marine
 2663 Ecology Progress Series, 395, 5-20, doi: 10.3354/meps08353.
 2664 1404
 2665 1405 Holler, P., Markert, E., Bartholoma, A., Capperucci, R., Hass, H.C., Kroncke, I., Mielck, F., Reimers,
 2666 H.C., 2017. Tools to evaluate seafloor integrity: comparison of multi-device acoustic seafloor
 2667 1406 classifications for benthic macrofauna-driven patterns in the German Bight, southern North
 2668 1407 Sea. Geo-Marine Letters, 37, 93-109.
 2669 1408
 2670 1409 ICES. 2015. Report of the workshop on guidance for the review of MSFD decision descriptor 3 -
 2671 commercial fish and shellfish II (WKGMSFDD3-II), 10-12 February 2015, ICES Headquarters,
 2672 1410 Denmark. ICES CM 2015/ACOM:48, 36 pp.
 2673 1411
 2674 1412 ICES. 2016. EU request to provide guidance on operational methods for the evaluation of the
 2675 MSFD Criterion D3C3. In Report of the ICES Advisory Committee, 2016. ICES Advice 2016,
 2676 1413 Book 1, Section 1.6.2.2.
 2677 1414
 2678 1415 ICES, 2017. Report of the Workshop on Guidance on Development of Operational Methods for the
 2679 1416 Evaluation of the MSFD Criterion D3.3 (WKIND3.3ii), 1-4 November 2016, Copenhagen,
 2680 1417 Denmark. ICES CM 2016/ACOM:44. 145 pp.
 2681 1418
 2682 1419 Iken, K., Brey, T., Wand, U., Voigt, J., Junghans, P., 2001. Food web structure of the benthic
 2683 community at the Porcupine Abyssal Plain (NE Atlantic): a stable isotope analysis. Progress in
 2684 1420 Oceanography, 50(1-4), 383-405.
 2685 1421
 2686 1422 Innocenti, G., Stasolla, G., Goren, M., Stern, N., Levitt-Barmats, Y., Diamant, A., Galil, B.S., 2017.
 2687 1423 Going down together: invasive host, *Charybdis longicollis* (Leene, 1938) (Decapoda:
 2688 Brachyura: Portunidae) and invasive parasite, *Heterosaccus dollfusi* Boschma, 1960
 2689 1424 (Cirripedia: Rhizocephala: Sacculinidae) on the upper slope off the Mediterranean coast of
 2690 Israel. Marine Biology Research, 13, 229-236.
 2691 1425
 2692 1426 Ioakeimidis, C., Zeri, C., Kaberi, E., Galatchi, M., Antoniadis, K., Streftaris, N., Galgani, F.
 2693 Papathanassiou, E., Papatheodorou, G., 2014. A comparative study of marine litter on the
 2694 1427 seafloor of coastal areas in the Eastern Mediterranean and Black Seas. Marine Pollution
 2695 1428 Bulletin, 89, 296-30.
 2696 1429
 2697 1430 Iwata, H., Tanabe, S., Sakai, N., Nishimura, A., Tatsukawa, R., 1994. Geographical distribution of
 2698 persistent organochlorines in air, water and sediments from Asia and Oceania, and their
 2699 1431
 2700
 2701
 2702
 2703
 2704
 2705
 2706
 2707
 2708
 2709
 2710
 2711
 2712
 2713
 2714

2715
2716
2717 1432 implications for global redistribution from lower latitudes. Environmental Pollution 85, 15-
2718 33.
2719 1433
2720
2721 1434 Jimenez, C., Patsalou, P., Andreou, V., Huseyinoglu, M.F., Çiçek, B.A., Hadjioannou, L., Petrou, A.,
2722 1435 2019. Out of sight, out of reach, out of mind: invasive lionfish *Pterois miles* in Cyprus at
2723 depths beyond recreational diving limits. 1st Mediterranean Symposium on the Non-
2724 1436 Indigenous Species (Antalya, Turkey, 17-18 January 2019). pp. 59-64.
2725
2726 1437
2727 1438 Junque, E., Gari, M., Arce, A., Torrent, M., Sunyer, J., Grimalt, J.O., 2017. Integrated assessment of
2728 1439 infant exposure to persistent organic pollutants and mercury via dietary intake in a central
2729 1440 western Mediterranean site (Menorca Island), Environmental Research 156, 714-724.
2730
2731 1441 Junqué, E., Garí M., Lull R.M., Grimalt J.O., 2018. Drivers of the accumulation of mercury and
2732 1442 organochlorine pollutants in Mediterranean lean fish and dietary significance. Science of
2733 1443 the total Environment 634, 170-180.
2734
2735
2736 1443
2737 1444 Katsanevakis, S., Verriopoulos, G., Nicolaidou, A., Thessalou-Legaki, M., 2007. Effect of marine
2738 1445 litter on the benthic megafauna of coastal soft bottoms: A manipulative field experiment.
2739 1446 Marine Pollution Bulletin, 54, 771-778.
2740
2741 1446
2742 1447 Kenny, A.J., Cato, I., Desprez, M., Fader, G., Schuttenhelm, R.T.E., Side, J., 2003. An overview of
2743 1448 seabed-mapping technologies in the context of marine habitat classification, Ices Journal of
2744 1449 Marine Science 60, 2, 411-418.
2745
2746 1449
2747 1450 Klein, B., Roether, W., Kress, N., Manca, B.B., d'Alcala, M.R., Souvermezoglou, E., Theocharis, A.,
2748 1451 Civitarese, G., Luchetta, A., 2003. Accelerated oxygen consumption in eastern
2749 1452 Mediterranean deep waters following the recent changes in thermohaline circulation,
2750 1453 Journal of Geophysical Research-Oceans 108, C9, doi: 10.1029/2002jc001454.
2751
2752
2753 1453
2754 1454 Klein, B., Roether, W., Manca, B.B., Bregant, D., Beitzel, V., Kovacevic, V., Luchetta, A., 1999. The
2755 1455 large deep water transient in the Eastern Mediterranean, Deep Sea Research, Part I, 46, 371-
2756 1456 414.
2757
2758 1456
2759 1457 Koenig, S., Fernandez, P., Sole, M., 2012. Differences in cytochrome P450 enzyme activities
2760 1458 between fish and crustacea: Relationship with the bioaccumulation patterns of
2761 1459 polychlorobiphenyls (PCBs). Aquatic Toxicology, 108, 11-17.
2762
2763 1459
2764 1460 Koenig, S., Fernandez, P., Company, J.B., Huertas, D., Sole, M., 2013a. Are deep-sea organisms
2765 1461 dwelling within a submarine canyon more at risk from anthropogenic contamination than
2766 1462 those from the adjacent open slope? A case study of Blanes canyon (NW Mediterranean),
2767 1463 Progress in Oceanography, 118, 249-259.
2768
2769
2770
2771
2772
2773

2774
2775
2776 1464 Koenig, S., Huertas, D., Fernández, P., 2013b. Legacy and emergent persistent organic pollutants
2777 (POPs) in NW Mediterranean deep-sea organisms, *Science of the Total Environment*, 443,
2778 358-366.
2779
2780 1466
2781 1467 Koenig, S., Sole, M., Fernandez-Gomez, C., Diez, S., 2013c. New insights into mercury
2782 bioaccumulation in deep-sea organisms from the NW Mediterranean and their human
2783 health implications, *Science of the Total Environment*, 442, 329-335.
2784
2785 1469
2786 1470 Koppelman, R., Bottger-Schnack, R., Mobius, J., Weikert, H., 2009. Trophic relationships of
2787 zooplankton in the eastern Mediterranean based on stable isotope measurements, *Journal*
2788 *of Plankton Research*, 31, 669-686.
2789
2790 1472
2791 1473 Koslow, J.A., Boehlert, G.W., Gordon, J.D.M., Haedrich, R.L., Lorange, P., Parin, N., 2000.
2792 [Continental slope and deep-sea fisheries: implications for a fragile ecosystem. ICES Journal of](#)
2793 [Marine Science](#), 57, 548-557.
2794
2795 1475
2796 1476 Kress, N., Hornung, H., Herut, B., 1998. Concentrations of Hg, Cd, Cu, Zn, Fe and Mn in deep sea
2797 benthic fauna from the southeastern Mediterranean Sea: A comparison study between
2798 fauna collected at a pristine area and at two waste disposal sites. *Marine Pollution Bulletin*,
2799 36, 911-921.
2800 1478
2801
2802 1479
2803 1480 Kress, N., Fainshtein, G., Hornung, H., 1996. Monitoring of Hg, Cd, Cu, Pb, Zn, Co, Be and V at a
2804 deep water coal fly ash dumping site. MAP Technical Reports Series N° 104, UNEP/FAO
2805 1481
2806 1482 Kress, N., Gertman, I., Herut, B., 2014. Temporal evolution of physical and chemical characteristics
2807 of the water column in the Easternmost Levantine basin (Eastern Mediterranean Sea) from
2808 2002 to 2010, *Journal of Marine Systems* 135, 6-13, doi: 10.1016/j.jmarsys.2013.11.016.
2809
2810 1484
2811 1485 Lawrence, M.J., Stemberger, H.I.J., Zolderdo, A.J., Struthers, D.P., Cooke S.J., 2015. The effects of
2812 [modern war and military activities on biodiversity and the environment. Environmental](#)
2813 [Reviews](#) 23(4), 443-460
2814
2815 1487
2816 1488 Lastras, G., Canals, M., Ballesteros, E., Gili, J.M., Sanchez-Vidal, A., 2016. Cold-water corals and
2817 anthropogenic impacts in La Fonera submarine canyon head. *PLoS One*, 16;11(5):e0155729.
2818
2819 1489
2820 1490 Lauria, V., Garofalo, G., Fiorentino, F., Massi, D., Milisenda, G., Piraino, S., Russo, T., Gristina, M.,
2821 2017. Species distribution models of two critically endangered deep-sea octocorals reveal
2822 fishing impacts on vulnerable marine ecosystems in central Mediterranean Sea. *Scientific*
2823 *Reports*, 7, 8049.
2824 1492
2825 1493
2826
2827
2828
2829
2830
2831
2832

2833
2834
2835
2836
2837
2838
2839
2840
2841
2842
2843
2844
2845
2846
2847
2848
2849
2850
2851
2852
2853
2854
2855
2856
2857
2858
2859
2860
2861
2862
2863
2864
2865
2866
2867
2868
2869
2870
2871
2872
2873
2874
2875
2876
2877
2878
2879
2880
2881
2882
2883
2884
2885
2886
2887
2888
2889
2890
2891

1494 Le Corre, G., Farrugio, H., 2011. Note sur la création par la CGPM d'une Zone de pêche
réglementée dans le golfe du Lion en mars 2009. RBE/HMT 2011-002.
<https://archimer.ifremer.fr/doc/00086/19688/>

1497 Lejeusne, C., Chevaldonné, P., Pergent-Martini, C., Boudouresque, C., Pérez, T., 2010. Climate
change effects on a miniature ocean: the highly diverse, highly impacted Mediterranean Sea.
Trends in Ecology & Evolution 1204: published online. doi 10.1016/j.tree.2009.1010.1009.

1500 Levin, L.A., 2003. Oxygen minimum zone benthos: Adaptation and community response to
hypoxia. Oceanography and Marine Biology: an Annual Review 2003, 41, 1-45.

1502 Levitus, S., Antonov, J.I., Boyer, T.P., Stephens, C., 2000. Warming of the World Ocean. Science,
287, 2225-2229

1504 Lipiatou, E., Marty, J.C., Saliot, A., 1993. Sediment trap fluxes of polycyclic aromatic hydrocarbons
in the Mediterranean Sea. Marine Chemistry, 44, 43-54.

1506 Livnat, M., 2014. Offshore safety in the Eastern Mediterranean energy sector. Implications of the
new EU directive. Mediterranean Paper Series 2014, The German Marshall Fund of the
United States, Washington DC, USA, 12 p.

1509 Llull, R.M., Garí, M., Canals, M., Rey-Maquieira, T., Grimalt, J.O., 2017. Mercury concentrations in
lean fish from the Western Mediterranean Sea: Dietary exposure and risk assessment in the
population of the Balearic Islands. Environmental Research, 158, 16-23.

1512 Lo Iacono, C., Gràcia, E., Bartolomé, R., Coiras, E., Dañobeitia, J.J., Acosta, J., 2012. The habitats of
the Chella Bank, Eastern Alboran Sea (Western Mediterranean). In: Harris P, Baker E
(eds.) Seafloor geomorphology as benthic habitat: GeoHab Atlas of seafloor geomorphic
features and benthic habitats. Elsevier, London, pp 681-687.

1516 Lo Iacono, C., Robert, K., Gonzalez-Villanueva, R., Gori, A., Gili, J.-M., Orejas, C., 2018. Predicting
cold-water coral distribution in the Cap de Creus Canyon (NW Mediterranean): implications
for marine conservation planning. Progress in Oceanography, 169, 169-180.

1519 Longhurst, A.R., 2017. Chapter 1 - Toward an ecological geography of the sea. In Ecological
Geography of the Sea (Second Edition), Editor A.R. Longhurst, Academic Press, 1-17.

1521 Luna, G.M., Bianchelli, S., Decembrini, F., De Domenico, E., Danovaro, R., Dell'Anno, A., 2012. The
dark portion of the Mediterranean Sea is a bioreactor of organic matter cycling. Global
Biogeochemical Cycles, 26(2) GB2017,doi:10.1029/2011GB004168

2892
2893
2894
2895 1524 Maglio, A., Pavan, G., Castellote, M., Frey, S., 2016. Overview of the noise hotspots in the
2896 1525 ACCOBAMS area – Part I, Mediterranean Sea. Agreement on the Conservation of Cetaceans
2897
2898 1526 in the Black Sea, Mediterranean Sea and Contiguous Area, 44 p.
2899
2900 1527 Martin, J., Palanques, A., Puig, P., 2006. Composition and variability of downward particulate
2901 1528 matter fluxes in the Palamos submarine canyon (NW Mediterranean), Journal of Marine
2902
2903 1529 Systems, 60, 75-97.
2904
2905 1530 Martin, J., Puig, P., Palanques, A., Masque, P., Garcia-Orellana, J., 2008. Effect of commercial
2906 1531 trawling on the deep sedimentation in a Mediterranean submarine canyon, Marine Geology,
2907
2908 1532 252, 150-155.
2909
2910 1533 Martin, J., Miquel, J.C., Khripounoff, A., 2010. Impact of open sea deep convection on sediment
2911 1534 remobilization in the western Mediterranean, Geophysical Research Letters, 37, doi:
2912
2913 1535 10.1029/2010gl043704.
2914
2915 1536 Martin, J., Puig, P., Masque, P., Palanques, A., Sanchez-Gomez, A., 2014. Impact of bottom trawling
2916 1537 on deep-sea sediment properties along the flanks of a submarine canyon, Plos One 9, 8, doi:
2917
2918 1538 e104536, 10.1371/journal.pone.0104536.
2919
2920 1539 Mascle, J., Mary, F., Praeg, D., Brosolo, L., Camera, L., Ceramicola, S., Dupré, S., 2014. Distribution
2921 1540 and geological control of mud volcanoes and other fluid/free gas seepage features in the
2922
2923 1541 Mediterranean Sea and nearby Gulf of Cadiz, Geo-Marine Letters, 34, 89-110.
2924
2925 1542 Massutí, E., Gordon, J. D., Moranta, J., Swan, S. C., Stefanescu, C., 2004. Mediterranean and
2926 1543 Atlantic deep-sea fish assemblages: differences in biomass composition and size-related
2927
2928 1544 structure. Scientia Marina, 68 (S3), 101-115.
2929
2930 1545 McCauley, R., Day, R.D., Swadling, K.M., Fitzgibbon, Q.P., Watson, R.A., 2017. Widely used marine
2931 1546 seismic survey air gun operations negatively impact zooplankton, Nature Ecology &
2932
2933 1547 Evolution, 1, 1-8, doi: 10.1038/s41559-017-0195.
2934
2935 1548 Mecho A., Aguzzi J., De Mol B., Lastras G., Ramirez-Ilodra E., Bahamon N., Company J.B., Canals M.
2936
2937 1549 2017. Visual faunistic exploration of geomorphological human-impacted deep-sea areas of
2938
2939 1550 the north-western Mediterranean Sea. J. Mar. Biol. Ass. UK, 98: 1241-1252.
2940
2941 1551 MEDIAS Handbook, 2015. Common protocol for the Pan-Mediterranean Acoustic Survey (MEDIAS).
2942 1552 Sète, France, March 2015, 21 pp. Available from: [http://www.medias-](http://www.medias-project.eu/medias/website/handbooks-menu/func-startdown/60/)
2943
2944 1553 [project.eu/medias/website/handbooks-menu/func-startdown/60/](http://www.medias-project.eu/medias/website/handbooks-menu/func-startdown/60/)
2945
2946 1554 MEDITS Handbook. Version n. 9, 2017, MEDITS Working Group: 106 pp. Available from:
2947 1555 <http://www.sibm.it/MEDITS%202011/principaledownload.htm>

2951
2952
2953 1556 Mercado J.M., Yebra L., Cortés D., Beken C., Simboura M., Moncheva S., Alonso A., Gómez F.,
2954
2955 1557 Salles S., Sánchez A., Valcarcel N., 2015. Designing joint monitoring programs for the MSFD
2956
2957 1558 Eutrophication assessment based on the monitoring strategy of UNEP/MAP (Barcelona
2958
2959 1559 Convention). In: Plans for the design of Joint Monitoring Programs in the Mediterranean and
2960
2961 1560 Black Sea regions adapted to MSFD requirements. - IRIS-SES project. Fransisco Alemany,
2962
2963 1561 Pagou Kalliopi, Giannoudi Louisa, Streftaris Nikos (eds.), IRIS-SES Project, May 2015.
2964
2965 1562 Migeon, S., Mascle, J., Coste, M., Rouillard, P., 2012. Mediterranean submarine canyons and
2966
2967 1563 channels: Morphological and geological backgrounds. In: Wurtz, M. (ed.) Mediterranean
2968
2969 1564 submarine canyons: Ecology and Governance, pp. 27-41, IUCN, Gland, Switzerland and
2970
2971 1565 Malaga, Spain.
2972
2973 1566 Moccia, D., Cau, A., Alvito, A., Canese, S., Cannas, R., Bo, M., Angiolillo, M., Follesa, M.C., 2019.
2974
2975 1567 New sites expanding the “Sardinian cold-water coral province” extension: A new potential
2976
2977 1568 cold-water coral network? Aquatic Conservation: Marine and Freshwater Ecosystems, 29,
2978
2979 1569 153-160.
2980
2981 1570 Molari M., Manini E., Dell’Anno A., 2013. Dark inorganic carbon fixation sustains the functioning of
2982
2983 1571 benthic deep-sea ecosystems. Global Biogeochemical Cycles, 27, 212-221.
2984
2985 1572 Monaco, A., Peruzzi, S., 2002. The Mediterranean Targeted Project MATER—a multiscale approach
2986
2987 1573 of the variability of a marine system—overview. Journal of Marine Systems, 33, 3-21.
2988
2989 1574 Montgomery, J. C., Radford, C. A. (2017). Marine bioacoustics. Current Biology, 27, R502-R507.
2990
2991 1575 Morley, N.H., Burton, J.D., Tankere, S.P.C., Martin, J.M., 1997. Distribution and behaviour of some
2992
2993 1576 dissolved trace metals in the western Mediterranean Sea. Deep-Sea Research Part II, 44, 3-4,
2994
2995 1577 675-691.
2996
2997 1578 Murray, F., Cowie, P., 2011. Plastic contamination in the decapod crustacean *Nephrops norvegicus*
2998
2999 1579 (Linnaeus, 1758). Marine Pollution Bulletin, 62, 1207–1217.
3000
3001 1580 Naumann, M.S., Imma Tolosa, I., Taviani, M., Grover, R., Christine Ferrier-Pagés, C., 2015. Trophic
3002
3003 1581 ecology of two cold-water coral species from the Mediterranean Sea revealed by lipid
3004
3005 1582 biomarkers and compound-specific isotope analyses. Coral Reefs, 34, 1165–1175.
3006
3007 1583 doi10.1007/s00338-015-1325-8.
3008
3009 1584 [NRC-National Research Council \(US\), 2003. Committee on Potential Impacts of Ambient Noise in
the Ocean on Marine Mammals. Washington \(DC\): National Academies Press \(US\).](#)
1585
1586 Nelms, S.E., Piniak, W.E.D., Weir, C.R., Godley, B.J., 2016. Seismic surveys and marine turtles: an
underestimated global threat? Biological Conservation, 193, 49–65.

3010
3011
3012
3013 1588 Newman, S., Watkins, E., Farmer, A., Ten Brink, P., & Schweitzer, J. P., 2015. The Economics of
3014 1589 marine litter. In: Bergmann M., Gutow L., Klages M. (eds) Marine Anthropogenic Litter.
3015
3016 1590 Springer, Cham.

3017
3018 1591 Nixon, S.W., 2009. Eutrophication and the macroscope. In Eutrophication in Coastal Ecosystems
3019 1592 (pp. 5-19). Springer, Dordrecht.

3020
3021 1593 Nowacek, D.P., Clark, C.W., Mann, D., Miller, P.J., Rosenbaum, H.C., Golden, J.S., Jasny, M., Kraska,
3022 1594 J. and Southall, B.L., 2015, Marine seismic surveys and ocean noise: time for coordinated and
3023
3024 1595 prudent planning. *Frontiers in Ecology and the Environment*, 13, 378-386.

3025
3026 1596 Oehlmann, J., Schulte-Oehlmann, U., Kloas, W., Jagnytsch, O., Lutz, I., Kusk, K., Wollenberger, L.,
3027
3028 1597 Santos, E., Paull, G., Van Look, K., Tyler, C., 2009. A critical analysis of the biological impacts
3029 1598 of plasticizers on wildlife. *Philosophical Transaction of the Royal Society Part B*, 364, 2047-
3030
3031 1599 2062.

3032
3033 1600 Olu-Le Roy, K., Sibuet, M., Fiala-Médioni, A., Gofas, S., Salas, C., Mariotti, A., Foucher, J.P.,
3034 1601 Woodside, J., 2004. Cold seep communities in the deep eastern Mediterranean Sea:
3035
3036 1602 composition, symbiosis and spatial distribution on mud volcanoes. *Deep Sea Research Part I*
3037
3038 1603 51, 1915-1936.

3039
3040 1604 Özcan, T., Ateş, A.S., Katağan, T., 2008. Expanding distribution and occurrence of the Indo-Pacific
3041 1605 Stomatopod, *Erugosquilla massavensis* (Kossmann, 1880) on the Aegean coast of Turkey.
3042
3043 1606 *Mediterranean Marine Science*, 9(2), 115-118.

3044
3045 1607 Özgür Özbek, E., Cardak, M., Kebapçioğlu, T., 2017. Spatio-temporal patterns of abundance,
3046 1608 biomass and length of the silver-cheeked toadfish *Lagocephalus sceleratus* in the Gulf of
3047
3048 1609 Antalya, *Turkish Journal of Fisheries and Aquatic Science*, 17, 725-733.

3049
3050 1610 Palanques, A., de Madron, X.D., Puig, P., Fabres, J., Guillen, J., Calafat, A., Canals, M., Heussner, S.,
3051 1611 Bonnin, J., 2006. Suspended sediment fluxes and transport processes in the Gulf of Lions
3052
3053 1612 submarine canyons. The role of storms and dense water cascading. *Marine Geology*, 234, 43-
3054
3055 1613 61.

3056
3057 1614 Palanques, A., Guillen, J., Puig, P., Durrieu de Madron, X., 2008. Storm-driven shelf-to-canyon
3058 1615 suspended sediment transport at the southwestern Gulf of Lions. *Continental Shelf*
3059
3060 1616 *Research*, 28, 1947-1956.

3061
3062 1617 Papiol, V., Cartes, J.E., Fanelli, E., Rumolo, P., 2013. Food web structure and seasonality of slope
3063 1618 megafauna in the NW Mediterranean elucidated by stable isotopes: Relationship with
3064
3065 1619 available food sources, *Journal of Sea Research*, 77, 53-69.

3069
3070
3071
3072
3073
3074
3075
3076
3077
3078
3079
3080
3081
3082
3083
3084
3085
3086
3087
3088
3089
3090
3091
3092
3093
3094
3095
3096
3097
3098
3099
3100
3101
3102
3103
3104
3105
3106
3107
3108
3109
3110
3111
3112
3113
3114
3115
3116
3117
3118
3119
3120
3121
3122
3123
3124
3125
3126
3127

Parravicini, V., Azzurro, E., Kulbicki, M., & Belmaker, J., 2015. Niche shift can impair the ability to predict invasion risk in the marine realm: an illustration using Mediterranean fish invaders. *Ecology Letters*, 18, 246-253.

Peng, C., Zhao, X.G., Liu, G.X., 2015. Noise in the sea and its impacts on marine organisms, *International Journal of Environmental Research and Public Health*, 12, 12304-12323.

Pham, C., Ramirez-Llodra, E., Claudia, H.S., Amaro, T., Bergmann, M., Canals, M., Company, J., Davies, J., Duineveld, G., Galgani, F., Howell, K., Huvenne, V.A., Isidro, E., Jones, D., Lastras, G., Morato, T., Gomes-Pereira, J., Purser, A., Stewart, H., Tojeira, I., Tubau, X., Van Rooij, D., Tyler, P., 2014. Marine litter distribution and density in European seas, from the shelves to deep basins. *PLoS One*, 9(4), e95839.

Piante, C., Ody, D., 2015. Blue Growth in the Mediterranean Sea: The Challenge of Good Environmental Status. *MedTrends Project*. WWF-France, 192 p.

Pierdomenico, M., Casalbore, D., Chiocci, F.L., 2019. Massive benthic litter funnelled to deep sea by flash-flood generated hyperpycnal flows. *Scientific Reports* 9: 5330.

Pinsky, M.L., Palumbi, S.R., 2014. [Meta-analysis reveals lower genetic diversity in overfished populations. *Molecular Ecology*, 23\(1\), 29-39.](#)

Piroddi, C., Coll, M., Liqueste, C., Macias, D., Greer, K., Buszowski, J., Steenbeek, J., Danovaro, R., Christensen, V., 2017. Historical changes of the Mediterranean Sea ecosystem: modelling the role and impact of primary productivity and fisheries changes over time. *Scientific Reports*, 7, 44491; doi: 10.1038/srep44491.

Popper, A.N., Hawkins, A.D., Fay, R.R., Mann, D.A., Bartol, S., Carlson, T.J., Coombs, S., Ellison, W.T., Gentry, R.L., Halvorsen, M.B., Lokkeborg, S., Rogers, P., Southall, B.L., Zeddies, D.G., Tavalga, W.N., 2014. Sound Exposure Guidelines for Fishes and Sea Turtles: A Technical Report prepared by ANSI-Accredited Standards Committee S3/SC1 and registered with ANSI. *Springer Briefs in Oceanography*. <http://www.springer.com/gp/book/9783319066585>

Porte, C., Escartin, E., Garcia, L.M., Solé, M., Albaigés, J., 2000. Xenobiotic metabolising enzymes and antioxidant defences in deep-sea fish: relationship with contaminant body burden. *Marine Ecology Progress Series*, 192, 259-266.

Poulos, S.E., Collins, M.B., Pattiaratchi, C., Cramp, A., Gull, W., Tsimplis, M., Papatheodorou, G., 1996. Oceanography and sedimentation in the semi-enclosed, deep-water Gulf of Corinth (Greece), *Marine Geology*, 134, 213-235.

3128
3129
3130
3131 1651 Puig, P., Canals, M., Company, J.B., Martin, J., Amblas, D., Lastras, G., Palanques, A., Calafat, A.M.,
3132 1652 2012. Ploughing the deep sea floor, *Nature* 489, 7415, 286-289, doi: 10.1038/nature11410.
3133
3134 1653 Pusceddu, A., Bianchelli, S., Martín, J., Puig, P., Palanques, A., Masqué, P., Danovaro, R., 2014.
3135 1654 Chronic and intensive bottom trawling impairs deep-sea biodiversity and ecosystem
3136 1655 functioning. *Proceedings of the National Academy of Sciences*, 111, 8861-8866.
3137
3138
3139 1656 Pusceddu, A., Bianchelli S., Gambi, C., Danovaro, R., 2011. Assessment of benthic trophic status of
3140 1657 marine coastal ecosystems: significance of meiofaunal rare taxa. *Estuarine, Coastal and Shelf*
3141 1658 *Science*, 93, 420-430.
3142
3143
3144 1659 Pusceddu, A., Dell'Anno, A., Fabiano, M., Danovaro, R., 2009. Quantity and bioavailability of
3145 1660 sediment organic matter as signature of benthic trophic state. *Marine Ecology Progress*
3146 1661 *Series* 375, 41-52.
3147
3148
3149 1662 Quetglas, A., Ordines, F., Gonzale, M., Zaragoza, N., Mallol, S., Valls, M., De Mesa, A., 2013.
3150 1663 Uncommon pelagic and deep-sea cephalopods in the Mediterranean: new data and
3151 1664 literature review. *Mediterranean Marine Science*, 14, 69-85.
3152
3153
3154 1665 Raicevich, S., Battaglia, P., Fortibuoni, T., Romeo, T., Giovanardi, O., Andaloro, F., 2017. Critical
3155 1666 Inconsistencies in Early Implementations of the Marine Strategy Framework Directive and
3156 1667 Common Fisheries Policy Objectives Hamper Policy Synergies in Fostering the Sustainable
3157 1668 Exploitation of Mediterranean Fisheries Resources. *Frontiers in Marine Science*, 4 : 316.
3158
3159
3160
3161 1669 Ramirez-Llodra, E., Company, J.B., Sardá, F., Rotllant, G., 2010. Megabenthic diversity patterns and
3162 1670 community structure of the Blanes submarine canyon and adjacent slope in the
3163 1671 Northwestern Mediterranean: A human overprint? *Marine Ecology*, 31, 167-182.
3164
3165
3166 1672 Ramirez-Llodra, E., Tyler, P.A., Baker, M.C., Bergstad, O.A., Clark, M.R., Escobar, E., Levin, L.A.,
3167 1673 Menot, L., Rowden, A.A., Smith, C.R., van Dover, C.L., 2011. Man and the last great
3168 1674 wilderness: human impact on the deep sea. *PLoS One* 6 (8):e22588.
3169
3170
3171 1675 Ramirez-Llodra, E., De Mol, B., Company, J.B., Coll, M., Sardà, F., 2013. Effects of natural and
3172 1676 anthropogenic processes in the distribution of marine litter in the deep Mediterranean Sea.
3173 1677 *Progress in Oceanography*, 118, 273-287.
3174
3175
3176 1678 Raoux, C., Bayona, J.M., Miquel, J.C., Teyssie, J.L., Fowler, S.W., Albaiges, J., 1999. Particulate
3177 1679 fluxes of aliphatic and aromatic hydrocarbons in near-shore waters to the northwestern
3178 1680 Mediterranean Sea, and the effect of continental runoff. *Estuarine, Coastal and Shelf*
3179 1681 *Science*, 48, 605-616.
3180
3181
3182
3183
3184
3185
3186

3187
3188
3189
3190 1682 Reid, P.C., Hari, R. E., Beaugrand, G., Livingstone, D.M., Marty, C., Straile, D., Barichivich, J.,
3191 1683 Goberville, E., Adrian, R., Aono, Y., Brown, R., Foster, J., Groisman, P., Hélaouët, P., Hsu, H.,
3192
3193 1684 Kirby, R., Knight, J., Kraberg, A., Li, J., Lo, T., Myneni, R.B., North, R.P., Pounds, J. A., Sparks,
3194
3195 1685 T., Stübi, R., Tian, Y., Wiltshire, K.H., Xiao, D. and Zhu, Z. (2016), Global impacts of the 1980s
3196 1686 regime shift. *Global change biology*, 22(2), 682-703.
3197
3198 1687 Rixen, M., Beckers, J.M., Levitus, S., Antonov, J., Boyer, T., Maillard, C., Fichaut, M., Balopoulos, E.,
3199
3200 1688 Iona, S., Dooley, H., Garcia, M.-J., Manca, B., Giorgetti, A., Manzella, G., Mikhailov, N.,
3201 1689 Pinardi, N., Zavatarelli, M., 2005. The Western Mediterranean Deep Water: A proxy for
3202
3203 1690 climate change, *Geophysical Research Letters* 32, 12, doi: 10.1029/2005gl022702.
3204
3205 1691 Roether, W., Well, R., 2001. Oxygen consumption in the Eastern Mediterranean, *Deep Sea*
3206 1692 *Research, Part I* , 48, 1535-1551.
3207
3208 1693 Rogers, A.D., 1999. The biology of *Lophelia pertusa* (Linnaeus 1758) and other deep-water reef-
3209
3210 1694 forming corals and impacts from human activities. *International Review of Hydrobiology*, 84
3211 1695 (4), 315-410.
3212
3213 1696 Rombouts, I., Beaugrand, G., Artigas, L.F., Dauvin, J.C., Gevaert, F., Goberville, E., Kopp, D.,
3214
3215 1697 Lefebvre, S., Luczak, C., Spilmont, N., Travers-Trolet, M., Villanueva, M.C., Kirby, R.R., 2013.
3216 1698 Evaluating marine ecosystem health: Case studies of indicators using direct observations and
3217
3218 1699 modelling methods. *Ecological Indicators* 24, 353-365.
3219
3220 1700 Rotllant, G., Abad, E., Sarda, F., Abalos, M., Company, J.B., Rivera, J., 2006. Dioxin compounds in
3221 1701 the deep-sea rose shrimp *Aristeus antennatus* (Risso, 1816) throughout the Mediterranean
3222
3223 1702 Sea, *Deep- Research, Part I*, 53, 1895-1906.
3224
3225 1703 Rountree, R., Aguzzi, J., Marini, S., Fanelli, E., De Leo, F.C., Del Rio, J. and Juanes, F., 2019. Towards
3226
3227 1704 an optimal design for ecosystem-level ocean observatories. *Advances in Marine Biology, An*
3228 1705 *Annual Review*, in press.
3229
3230 1706 Ryan, P. G., Moore, C. J., van Franeker, J. A., Moloney, C. L., 2009. Monitoring the abundance of
3231
3232 1707 plastic debris in the marine environment. *Philosophical Transactions of the Royal Society B:*
3233 1708 *Biological Sciences*, 364, 1999-2012.
3234
3235 1709 Salvadó, J.A., Grimalt, J.O., López, J.F., Palanques, A., Heussner, S., Pasqual, C., Sanchez-Vidal A.,
3236
3237 1710 Canals, M., 2017. Transfer of lipid molecules and polycyclic aromatic hydrocarbons to open
3238
3239 1711 marine waters by dense water cascading events. *Progress in Oceanography*, 159, 178-194.
3240 1712 Sanchez-Vidal, A., Pasqual, C., Kerherve, P., Calafat, A., Heussner, S., Palanques, A., Durrieu de
3241
3242 1713 Madron, X., Canals, M., Puig, P., 2008. Impact of dense shelf water cascading on the transfer
3243
3244
3245

3246
3247
3248
3249 1714 of organic matter to the deep western Mediterranean basin. Geophysical Research Letters
3250 1715 35, L05605, doi:10.1029/2007GL032825.
3251
3252 1716 Sanchez-Vidal, A., Canals, M., Calafat, A.M., Lastras, G., Pedrosa-Pàmies, R., Menéndez, M.,
3253 1717 Medina, R., Company, J.B., Hereu, B., Romero, J., Alcoverro, T., 2012. Impacts on the deep-
3254 1718 sea ecosystem by a severe coastal storm; PLoS one, 7, e30395. doi:
3255 1718 10.1371/journal.pone.0030395.
3256 1719
3257 1720 Sanchez-Vidal, A., Thompson, R.C., Canals, M., de Haan, W.P., 2018. The imprint of microfibrils in
3258 1721 southern European deep seas. PLoS ONE 13 (11): e0207033.
3259 1721
3260 1722 Sardá, F., Calafat, A., Flexas, M., Tselepidis, A., Canals, M., Espino, M., Tursi, A., 2004. An
3261 1723 introduction to Mediterranean deep-sea biology. Scientia Marina, 68 (suppl. 3), 7-38.
3262 1723
3263 1724 Schembri PJ, Dimech M, Camilleri M, Page, R., 2007. Living deep-water *Lophelia* and *Madrepora*
3264 1725 corals in Maltese waters (Strait of Sicily, Mediterranean Sea). Cahiers de Biologie Marine, 48,
3265 1726 77-83.
3266 1726
3267 1727 Schroeder, K., Gasparini, G.P., Borghini, M., Ribotti, A., 2009. Experimental evidences of the recent
3268 1728 abrupt changes in the deep Western Mediterranean Sea. Dynamics of Mediterranean Deep
3269 1729 Waters, n° 38, Qwara, Malta.
3270 1730 Schroeder, K., Millot, C., Bengara, L., Ben Ismail, S., Bensi, M., Borghini, M., Budillon, G., Cardin, V.,
3271 1731 Coppola, L., Curtil, C., Drago, A., El Moumni, B., Font, J., Fuda, J.L., Garcia-Lafuente, J.,
3272 1732 Gasparini, G.P., Kontoyiannis, H., Lefevre, D., Puig, P., Raimbault, P., Rougier, G., Salat, J.,
3273 1733 Sammari, C., Sanchez Garrido, J. C., Sanchez-Roman, A., Sparnocchia, S., Tamburini, C.,
3274 1734 Taupier-Letage, I., Theocharis, A., Vargas-Yanez, M., Vetrano, A., 2013. Long-term
3275 1735 monitoring programme of the hydrological variability in the Mediterranean Sea: a first
3276 1736 overview of the HYDROCHANGES network. Ocean Science 9, 301-324.
3277 1737 Schroeder, K., Chiggiato, J., Bryden, H.L., Borghini, M., Ben Ismail, S., 2016. Abrupt climate shift in
3278 1738 the Western Mediterranean Sea, Scientific Reports 6, 23009, doi: 10.1038/srep23009.
3279 1739 Semprucci, F., Sbrocca, C., Rocchi, M., Balsamo, M., 2014. Temporal changes of the meiofaunal
3280 1740 assemblage as a tool for the assessment of the ecological quality status. Journal of the
3281 1741 Marine Biological Association of the United Kingdom, 95, 247-254.
3282 1742 Shaltout M., Omstedt A., 2014. Recent sea surface temperature trends and future scenarios for
3283 1743 the Mediterranean Sea, Oceanologia, 56, 411-443.
3284 1743
3285
3286
3287
3288
3289
3290
3291
3292
3293
3294
3295
3296
3297
3298
3299
3300
3301
3302
3303
3304

3305
3306
3307 1744 Simoncelli, S., Coatanoan, C., Myroshnychenko, V., Sagen, H., BÄck, Ö., Scory, S., Grandi, A.,
3308
3309 1745 Schlitzer, R., Fichaut, M., 2015. Second release of the SeaDataNet aggregated data sets
3310
3311 1746 products. SeaDataNet, 10.13155/50382, <http://archimer.ifremer.fr/doc/00392/50382/>
3312
3313 1747 Simpson, S.D., Meekan, M., Montgomery, J., McCauley, R., Jeffs, A., 2005. Homeward sound,
3314 1748 Science 308, 5719, 221-221, doi: 10.1126/science.1107406.
3315
3316 1749 Smith CR, De Leo FC, Bernardino AF, Sweetman AK, Arbizu PM., 2008. Abyssal food limitation,
3317
3318 1750 ecosystem structure and climate change. Trends in Ecology & Evolution, 23, 518-28.
3319
3320 1751 Snelgrove, P.V.R., Soetaert, K., Solan, M., Thrush, S., Wei, C.-L., Danovaro, R, Fulweiler, R.W.,
3321 1752 Kitazato, H., Ingole, B., Norkko, A., Parkes, R.J., Volkenborn, N., 2018. Contrasting
3322
3323 1753 biogeochemical and biological estimates of carbon turnover on the global seafloor. Trends in
3324 1754 Ecology & Evolution, 33, 96-105.
3325
3326 1755 Solé, M., Porte, C., Albaiges, J., 2001. Hydrocarbons, PCBs and DDT in the NW Mediterranean
3327
3328 1756 deep-sea fish *Mora moro*. Deep Sea Research, Part I, 48, 495-513.
3329
3330 1757 Stefanescu, C., Moralesnin, B., Massuti, E., 1994. Fish assemblages on the slope in the Catalan sea
3331 1758 (western Mediterranean) - influence of a submarine-canyon. Journal of the Marine Biological
3332
3333 1759 Association of the United Kingdom, 74, 499-512.
3334
3335 1760 Stephens, D., Diesing, M., 2014. A comparison of supervised classification methods for the
3336 1761 prediction of substrate type using Multibeam acoustic and legacy grain-size data. Plos One 9,
3337
3338 1762 4, doi: e93950, 10.1371/journal.pone.0093950.
3339
3340 1763 Storelli, M.M., Storelli, A., D'Addabbo, R., Barone, G., Marcotrigiano, G.O., 2004b. Polychlorinated
3341 1764 biphenyl residues in deep-sea fish from Mediterranean Sea. Environment International, 30,
3342
3343 1765 343-349.
3344
3345 1766 Storelli, M.M., Perrone, V.G., Marcotrigiano, G.O., 2007. Organochlorine contamination (PCBs and
3346 1767 DDTs) in deep-sea fish from the Mediterranean SeaseaSea, Marine Pollution Bulletin, 54,
3347
3348 1768 1968-1971.
3349
3350 1769 Storelli, M.M., Losada, S., Marcotrigiano, G.O., Roosens, L., Barone, G., Neels, H., Covaci, A., 2009.
3351 1770 Polychlorinated biphenyl and organochlorine pesticide contamination signatures in deep-sea
3352
3353 1771 fish from the Mediterranean Sea, Environmental Research, 109, 851-856.
3354
3355 1772 Stramma, L., Johnson, G.C., Sprintall, J., Mohrholz, V., 2008. Expanding oxygen-minimum zones in
3356 1773 the tropical oceans, Science 320, 5876, 655-658.
3357
3358
3359
3360
3361
3362
3363

3364
3365
3366 1774 Tamburini, C., Canals, M., Durrieu de Madron, X., Houpert, L., Lefèvre, D., Martini, S., and
3367
3368 1775 ANTARES consortium (2013) Deep-Sea Bioluminescence Blooms after Dense Water
3369
3370 1776 Formation at the Ocean Surface. PLoS ONE 8(7): e67523.
3371
3372 1777 Taviani, M., 2002. The Mediterranean benthos from late Miocene up to present: ten million years
3373 1778 of dramatic climatic and geological vicissitudes. *Biologia Marina Mediterranea*, 9, 445–463.
3374
3375 1779 Taviani, M., 2003. Shaping the biogeography of the Mediterranean basin: one geologist's
3376
3377 1780 perspective: Marine biogeography of the Mediterranean Sea: Patterns and dynamics of
3378 1781 biodiversity. *Biogeographia*, 24, 15-22.
3379
3380 1782 Taviani, M., Freiwald, F., Zibrowius, H., 2005. Deep coral growth in the Mediterranean Sea: an
3381
3382 1783 overview. In: Freiwald A, Roberts JM (eds. *Cold-water Corals and Ecosystems*. Springer-
3383 1784 Verlag, Berlin, 137–156
3384
3385 1785 Taviani, M., 2011. The deep-sea chemoautotroph microbial world as experienced by the
3386
3387 1786 Mediterranean metazoans through time. *Lecture Notes in Earth Sciences*, 131, 277–295.
3388
3389 1787 Taviani, M., Angeletti, L., Antolini, B., Ceregato, A., Froglià, C., Lopez Correa, M., Montagna, P.,
3390 1788 Remia, A., Trincardi, F., Vertino, A., 201). *Geo-biology of Mediterranean deep-water coral*
3391
3392 1789 *ecosystems*. *Marine Research at CNR*, 6, pp.705-719.
3393
3394 1790 Taviani, M., 2014. Marine Chemosynthesis in the Mediterranean Sea. In: Goffredo S, Dubinsky Z
3395 1791 (eds.), *The Mediterranean Sea: its history and present challenges* Springer Science+Business
3396
3397 1792 Media Dordrecht 2014, 69-83. DOI 10.1007/978-94-007-6704-1_5,
3398
3399 1793 Taviani, M., Angeletti, L., Beuck, L., Campiani, E., Canese, S., Foglini, F., Freiwald, A., Montagna, P.
3400 1794 and Trincardi, F., 2016. Reprint of 'On and off the beaten track: Megafaunal sessile life and
3401
3402 1795 Adriatic cascading processes'. *Marine Geology*, 375, 146-160.
3403
3404 1796 Taviani, M., Angeletti, L., Canese, S., Cannas, R., Cardone, F., Cau, A., Cau, A.B., Follesa, M.C.,
3405 1797 Marchese, F., Montagna, P., Tessarolo, C., 2017. The "Sardinian cold-water coral province" in
3406
3407 1798 the context of the Mediterranean coral ecosystems. *Deep Sea Research, Part II, Topical*
3408
3409 1799 *Studies in Oceanography*, 145, 61-78.
3410
3411 1800 Taviani, M., Angeletti, L., Cardone, F., Montagna, P., Danovaro, R. (2019) A unique and threatened
3412 1801 deep water coral-bivalve biotope new to the Mediterranean Sea offshore the Naples
3413
3414 1802 megalopolis. *Scientific Reports*, 9, 3411. doi.org/10.1038/s41598-019-39655-8
3415
3416 1803 Taylor, M. L., Gwinnett, C., Robinson, L. F., Woodall, L. C., 2016. Plastic microfibre ingestion by
3417 1804 deep-sea organisms. *Scientific Reports*, 6, 33997.
3418
3419
3420
3421
3422

3423
3424
3425
3426
3427
3428
3429
3430
3431
3432
3433
3434
3435
3436
3437
3438
3439
3440
3441
3442
3443
3444
3445
3446
3447
3448
3449
3450
3451
3452
3453
3454
3455
3456
3457
3458
3459
3460
3461
3462
3463
3464
3465
3466
3467
3468
3469
3470
3471
3472
3473
3474
3475
3476
3477
3478
3479
3480
3481

1805 Tecchio, S., van Oevelen, D., Soetaert, K., Navarro, J., Ramirez-Llodra, E., 2013. Trophic dynamics of
1806 deep-sea megabenthos are mediated by surface productivity, Plos One 8, 5, e63796
1807 10.1371/journal.pone.0063796.

1808 The Petroleum Economist Ltd. (2013). World Energy Atlas. 7th Edition. ISBN 1 86186 343 8. SC
1809 (Sang Choy). International Pte Ltd, Singapore.

1810 Thomsen, L., Aguzzi, J., Costa, C., De Leo, F., Ogston, A., Purser, A., 2017. The oceanic biological
1811 pump: Rapid carbon transfer to the Deep Sea during winter. Scientific Report, 7, 10763.

1812

1813 Trudel, M., Rasmussen, J.B., 2001. Predicting mercury concentration in fish using mass balance
1814 models. Ecological Applications 11, 517-529.

1815 Tsapakis, M., Apostolaki, M., Eisenreich, S., Stephanou, E.G., 2006. Atmospheric deposition and
1816 marine sedimentation fluxes of polycyclic aromatic hydrocarbons in the eastern
1817 Mediterranean basin. Environmental Science & Technology, 40, 4922-4927.

1818 Tubau, X., Canals, M., Lastras, G., Rayo, X., Rivera, J., Amblas, D., 2015. Marine litter on the floor of
1819 deep submarine canyons of the Northwestern Mediterranean Sea: The role of hydrodynamic
1820 processes. Progress in Oceanography, 134, 379-403.

1821 Ulses, C., Estournel, C., Puig, P., Durrieu de Madron, X., Marsaleix P., 2008. Dense water cascading
1822 in the northwestern Mediterranean during the cold winter 2005. Quantification of the
1823 export through the Gulf of Lion and the Catalan margin. Geophysical Research Letters, 35,
1824 L07610, doi:10.1029/2008GL03325.

1825 UNEP(DEPI)/MED, 2007. Eutrophication Monitoring Strategy for the MED POL (REVISION)
1826 UNEP/MAP; WG.321/Inf.5, 9 November 2007.

1827 UNEP, 2012. Implementation of the Ecosystem Approach (EcAp) in the Mediterranean by the
1828 contracting parties in the context of the Barcelona Convention for the Protection of the
1829 Marine Environment and the Coastal region of the Mediterranean and its Protocols.
1830 http://www.rac-spa.org/ecapmed_i

1831 Valls, M., Sweeting, C.J., Olivar, M.P., de Puellas, M.F., Pasqual, C., Polunin, N.V.C., Quetglas, A.,
1832 2014. Structure and dynamics of food webs in the water column on shelf and slope grounds
1833 of the western Mediterranean. Journal of Marine Systems, 138, 171-181.

1834 Van Cauwenberghe, L., Vanreusel, A., Maes, J.C., 2013. Microplastic pollution in deep sea
1835 sediments. Environmental Science and Pollution Research, 182, 495-499.

3482
3483
3484
3485 1836 van Oevelen, D., Bergmann, M., Soetaert, K., Bauerfeind, E., Hasemann, C., Klages, M., Schewe I.,
3486 1837 Soltwedel T., Budaeva, N. E., 2011. Carbon flows in the benthic food web at the deep-sea
3487
3488 1838 observatory HAUSGARTEN (Fram Strait). Deep Sea Research, Part I, 58, 1069-1083.
3489
3490 1839 Varnavas, S.P., Achilleopoulos, P.P., 1995. Factors controlling the vertical and spatial transport of
3491 1840 metal-rich particulate matter in seawater at the outfall of bauxitic red mud toxic waste,
3492
3493 1841 Science of the Total Environment, 175, 199-205.
3494
3495 1842 Varnavas, S., Ferentinos, G., Collins, M., 1986. Dispersion of Bauxitic red mud in the Gulf-of-
3496 1843 Corinth, Greece, Marine Geology 70, 3-4, 211-222, doi: 10.1016/0025-3227(86)90003-4.
3497
3498 1844 Vasilakopoulos, P., Maravelias, C.D., Tserpes, G., 2014. The alarming decline of Mediterranean fish
3499
3500 1845 stocks. Current Biology, 24, 1643-1648.
3501
3502 1846 Vasilis, G., Smith, C. J., Kiparissis, S., Stamouli, C., Dounas, C., Mytilineou, C., 2019. Updating the
3503 1847 distribution status of the critically endangered bamboo coral *Isidella elongata* (Esper, 1788)
3504
3505 1848 in the deep Eastern Mediterranean Sea. Regional Studies in Marine Science, 28, 100610,
3506 1849 doi.org/10.1016/j.rsma.2019.100610.
3507
3508 1850 Wartzok, D., Ketten, D.R., 1999. Marine mammal sensory systems. In: Reynolds, J., Rommel, S.
3509
3510 1851 (eds.) Biology of marine mammals, pp. 117-175, Smithsonian Institution Press, Washington,
3511 1852 DC. ISBN: 1560983752
3512
3513 1853 Weilgart, L., 2017. The impact of ocean noise pollution on fish and invertebrates. Report for
3514
3515 1854 OceanCare, OCEANCARE & DALHOUSIE UNIVERSITY, [https://www.oceancare.org/wp-](https://www.oceancare.org/wp-content/uploads/2017/10/noise_and_fish_review_paper-20171016.pdf)
3516
3517 1855 [content/uploads/2017/10/noise_and_fish_review_paper-20171016.pdf](https://www.oceancare.org/wp-content/uploads/2017/10/noise_and_fish_review_paper-20171016.pdf)
3518
3519 1856 Wenz, G.M., 1962. Acoustic Ambient Noise in the Ocean: Spectra and Sources, The Journal of the
3520 1857 Acoustical Society of America, 34, 1936-1956.
3521
3522 1858 Woodall, L. C., Sanchez-Vidal, A., Canals, M., Paterson, G. L. J., Coppock, R., Sleight, V., Calafat, A.,
3523 1859 Rogers, A. D., Narayanaswamy, B. E., Thompson, R. C., 2014. The deep sea is a major sink for
3524
3525 1860 microplastic debris. Royal Society Open Science, 1, 140317.
3526
3527 1861 Wurtz, M., Rovere, M., 2015. Atlas of the Mediterranean Seamounts and Seamount-like
3528 1862 Structures. IUCN, Gland, Switzerland and Malaga, Spain, pp. 276.
3529
3530 1863 WWF/IUCN, 2004. The Mediterranean deep-sea ecosystems: an overview of their diversity,
3531
3532 1864 structure, functioning and anthropogenic impacts, with a proposal for conservation. IUCN,
3533 1865 Malaga and WWF, Rome.
3534
3535
3536
3537
3538
3539
3540

3541
3542
3543
3544
3545
3546
3547
3548
3549
3550
3551
3552
3553
3554
3555
3556
3557
3558
3559
3560
3561
3562
3563
3564
3565
3566
3567
3568
3569
3570
3571
3572
3573
3574
3575
3576
3577
3578
3579
3580
3581
3582
3583
3584
3585
3586
3587
3588
3589
3590
3591
3592
3593
3594
3595
3596
3597
3598
3599

1866 Xiao, Y.J., Ferreira, J.G., Bricker, S.B., Nunes, J.P., Zhu, M.Y., Zhang, X.L., 2007. Trophic
assessment in Chinese coastal systems - Review of methods and application to the
Changjiang (Yangtze) Estuary and Jiaozhou Bay, Estuaries and Coasts, 30, 901-918.
1869 Yağlioğlu, D., Ayas, D., 2016. New occurrence data of four alien fishes (*Pisodonophis*
semicinctus, *Pterois miles*, *Scarus ghobban* and *Parupeneus forsskali*) from the north
eastern Mediterranean (Yeşilovacik Bay, Turkey). Biharean Biologist, 10, 150–152.
1872 Zenetos, A., 2019. Mediterranean Sea: 30 years of biological invasions (1988-2017). In: 1st
Mediterranean Symposium on the Non-Indigenous Species, 2018, p. 13.
1874 Zuñiga, D., Flexas, M.M., Sanchez-Vidal, A., Coenjaerts, J., Calafat, A., Jorda, G., Garcia-
Orellana, J., Puigdefabregas, J., Canals, M., Espino, M., et al., 2009. Particle fluxes
dynamics in Blanes submarine canyon (Northwestern Mediterranean), Progress in
Oceanography 82, 4, 239-251.

Table 1. List of the descriptors, their definition if GES is achieved and if they are a state or pressure descriptors. Only D3 is both a state and pressure descriptor as it related to aspects such as the level of fishing activity (pressure) and population age, size distribution and biomass indices (state).

Descriptor	GES	State	Pressure
1	Biodiversity is maintained	X	
2	Non-indigenous species do not adversely alter the ecosystem		X
3	The population of commercial fish species is healthy	X	X
4	Elements of food webs ensure long-term abundance and reproduction	X	
5	Eutrophication is minimised		X
6	The sea floor integrity ensures functioning of the ecosystem	X	
7	Permanent alteration of hydrographical conditions does not adversely affect the ecosystem		X
8	Concentrations of contaminants give no effects		X
9	Contaminants in seafood are below safe levels		X
10	Marine litter does not cause harm		X
11	Introduction of energy (including underwater noise) does not adversely affect the ecosystem		X


3659
3660
3661 **Figure captions**
3662
3663

3664 **Figure 1.** Jurisdictional Continental Shelf and deep Mediterranean Sea, with indication of
3665 jurisdictional continental shelf per each country (including non-EU countries).
3666

3667 **Figure 2.** Location of GFCM Geographical Sub-Areas (GSAs) and MSFD Mediterranean national sub-
3668 regions. FAO divisions are shown in thumbnail map.
3669
3670

3671
3672
3673
3674
3675
3676
3677
3678
3679
3680
3681
3682
3683
3684
3685
3686
3687
3688
3689
3690
3691
3692
3693
3694
3695
3696
3697
3698
3699
3700
3701
3702
3703
3704
3705
3706
3707
3708
3709
3710
3711
3712
3713
3714
3715
3716
3717

Jurisdictional Continental Shelf and Deep Mediterranean Sea



Approximative percentages of Jurisdictional Continental Shelf per Country related to the total area of deep sea, calculated with the 200 meters isobath limit, in the Mediterranean Sea: Albania (AL): 0,3%; Algeria (DZ): 6,1%; Croatia (HR): 0,5%; Cyprus (CY): 4,8%; Egypt (EG): 7%; France (FR): 3,5%; Greece (EL): 20,6%; Israel (IL): 1,1%; Italy (IT): 21,2%; Lebanon (LB): 0,95%; Lybia (LY): 15,1%; Malta (MT): 2,3%; Montenegro (ME): 0,15%; Morocco (MA): 0,6%; Spain (ES): 11,1%; Syria (SY): 0,5%; Tunisia (TN): 1,5%; Turkey (TR): 2,5%.

