



UNIVERSITÀ POLITECNICA DELLE MARCHE  
Repository ISTITUZIONALE

Integrated characterization and risk management of marine sediments: The case study of the industrialized Bagnoli area (Naples, Italy)

This is the peer reviewed version of the following article:

*Original*

Integrated characterization and risk management of marine sediments: The case study of the industrialized Bagnoli area (Naples, Italy) / Morroni, L.; D'Errico, G.; Sacchi, M.; Molisso, F.; Armiento, G.; Chiavarini, S.; Rimauro, J.; Guida, M.; Siciliano, A.; Ceparano, M.; Aliberti, F.; Tosti, E.; Gallo, A.; Libralato, G.; Patti, F. P.; Gorbi, S.; Fattorini, D.; Nardi, A.; Di Carlo, M.; Mezzelani, M.; Benedetti, M.; Pellegrini, D.; Musco, L.; Danovaro, R.; Dell'Anno, A.; Regoli, F.. - In: MARINE ENVIRONMENTAL RESEARCH. - ISSN 0141-1136. - STAMPA. - 160:(2020). [10.1016/j.marenvres.2020.104984]

*Availability:*

This version is available at: 11566/277413 since: 2024-04-11T14:05:41Z

*Publisher:*

*Published*

DOI:10.1016/j.marenvres.2020.104984

*Terms of use:*

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. The use of copyrighted works requires the consent of the rights' holder (author or publisher). Works made available under a Creative Commons license or a Publisher's custom-made license can be used according to the terms and conditions contained therein. See editor's website for further information and terms and conditions.

This item was downloaded from IRIS Università Politecnica delle Marche (<https://iris.univpm.it>). When citing, please refer to the published version.

(Article begins on next page)



# Integrated characterization and risk management of marine sediments: the case study of the industrialized Bagnoli area (Naples, Italy)

Morroni L.<sup>1±</sup>, d'Errico G.<sup>2±</sup>, Sacchi M.<sup>3</sup>, Molisso F.<sup>3</sup>, Armiento G.<sup>4</sup>, Chiavarini, S.<sup>4</sup>, Rimauro, J.<sup>5</sup>, Guida M.<sup>6</sup>, Siciliano A.<sup>6</sup>, Ceparano M.<sup>6</sup>, Aliberti F.<sup>6</sup>, Tosti, E.<sup>7</sup>, Gallo A.<sup>7</sup>, Libralato G.<sup>6</sup>, Patti F.P.<sup>7</sup>, Gorbi S.<sup>2</sup>, Fattorini D.<sup>2</sup>, Nardi A.<sup>2</sup>, Di Carlo M.<sup>2</sup>, Mezzelani M.<sup>2</sup>, Benedetti M.<sup>2</sup>, Pellegrini D.<sup>1</sup>, Musco L.<sup>7</sup>, Danovaro R.<sup>2,7</sup>, Dell'Anno A.<sup>2</sup>, Regoli F.<sup>2</sup>

1: Istituto Superiore per la Protezione e la Ricerca Ambientale (ISPRA), Via del Cedro 38, 57122 Livorno, Italy.

2: Dipartimento di Scienze della Vita e dell'Ambiente, Università Politecnica delle Marche, Via Brecce Bianche, 60131, Ancona (Italy)

3: Istituto di Scienze Marine CNR-ISMAR, Calata Porta di Massa 80, 80133, Napoli (Italy)

4: ENEA, Dipartimento Sostenibilità, CR Casaccia, via Anguillarese 301, 00123, Roma (Italy)

5: ENEA, Dipartimento Sostenibilità, CR Portici, P.le Enrico Fermi 1, 80055, Portici (Naples, Italy)

6: Department of Biology, Università di Napoli Federico II, via Cinthia, 80126 Naples (Italy)

7: Stazione Zoologica Anton Dohrn, 80121 Napoli (Italy)

±: equally contributed

Corresponding author: Lorenzo Morroni

**Keywords:** Ecological Risk Assessment; Weight Of Evidence (WOE); Sediments; Contamination; Bioavailability; Bioassays; Biomarkers; Benthic Communities

## Abstract

The aim of the present work is to demonstrate the practical importance of a multidisciplinary approach and weighted criteria to synthesize and integrate different typologies of data (or lines of evidence, LOEs), including chemical levels in marine sediments, their bioavailability to specific indicator species, ecotoxicological effects measured through subcellular biomarkers and batteries of bioassays, and potential impacts of pollution on local benthic communities. The area of Bagnoli (Gulf of Naples, Southern Italy) was selected as a model case-study, as it is a coastal area chronically impacted by massive industrial contamination (trace metals and hydrocarbons), and dismissed decades ago without any subsequent remediation or habitat restoration. The results of each LOE were elaborated to provide specific hazard indices before their overall integration in a weight of evidence (WOE) evaluation. Levels of some trace metals and PAHs revealed a severe contamination in the entire study area. Bioavailability of hydrocarbons was evident particularly for high molecular weight PAHs, which also caused significant variations of cellular biomarkers, such as cytochrome P450 metabolization in fish, lysosomal membrane destabilization in mussels,

genotoxic effects both in fish and molluscs. The results of a battery of bioassays indicated less marked responses compared to those obtained from chemical and biomarkers analyses, with acute toxicity still present in sediments close to the source of contamination. The analysis of benthic assemblages showed limited evidence of impact in the whole area, indicating a good functioning of local ecosystems at chronic contamination. Overall, the results of this study confirm the need of combining chemical and biological data, the quantitative characterization of various typologies of hazard and the importance of assessing an integrated environmental WOE risk, to orientate specific and scientifically-supported management options in industrialized areas.

## **1. Introduction**

Since the first chemical factory built in 1854, the Bagnoli-Coroglio industrial area (Gulf of Naples, Italy) rapidly become a key site for Italian economic growth, with several industrial plants producing steel, cement and asbestos, using fossil coal, iron ores and limestone as raw materials, transported by vessels and processed on site. In the mid-80s, the environmental risk of such activities was recognized, leading to a progressive dismantling of the industrial area, which ended in the mid-90s. However, the drastic impact of industrial activities was never remediated after the plant dismissal, with extremely high concentrations of polycyclic aromatic hydrocarbons (PAHs) and trace metals in sediments, especially close to the piers of the plant (Romano et al. 2004).

Nowadays it is widely recognized that the impact of chemical pollution should be evaluated by giving increasing importance to the assessment of biological effects of contaminants, and using an integrated approach with chemical data. The first example of integrated assessment was the Sediment Quality Triad (SQT), which considered chemical analyses, ecotoxicological testing and benthic communities as different Lines of Evidence (LOEs) to describe environmental quality of marine sediments (Chapman, 2007). Integrated strategies have also been proposed by various

63 international agencies, e.g. OSPAR, HELCOM, MEDPOL, ICES. An important advantage of  
64 these approaches is the added interpretative value derived from the integration of multiple  
65 typologies of studies, thus improving our ability to describe and interpret variations of  
66 environmental conditions (Regoli et al. 2019). The chemical approach by itself does not provide  
67 information on real bioavailability and biological risk of measured pollutants, often resulting in  
68 overestimated and costly management decisions (Bradham et al. 2006). Ecotoxicological batteries  
69 of bioassays have progressively been applied to quantify the potential biological hazard caused by  
70 bioavailable multi-factorial contamination, thus providing a more relevant response not restricted  
71 by a predetermined list of contaminants (Volpi Ghirardini et al. 2005). The benthic studies add  
72 information on the functioning of local communities and Ecological Quality Status Descriptors  
73 have been developed from these results (Dauvin 2015, Borja et al. 2016). In recent years,  
74 additional LOEs have been integrated in a weight of evidence (WOE) framework, such as  
75 bioaccumulation and biomarkers investigations. The bioaccumulation LOE quantifies the  
76 bioavailable fraction of contaminants, which can be transferred to aquatic organisms, being  
77 responsible for potential onset of adverse effects. Biomarkers reflect sub-lethal alterations at  
78 molecular and cellular level, representing a sensitive and early warning method to better  
79 understand the toxic effects and mechanism of action of environmental contaminants (Regoli &  
80 Giuliani 2014, Benedetti et al. 2015, Regoli et al. 2019). The WOE integration of chemical  
81 analysis, bioaccumulation, biomarkers, bioassays and analysis of benthic communities provides a  
82 more robust basis for environmental control and management in respect to the first SQT. In recent  
83 years this approach was synthesized in a quantitative model (SediquaSoft), validated in several  
84 case studies for environmental risk assessment associated with polluted sediments, harbor areas,  
85 or complex natural and anthropic impacts on the marine environment (Piva et al. 2011, Benedetti  
86 et al. 2012, 2014, Regoli et al. 2014, 2019, Bebianno et al. 2015, Mestre et al. 2017, Pittura et al.  
87 2018). In SediquaSoft different LOEs are independently elaborated, using specific criteria for  
88 each data, which weight typology of chemical pollutants and toxicological relevance of measured

endpoints, as well as the number and magnitude of observed variations normalized toward specific thresholds. Synthetic and quantitative hazard indices are calculated for each LOE, before their overall integration in the WOE assessment: the calculated level of risk is assigned to 1 of 5 classes, ranging from absent to severe (Piva et al. 2011, Regoli et al. 2019), which is the basis for different management options associated to each class of environmental risk. Weighted criteria for elaboration of chemical data and ecotoxicological bioassays have been incorporated in the last Italian law for determining quality class and management options for dredged marine sediments, based on the weighted elaboration and integration of their chemical and ecotoxicological characteristics (DM 173/2016).

As part of an extensive research project aimed to characterize the environmental quality and a possible remediation strategy of Bagnoli industrial site (ABBaCo 2018), the objective of the present work was to demonstrate the practical efficacy of the WOE approach to elaborate and integrate huge datasets of heterogeneous results. The combination of rigorous mathematical algorithms with the user-friendly outputs of the Sediqualssoft model was expected to represent an important tool to facilitate site-oriented and scientifically supported management options for sediments of such a polluted area.

## **2. Materials and methods**

### *2.1 Experimental design and sampling activities*

The study area is located in the eastern part of the Bay of Pozzuoli, within the Gulf of Naples. According to knowledge from previous studies (ISPRA-ICRAM 2005; Arienzo et al. 2017, 2019; Romano et al. 2018; Trifuoggi et al. 2017, 2018), it was divided in 11 sub-areas, each containing from 5 to 23 sampling points, depending on the distance from the industrial plant and the expected level of contamination (Figure 1). According to requirements of Italian Ministry of Environment and within “ABBaCo project”, 118 sediments samples were collected in the study

area between May 2017 and December 2017 for chemical analyses, ecotoxicological bioassays and analyses of benthic communities.

Bioaccumulation and biomarker responses were carried out on mussels, *Mytilus galloprovincialis*, and on the fish species, *Mullus barbatus*, *Pagellus erythrinus* and *Diplodus vulgaris*, sampled in December 2017. These fish species are commonly used in the biomonitoring of marine environment (Regoli et al. 2002; Bonsignore et al. 2013), and they were selected to highlight the potential influence of their different mobility, contact with sediments and feeding behavior on bioaccumulation and responsiveness to chemical pollutants. Organisms were sampled from different sites in the same period to avoid that comparisons of bioaccumulation and biomarkers responses could be differently influenced by seasonal variations (Bocchetti et al. 2006; Fattorini et al. 2014). Mussels (shell length  $5.5 \pm 0.5$  cm) were collected in 3 points along the 2 industrial piers (P2, PGT, PGP) and in additional 4 sites at different distances from the plant (Figure 1). Fish were sampled by local fishermen both in area of Bagnoli (INSIN) and in a reference site outside the bay (OUTSIN). Mean lengths were recorded both in fish from INSIN (*M. barbatus*:  $17.5 \pm 1.4$  cm; *P. erythrinus*:  $15.7 \pm 1.5$  cm; *D. vulgaris*:  $18.7 \pm 2.1$  cm) and from OUTSIN (*M. barbatus*:  $11.5 \pm 2.1$  cm; *P. erythrinus*:  $22.6 \pm 4.6$  cm; *D. vulgaris*:  $16.7 \pm 0.9$  cm). Analyses on fish samples (n=5) were performed using tissues of one individual for each replicate, while mussels samples (n=5) were constituted each by pooling tissues of 3 individuals.

## 2.2 Chemical characterization of sediments

After collection, sediment samples for chemical analyses were stored at -20 °C, until analysed for grain-size distribution (gravel, sand, silt, and clay), organic matter (OM), trace metals and metalloids (Al, As, Cd, Cr, Cu, Fe, Hg, Ni, Pb, V, and Zn), hydrocarbons with C>12, polycyclic aromatic polycyclic aromatic hydrocarbons (PAHs), polychlorobiphenyls (PCBs), organotin compounds, organochlorine pesticides, dioxin (PCDDs) and furan (PCDFs). Measurements were carried out through validated methods by sieves and laser particle sizer, inductively coupled

plasma-mass spectrometry (ICP-MS), inductively coupled plasma-optical emission spectroscopy (ICP-OES), Atomic Absorption Spectrophotometry, gas-chromatography with mass spectrometry and flame ionization detector (GC-MS, GC-FID), high resolution DFS mass spectrometer Dioxine B Thermo Fischer. For the determination of PCDDs and PCDFs the isotopic dilution procedures reported by USEPA 1613B (1994) were performed. Detailed analytical protocols, including QA/QC procedures, have been given elsewhere (Armiento et al., 2020; Molisso et al. 2020).

### 2.3 Bioaccumulation analyses

Bioaccumulation of trace elements (Al, As, Cd, Cu, Cr, Hg, Ni, Pb and Zn) and polycyclic aromatic hydrocarbons (PAHs) were carried out on the whole soft tissues of wild specimens of *M. galloprovincialis* and in the liver tissues of *M. barbatus*, *P. erythrinus* and *D. vulgaris*, following previously validated procedures based on atomic absorption spectrophotometry (AAS) and high performance liquid chromatography (HPLC) with diode array (DAD) and fluorimetric detection (Benedetti et al. 2014, Regoli et al. 2019). All the analytical determinations were performed by analysing five replicates, carefully checking for accuracy, precision and recovery by testing a series of blank solutions (reagents only), reference standards and selected certified standard materials.

### 2.4 Biomarkers analyses

Lysosomal membrane stability was measured in mussels hemocytes through the Neutral Red Retention Time (NRRT) assay, while metallothioneins (MTs) were determined in digestive glands by spectrophotometric analysis after acidic ethanol/chloroform fractionation using GSH as standard (Viarengo et al., 1997). Acetylcholinesterase enzymatic activity (AChE) was spectrophotometrically assayed in mussels haemolymph and fish brain using the Ellman's reaction (Gorbi et al. 2008). Ethoxyresorufin O-deethylase (EROD) was spectrofluorimetrically determined in individual fish livers measuring resorufin formation from 7-ethoxyresorufin



(Regoli et al. 2003). Aromatic metabolites in fish bile were measured by fixed fluorescence (FF) spectrofluorimetry and semi-quantitatively assessed as naphthalene-like, pyrene-like and benzo[a]pyrene-like metabolites (Gorbi and Regoli 2004). Micronuclei (MN) frequency was microscopically measured in mussels hemocytes and fish gills, observing 2000 cells with preserved cytoplasm: such genotoxic alterations were defined as round structures, smaller than 1/3 of the main nucleus diameter, on the same optical plan and clearly (Gorbi et al. 2008). Detailed analytical procedure have been previously reported (Benedetti et al., 2014).

## 2.5 Ecotoxicological bioassays

A battery of ecotoxicological bioassays was applied to sediment samples following standardized procedures. The bioluminescence test with *Vibrio fischeri* (Doe et al. 2005) was selected for the solid phase, while the algal growth inhibition of *Skeletonema costatum* (ISO 10253: 2006) and the sea urchin embryotoxicity assay with *Paracentrotus lividus* (ISPRA, 2017) were used to test elutriates, prepared with standard procedure (USEPA 503/8-91/001: 1991). Detailed procedures for sediments treatment and ecotoxicological bioassays have been reported elsewhere (Morrone et al. 2018; Gallo et al. 2020).

## 2.6 Benthic communities

Sediment samples were sieved through a mesh net of 1 mm and sorted under the stereomicroscope. All taxa were identified and the main taxa classified to the species level or to the lower possible taxonomic level. For each species, whenever possible, the corresponding biocoenosis was identified. From the list of species, the WOE SediquaSoft elaborated the available community descriptors, diversity indices and ecological indicators (including abundance, richness, Margalef, Shannon, Pielou, AMBI, BENTHIX, BOPA, BITS, mAMBI) which are reported in Table S4b. The results and the methodological details of these analyses have been reported in Hay Mele et al. (2020).

## 2.7 Statistical analyses and WOE elaboration

Analysis of variance (ANOVA) was applied to chemical data and biomarker responses of *M. galloprovincialis*, to test the significance of the differences between areas (level of significance at the 95% of confidence interval,  $\alpha=0.05$ ); homogeneity of variance was tested by Cochran C, and post-hoc comparisons (Student-Newman-Keuls) were used to compare means of values. Student's t-test was used to test for statistical significance at the 95% of confidence interval ( $\alpha=0.05$ ) between INSIN and OUTSIN areas for chemical data and biomarker responses on each of the three fish species.

All results, for various typologies of data, have been elaborated within the quantitative WOE, Sediqualsoft model, which consists in various modules to summarize specific hazard indices for individual LOEs, before their overall integration in the final WOE assessment (Piva et al. 2011, Benedetti et al. 2012, Regoli et al. 2014, 2019). Logical flow charts, based on expert judgment and legislative constraints, were converted into algorithms for weighted elaboration of data from sediment chemistry, bioavailability of chemicals in bioindicator species, ecotoxicological effects measured at subcellular level (biomarkers), toxicity at organism level (laboratory bioassays) and at the community level (benthic communities): the individual LOEs have been finally integrated for the WOE evaluation (see below).

#### *LOE 1: Chemical characterization of sediments*

The evaluation of chemical hazard (LOE-1) is initially based on the calculation for each pollutant of Ratio to Reference (*RTR*), i.e. the ratio between concentration measured in sediments and those indicated by a sediment quality guideline (SQG); in the present investigation, reference limits were those indicated by the SQG-L2 of the Italian decree for determining quality class and management options for dredged marine sediments (DM 173/2016). The *RTR* is corrected by a factor (*w*) which depend on the typology of chemicals (i.e. non priority  $w=1$ , priority  $w=1.1$ , priority and hazardous pollutants  $w=1.3$ ). In the calculation of the specific Hazard Quotient

217 ( $HQ_C$ ), an average  $RTR_w$  is obtained for all of the parameters with  $RTR \leq 1$  (i.e. values below the  
 218 SQG), while for those with  $RTR > 1$ , the  $RTR_w$  are individually added into the summation  $\Sigma$ :

$$219 \quad HQ_C = \frac{\sum_{j=1}^N RTR_w(j)_{RTR(j) \leq 1}}{N} + \sum_{k=1}^M RTR_w(k)_{RTR(k) > 1}$$

220 Based on expert judgment, the values of  $HQ_C$  are assigned to one of six classes of chemical  
 221 hazard, absent, negligible, slight, moderate, major and severe depending on the number, typology  
 222 and magnitude of exceeding chemicals (Regoli et al., 2019).

### 223 *LOE 2: Bioavailability of chemicals*

224 The results on bioaccumulation of chemicals in tissues of mussels and fish (LOE2) are elaborated  
 225 calculating, for each parameter, the increase of concentration compared to control organisms,  
 226 corrected for the typology of pollutant and the statistical significance of the difference. The  
 227 cumulative  $HQ_{BA}$  does not consider parameters with  $RTR_w < 1.3$ , calculates the average for those  
 228 with  $RTR_w$  ranging between 1.3 and 2.6, and adds the summation of all those with  $RTR_w \geq 2.6$ ):

$$229 \quad HQ_{BA} = \frac{\sum_{n=1}^j RTR_w(n)_{1.3 \leq RTR_w(j) < 2.6}}{j} + \sum_{n=1}^K RTR_w(n)_{RTR(k) \geq 2.6}$$

230 The  $HQ_{BA}$  is assigned to one of five classes of hazard for bioavailability, from Absent to Severe  
 231 (Regoli et al., 2019).

232

233 *LOE 3: Sublethal effects: Biomarkers* The module for the elaboration of biomarkers (LOE3)  
 234 contains a wide battery of responses, each assigned with a weight (based on the relevance of  
 235 biological endpoint) and a threshold indicative of changes of biological relevance. For each  
 236 biomarker, the measured variation is compared to the threshold (Table S5), corrected for

statistical significance and importance of biomarker (weight), and assigned to 1 of 5 classes of effect which are then differently weighted in the calculation of cumulative  $HQ_{BM}$ .

$$HQ_{BM} = \left( \frac{\sum_{j=1}^N Effect_w(j)_{1.5 < Effect(j) \leq 2.5}}{num\ biomarker_{1.5 < Effect(j) \leq 2.5}} + \sum_{k=1}^M Effect_w(k)_{Effect(j) > 2.5} \right)$$

According to the % distribution of biomarkers in the 5 classes, the level of cumulative  $HQ_{BM}$  is assigned to 1 of 5 classes of hazard: all the more relevant information are given in the model output (Regoli et al., 2019).

#### *LOE 4: Ecotoxicological Bioassays*

Weighted criteria to elaborate results from standardized ecotoxicological bioassays (LOE-4) are based on specific thresholds and weights assigned to each bioassay depending on the biological endpoint, tested matrix, time of exposure, and the possibility of hormetic responses.

In the module for ecotoxicological bioassays, the cumulative hazard quotient ( $HQ_{Battery}$ ) is obtained by the summation ( $\Sigma$ ) of the weighted effects ( $Ew$ ), i.e., the variations measured for each test compared to specific thresholds, corrected for the statistical significance of the difference ( $w$ ), biological importance of the endpoint and exposure conditions ( $w_2$ ):

$$HQ_{BATTERY} = \sum_{k=1}^N Effect_w(k) \cdot w_2$$

The  $HQ_{Battery}$  is normalized to a scale ranging from 0 to 10, where 1 is the battery threshold (when all the measured bioassays exhibit an effect equal to the threshold, 10 when all the assays exhibit 100% of effect); the  $HQ_{Battery}$  is then assigned to one of five classes of hazard, from Absent to Severe (Regoli et al., 2019).

#### *LOE 5: Benthic Communities*

Data on benthic communities are elaborated within a specific module (LOE-5), which converts the list of identified species in several available univariate and multivariate indices for the classification of ecological quality (Vincent 2002, Dauvin and Ruellet 2007, Muxika et al. 2007, Anderson et al. 2008, Mistri and Munari 2008, Sigovini et al. 2013). Such elaborated indices include total abundance (N), species richness (S), Shannon-Weaver Diversity Index ( $H'$ ), Margalef index (D), Pielou's evenness index (J), AZTI' Marine Biotic Index (AMBI), multimetric-AZTI Marine Biotic Index (m-AMBI), Benthic Index (BENTIX), Benthic Index based on Taxonomic Sufficiency (BITS) and Benthic Opportunistic Polychaetes Amphipods (BOPA index) (Regoli et al., 2019). In this work, the AMBI index was chosen for the integration with other LOEs in the final WOE elaboration of ecological risk.

#### *WOE integration*

The huge datasets of results elaborated from the 5 LOEs have been finally integrated through a WOE approach based on the quantitative model SediquaSoft. The quantitative hazard quotients (HQs) obtained for each LOEs are normalized to a common scale and given a different weight according to previously validated procedures (Piva et al., 2011; Lethonen et al., 2019; Regoli et al., 2019). LOE-2, summarizing bioavailability of chemicals in mussels and fish had a greater weight (w: 1.2) compared to LOE-1 assessing the presence of such compounds in the sediments (w: 1.0); at the biological level, a greater ecological relevance was assigned to LOE-5 on benthic communities (w: 1.3) compared to LOE-4 reflecting acute ecotoxicological effect at an organismal level (w: 1.2), or LOE-3 on sublethal effects at the cellular level (w: 1). An overall WOE level of risk is thus calculated and assigned to 1 of 5 classes of risk from Absent to Severe (Piva et al., 2011). Scientific criteria, validation of weights and thresholds, expert judgment evaluations and specific flow-charts of each LOE have been validated elsewhere (Piva et al., 2011; Benedetti et al., 2012, 2014; Lethonen et al., 2019; Regoli et al., 2019).

### **3. Results**

### 3.1 Chemical characterization of sediments

Chemical analyses were performed on sediments from the 11 sub-areas (Figure 1) and measured values are reported by Armiento et al (2020). Several critical values were obtained, especially for PAHs and trace metals (As, Zn, Pb, V and to a lesser extent Cd Cu and Hg). The highest concentrations of  $\Sigma$  PAHs (approximately 2800 mg/kg) were measured in some samples collected within the sub-area 7.

When data were elaborated according to weighted criteria, 100 samples on 118 exhibited a “Severe” chemical Hazard Quotient (HQ) indicating a widespread contamination in the whole investigated area (Figure 2a).

### 3.2 Bioaccumulation in mussels and fish

Concentrations of trace metals and PAHs measured in whole tissues of mussels *M. galloprovincialis* sampled from 7 sampling stations are shown in Supplementary Materials (Table S1a). Among inorganic elements, concentrations of As, Cd, Fe, Mn, Pb and V were typically higher in mussels sampled from the industrial piers (P2, PGT and PGP), exhibiting a gradual decrease in organisms collected at increasing distances from the industrial plant. .

Mussels from the piers (P2, PGT and PGP) were also characterized by elevated concentrations of high molecular weight (HMW) PAHs, which confirm the industrial origin of such bioavailable organic chemicals. In contrast, low molecular weight (LMW) PAHs did not exhibit significant differences between organisms from the piers or the other sampling sites (BRM, BBM and BNF). Results on bioaccumulation of chemicals in fish are given in the Supplementary Materials (Table S1b). *M. barbatus* and *D. vulgaris* exhibited comparable concentrations of trace metals and PAHs in specimens collected in the area of Bagnoli and in those from the reference site, while concentrations of HMW-PAHs (*M. barbatus*), LMW-PAHs and total PAHs (*D. vulgaris*) were significantly higher in organisms from the industrial area ( $p < 0.05$ ). Bioaccumulation of chemicals

in *P. erythrinus* did not reveal significant variations for all the analysed parameters comparing the specimens collected in the industrial and reference areas (Table S1b).

The weighted elaboration of these results summarized as “Major” the HQ for bioavailability in *M. galloprovincialis* from sub-area 1, essentially due to concentrations of HMW-PAHs (supplementary Table S1), while “Moderate” in mussels collected from the sub-areas 7 and 9 (Figure 2b); a lower level of HQ was assigned to bioavailability for mussels in sub-areas 2, 6, 10, 11 (Figure 2b).

Considering results obtained in fish species, the higher bioavailability HQ was elaborated for *M. barbatus*, appearing as “Moderate” in fish collected within the industrial sub-areas 3, 4, 5 and 8.

### 3.3 Biomarkers responses in mussels and fish

Biomarkers analyzed in native mussels revealed a higher sensitivity of mussels collected in sites PGP and PGT (included in sub-areas 1, 7 and 9) which exhibited a decreased lysosomal membrane stability and increased micronuclei frequency compared to organisms from other areas (Supplementary Table S2a). No variations were observed for the enzymatic activity of acetylcholinesterase in hemolymph, nor for metallothioneins in digestive gland of mussels from various sites (Supplementary Table S2a).

Results on acetylcholinesterase in brain, EROD enzymatic activity in liver, aromatic bile metabolites and frequency of branchial micronuclei in fish species (*M. barbatus*, *P. erythrinus* and in *D. vulgaris*) are reported in supplementary Table S2b. Acetylcholinesterase enzymatic activity was not affected in *M. barbatus* sampled in the industrial area while a decrease of this biomarker was observed in *P. erythrinus* and *D. vulgaris*. The EROD enzymatic activity was significantly induced in *M. barbatus* ( $p<0.05$ ) and *P. erythrinus* ( $p<0.05$ ) sampled in the industrial area compared to specimens from the reference site, while the cytochrome P450 biotransformation pathway was unaffected in *D. vulgaris*. At the same time, all the fish species exhibited higher levels of aromatic metabolites, particularly B[a]P-like and pyrene-like, in

organisms sampled in the industrial compared to reference area. The frequency of micronuclei significantly ( $p < 0.05$ ) increased in gills of *M. barbatus* from the industrial area.

Combining the weighted elaboration of biomarker results obtained in mussels and fish, it was possible to assign a hazard index in all the investigated sub-areas: such HQ resulted as “Major” in sub-areas 1, 3, 4, 5 and 8, “Moderate” in 9, “Slight” or “Absent” in the remaining sub-areas (Figure 2c).

### 3.4 Ecotoxicological bioassays

Ecotoxicological characteristics of the sediments, evaluated through a battery of three bioassays (*V. fischeri*, *S. costatum* and *P. lividus*), are detailed in Supplementary Table S3. The weighted elaboration revealed a “Slight” or “Absent” toxicity for most samples; a “Moderate” HQ was summarized in sub-areas 1, 2 and 7, where some individual samples exhibited a “Major” level of hazard (Figure 2d). The embryotoxicity of *P. lividus* was the most sensitive bioassay, often in combination with the inhibition of the algal growth in *S. costatum* (Supplementary Table S3).

### 3.5 Benthic communities

The analyses of benthic communities carried out in 15 stations allowed to identify 1796 organisms belonging to 164 taxonomic groups (Supplementary Table S4). The AMBI index was selected as the most appropriate for the study area and the results indicated a “Slight” or “Absent” level of HQ at this level of biological organization (Table 1, Figure 2e).

### 3.6 Weight of evidence integration

The elaborated WOE risk indices were “Moderate” for the majority of sub-areas (1-5, 7-9 with calculated values between 40.48 and 54.61), and “Slight” in sub-areas 6, 10, 11 (WOE values between 29.61 and 37.03; Table 1 and Figure 3).



#### 4. Discussion

The weight of evidence (WOE) approach, integrating individual lines of evidence through qualitative or quantitative methods, has been widely used in ecological and risk assessments to draw conclusions and justify selection of regulatory benchmarks (Linkov et al., 2009, 2015). Procedures for integration of different typologies of data must be quantitative and transparent for their acceptance in regulatory normative. As part of a decision-making process, various WOE methodologies have been recently formalized in different fields, e.g. by US-EPA (Linkov et al., 2009, 2011, 2015), European Food Safety Authority (Suter et al., 2017), or by the last Italian law on management of dredged sediments (DM 173/2016). The latter is based on the same weighting criteria for chemical analyses and ecotoxicological bioassays presented in this work. The combination of chemical and biological analyses, is recognized as an added value to the use of individual lines of evidence (LOEs), and in line with European Directives which recommend the use of multiple quality indicators for aquatic ecosystems (Lyons et al. 2010; Lethoten et al. 2014). The application of quantitative weighted criteria to process and integrate huge amounts of heterogeneous data from different LOEs allowed to summarize complex scientific information for an easier interpretation by policymakers or environmental managers (Piva et al. 2011; Borja et al. 2017; Regoli et al. 2019).

In the present study, a WOE assessment has been carried out in Bagnoli industrial site, chosen as model area to demonstrate the practical applicability in a complex environmental scenario of a model integrating chemical characterization of sediments, bioavailability of pollutants to key bioindicator species and the onset of effects at different levels of biological organization, from cellular responses to status of benthic communities.

Concentrations of trace metals and especially PAHs in sediments revealed a severe chemical contamination still present in the entire study area (Armiento et al 2020), with concentrations extremely higher than baseline levels and maximum limits indicated by Italian legislation for harbor-dredged sediments (L2 values, DM 173/2016).

386 The integrative approach of this study showed that toxicity of sediments, measured through  
387 standardized batteries of ecotoxicological bioassays, was often not in accordance with chemical  
388 characterization. In fact, despite an evident contamination, sediments from the area of Bagnoli  
389 inlet as well as those from the southern and the northern stations, showed low levels of acute  
390 toxicity; only some samples, particularly those collected close to the industrial plant, revealed  
391 evidence of a major acute toxicity, but the overall HQ for ecotoxicological bioassays resulted as  
392 “Moderate” in these sub-areas.

393 Despite the lack of an elevated acute toxicity, the bioavailability of contaminants was evident in  
394 terms of bioaccumulation of PAHs, and of HMW hydrocarbons, both in mussels and fish from the  
395 industrialized area. For this reason, the level of hazard for bioavailability was calculated as  
396 “Moderate” in the greatest part of the study area, with major effects in proximity of industrial  
397 piers. The significant accumulation of HMW-PAHs confirmed a pyrolytic origin of such  
398 pollution, related to combustion processes of the plant. A significant accumulation of these PAHs  
399 was observed also in liver of the benthic *M. barbatus*, indicating sediments as a still active source  
400 of these chemical contaminants to local biota.

401 Bioavailability of PAHs was further confirmed by biotransformation-related biomarkers, such as  
402 the induction of EROD enzymatic activity and the accumulation of aromatic metabolites in the  
403 bile of *M. barbatus* and *P. erythrinus*. The latter species confirmed the higher sensitivity of these  
404 biomarkers compared to tissue concentrations, in revealing bioavailability and metabolism of  
405 PAHs in fish. The comparison of the three fish species highlighted *M. barbatus* as the most  
406 affected by PAHs, both in terms of accumulation and of cellular responses related to their  
407 biotransformation; in *P. erythrinus* the lack of PAHs accumulation did not reflect a lack of  
408 exposure but the active metabolization and excretion of these chemicals, as indicated by the  
409 significant EROD induction and accumulation of aromatic bile metabolites. Finally, the slight  
410 PAHs effects in the more pelagic *D. vulgaris* further confirm the significant role of sediments in  
411 direct transfer of chemicals to benthic biota and, indirectly, to trophic webs.

Additional adverse effects of PAHs were revealed by lysosomal membrane destabilization in mussels, and the onset of genotoxic effects in both fish and mussels. The enhancement of micronuclei frequency is a well-known effect of PAHs, partly related to increased formation of oxyradicals during their metabolization pathway (Benedetti et al., 2015).

The overall biological significance of observed biomarkers responses in biota corresponded to a “Major” level of subcellular hazard in organisms collected from the industrial sub-areas.

Concerning the ecological effects on benthic communities, the application of AMBI index (Borja et al. 2000) allowed to classify as “Absent” the hazard in the samples collected in the majority of study area, except in sub-areas 4, 5, 10 and 7, located near the plant, where the impact was summarized as “Slight”. Such evaluation is in accordance with Fasciglione et al. (2016), who reported an unexpected evidence of biodiversity in this area, indicating a good functioning of local ecosystems at high levels of chemical contamination.

The overall integration of hazard quotients elaborated from sediment chemistry, bioassays, bioaccumulation, biomarkers and benthic communities provided a more holistic assessment of the environmental quality in the investigated area. The results indicate a clear pollution, especially near the piers. Beside a “Slight” level of hazard summarized in sub-areas 6, 10 and 11, the WOE elaboration increased to a “Moderate” level in all the other zones: interestingly the highest hazard coefficients were obtained near the plant: values close to the limit between “Moderate” and “Major” level of risk were obtained in sub-areas 1 and 7 (Table 1), further confirming the impact of discharged material from the industrial activities.

The possibility of converting complex scientific information into simple hazard indices easily understandable from policy makers and environmental managers could facilitate and orientate site-specific decisions on environmental sediment management. This study confirm the need of performing multidisciplinary approaches to assess the health status of marine ecosystems. In this perspective the use of the WOE integration, which combine and weight different kinds of data and analyses allows to better discriminate the presence of contaminants and their short or long-

term consequences. The importance of WOE models is particularly evident in complex environmental scenarios where apparently contrasting results are provided by various LOEs. The present study also highlights that different conclusions (with consequent different management scenarios) can be made on the basis of the analyses chosen: from a severe hazard derived from the use of the sediment contamination alone to the lack of hazard if we consider only the data from the analyses of the benthic assemblages. The multidisciplinary approach used here confirms that different typologies of pollutants are bioavailable and selectively transferred from sediments to local biota; such chemicals do not always exert acute toxicological effects, but can induce cellular responses that, being highly sensitive, might be prognostic of future adverse effects. Although the analysis of local sentinel organisms confirms that the contaminants of the sediments are still transferred to biota, the results of the WOE approach suggests that the management of the sediments from the industrial area of Bagnoli should not be based only on its chemical characterization.

## **Acknowledgments**

This study was supported by the project ABBaCo funded by the Italian Ministry for Education, University and Research (grant number C62F16000170001) and by the project MERCES (Marine Ecosystem Restoration in Changing European Seas), funded by the European Union's Horizon 2020 research and innovation program (grant agreement no. 689518).

## **6. References**

ABBaCo project: “Sperimentazioni pilota finalizzate al restauro ambientale e balneabilità del SIN Bagnoli-Coroglio” (2018)..Italian Ministry for Education, University and Research Grant number C62F16000170001. <http://www.szn.it/index.php/en/research/integrative-marine-ecology/research-projects-emi/abbaco>.

464 Anderson MJ., Gorley RN., Clarke KR. (2008). PERMANOVA for PRIMER: guide to software  
 465 and statistical methods. PRIMER-E Ltd., Plymouth, United Kingdom: 214 pp.

466 Armiento G., Caprioli R., Cerbone A., Chiavarini S., Crovato C., De Cassan M., De Rosa L.,  
 467 Montereali M.R., Nardi E, Nardi L., Pezza M., Proposito M., Rimauro J., Salerno A.,  
 468 Salluzzo A. Spaziani F. (2020). Current status of coastal sediments contamination in the  
 469 former industrial area of Bagnoli-Coroglio (Naples, Italy), G. Budillon, R. Delfanti and D.  
 470 Vicinanza eds. Special Issue of Chemistry and Ecology: Multidisciplinary Approach to the  
 471 Characterization of Marine Coastal Areas Subjected to Chronic Industrial Contamination.  
 472 Submitted.

473 Arienzo M., Donadio C., Mangoni O., Bolinesi F., Stanislao C., Trifuoggi M., Toscanesi M., Di  
 474 Natale G., Ferrara L. (2017). Characterization and source apportionment of polycyclic  
 475 aromatic hydrocarbons (PAHs) in the sediments of gulf of Pozzuoli (Campania, Italy). Mar  
 476 Poll Bull, 124:480–487.

477 Arienzo M., Toscanesi M., Trifuoggi M., Ferrara L., Stanislao C., Donadio C., Villari G., De  
 478 Vico G., Carella F. (2019). Contaminants bioaccumulation and pathological assessment in  
 479 *Mytilus galloprovincialis* in coastal waters facing the brownfield site of Bagnoli, Italy. Mar  
 480 Poll Bull 140:341-352.

481 Bebianno MJ, Pereira CG, Rey F, Cravo A, Duarte D, D’Errico G, Regoli F. (2015). Integrated  
 482 approach to assess ecosystem health in harbor areas. Sci Total Environ 514:92–107.

483 Benedetti, M., Bocchetti, R., Di Mento, R., Fattorini, D., Gorbi, S., Machella, N., Moltedo, G.,  
 484 Notti, A., Scarpato, A., Virno-Lamberti, C., Cicero, A.M., Regoli, F. (2006). Multimaker  
 485 approach with caged mussels *Mytilus galloprovincialis*, for monitoring the impact of  
 486 produced waters in the Adriatic sea. Mar Environ Res, 62: S377-S378.

487 Benedetti, M., Ciapriani, F., Piva, F., Onorati, F., Fattorini, D., Notti, A., Ausili, A., Regoli, F.  
 488 (2012). A multidisciplinary weight of evidence approach for classifying polluted sediments:

489 Integrating sediment chemistry, bioavailability, biomarkers responses and bioassays.  
 490 Environ Int 38:17–28.

491 Benedetti, M., Giuliani, M.E., Regoli, F. (2015). Oxidative metabolism of chemical pollutants in  
 492 marine organisms: Molecular and biochemical biomarkers in environmental toxicology.  
 493 Ann N Y Acad Sci 1340:8–19.

494 Benedetti, M., Gorbi, S., Fattorini, D., D’Errico, G., Piva, F., Pacitti, D., Regoli, F. (2014).  
 495 Environmental hazards from natural hydrocarbons seepage: Integrated classification of risk  
 496 from sediment chemistry, bioavailability and biomarkers responses in sentinel species.  
 497 Environ Pollut 185:116–126.

498 Bocchetti, R., Regoli, F. (2006). Seasonal variability of oxidative biomarkers, lysosomal  
 499 parameters, metallothioneins and peroxisomal enzymes in the Mediterranean mussel  
 500 *Mytilus galloprovincialis* from Adriatic Sea. Chemosphere 65, 913–921.

501 Bonsignore, M., Salvagio Manta, D., Oliveri, E., Sprovieri, M., Basilone, G., Bonanno, A., Falco,  
 502 F., Traina, A., Mazzola, S. (2013). Mercury in fish from Augusta Bay (southern Italy): risk  
 503 assessment and health implication. Food Chem. Toxicol. 56, 184-194.

504 Borja, A., Franco, J., and Pérez, V. (2000). A marine Biotic Index to establish the ecological  
 505 quality of soft-bottom benthos within European estuarine and coastal environments. Mar  
 506 Pollut Bull 40:1100–1114.

507 Borja, A., Elliott, M., Andersen, J.H., Berg, T., Carstensen, J., Halpern, B.S., Heiskanen, A.S.,  
 508 Korpinen, S., Stewart Lowndes, J.S., Martin, G., Rodriguez-Ezpeleta, N. (2016). Overview  
 509 of integrative assessment of marine systems: The ecosystem approach in practice. Front  
 510 Mar Sci 3:1–20.

511 Borja, A., Elliott, M., Uyarra, M.C., Carstensen, J., Mea, M. (2017). Editorial: Bridging the gap  
 512 between policy and science in assessing the health status of marine ecosystems. Front Mar  
 513 Sci 4:1–3.

514 Bradham, C., Foltz, K.R., Beane, W.S., Arnone, M.I., Rizzo, F., Coffman, J., Mushegian, A.,  
515 Goel, M., Morales, J., Geneviere, A.M., Lapraz, F., Robertson, A.J., Kelkar, H., Loza-Coll,  
516 M., Townley, I.K., Raisch, M., Roux, M.M., Lepage, T., Gache, C., McClay, D.R.,  
517 Manning, G. (2006). The sea urchin kinome: a first look. *Dev Biol* 300:180–93.

518 Chapman, P.M. (2007). Determining when contamination is pollution — Weight of evidence  
519 determinations for sediments and effluents. *Environ Int* 33:492–501.

520 Dauvin, J. C., and Ruellet, T. (2007) Polychaete/amphipod ratio revisited. *Mar Pollut Bull*  
521 55:215–224.

522 Dauvin, J.C. (2015). History of benthic research in the English Channel: From general patterns of  
523 communities to habitat mosaic description. *J Sea Res* 100:32–45.

524 DM 173/2016. Ministero dell’Ambiente e della Tutela del Territorio e del Mare, Supplemento  
525 ordinario alla Gazzetta Ufficiale, n. 208 del 6 settembre 2016-Serie generale. Regolamento  
526 recante modalità e criteri tecnici per l’autorizzazione all’immersione in mare dei materiali di  
527 escavo di fondali marini.

528 Doe, K., Jackman, P, Scroggins, R, McLeay, D, Wohlgeschaffen, G. (2005). Solid-Phase Test for  
529 Sediment Toxicity Using the Luminescent Bacterium, *Vibrio Fischeri*. In: Blaise C., Férard  
530 JF. (eds) *Small-scale Freshwater Toxicity Investigations*. Springer, Dordrecht.

531 Fasciglione, P., Barra, M., Santucci, A., Ciancimino, S., Mazzola, S., Passaro, S. (2016).  
532 Macrobenthic community status in highly polluted area: a case study from Bagnoli,  
533 Naples Bay, Italy. *Rend. Fis. Acc. Lincei* 27:229-239.

534 Fattorini, D, Sarkar, S.K., Regoli, F, Bhattacharya, B.D., Rakshit, D. et al. (2013). Levels and  
535 chemical speciation of arsenic in representative biota and sediments of a tropical mangrove  
536 wetland, India. *Environ Sci Process Impacts* 15: 773-782.

537 Gallo, A., Guida, M., Armiento, G., Siciliano, A., Carraturo, F, Mormile, N, Pellegrini, D,  
538 Morroni, L, Tosti, E., Ferrante, M., Montresor, M., Molisso, F., Sacchi, M., Danovaro, R.,

539 Libralato, G. (2020). Species-specific sensitivity of three microalgae to sediment elutriates.  
 540 Mar Environ Res, this issue.

541 Gorbi, S., Regoli, F. (2004). Induction of cytochrome P4501A and biliary PAH metabolites in  
 542 European eel *Anguilla anguilla*: Seasonal, dose- and time-response variability in field and  
 543 laboratory conditions. Mar Environ Res 58 (2-5):511-515.

544 Gorbi, S., Virno Lamberti, C., Notti, A., Benedetti, M., Fattorini, D., Molledo, G., Regoli, F.  
 545 (2008). An ecotoxicological protocol with caged mussels, *Mytilus galloprovincialis*, for  
 546 monitoring the impact of an offshore platform in the Adriatic sea. Mar Environ Res 65  
 547 (1):34-49.

548 Hay Mele, B., Russo, B., Crocetta, F., Gambi, C., Dell'Anno, A., Danovaro, R., Guglielmo, R.,  
 549 Musco, L., Patti, F.P., Riginella, E., Tangherlini, M., Ribera d'Alcala', M., D'Alelio, D.  
 550 (2020). Ecological assessment of anthropogenic impact in marine ecosystems: the case of  
 551 Bagnoli Bay. Mar Environ Res, this issue.

552 ISO, 2006. Water quality — Marine algal growth inhibition test with *Skeletonema* sp. and  
 553 *Phaeodactylum tricornutum*. ISO 10253:2006.

554 ISPRA-ICRAM (2005). Caratterizzazione ambientale dei fondali e degli arenili inclusi nella  
 555 perimetrazione del sito di bonifica di Bagnoli–Coroglio. BoI-Pr-CA-BA-relazione-02.04.

556 ISPRA (2011). Procedura per la determinazione degli idrocarburi con C>12 nei suoli contaminati.  
 557 Manuali e linee guida, 75/2011.

558 ISPRA (2017). Saggio di fecondazione e saggio di sviluppo embrionale con il riccio di mare  
 559 *Paracentrotus lividus* (Lamarck) (Echinodermata: Echinoidea). Rome: ISPRA.

560 Lehtonen, K. K., Sundelin, B., Lang, T., and Strand, J. (2014). Development of tools for  
 561 integrated monitoring and assessment of hazardous substances and their biological effects  
 562 in the Baltic Sea. AMBIO 43:69–81.



563 Linkov, I., Long, E.B., Cormier, S.M., Satterstrom, F.K., Bridges, T. (2009). Weightof-evidence  
 564 evaluation in environmental assessment: Review of qualitative and quantitative approaches.  
 565 Sci Total Environ 497:5199–5205.

566 Linkov, I., Welle, P., Loney, D., Tkachuk, A., Canis, L., Kim, J.B., Bridges, T. (2011). Use of  
 567 multicriteria decision analysis to support weight of evidence evaluation. Risk Anal  
 568 doi:10.1111/j.1539-6924.2011.01585.

569 Linkov, I., Massey, O., Keisler, J., Rusyn, I., Hartung, T. (2015). From “ weight of evidence to  
 570 quantitative data integration using multicriteria decision analysis and Bayesian methods”.  
 571 Altex 32:3–8.

572 Lyons, B.P., Thain, J.E., Stentiford, G.D., Hylland, K., Davies, I.M., Vethaak, A.D. (2010). Using  
 573 biological effects tools to define Good Environmental Status under the European Union  
 574 Marine Strategy Framework Directive. Mar Pollut Bull 60:1647-1651.

575 Lowe, D.M., Pipe, R.K. (1994). Contaminant induced lysosomal membrane damage in marine  
 576 mussel digestive cells: an in vitro study. Aquat Toxicol, 30(4):357-365.

577 Mestre, N.C., Rocha, T.L., Canals, M., Cardoso, C., Danovaro, R., Dell’Anno, A., Gambi, C.,  
 578 Regoli, F., Sanchez-Vidal, A., Bebianno, M.J. (2017). Environmental hazard assessment of  
 579 a marine mine tailings deposit site and potential implications for deep-sea mining. Environ  
 580 Pollut 228:169–178.

581 Mistri, M., and Munari, C. (2008). BITS: a SMART indicator for soft-bottom, nontidal lagoons.  
 582 Mar Pollut Bull 56:587–599.

583 Molisso, F., Capodanno, M., Di Gregorio, C., Gilardi, M., Guarino, A., Oliveri, E., Tamburrino  
 584 S., Sacchi, M. (2020). Sedimentological analysis of marine deposits off the Bagnoli-  
 585 Coroglio Site of National Interest (SNI), Pozzuoli (Napoli) Bay. G. Budillon, R. Delfanti  
 586 and D. Vicinanza eds. Special Issue of Chemistry and Ecology: Multidisciplinary Approach  
 587 to the Characterization of Marine Coasta Areas Subjected to Chronic Industrial  
 588 Contamination. Submitted.

589 Morroni, L., Pinsino, A., Pellegrini, D., Regoli, F. (2018). Reversibility of metal induced  
590 malformations in sea urchin embryos. *Ecotoxicol Environ Saf* 148:923-929.

591 Muxika, I., Borja, Á., and Bald, J. (2007). Using historical data, expert judgement and  
592 multivariate analysis in assessing reference conditions and benthic ecological status,  
593 according to the European Water Framework Directive. *Mar Pollut Bull* 55:16–29.

594 Pittura, L., Avio, C.G., Giuliani, M.E., D’Errico, G., Keiter, S.H., Cormier, B., Gorbi, S., Regol,  
595 F. (2018). Microplastics as vehicles of environmental PAHs to marine organisms:  
596 Combined chemical and physical hazards to the mediterranean mussels, *Mytilus*  
597 *galloprovincialis*. *Front Mar Sci* 5(103).

598 Piva, F., Ciaprin, F., Onorati, F., Benedetti, M., Fattorini, D., Ausili, A., Regoli, F. (2011).  
599 Assessing sediment hazard through a weight of evidence approach with bioindicator  
600 organisms: A practical model to elaborate data from sediment chemistry, bioavailability,  
601 biomarkers and ecotoxicological bioassays. *Chemosphere* 83:475–485.

602 Regoli, F., Pellegrini, D., Winston, G.W., Gorbi, S., Giuliani, S., Virno-Lamberti, C., Bompadre,  
603 S., (2002). Application of biomarkers for assessing the biological impact of dredged  
604 materials in the Mediterranean: the relationship between antioxidant responses and  
605 susceptibility to oxidative stress in the red mullet (*Mullus barbatus*). *Mar. Pollut. Bull.* 44,  
606 912–922

607 Regoli, F., Winston, G.W., Gorbi, S., Frenzilli, G., Nigro, M., Corsi, I., Focardi, S. (2003).  
608 Integrating enzymatic responses to organic chemical exposure with total oxyradical  
609 absorbing capacity and DNA damage in the European eel *Anguilla anguilla*. *Environ*  
610 *Toxicol Chem* 22 (9):2120-2129.

611 Regoli, F. and Giuliani, M.E. (2014). Oxidative pathways of chemical toxicity and oxidative  
612 stress biomarkers in marine organisms. *Mar Environ Res* 93:106–117.

613 Regoli, F., Pellegrini, D., Cicero, A.M., Nigro, M., Benedetti, M., Gorbi, S., Fattorini, D.,  
614 D’Errico, G., Di Carlo, M., Nardi, A., Gaion, A., Scuderi, A., Giuliani, S., Romanelli, G.,

615 Berto, D., Trabucco, B., Guidi, P., Bernardeschi, M., Scarcelli, V., Frenzilli, G. (2014). A  
616 multidisciplinary weight of evidence approach for environmental risk assessment at the  
617 Costa Concordia wreck: Integrative indices from Mussel Watch. *Mar Environ Res* 96:92–  
618 104.

619 Regoli, F., D’Errico, G., Nardi, A., Mezzelani, M., Fattorini, D., Benedetti, M., Di Carlo, M.,  
620 Pellegrini, D., Gorbi, S. (2019). Application of a Weight of Evidence Approach for  
621 Monitoring Complex Environmental Scenarios: the Case-Study of Off-Shore Platforms.  
622 *Front Mar Sci* 6:1–15.

623 Romano, E., Ausili, A., Zharova, N., Celia Magno, M., Pavoni, B., Gabellini, M. (2004). Marine  
624 sediment contamination of an industrial site at Port of Bagnoli, Gulf of Naples, Southern  
625 Italy. *Mar Pollut Bull* 49:487–495.

626 Romano E., Bergamin L., Pierfranceschi G., Ausili A. (2018). Temporal changes of metal and  
627 trace element contamination in marine sediments due to a steel plant: The case study of  
628 Bagnoli (Naples, Italy). *Appl Geochem* 88:85-94.

629 Sigovini, M., Keppel, E., and Tagliapietra, D. (2013). M-AMBI revisited: looking inside a  
630 widely-used benthic index. *Hydrobiologia* 717:41–50.

631 Suter, G.W., Cormier, S.M., Barron, M.G. (2017). A weight of evidence framework for  
632 environmental assessments: Inferring qualities. *Integr Environ Assess Manag* 13:1038–  
633 1044.

634 Trifuoggi M., Donadio C., Ferrara L., Stanislao C., Toscanesi M., Arienzo M. (2018). Levels of  
635 pollution of rare earth elements in the surface sediments from the Gulf of Pozzuoli  
636 (Campania, Italy). *Mar Poll Bull* 136:374–384.

637 Trifuoggi M., Donadio C., Mangoni O., Ferrara L., Bolinesi F., Nastro R.A., Stanislao C.,  
638 Toscanesi M., Di Natale G., Arienzo M. (2017). Distribution and enrichment of trace metals  
639 in surface marine sediments in the Gulf of Pozzuoli and off the coast of the brownfield  
640 metallurgical site of Ilva of Bagnoli (Campania, Italy). *Mar Poll Bull* 124:502–511.

641 USEPA (1991). Evaluation of dredged material proposed for ocean disposal testing manual. EPA  
642 503/8-91/001.

643 USEPA (1994). Tetra-through Octa-Chlorinated Dioxins and Furans by Isotope Dilution  
644 HRGC/HRMS. 1613.

645 Viarengo, A., Ponzano, E., Dondero, F., Fabbri, R. (1997). A simple spectrophotometric method  
646 for metallothionein evaluation in marine organisms: An application to Mediterranean and  
647 Antarctic molluscs. *Mar Environ Res* 44 (1):69-84.












648 Vincent, C., Heinrich, H., Edwards, A., Nygaard, K., Haythornthwaite, J. (2002). Guidance on  
649 typology, reference conditions and classification systems for transitional and coastal waters,  
650 CIS Working Group 2.4(Coast) Common Implementation Strategy of the Water  
651 Framework Directive, European Commission.

652 Volpi Ghirardini, A, Arizzi Novelli, A., Tagliapietra, D. (2005). Sediment toxicity assessment in  
653 the Lagoon of Venice (Italy) using *Paracentrotus lividus* (Echinodermata: Echinoidea)  
654 fertilization and embryo bioassays. *Environ Int* 31:1065–77.

655

## Tables

**Table 1** - Classification of sub-areas within Bagnoli-Coroglio industrial site, according to the weight of evidence elaboration. Levels of hazard are reported for each LOE and for their overall WOE integration.

Area	LOE1 (Level of chemical hazard)	LOE2 (Level of hazard for bioavailability)	LOE3 (Level of hazard for biomarkers)	LOE4 (Level of hazard for bioassays)	LOE5 (Level of hazard for Benthic Communities)	WOE (Weight of Evidence integration)	
Area 1	SEVERE	MAJOR	MAJOR	MODERATE	ABSENT	MODERATE (54.61)	
Area 2	SEVERE	SLIGHT	SLIGHT	MODERATE	ABSENT	MODERATE (40.48)	
Area 3	SEVERE	MODERATE	MAJOR	ABSENT	ABSENT	MODERATE (43.40)	
Area 4	SEVERE	MODERATE	MAJOR	ABSENT	SLIGHT	MODERATE (45.36)	
Area 5	SEVERE	MODERATE	MAJOR	ABSENT	SLIGHT	MODERATE (46.54)	
Area 6	SEVERE	ABSENT	ABSENT	ABSENT	ABSENT	SLIGHT (29.61)	
Area 7	SEVERE	MODERATE	MAJOR	MODERATE	SLIGHT	MODERATE (53.56)	
Area 8	SEVERE	MODERATE	MAJOR	ABSENT	ABSENT	MODERATE (41.30)	
Area 9	SEVERE	MODERATE	MODERATE	ABSENT	ABSENT	MODERATE (41.73)	
Area 10	SEVERE	SLIGHT	SLIGHT	ABSENT	ABSENT	SLIGHT (34.83)	
Area 11	SEVERE	ABSENT	ABSENT	SLIGHT	ABSENT	SLIGHT (37.03)	

670 **Legends of Figures**

671 **Figure 1** - Localization of sampling sites in the Bagnoli-Coroglio industrial area. In each sub-area  
672 were performed: chemical characterization of sediments; bioavailability of trace metals and  
673 organic pollutants in mussels and fish, biomarkers in mussels and fish, benthic communities  
674 analysis; ecotoxicological bioassays.

675

676 **Figure 2** - Classification of sub-areas within Bagnoli-Coroglio industrial area for each LOE  
677 according to the weight of evidence elaboration. Levels of hazard are reported in different colors  
678 on the areas, as detailed in the legend, considering as LOE: chemical characterization of  
679 sediments (A); bioavailability of trace metals and organic pollutants in mussels and fish (B),  
680 biomarkers in mussels and fish (C); ecotoxicological bioassays (D); benthic communities analysis  
681 (E).


682

683 **Figure 3** – Classification of areas within Bagnoli-Coroglio industrial area considering the overall  
684 weight of evidence integration. Levels of hazard are reported in different colors on the areas as  
685 detailed in the legend.

686

687 **Figure 1**


688



689

690


691 **Figure 2**  
692



693  
694  
695



696 **Figure 3**  
697



698  
699

## SUPPLEMENTARY MATERIAL

**Table S1a.** Bioaccumulation of trace metals and polycyclic aromatic hydrocarbons (PAHs) concentrations in tissues of mussels sampled. Data expressed as mean  $\pm$  standard deviation. Different letters (a, b, c) indicate significant differences ( $p < 0.05$ ) between sites, as determined by one way analyses of variance (ANOVA) and Student Newman Keuls post-hoc test.

		PGT (Area 1;9)	PGP (Area 1;7)	P2 (Area 2;10)	BBM (Area 6;9;11)	BNF (Area 8)	BRM (Reference area)	Sig.
Al	$\mu\text{g/g (dw)}$	$85 \pm 15.4$	$61.6 \pm 5.48$	$52.6 \pm 15.4$	$58.2 \pm 17$	$49.2 \pm 47.5$	$46.5 \pm 14.7$	<i>n.s.</i>
As		$22.4 \pm 5.87$ <i>bc</i>	$25.6 \pm 2.29$ <i>c</i>	$22.2 \pm 0.783$ <i>bc</i>	$17.1 \pm 1.81$ <i>ab</i>	$11.9 \pm 1.86$ <i>a</i>	$19.8 \pm 0.827$ <i>bc</i>	$p < 0.0001$
Cd		$0.873 \pm 0.031$ <i>c</i>	$0.83 \pm 0.0943$ <i>c</i>	$0.671 \pm 0.0924$ <i>b</i>	$0.579 \pm 0.107$ <i>b</i>	$0.394 \pm 0.0692$ <i>a</i>	$0.519 \pm 0.014$ <i>ab</i>	$p < 0.0001$
Cr		$2.08 \pm 0.315$ <i>c</i>	$1.65 \pm 0.263$ <i>b</i>	$1.17 \pm 0.0242$ <i>a</i>	$1.17 \pm 0.127$ <i>a</i>	$1.1 \pm 0.221$ <i>a</i>	$0.944 \pm 0.0518$ <i>a</i>	$p < 0.0001$
Cu		$6.88 \pm 0.872$ <i>bc</i>	$4.07 \pm 1.95$ <i>ab</i>	$7.88 \pm 2.89$ <i>bc</i>	$6.93 \pm 1$ <i>c</i>	$1.19 \pm 0.712$ <i>a</i>	$8.68 \pm 1.16$ <i>c</i>	$p < 0.005$
Fe		$626 \pm 174$ <i>b</i>	$604 \pm 96.5$ <i>b</i>	$362 \pm 95.5$ <i>a</i>	$340 \pm 103$ <i>a</i>	$230 \pm 118$ <i>a</i>	$218 \pm 43.4$ <i>a</i>	$p < 0.005$
Hg		$0.171 \pm 0.011$ <i>b</i>	$0.17 \pm 0.00796$ <i>b</i>	$0.157 \pm 0.0326$ <i>b</i>	$0.146 \pm 0.0123$ <i>b</i>	$0.11 \pm 0.0115$ <i>a</i>	$0.139 \pm 0.0112$ <i>b</i>	$p < 0.005$
Mn		$68.1 \pm 11.7$ <i>c</i>	$62 \pm 9$ <i>c</i>	$42.2 \pm 12.2$ <i>b</i>	$33.7 \pm 1.83$ <i>ab</i>	$14.8 \pm 4.53$ <i>a</i>	$31.7 \pm 4.04$ <i>ab</i>	$p < 0.0001$
Ni		$1.1 \pm 0.0932$ <i>b</i>	$0.997 \pm 0.0895$ <i>ab</i>	$0.718 \pm 0.0991$ <i>a</i>	$1.32 \pm 0.259$ <i>b</i>	$0.725 \pm 0.164$ <i>a</i>	$1.22 \pm 0.0497$ <i>b</i>	$p < 0.0001$
Pb		$10.4 \pm 0.585$ <i>c</i>	$10.5 \pm 0.484$ <i>c</i>	$6.16 \pm 0.655$ <i>b</i>	$4.1 \pm 0.705$ <i>a</i>	$3.57 \pm 2.32$ <i>a</i>	$2.9 \pm 0.569$ <i>a</i>	$p < 0.0001$
V		$2.65 \pm 0.531$ <i>c</i>	$2.42 \pm 0.218$ <i>bc</i>	$1.68 \pm 0.326$ <i>ab</i>	$1.52 \pm 0.472$ <i>a</i>	$1.33 \pm 0.387$ <i>a</i>	$1.43 \pm 0.259$ <i>a</i>	$p < 0.005$
Zn		$53.2 \pm 11$	$90.7 \pm 74.5$	$41.1 \pm 6.91$	$123 \pm 116$	$38.7 \pm 5.54$	$42.2 \pm 20.4$	<i>n.s.</i>

n.s. = not significant

Table S1a. Continues.

		PGT	PGP	P2	BBM	BNF	BRM	Sig.
		(Area 1;9)	(Area 1;7)	(Area 2;10)	(Area 6;9;11)	(Area 8)	(Reference area)	
Naphthalene	ng/g (dw)	18.7 ± 3.19	35.5 ± 24.1	18.2 ± 5.99	14.1 ± 0.932	16.1 ± 0.508	16.6 ± 0.965	<i>n.s.</i>
Acenaphthylene		< 0.05	< 0.05	< 0.05	< 0.05	< 0.05	33.8 ± 58.5	<i>n.s.</i>
1-methylnaphthalene		< 0.05	< 0.05	1.96 ± 3.3 <i>a</i>	16 ± 1.88 <i>b</i>	17.3 ± 1.94 <i>b</i>	18.4 ± 2.57 <i>b</i>	<i>p</i> < 0.0001
2-methylnaphthalene		< 0.05	< 0.05	2.58 ± 4.38 <i>a</i>	10.8 ± 1.67 <i>b</i>	8.14 ± 0.336 <i>b</i>	9.42 ± 0.391 <i>b</i>	<i>p</i> < 0.0001
Acenaphthene		< 0.01	< 0.01	< 0.01	< 0.01	0.981 ± 0.0632 <i>b</i>	< 0.01	<i>p</i> < 0.005
Fluorene		0.778 ± 0.138 <i>ab</i>	0.832 ± 0.186 <i>ab</i>	1.13 ± 0.229 <i>b</i>	0.596 ± 0.136 <i>a</i>	0.68 ± 0.204 <i>ab</i>	0.684 ± 0.206 <i>ab</i>	<i>p</i> < 0.05
Phenanthrene		< 0.01	< 0.01	4.31 ± 4.51	4.04 ± 0.723	4.46 ± 0.689	4.18 ± 0.381	<i>n.s.</i>
Anthracene		2.74 ± 2.67	0.775 ± 1.32	0.768 ± 0.694	0.057 ± 0.0567	< 0.01	< 0.01	<i>n.s.</i>
Fluoranthene		10 ± 17.3	4.63 ± 8.01	0.225 ± 0.372	< 0.01	< 0.01	< 0.01	<i>n.s.</i>
Pyrene		34.5 ± 28.6	10 ± 16.2	7.11 ± 6.13	1.39 ± 0.433	0.379 ± 0.082	0.342 ± 0.291	<i>n.s.</i>
Benzo(a)anthracene		13.2 ± 9.95 <i>b</i>	4.02 ± 6.95 <i>a</i>	1.64 ± 2.71 <i>a</i>	< 0.01	0.016 ± 0.0103 <i>a</i>	0.0766 ± 0.115 <i>a</i>	<i>p</i> < 0.05
Chrysene		9.93 ± 7.03 <i>b</i>	2.95 ± 5.1 <i>a</i>	< 0.01	< 0.01	< 0.01	0.354 ± 0.119 <i>a</i>	<i>p</i> < 0.05
7,12-dimethylbenzo(a)anthracene		10.9 ± 15.7	4.88 ± 8.38	2.29 ± 1.95	3.68 ± 4.02	1.59 ± 0.779	1.21 ± 1.04	<i>n.s.</i>
Benzo(b)fluoranthene		13.6 ± 9.71	10.1 ± 9.37	0.0446 ± 0.0756	0.306 ± 0.0459	0.179 ± 0.01	0.321 ± 0.106	<i>n.s.</i>
Benzo(k)fluoranthene		8.78 ± 9.5	7.05 ± 6.11	0.0273 ± 0.0275	0.401 ± 0.316	0.0706 ± 0.0366	0.307 ± 0.0646	<i>n.s.</i>
Benzo(a)pyrene		5.99 ± 3.88 <i>b</i>	2.66 ± 2.36 <i>a</i>	1.13 ± 0.893 <i>a</i>	0.321 ± 0.0481 <i>a</i>	0.0946 ± 0.0465 <i>a</i>	0.142 ± 0.0989 <i>a</i>	<i>p</i> < 0.05
Dibenzo(ah)anthracene		0.677 ± 0.391	14 ± 18.8	0.454 ± 0.327	0.094 ± 0.161	0.189 ± 0.0267	0.236 ± 0.0669	<i>n.s.</i>
Benzo(ghi)perylene		0.227 ± 0.0937	0.158 ± 0.271	0.105 ± 0.0136	0.032 ± 0.0536	0.0346 ± 0.0583	< 0.001	<i>n.s.</i>
Indeno(123cd)pyrene		1.09 ± 1.8	0.043 ± 0.00608	< 0.05	0.0426 ± 0.0234	0.0393 ± 0.0184	< 0.05	<i>n.s.</i>
Low MW PAHs		43.5 ± 9.17 <i>a</i>	72.1 ± 47.4 <i>a</i>	57.8 ± 24.1 <i>a</i>	99.3 ± 5.56 <i>a</i>	99.4 ± 5.76 <i>a</i>	170 ± 113 <i>b</i>	<i>p</i> < 0.05
High MW PAHs		172 ± 123 <i>b</i>	121 ± 30.5 <i>b</i>	32.7 ± 10.5 <i>a</i>	13.3 ± 8.24 <i>a</i>	5.31 ± 1.67 <i>a</i>	6.07 ± 2.23 <i>a</i>	<i>p</i> < 0.05
Total PAHs		216 ± 129	194 ± 67.6	90.5 ± 27	112 ± 11.4	104 ± 6.33	176 ± 115	<i>n.s.</i>

n.s. = not significant

**Table S1b.** Bioaccumulation of trace metals and polycyclic aromatic hydrocarbons (PAHs) concentrations in tissues of fish sampled. Data expressed as mean  $\pm$  standard deviation. Different letters (a, b, c) indicate significant differences ( $p < 0.05$ ) between sites, as determined by one way analyses of variance (ANOVA) and Student Newman Keuls post-hoc test. Significant differences between areas and reference area ( $p < 0.05$ ), as determined by t-student test.

		<i>Mullus barbatus</i>			<i>Diplodus vulgaris</i>			<i>Pagellus sp.</i>		
		OUTSIN	INSIN	Sig.	OUTSIN	INSIN	Sig.	OUTSIN	INSIN	Sig.
Al	$\mu\text{g/g (dw)}$	n/a	22.3 $\pm$ 14.4	n.t.	12.6 $\pm$ 7.34	18 $\pm$ 7.43	n.s.	24.7 $\pm$ 26.9	36.6 $\pm$ 32.5	n.s.
As		61.1 $\pm$ 8.3	58.6 $\pm$ 15.9	n.s.	17.2 $\pm$ 0.936	20.6 $\pm$ 6.46	n.s.	13.7 $\pm$ 3.74	9.05 $\pm$ 1.48	$p < 0.05$
Cd		0.272 $\pm$ 0.156	0.4 $\pm$ 0.161	n.s.	0.313 $\pm$ 0.0418	1.23 $\pm$ 0.617	$p < 0.05$	0.657 $\pm$ 0.302	0.671 $\pm$ 0.571	n.s.
Cr		0.0573 $\pm$ 0.0279	0.647 $\pm$ 0.323	$p < 0.05$	1.39 $\pm$ 0.808	0.674 $\pm$ 0.45	$p < 0.05$	0.585 $\pm$ 0.274	0.633 $\pm$ 0.324	n.s.
Cu		16.9 $\pm$ 3.11	9.89 $\pm$ 2.61	$p < 0.05$	25.7 $\pm$ 6.17	32.1 $\pm$ 7.42	n.s.	25.6 $\pm$ 11.9	26.1 $\pm$ 13	n.s.
Fe		366 $\pm$ 150	646 $\pm$ 223	$p < 0.05$	1148 $\pm$ 357	1069 $\pm$ 328	n.s.	536 $\pm$ 111	538 $\pm$ 255	n.s.
Hg		n/a	0.372 $\pm$ 0.216	n.t.	0.611 $\pm$ 0.215	0.9 $\pm$ 0.552	n.s.	0.725 $\pm$ 0.208	0.62 $\pm$ 0.335	n.s.
Mn		n/a	13.3 $\pm$ 2.25	n.t.	10.1 $\pm$ 2.2	18.1 $\pm$ 7.1	$p < 0.05$	15.3 $\pm$ 3.89	13.1 $\pm$ 5.2	n.s.
Ni		n/a	0.533 $\pm$ 0.39	n.t.	0.392 $\pm$ 0.235	0.744 $\pm$ 0.379	$p < 0.05$	0.417 $\pm$ 0.147	0.386 $\pm$ 0.0409	n.s.
Pb		0.786 $\pm$ 0.376	2.28 $\pm$ 2.27	$p < 0.05$	1.56 $\pm$ 0.47	4.19 $\pm$ 0.852	$p < 0.05$	4.68 $\pm$ 1.22	7.14 $\pm$ 3.21	n.s.
V		n/a	0.315 $\pm$ 0.194	n.t.	0.166 $\pm$ 0.0422	6.92 $\pm$ 3.63	$p < 0.05$	0.292 $\pm$ 0.184	0.497 $\pm$ 0.244	$p < 0.05$
Zn		61 $\pm$ 7.58	98.3 $\pm$ 13.6	$p < 0.05$	142 $\pm$ 0.473	171 $\pm$ 48	$p < 0.05$	129 $\pm$ 12	126 $\pm$ 13.9	n.s.

n.s. = not significant

n.t. = not tested

Table S1b. Continues.

		<i>Mullus barbatus</i>			<i>Diplodus vulgaris</i>			<i>Pagellus sp.</i>		
		OUTSIN	INSIN	Sig.	OUTSIN	INSIN	Sig.	OUTSIN	INSIN	Sig.
Naphthalene	ng/g (dw)	88.7 ± 36.1	89.1 ± 32.8	<i>n.s.</i>	196 ± 42.9	65.8 ± 13.7	<i>p</i> < 0.05	92.5 ± 44.3	129 ± 42.7	<i>n.s.</i>
Acenaphthylene		< 0.05	< 0.05	<i>n.t.</i>	< 0.05	< 0.05	<i>n.t.</i>	< 0.05	< 0.05	<i>n.t.</i>
1-methylnaphthalene		80.7 ± 20.7	50.7 ± 10.6	<i>n.s.</i>	115 ± 18.7	38.4 ± 7.8	<i>p</i> < 0.05	66.6 ± 35.1	79.9 ± 33	<i>n.s.</i>
2-methylnaphthalene		62.8 ± 16.1	39.3 ± 11.9	<i>n.s.</i>	105 ± 6.69	40.2 ± 9.11	<i>p</i> < 0.05	42.4 ± 22.3	59.5 ± 21.9	<i>n.s.</i>
Acenaphthene		3.79 ± 1.15	9.77 ± 1.36	<i>p</i> < 0.05	< 0.01	12.7 ± 1.58	<i>n.t.</i>	< 0.01	8.35 ± 3.23	<i>n.t.</i>
Fluorene		13 ± 2.8	6.11 ± 0.385	<i>p</i> < 0.05	7.68 ± 0.0539	6.47 ± 0.81	<i>n.s.</i>	5.83 ± 1.87	6.46 ± 1.62	<i>n.s.</i>
Phenanthrene		36.6 ± 9.98	23.5 ± 18.4	<i>n.s.</i>	31.2 ± 26.7	3.16 ± 1.18	<i>p</i> < 0.05	27 ± 21.8	38.4 ± 18.8	<i>n.s.</i>
Anthracene		0.38 ± 0.489	0.728 ± 0.121	<i>p</i> < 0.05	< 0.01	1.08 ± 0.0569	<i>n.t.</i>	0.473 ± 0.329	0.52 ± 0.079	<i>n.s.</i>
Fluoranthene		4.05 ± 1.76	< 0.01	<i>n.t.</i>	< 0.01	< 0.01	<i>n.t.</i>	< 0.01	7.96 ± 2.82	<i>n.t.</i>
Pyrene		1.81 ± 0.858	2.32 ± 1.54	<i>n.s.</i>	6.77 ± 1.12	3.26 ± 1.67	<i>n.s.</i>	9.23 ± 12.4	2.69 ± 1.58	<i>p</i> < 0.05
Benzo(a)anthracene		< 0.01	< 0.01	<i>n.t.</i>	< 0.01	< 0.01	<i>n.t.</i>	< 0.01	< 0.01	<i>n.t.</i>
Chrysene		< 0.05	0.795 ± 0.722	<i>n.t.</i>	< 0.01	0.456 ± 0.4	<i>n.t.</i>	4.18 ± 6.44	0.473 ± 0.287	<i>p</i> < 0.05
7,12-dimethylbenzo(a)anthracene		< 1	20.8 ± 11.2	<i>n.t.</i>	< 0.05	16.2 ± 12.3	<i>n.t.</i>	< 0.05	7.91 ± 2.05	<i>n.t.</i>
Benzo(b)fluoranthene		0.944 ± 0.765	2.07 ± 2.27	<i>n.s.</i>	2.76 ± 0.342	0.85 ± 0.299	<i>p</i> < 0.05	5.13 ± 7.96	1.12 ± 0.84	<i>n.s.</i>
Benzo(k)fluoranthene		0.611 ± 0.684	1.71 ± 1.99	<i>n.s.</i>	1.48 ± 1.75	1.66 ± 0.849	<i>n.s.</i>	2.69 ± 3.95	0.789 ± 0.657	<i>n.s.</i>
Benzo(a)pyrene		0.872 ± 0.627	1.06 ± 1.01	<i>n.s.</i>	1.27 ± 0.00229	1.02 ± 0.459	<i>n.s.</i>	2.3 ± 3.67	0.73 ± 0.461	<i>n.s.</i>
Dibenzo(ah)anthracene		< 0.001	2,391	<i>n.t.</i>	2.92 ± 0.00591	< 0.001	<i>n.t.</i>	1.12 ± 0.749	1.79 ± 1.33	<i>n.s.</i>
Benzo(ghi)perylene		0.631 ± 0.297	2,648	<i>n.t.</i>	0.513 ± 0.725	0.376 ± 0.0947	<i>n.s.</i>	0.858 ± 0.464	0.251 ± 0.434	<i>p</i> < 0.05
Indeno(123cd)pyrene		< 0.001	0,084	<i>n.t.</i>	< 0.05	< 0.05	<i>n.t.</i>	< 0.05	0.192 ± 0.333	<i>n.t.</i>
Low MW PAHs		286 ± 79.1	219 ± 71.2	<i>n.s.</i>	456 ± 41.7	167 ± 28.9	<i>p</i> < 0.05	234 ± 120	319 ± 119	<i>n.s.</i>
High MW PAHs		8.39 ± 0.742	30.1 ± 16.1	<i>p</i> < 0.05	15.7 ± 2.5	23.3 ± 15.9	<i>n.s.</i>	25 ± 34.3	23.3 ± 8.57	<i>n.s.</i>
Total PAHs		294 ± 79.4	249 ± 86.6	<i>n.s.</i>	472 ± 44.2	190 ± 44.8	<i>p</i> < 0.05	259 ± 125	343 ± 127	<i>n.s.</i>

n.s. = not significant

n.t. = not tested

**Table S2a.** Results of biomarkers analyzed in mussels sampled. Data expressed as mean  $\pm$  standard deviation (n=5). Asterisks (\*) indicate significant differences between areas and reference area ( $p < 0.05$ ), as determined by t-student test.

		PGT (Area 1;9)	PGP (Area 1;7)	P2 (Area 2;10)	BB (Area 6;9;11)	BNF (Area 8)	BRM (Reference area)	Sig.
Lysosomal membranes stability	(min)	53.1 $\pm$ 8.0		55.2 $\pm$ 19.8	76.8 $\pm$ 5.0	61.8 $\pm$ 15.5	73.5 $\pm$ 28.9	n.s.
Acetylcholinesterase enzyme activity	(nmol/min/mg prt)	88.1 $\pm$ 10.1	94.5 $\pm$ 21.7	99.9 $\pm$ 23.8	82.7 $\pm$ 19.4	99.1 $\pm$ 30.5	105.1 $\pm$ 25.1	n.s.
Metallothioneins	(nmol eq.(G)SH/mg prt)	3.2 $\pm$ 1.7	1.7 $\pm$ 0.5	2.4 $\pm$ 0.6	2.00 $\pm$ 1.2	2.8 $\pm$ 0.5	2.3 $\pm$ 0.7	n.s.
Micronuclei frequency	(‰)	1.3 $\pm$ 0.3	0.8 $\pm$ 0.1	0.5 $\pm$ 0.3	0.4 $\pm$ 0.2	0.3 $\pm$ 0.3	0.3 $\pm$ 0.1	n.s.

n.s. = not significant

**Table S2b.** Results of biomarkers analyzed in fish species. Data expressed as mean  $\pm$  standard deviation (n=5). Asterisks (\*) indicate significant differences between areas and reference area ( $p < 0.05$ ), as determined by t-student test.

		<i>Mullus barbatus</i>			<i>Diplodus vulgaris</i>			<i>Pagellus sp.</i>		
		OUTSIN (Reference area)	INSIN (Area 3;4;5;8)	Sig.	OUTSIN (Reference area)	INSIN (Area 3;4;5;8)	Sig.	OUTSIN (Reference area)	INSIN (Area 3;4;5;8)	Sig.
Acetylcholinesterase enzyme activity	(nmol/min/mg prt)	63.3 $\pm$ 14.3	64.9 $\pm$ 13.1	n.s.	85.5 $\pm$ 10.6	61.2 $\pm$ 12.9	n.s.	91.4 $\pm$ 16.3	66.5 $\pm$ 11.8	n.s.
EROD enzyme activity	(pmol/min/mg prt)	85.3 $\pm$ 23.4	162.6 $\pm$ 47	$p < 0.05$	7.3 $\pm$ 4.1	4.3 $\pm$ 2.6	n.s.	27.2 $\pm$ 20.1	67.7 $\pm$ 15.3	$p < 0.05$
Pyrene-like metabolites	( $\mu$ g/ $\mu$ mol biliverdina)	0.4 $\pm$ 0.3	6.7 $\pm$ 2.3	$p < 0.01$	1.3 $\pm$ 1.0	5.4 $\pm$ 5.9	n.s.	0.1 $\pm$ 0.1	3.8 $\pm$ 2.7	$p < 0.05$
B[a]P-like metabolites	( $\mu$ g/ $\mu$ mol biliverdina)	3.6 $\pm$ 1.7	20.2 $\pm$ 4.7	$p < 0.01$	7.1 $\pm$ 3.9	10.5 $\pm$ 7.8	n.s.	0.6 $\pm$ 0.5	11.8 $\pm$ 3.6	$p < 0.01$
Naphtalene-like metabolites	(mg/ $\mu$ mol biliverdina)	9.0 $\pm$ 5.9	8.6 $\pm$ 1.8	n.s.	3.9 $\pm$ 2.0	8.2 $\pm$ 3.4	n.s.	1.1 $\pm$ 0.6	6.5 $\pm$ 4.0	$p < 0.05$
Micronuclei frequency	(‰)	6.3 $\pm$ 0.5	10.6 $\pm$ 1.8	$p < 0.05$	4.8 $\pm$ 0.6	5.3 $\pm$ 1.1	n.s.	5.5 $\pm$ 0.8	6.3 $\pm$ 0.8	n.s.

n.s. = not significant

**Table S3a.** Result of the bioassay with *Vibrio fischeri*. Data are expressed in toxic units (TU), reporting mean  $\pm$  standard deviation.

Area	Sample	CONTROL T.U.	TREATED T.U.
Area 1	18	100	100
	19	100	100
	20	100	100
	23	100	100
	24	100	100
	25	100	100
	26	81.2 $\pm$ 1.1	240.8 $\pm$ 18.6
	27	63 $\pm$ 0.9	36.7 $\pm$ 4.7
	28	108.8 $\pm$ 2.5	118.6 $\pm$ 41.4
	29	100	100
	30	100	100
	31	100	100
	32	100	100
	33	92.2 $\pm$ 2	4799.1 $\pm$ 824.1
	34	100	100
	35	67.8 $\pm$ 3.9	108.2 $\pm$ 12
	36	100	100
	37	100	100
	41	100	100
	42	117.5 $\pm$ 2.6	3006.3 $\pm$ 4338.4
Area 2	43	130 $\pm$ 7.5	366.3 $\pm$ 77
	44	100	100
	48	71.5 $\pm$ 8	172.9 $\pm$ 14.5
	59	100	100
	60	63.8 $\pm$ 4.8	93.4 $\pm$ 22.8
	61	100	100
	62	100	100
	69	100	100
Area 3	70	100	100
	71	100	100
	96	96 $\pm$ 1.9	69.8 $\pm$ 3.5
	97	96 $\pm$ 1.9	80.3 $\pm$ 2
	98	96 $\pm$ 1.9	74 $\pm$ 11.3
	99	96 $\pm$ 1.9	64.2 $\pm$ 1.5
	100	96 $\pm$ 1.9	86.8 $\pm$ 1.4
	104	96 $\pm$ 1.9	37.6 $\pm$ 0.2
Area 4	105	96 $\pm$ 1.9	52.8 $\pm$ 1.7
	106	96 $\pm$ 1.9	37 $\pm$ 0.4
	108	96 $\pm$ 1.9	17.9 $\pm$ 3
	109	93.9 $\pm$ 1.4	67.7 $\pm$ 0.9
	110	93.9 $\pm$ 1.4	78.8 $\pm$ 1.5
	111	96 $\pm$ 1.9	10.6 $\pm$ 0.2
	112	91.5 $\pm$ 2.1	7.5 $\pm$ 0.7
	113	91.5 $\pm$ 2.1	2.5 $\pm$ 0.7
	114	106 $\pm$ 8.4	58 $\pm$ 4.2
	115	91.5 $\pm$ 2.1	7 $\pm$ 2.8
	116	91.5 $\pm$ 0.7	12.5 $\pm$ 3.5

**Table S3a.** Continues.

Area	Sample	CONTROL T.U.	TREATED T.U.
Area 5	117	99 ± 1.4	34.5 ± 6.3
	118	99 ± 1.4	21.5 ± 0.7
	119	91.5 ± 0.7	5.5 ± 0.7
	120	94.5 ± 0.7	56 ± 1.4
	121	95	15.5 ± 0.7
	122	94.5 ± 0.7	14 ± 1.4
	123	94.5 ± 0.7	25 ± 1.4
Area 6	101	96 ± 1.9	65.5 ± 2.9
	102	96 ± 1.9	83 ± 1.9
	103	96 ± 1.9	83 ± 2.1
	107	96 ± 1.9	61.8 ± 2.7
Area 7	38	100	100
	39	100	100
	40	100	100
	45	216.2 ± 20.5	293.3 ± 45.2
	46	100	100
	47	75.9 ± 8.5	78.8 ± 8.7
	49	138.6 ± 1	1269.7 ± 358.7
	50	158.4 ± 1.2	6641.8 ± 1243.6
	51	151.3 ± 0	9510.6 ± 1182.3
	52	199.9 ± 22.8	3434.2 ± 663.3
	53	209.7 ± 7.7	1263 ± 426.4
	54	239.3 ± 19.4	228.7 ± 20.3
	55	214.4 ± 6.5	18384.7 ± 1613.5
	56	165.5 ± 2.4	335.3 ± 35
	57	100	100
	58	197.7 ± 10.4	1058.7 ± 323.1
	64	216.6 ± 15.7	77.2 ± 4.7
	65	220.7 ± 13.2	2401.4 ± 563.3
	66	243.7 ± 14.6	168.4 ± 11.2
	67	100	100
	68	100	100
	73	217.1 ± 12.7	1866.9 ± 688.8
	74	171.9 ± 7.5	881.5 ± 166.7
	75	156.2 ± 12.7	298.9 ± 62.1
	77	100	100
Area 8	124	94.5 ± 0.7	50 ± 1.4
	125	94.5 ± 0.7	65 ± 2.8
	126	94.5 ± 0.7	47 ± 2.8
	127	94.5 ± 0.7	41 ± 1.4



**Table S3a.** Continues.

Area	Sample	CONTROL	TREATED
		T.U.	T.U.
Area 9	1	100	100
	2	100	100
	3	100	100
	4	100	100
	5	100	100
	6	100	100
	7	100	100
	8	100	100
	9	100	100
	10	100	100
	11	100	100
	12	100	100
	13	100	100
	14	100	100
	15	100	100
Area 10	63	100	100
	72	100	100
	80	92 .4 $\pm$ 0 .6	133 .9 $\pm$ 13 .7
	81	100	100
	82	100	100
	83	100	100
	84	100	100
	85	100	100
	91	100	100
	92	100	100
	94	71 .3 $\pm$ 2 .6	109 .1 $\pm$ 17 .1
	95	101 .2 $\pm$ 3 .7	108 .3 $\pm$ 6 .1
Area 11	16	100	100
	17	100	100
	21	100	100
	22	100	100

**Table S3b.** Result of the bioassay with *Skeletonema costatum*. Algal growth values are expressed as mean cell density (cells/mL)  $\pm$  standard deviation.

Area	Sample	CONTROL cell/ml	TREATED cell/ml
Area 1	18	19563 .4 $\pm$ 247849 .7	27307 .3 $\pm$ 68014 .5
	19	14653 .7 $\pm$ 210384 .1	15619 .4 $\pm$ 98947 .7
	20	19971 $\pm$ 226955 .4	1222 .7 $\pm$ 166722 .2
	23	19971 $\pm$ 226955 .4	5909 .8 $\pm$ 165425 .3
	24	33828 .5 $\pm$ 222488 .4	4483 .2 $\pm$ 45343
	25	14653 .7 $\pm$ 210384 .1	8312 .8 $\pm$ 87900 .2
	26	14653 .7 $\pm$ 210384 .1	815 .1 $\pm$ 213266
	27	8359 .3 $\pm$ 189441 .7	8287 .3 $\pm$ 312406
	28	19563 .4 $\pm$ 247849 .7	10153 .9 $\pm$ 74162 .8
	29	23948 .7 $\pm$ 169844 .3	4991 .7 $\pm$ 65709
	30	14653 .7 $\pm$ 210384 .1	29751 .8 $\pm$ 193572 .6
	31	19971 $\pm$ 226955 .4	3750 .2 $\pm$ 110975 .2
	32	19971 $\pm$ 226955 .4	27307 .3 $\pm$ 245928 .4
	33	23948 .7 $\pm$ 169844 .3	11699 .5 $\pm$ 132955
	34	14653 .7 $\pm$ 210384 .1	53828 .4 $\pm$ 153705 .2
	35	2512 .4 $\pm$ 182601 .8	8438 .4 $\pm$ 176146 .2
	36	19563 .4 $\pm$ 247849 .7	14471 .2 $\pm$ 350832 .3
	37	19563 .4 $\pm$ 247849 .7	4075 .7 $\pm$ 141793 .1
	41	2512 .4 $\pm$ 182601 .8	3668 .1 $\pm$ 244391 .4
	42	19563 .4 $\pm$ 247849 .7	12763 .3 $\pm$ 121811 .4
Area 2	43	13042 .3 $\pm$ 212497 .5	13449 .8 $\pm$ 136413 .4
	44	13042 .3 $\pm$ 212497 .5	25946 .3 $\pm$ 458041 .7
	48	2512 .4 $\pm$ 182601 .8	10049 .7 $\pm$ 135568
	59	33828 .5 $\pm$ 222488 .4	6647 .3 $\pm$ 55333 .9
	60	19563 .4 $\pm$ 247849 .7	14547 .5 $\pm$ 401170 .8
	61	6349 $\pm$ 174455 .5	9246 .3 $\pm$ 136413 .4
	62	2512 .4 $\pm$ 182601 .8	3260 .5 $\pm$ 229635 .7
Area 3	69	33828 .5 $\pm$ 222488 .4	1199 .8 $\pm$ 39963 .3
	70	2512 .4 $\pm$ 182601 .8	8823 .4 $\pm$ 212113 .3
	71	6349 $\pm$ 174455 .5	18650 .6 $\pm$ 216724 .4
	96	4890 .8 $\pm$ 204966	10316 .2 $\pm$ 199048 .4
	97	4890 .8 $\pm$ 204966	11412 $\pm$ 232863 .5
Area 4	98	4890 .8 $\pm$ 204966	5598 .2 $\pm$ 172534 .2
	99	4890 .8 $\pm$ 204966	8671 .5 $\pm$ 201353 .9
	100	4890 .8 $\pm$ 204966	4048 .4 $\pm$ 181948 .6
	104	4890 .8 $\pm$ 204966	20760 .8 $\pm$ 221527 .8
	105	4890 .8 $\pm$ 204966	6113 .5 $\pm$ 204043 .8
	106	4890 .8 $\pm$ 204966	18748 .3 $\pm$ 205580 .8
	108	4890 .8 $\pm$ 204966	14986 .2 $\pm$ 205773
Area 4	109	4890 .8 $\pm$ 204966	17862 $\pm$ 188289
	110	4890 .8 $\pm$ 204966	17118 $\pm$ 242278
	111	4890 .8 $\pm$ 204966	10596 .8 $\pm$ 186137 .1
	112	4890 .8 $\pm$ 204966	7743 .8 $\pm$ 189518 .6
	113	4890 .8 $\pm$ 204966	2512 .4 $\pm$ 130649 .5
	114	4890 .8 $\pm$ 204966	15935 .3 $\pm$ 202276 .2
	115	4890 .8 $\pm$ 204966	9046 .5 $\pm$ 235553 .4
	116	4890 .8 $\pm$ 204966	15690 .2 $\pm$ 190786 .7

**Table S3b.** Continues.

Area	Sample	CONTROL cell/ml	TREATED cell/ml
Area 5	117	4890 .8 ± 204966	13449 .9 ± 221912
	118	4890 .8 ± 204966	9374 .1 ± 231518 .6
	119	4890 .8 ± 204966	10507 .6 ± 176569
	120	4890 .8 ± 204966	5763 .9 ± 192515 .9
	121	4890 .8 ± 204966	16152 .7 ± 205580 .8
	122	4890 .8 ± 204966	4075 .7 ± 180699 .8
	123	4890 .8 ± 204966	16387 .6 ± 218645 .8
Area 6	101	4890 .8 ± 204966	13857 .4 ± 186944 .1
	102	4890 .8 ± 204966	17933 .2 ± 181948 .6
	103	4890 .8 ± 204966	14494 .1 ± 172918 .5
	107	4890 .8 ± 204966	4924 .7 ± 180988
Area 7	38	8359 .3 ± 189441 .7	18528 .5 ± 301646 .6
	39	2512 .4 ± 182601 .8	7321 .1 ± 178912 .9
	40	2512 .4 ± 182601 .8	2853 ± 127267 .9
	45	19563 .4 ± 247849 .7	5706 ± 300109 .5
	46	8359 .3 ± 189441 .7	9719 .2 ± 307410 .5
	47	19563 .4 ± 247849 .7	58690 .4 ± 236321 .8
	49	23948 .7 ± 169844 .3	14721 .5 ± 258609 .1
	50	19971 ± 226955 .4	40443 .7 ± 77813 .3
	51	6349 ± 174455 .5	38311 .8 ± 264373 .1
	52	14653 .7 ± 210384 .1	24454 .3 ± 271385 .9
	53	14653 .7 ± 210384 .1	25946 .3 ± 402707 .8
	54	13042 .3 ± 212497 .5	7043 .6 ± 208270 .6
	55	8359 .3 ± 189441 .7	8559 ± 169844 .3
	56	14653 .7 ± 210384 .1	3174 .5 ± 127767 .5
	57	14653 .7 ± 210384 .1	47278 .4 ± 340553 .2
	58	14653 .7 ± 210384 .1	4483 .2 ± 181564 .3
	64	6349 ± 174455 .5	5198 .2 ± 209807 .7
	65	33828 .5 ± 222488 .4	1222 .7 ± 6916 .7
	66	6349 ± 174455 .5	7411 .4 ± 294345 .6
	67	2512 .4 ± 182601 .8	4483 .2 ± 262836
	68	13042 .3 ± 212497 .5	6099 .9 ± 504921 .8
	73	13042 .3 ± 212497 .5	25946 .3 ± 402707 .8
	74	2512 .4 ± 182601 .8	18340 .7 ± 209346 .6
	75	6349 ± 174455 .5	8372 .6 ± 396559 .6
	77	2512 .4 ± 182601 .8	9374 .1 ± 177760 .1
Area 8	124	4890 .8 ± 204966	15487 .7 ± 174455 .5
	125	4890 .8 ± 204966	15487 .7 ± 177913 .9
	126	4890 .8 ± 204966	14744 .1 ± 182332 .9
	127	4890 .8 ± 204966	20378 .6 ± 173610 .1

**Table S3b.** Continues.

Area	Sample	CONTROL cell/ml	TREATED cell/ml
Area 9	1	13042 .3 ± 212497 .5	8873 .4 ± 397328 .1
	2	23948 .7 ± 169844 .3	33417 .6 ± 269368 .5
	3	19563 .4 ± 247849 .7	2445 .4 ± 152168 .2
	4	6349 ± 174455 .5	3049 .9 ± 175224
	5	14653 .7 ± 210384 .1	9228 .3 ± 191170 .9
	6	19563 .4 ± 247849 .7	18492 .6 ± 157547 .9
	7	6349 ± 174455 .5	37806 .9 ± 199048 .3
	8	8359 .3 ± 189441 .7	15640 .7 ± 143714 .4
	9	6349 ± 174455 .5	7500 .5 ± 174455 .5
	10	23948 .7 ± 169844 .3	40660 .6 ± 278975 .1
	11	19563 .4 ± 247849 .7	45847 .8 ± 141793 .1
	12	14653 .7 ± 210384 .1	6555 ± 249194 .7
	13	8359 .3 ± 189441 .7	6730 ± 309716 .1
	14	19971 ± 226955 .4	20582 .4 ± 205556 .8
	15	13042 .3 ± 212497 .5	14944 .3 ± 428069 .2
Area 10	63	2512 .4 ± 182601 .8	1729 .1 ± 196435 .3
	72	6349 ± 174455 .5	6754 .7 ± 142945 .9
	80	6349 ± 174455 .5	14270 .8 ± 230557 .9
	81	2512 .4 ± 182601 .8	3260 .5 ± 202890 .9
	82	6349 ± 174455 .5	2641 .3 ± 262836
	83	2512 .4 ± 182601 .8	2037 .8 ± 172918 .4
	84	2512 .4 ± 182601 .8	20378 .6 ± 269291 .6
	85	6349 ± 174455 .5	22303 .8 ± 389642 .9
	91	6349 ± 174455 .5	21661 .5 ± 185983 .4
	92	6349 ± 174455 .5	16475 .2 ± 319322 .7
	94	6349 ± 174455 .5	9922 .2 ± 219030
	95	6349 ± 174455 .5	23472 .3 ± 276669 .5
Area 11	16	14653 .7 ± 210384 .1	12789 .3 ± 76852 .6
	17	19971 ± 226955 .4	8966 .5 ± 213611 .9
	21	33828 .5 ± 222488 .4	11819 .6 ± 192285 .3
	22	33828 .5 ± 222488 .4	3923 .4 ± 226715 .3

**Table S3c.** Result of bioassay with *Paracentrotus lividus*. Values are expressed as mean % of normal embryos  $\pm$  standard deviation.

Area	Sample	CONTROL %	TREATED %
Area 1	18	84.6 $\pm$ 1.5	82.6 $\pm$ 2.5
	19	84.6 $\pm$ 1.5	3.3 $\pm$ 1.5
	20	84.6 $\pm$ 1.5	68 $\pm$ 2
	23	84.6 $\pm$ 0.5	84.6 $\pm$ 0.5
	24	84.6 $\pm$ 1.5	82.3 $\pm$ 0.5
	25	81.3 $\pm$ 1.5	59.6 $\pm$ 1.5
	26	81.3 $\pm$ 1.5	18.6 $\pm$ 2
	27	81.3 $\pm$ 1.5	70.3 $\pm$ 0.5
	28	84.6 $\pm$ 0.5	70 $\pm$ 2.6
	29	84.6 $\pm$ 0.5	85.3 $\pm$ 2.5
	30	84.6 $\pm$ 0.5	9 $\pm$ 1
	31	81.3 $\pm$ 1.5	80.6 $\pm$ 1.1
	32	81.3 $\pm$ 1.5	62.3 $\pm$ 2.5
	33	84.6 $\pm$ 0.5	61.6 $\pm$ 1.5
	34	84.6 $\pm$ 0.5	38.3 $\pm$ 3.5
	35	79.6 $\pm$ 3	79.6 $\pm$ 3.5
	36	84.6 $\pm$ 0.5	83.3 $\pm$ 2.8
	37	79.6 $\pm$ 3	79.3 $\pm$ 1.1
	41	84.6 $\pm$ 0.5	16.6 $\pm$ 2.8
	42	81.3 $\pm$ 1.5	0
Area 2	43	81.3 $\pm$ 1.5	0.6 $\pm$ 0.5
	44	79.6 $\pm$ 3	80.3 $\pm$ 3.5
	48	84.6 $\pm$ 0.5	5.3 $\pm$ 0.5
	59	84.6 $\pm$ 0.5	23 $\pm$ 1
	60	84.6 $\pm$ 1.5	32.6 $\pm$ 2
	61	84.6 $\pm$ 0.5	84.3 $\pm$ 0.5
	62	84.6 $\pm$ 0.5	61.6 $\pm$ 2.8
	69	81.3 $\pm$ 1.5	80.6 $\pm$ 1.1
Area 3	70	79.6 $\pm$ 3	78.6 $\pm$ 1.1
	71	79.6 $\pm$ 3	61 $\pm$ 3.6
	96	89 $\pm$ 2	0.6 $\pm$ 1.1
	97	89 $\pm$ 2	82 $\pm$ 2.6
	98	89 $\pm$ 2	88.6 $\pm$ 0.5
	99	89 $\pm$ 2	0
	100	89 $\pm$ 2	0
	104	89 $\pm$ 2	73 $\pm$ 2.6
	105	89 $\pm$ 2	23 $\pm$ 5.2
Area 4	106	89 $\pm$ 2	89 $\pm$ 1
	108	89 $\pm$ 2	89.3 $\pm$ 0.5
	109	89 $\pm$ 2	17.3 $\pm$ 6.8
	110	89 $\pm$ 2	88.3 $\pm$ 1.5
	111	89 $\pm$ 2	89 $\pm$ 1
	112	89 $\pm$ 2	88.3 $\pm$ 2
	113	89 $\pm$ 2	46.3 $\pm$ 1.5
	114	89 $\pm$ 2	88.6 $\pm$ 0.5
	115	89 $\pm$ 2	89.3 $\pm$ 0.5
	116	89 $\pm$ 2	87.6 $\pm$ 0.5

**Table S3c.** Continues.

Area	Sample	CONTROL %	TREATED %
Area 5	117	89 ± 2	88.3 ± 2.5
	118	89 ± 2	0
	119	89 ± 2	88 ± 1
	120	89 ± 2	21.3 ± 11.7
	121	89 ± 2	77.3 ± 2.5
	122	89 ± 2	89 ± 1
	123	89 ± 2	15.3 ± 0.5
Area 6	101	89 ± 2	82.3 ± 2.5
	102	89 ± 2	80.6 ± 1.1
	103	89 ± 2	87 ± 1.7
	107	89 ± 2	89.6 ± 1.1
Area 7	38	84.6 ± 0.5	84.6 ± 0.5
	39	84.6 ± 0.5	73.3 ± 2.8
	40	79.6 ± 3	79.6 ± 1.5
	45	79.6 ± 3	74 ± 5.2
	46	79.6 ± 3	79.3 ± 2
	47	84.6 ± 1.5	65 ± 3
	49	84.6 ± 1.5	0
	50	81.3 ± 1.5	0
	51	84.6 ± 1.5	17.6 ± 2.5
	52	84.6 ± 0.5	7.3 ± 1.5
	53	81.3 ± 1.5	0.6 ± 0.5
	54	79.6 ± 3	0
	55	79.6 ± 3	0
	56	84.6 ± 0.5	52.3 ± 2.5
	57	81.3 ± 1.5	70.6 ± 1.1
	58	81.3 ± 1.5	72 ± 2.6
	64	84.6 ± 0.5	84.6 ± 0.5
	65	84.6 ± 0.5	53.6 ± 1.5
	66	84.6 ± 1.5	0
	67	84.6 ± 0.5	36 ± 1.7
	68	81.3 ± 1.5	81.3 ± 1.1
	73	84.6 ± 1.5	77.6 ± 2.5
	74	84.6 ± 1.5	86.6 ± 1.5
	75	84.6 ± 1.5	84.6 ± 0.5
	77	79.6 ± 3	81.3 ± 3.2
Area 8	124	89 ± 2	58.6 ± 4
	125	89 ± 2	0
	126	89 ± 2	89.3 ± 1.1
	127	89 ± 2	88.3 ± 1.1

**Table S3c.** Continues.

Area	Sample	Area	CONTROL %	TREATED %
Area 9	1	Area 9	84.6 ± 0.5	84.6 ± 1.5
	2	Area 9	84.6 ± 0.5	84.6 ± 1.5
	3	Area 9	84.6 ± 0.5	84.3 ± 0.5
	4	Area 9	84.6 ± 0.5	84.6 ± 0.5
	5	Area 9	84.6 ± 0.5	84.6 ± 1.1
	6	Area 9	84.6 ± 0.5	84.6 ± 0.5
	7	Area 9	84.6 ± 0.5	84.3 ± 3
	8	Area 9	84.6 ± 0.5	85.3 ± 1.5
	9	Area 9	84.6 ± 0.5	84.6 ± 0.5
	10	Area 9	84.6 ± 1.5	83.6 ± 1.1
	11	Area 9	84.6 ± 0.5	84.6 ± 0.5
	12	Area 9	84.6 ± 0.5	84.6 ± 4.1
	13	Area 9	84.6 ± 0.5	80.3 ± 1.5
	14	Area 9	84.6 ± 0.5	84.6 ± 0.5
	15	Area 9	84.6 ± 1.5	54.6 ± 2.5
Area 10	63	Area 10	79.6 ± 3	68.6 ± 1.5
	72	Area 10	79.6 ± 3	62.6 ± 4.6
	80	Area 10	81.3 ± 1.5	80.6 ± 1.1
	81	Area 10	79.6 ± 3	80.6 ± 4
	82	Area 10	79.6 ± 3	78.6 ± 4.1
	83	Area 10	79.6 ± 3	79.3 ± 0.5
	84	Area 10	84.6 ± 1.5	87 ± 1.7
	85	Area 10	79.6 ± 3	6.3 ± 1.5
	91	Area 10	79.6 ± 3	63.6 ± 3.2
	92	Area 10	79.6 ± 3	80.6 ± 1.1
	94	Area 10	79.6 ± 3	79.6 ± 2.5
	95	Area 10	79.6 ± 3	80.6 ± 1.1
Area 11	16	Area 11	84.6 ± 0.5	60.3 ± 0.5
	17	Area 11	84.6 ± 0.5	84.6 ± 3.5
	21	Area 11	84.6 ± 0.5	84.6 ± 0.5
	22	Area 11	84.6 ± 1.5	0

**Table S4a.** List of observed species in benthic communities analyses.

Phylum	Class	Order	Family	Species
Annelida	Polychaeta	Eunicida	Dorvilleidae	<i>Protodorvillea kefersteini</i> (McIntosh, 1869)
			Eunicidae	<i>Eunice vittata</i> (Delle Chiaje, 1828)
			Eunicidae	<i>Lysidice unicornis</i> (Grube, 1840)
			Eunicidae	<i>Marphysa bellii</i> (Audouin & Milne-Edwards, 1833)
			Lumbrineridae	<i>Lumbrineris latreilli</i> Audouin & Milne-Edwards, 1834
			Lumbrineridae	<i>Ninoe armoricana</i> Glémarec, 1968
			Oeonidae	<i>Drilonereis filum</i> (Claparède, 1868)
			Onuphidae	<i>Aponuphis bilineata</i> (Baird, 1870)
			Onuphidae	<i>Hyalinoecia tubicola</i> (O.F. Müller, 1776)
			Onuphidae	<i>Onuphis eremita</i> Audouin & Milne Edwards, 1833
		Phyllodocida	Aphroditidae	<i>Pontogenia chrysocoma</i> (Baird, 1865)
			Glyceridae	<i>Glycera tridactyla</i> Schmarda, 1861
			Glyceridae	<i>Glycera unicornis</i> Lamarck, 1818
			Goniadidae	<i>Goniada maculata</i> Örsted, 1843
			Hesionidae	<i>Psamathe fusca</i> Johnston, 1836
			Nephtyidae	<i>Nephtys hombergii</i> Savigny in Lamarck, 1818
			Nereididae	<i>Nereis rava</i> Ehlers, 1868
			Paralacydoniidae	<i>Paralacydonia paradoxa</i> Fauvel, 1913
			Phyllodocidae	<i>Mysta picta</i> (Quatrefages, 1866)
			Phyllodocidae	<i>Nereiphylla rubiginosa</i> (Saint-Joseph, 1888)
			Phyllodocidae	<i>Phyllodoce lineata</i> (Claparède, 1870)
			Phyllodocidae	Phyllodocidae indet.
			Pilargidae	<i>Sigambra tentaculata</i> (Treadwell, 1941)
			Polynoidae	<i>Harmothoe antilopes</i> McIntosh, 1876
			Polynoidae	<i>Harmothoe longisetis</i> (Grube, 1863)
			Polynoidae	<i>Harmothoe</i> sp.
			Polynoidae	Polynoidae indet.
			Sigalionidae	<i>Sigalion mathildae</i> Audouin & Milne Edwards in Cuvier, 1830
			Sigalionidae	<i>Sthenelais boa</i> (Johnston, 1833)
			Syllidae	<i>Exogone</i> sp.
			Syllidae	Syllidae indet.
		Sabellida	Oweniidae	<i>Owenia fusiformis</i> Delle Chiaje, 1844
			Sabellidae	<i>Acromegalomma claparedei</i> (Gravier, 1906)
			Sabellidae	<i>Dialychone acustica</i> Claparède, 1870
			Sabellidae	<i>Dialychone arenicola</i> (Langerhans, 1881)
			Serpulidae	<i>Ditrupa arietina</i> (O. F. Müller, 1776)
		Spionida	Magelonidae	<i>Magelona allenii</i> Wilson, 1958
			Magelonidae	<i>Magelona johnstoni</i> Fiege, Licher & Mackie, 2000
			Poecilochaetidae	<i>Poecilochaetus serpens</i> Allen, 1904
			Spionidae	<i>Dipolydora coeca</i> (Örsted, 1843)
			Spionidae	<i>Paraprionospio pinnata</i> (Ehlers, 1901)
			Spionidae	<i>Prionospio ehlersi</i> Fauvel, 1928
			Spionidae	<i>Prionospio fallax</i> Soderstrom, 1920
			Spionidae	<i>Pseudopolydora antennata</i> (Claparède, 1869)
			Spionidae	<i>Scolecopsis (Scolecopsis) squamata</i> (O.F. Muller, 1806)
			Spionidae	<i>Spio filicornis</i> (Müller, 1776)



Table S4a. Continues.

Phylum	Class	Order	Family	Species
Annelida	Polychaeta	Spionida	Spionidae	<i>Spio multioculata</i> (Rioja, 1918)
			Spionidae	<i>Spiophanes reyssi</i> Laubier, 1964
		Terebellida	Ampharetidae	<i>Adercodon pleijeli</i> Mackie, 1994
			Ampharetidae	<i>Adercodon pleijeli</i> Mackie, 1994
			Ampharetidae	<i>Ampharete acutifrons</i> (Grube, 1860)
			Ampharetidae	<i>Melinna palmata</i> Grube, 1870
			Cirratulidae	<i>Aphelochaeta marioni</i> (Saint-Joseph, 1894)
			Cirratulidae	<i>Chaetozone caputesocis</i> (Saint-Joseph, 1894)
			Cirratulidae	Cirratulidae indet.
			Cirratulidae	<i>Kirkegaardia dorsobranchialis</i> (Kirkegaard, 1959)
			Flabelligeridae	<i>Diplocirrus glaucus</i> (Malmgren, 1867)
			Sternaspidae	<i>Sternaspis scutata</i> (Ranzani, 1817)
			Terebellidae	<i>Pista cretacea</i> (Grube, 1860)
			Trichobranchidae	<i>Terebellides stroemii</i> Sars, 1835
			Capitellidae	<i>Leiocapitella dollfusi</i> (Fauvel, 1936)
			Capitellidae	<i>Notomastus latericeus</i> Sars, 1851
			Capitellidae	<i>Pseudoleiocapitella fauveli</i> Harmelin, 1964
			Chaetopteridae	<i>Phyllochaetopterus socialis</i> Claparède, 1868
			Cossuridae	<i>Cossura soyeri</i> Laubier, 1962
			Maldanidae	<i>Chirimia biceps</i> (M. Sars, 1861)
			Maldanidae	<i>Euclymene lombricoides</i> (Quatrefages, 1866)
			Maldanidae	<i>Euclymene oerstedii</i> (Claparède, 1863)
			Maldanidae	<i>Leiochone leiopygos</i> (Grube, 1860)
			Maldanidae	<i>Metasychis gotoi</i> (Izuka, 1902)
			Maldanidae	<i>Praxillella</i> sp.
			Opheliidae	<i>Armandia cirrhosa</i> Filippi, 1861
			Opheliidae	<i>Ophelia</i> sp.
			Orbiniidae	<i>Phylo foetida ligustica</i> (Orlandi, 1896)
			Paraonidae	<i>Aricidea (Acmira) catherinae</i> Laubier, 1967
			Paraonidae	<i>Levinsonia gracilis</i> (Tauber, 1879)
Arthropoda	Hexanauplia	Sessilia	Balanidae	<i>Balanus trigonus</i> Darwin, 1854
	Malacostraca	Amphipoda	Ampeliscidae	<i>Ampelisca brevicornis</i> (Costa, 1853)
			Ampeliscidae	<i>Ampelisca ledoyeri</i> Bellan-Santini & Kaim-Malka, 1977
			Ampeliscidae	<i>Ampelisca ruffoi</i> Bellan-Santini & Kaim-Malka, 1977
			Ampeliscidae	<i>Ampelisca</i> sp.
			Ampeliscidae	<i>Ampelisca spinifer</i> Reid, 1951
			Ampeliscidae	<i>Ampelisca typica</i> (Spence Bate, 1856)
			Aoridae	<i>Autonoe spiniventris</i> Della Valle, 1893
			Aoridae	<i>Microdeutopus versiculatus</i> (Spence Bate, 1857)
			Bathyporeiidae	<i>Bathyporeia lindstromi</i> Stebbing, 1906
			Cheirocratidae	<i>Cheirocratus sundevallii</i> (Rathke, 1843)
			Dexaminidae	<i>Dexamine spinosa</i> (Montagu, 1813)
			Dexaminidae	<i>Guernea (Guernea) coalita</i> (Norman, 1868)
			Leucothoidae	<i>Leucothoe pachycera</i> Della Valle, 1893
			Maeridae	<i>Othomaera schmidtii</i> (Stephensen, 1915)
			Oedicerotidae	<i>Deflexilodes gibbosus</i> (Chevreux, 1888)

Table S4a. Continues.

Phylum	Class	Order	Family	Species
Arthropoda	Malacostraca	Amphipoda	Oedicerotidae	<i>Kroyera carinata</i> Spence Bate, 1857
			Oedicerotidae	<i>Periocolodes longimanus</i> (Spence Bate & Westwood, 1868)
			Oedicerotidae	<i>Synchelidium haplocheles</i> (Grube, 1864)
			Photidae	<i>Photis longicaudata</i> (Spence Bate & Westwood, 1862)
			Phoxocephalidae	<i>Harpinia antennaria</i> Meinert, 1890
			Phoxocephalidae	<i>Harpinia truncata</i> Sars, 1891
			Phoxocephalidae	<i>Metaphoxus gruneri</i> Karaman, 1986
			Tryphosidae	<i>Hippomedon ambiguus</i> Ruffo, 1946
			Tryphosidae	<i>Hippomedon massiliensis</i> Bellan-Santini, 1965
			Urothoidae	<i>Urothoe elegans</i> (Spence Bate, 1857)
		Cumacea	Bodotriidae	<i>Bodotria scorpioides</i> (Montagu, 1804)
			Bodotriidae	<i>Iphinoe serrata</i> Norman, 1867
		Decapoda	Alpheidae	<i>Alpheus glaber</i> (Olivi, 1792)
			Callianassidae	<i>Callianassa</i> sp.
			Carcinidae	<i>Xaiva biguttata</i> (Risso, 1816)
			Diogenidae	<i>Diogenes pugilator</i> (Roux, 1829)
			Solenoceridae	<i>Solenocera membranacea</i> (Risso, 1816)
			Decapoda indet.	
		Isopoda	Sphaeromatidae	<i>Cymodoce tuberculata</i> Costa in Hope, 1851
		Mysida		Mysida indet.
		Stomatopoda	Nannosquillidae	<i>Platysquilla eusebia</i> (Risso, 1816)
		Tanaidacea	Apseudidae	<i>Apseudopsis latreillii</i> (Milne Edwards, 1828)
			Leptocheliidae	<i>Chondrochelia savignyi</i> (Kroyer, 1842)
Echinodermata	Echinoidea	Clypeasteroidea	Echinocyamidae	<i>Echinocyamus pusillus</i> (O.F. Müller, 1776)
	Ophiuroidea	Ophiurida	Amphiuridae	<i>Amphipholis squamata</i> (Delle Chiaje, 1828)
			Amphiuridae	<i>Amphiura chiajei</i> Forbes, 1843
			Amphiuridae	<i>Amphiura filiformis</i> (O.F. Müller, 1776)
			Ophiotrichidae	<i>Ophiotrix</i> sp.
			Ophiuridae	<i>Ophiura ophiura</i> (Linnaeus, 1758)
		Spatangoida	Loveniidae	<i>Echinocardium cordatum</i> (Pennant, 1777)
Mollusca	Bivalvia	Arcida	Glycymerididae	<i>Glycymeris bimaculata</i> (Poli, 1795)
			Noetiidae	<i>Striarca lactea</i> (Linnaeus, 1758)
		Cardiida	Cardiidae	<i>Acanthocardia echinata</i> (Linnaeus, 1758)
			Cardiidae	<i>Fulvia australis</i> (G. B. Sowerby II, 1834)
			Cardiidae	<i>Laevicardium oblongum</i> (Gmelin, 1791)
			Cardiidae	<i>Papillicardium papillosum</i> (Poli, 1791)
			Cardiidae	<i>Parvicardium exiguum</i> (Gmelin, 1791)
			Donacidae	<i>Donax venustus</i> Poli, 1795
			Tellinidae	<i>Fabulina fabula</i> (Gmelin, 1791)
			Tellinidae	<i>Moerella pulchella</i> (Lamarck, 1818)
			Astartidae	<i>Astarte fusca</i> (Poli, 1791)
		Littorinimorpha	Naticidae	<i>Neverita josephinia</i> Risso, 1826
		Lucinida	Lucinidae	<i>Lucinella divaricata</i> (Linnaeus, 1758)
			Thyasiridae	<i>Thyasira biplicata</i> (Philippi, 1836)
		Myida	Corbulidae	<i>Corbula gibba</i> (Olivi, 1792)
		Nuculanida	Nuculanidae	<i>Saccella commutata</i> (Philippi, 1844)

**Table S4a.** Continues.

Phylum	Class	Order	Family	Species
Mollusca	Bivalvia	Nuculida	Nuculidae	<i>Nucula nitidosa</i> Winckworth, 1930
		Venerida	Veneridae	<i>Callista chione</i> (Linnaeus, 1758)
			Veneridae	<i>Chamelea gallina</i> (Linnaeus, 1758)
			Veneridae	<i>Chamelea striatula</i> (da Costa, 1778)
			Veneridae	<i>Clausinella fasciata</i> (da Costa, 1778)
			Veneridae	<i>Dosinia lupinus</i> (Linnaeus, 1758)
			Veneridae	<i>Pitar rudis</i> (Poli, 1795)
			Macluridae	<i>Spisula subtruncata</i> (da Costa, 1778)
			Thraciidae	<i>Thracia phaseolina</i> (Lamarck, 1818)
			Ungulinidae	<i>Diplodonta trigona</i> (Scacchi, 1835)
	Gastropoda	Cephalaspidea	Philinidae	<i>Philine</i> sp.
		Littorinimorpha	Naticidae	<i>Naticarius hebraeus</i> (Martyn, 1786)
		Neogastropoda	Mangeliidae	<i>Mangelia costata</i> (Pennant, 1777)
			Muricidae	<i>Hexaplex trunculus</i> (Linnaeus, 1758)
			Nassariidae	<i>Tritia mutabilis</i> (Linnaeus, 1758)
			Nassariidae	<i>Tritia pygmaea</i> (Lamarck, 1822)
			Ringiculidae	<i>Ringicula auriculata</i> (Ménard de la Groye, 1811)
		Polyplacophora		Polyplacophora indet.
	Scaphopoda	Dentaliida	Dentaliidae	<i>Antalis inaequicostata</i> (Dautzenberg, 1891)
	Bivalvia	Arcida	Arcidae	<i>Anadara gibbosa</i> (Reeve, 1844)
		Cardiida	Tellinidae	<i>Peronidia albicans</i> (Gmelin, 1791)
		Lucinida	Lucinidae	<i>Loripinus fragilis</i> (Philippi, 1836)
			Macluridae	<i>Lutraria lutraria</i> (Linnaeus, 1758)
Sipuncula	Phascolosomatidea	Aspidosiphonida	Aspidosiphonidae	<i>Aspidosiphon (Aspidosiphon) muelleri muelleri</i> Diesing, 1851
	Sipunculidea	Golfingiida	Golfingiidae	<i>Golfingia (Golfingia) elongata</i> (Keferstein, 1862)
			Phascolionidae	<i>Phascolion (Phascolion) strombus strombus</i> (Montagu, 1804)

**Table S4b.** Ecological quality indices values: organisms abundance (N), species richness (S), Margalef index (d), Shannon index (H'), Pielou index (J'), AMBI index, BENTHIX index, BOPA index, BITS index, m AMBI index, HQ and level of hazard for Benthic Communities. Quality classes of AMBI index are expressed through conventional colours (blue: elevated; green: good; yellow: sufficient; orange: scarce; red: bad)

Area	Sample	N	H'	S	D	J'	AMBI	BENTHIX	BOPA	BITS	m-AMBI	HQ	Level of hazard
Area 1	19	37.7	3.2	13.3	3.4	1.3	0.8	2.6	0.1	1.1	0.7	13.2	Absent
Area 1 - 2 - 10	44	42.3	3.1	14.3	3.6	1.2	0.6	2.9	0.1	1.3	0.7	10.5	Absent
Area 3	98	23	3.1	10.7	3.1	1.3	0.6	3.9	0.2	1.2	0.7	10.8	Absent
Area 3	99	51.3	3.7	21.3	5.2	1.2	0.8	3.8	0.2	1.2	0.9	12.7	Absent
Area 3	104	57	3.9	22	5.2	1.3	0.8	2.5	0.2	1.1	0.9	13.7	Absent
Area 4	111	24	3.7	15.7	4.6	1.4	1.6	2.5	0.3	0.6	0.7	23.6	Slight
Area 4	113	42	3.5	17.7	4.5	1.2	2.6	2.6	0.3	0.6	0.7	32.9	Slight
Area 5	117	17.7	3.4	12	3.8	1.4	1.6	3.7	0.3	0.9	0.7	23.7	Slight
Area 5	122	18.7	2.7	9.3	2.9	1.2	3	2.7	0.2	0.6	0.5	36.8	Slight
Area 6	107	43.7	3.3	17.7	4.5	1.2	0.5	3.6	0.2	0.8	0.8	8.5	Absent
Area 7 - 10	67	51	3.9	21	5.1	1.3	1.2	3.1	0.2	1.1	0.8	20.3	Slight
Area 8	126	74	3.4	25	5.6	1.1	0.5	4	0.2	1	0.9	8.7	Absent
Area 8	127	49.3	3.5	15.7	3.8	1.3	0.4	4.1	0.1	1.2	0.8	6.8	Absent
Area 9	12	34.7	3.5	13.7	3.6	1.3	0.9	2.8	0.2	0.9	0.7	15.5	Absent
Area 11	21	32.3	3.6	17	4.6	1.3	1.1	3.1	0.2	0.8	0.8	18.1	Absent

7 **Table S5.** Weight and threshold (%) of biomarkers used for elaboration LOE3: Sublethal effects.

8

Biomarkers	Species	Weight	Inhibition threshold	Induction threshold
Lysosomal membranes stability	<i>M. galloprovincialis</i>	1.2	25	
Acetylcholinesterase enzyme activity		1.5	25	60
Metallothioneins		1		40
Micronuclei frequency		1.9		50
Acetylcholinesterase enzyme activity	<i>Fish species</i>	1.5	25	60
EROD enzyme activity		1.5		200
Pyrene-like metabolites		1		150
B[a]P-like metabolites		1		100
Naphtalene-like metabolites		1		200
Micronuclei frequency		1.9		50

9

10