



UNIVERSITÀ POLITECNICA DELLE MARCHE
Repository ISTITUZIONALE

Chitosan and postharvest decay of fresh fruit: Meta-analysis of disease control and antimicrobial and eliciting activities

This is a pre print version of the following article:

Original

Chitosan and postharvest decay of fresh fruit: Meta-analysis of disease control and antimicrobial and eliciting activities / Rajestary, R., Landi, L., Romanazzi, G.. - In: COMPREHENSIVE REVIEWS IN FOOD SCIENCE AND FOOD SAFETY. - ISSN 1541-4337. - 20:1(2021), pp. 563-582. [10.1111/1541-4337.12672]

Availability:

This version is available at: 11566/286700 since: 2024-05-24T12:26:34Z

Publisher:

Published

DOI:10.1111/1541-4337.12672

Terms of use:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. The use of copyrighted works requires the consent of the rights' holder (author or publisher). Works made available under a Creative Commons license or a Publisher's custom-made license can be used according to the terms and conditions contained therein. See editor's website for further information and terms and conditions.

This item was downloaded from IRIS Università Politecnica delle Marche (<https://iris.univpm.it>). When citing, please refer to the published version.

Publisher copyright:

Wiley - Preprint/ Author's submitted Manuscript

This is the pre-peer reviewed version of the above quoted article which has been published in final form at 10.1111/1541-4337.12672. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions <https://authorservices.wiley.com/author-resources/Journal-Authors/licensing/self-archiving.html>.

(Article begins on next page)

15 July 2026

1	Table of contents
2	Title: Chitosan and Postharvest Decay of Fresh Fruit: Meta-Analysis of Disease Control
3	and Antimicrobial and Eliciting Activities
4	Abstract
5	1. INTRODUCTION
6	2. METHODS
7	2.1 Search strategy and study selection
8	2.2 Data extraction
9	2.3 Data analysis
10	3 RESULTS OF THE REVIEW
11	3.1 Chitosan-microbe interactions
12	3.2 Chitosan-plant interactions
13	3.3 Description of included studies
14	3.4 Effects of 1% chitosan on disease incidence
15	3.5 Effects of 1% chitosan on in-vitro mycelium growth
16	3.6 Effects of 1% chitosan on enzyme activities associated with host defence
17	4 DISCUSSION
18	5 CONCLUSIONS
19	References
20	Acknowledgements
21	Author Contributions
22	Conflicts of Interest
23	

Chitosan and Postharvest Decay of Fresh Fruit: Meta-Analysis of Disease Control and Antimicrobial and Eliciting Activities

Razieh Rajestary†, Lucia Landi† & Gianfranco Romanazzi*

Department of Agricultural, Food and Environmental Sciences, Marche Polytechnic University, Via Brecce Bianche 10, 60131 Ancona, Italy

***Corresponding author: Gianfranco Romanazzi**

Tel: +39-071-2204336

Email: g.romanazzi@univpm.it

†Razieh Rajestary and Lucia Landi contributed equally to the paper

Word count of text: 11401

Short version of title: **Chitosan and Postharvest Decay of Fruit**

Abstract

Consumers are increasingly aware of the importance of regular consumption of fresh fruit in their diet. Since fresh fruit are highly sensitive to postharvest decay, several investigations focused on the study natural compounds alternative to synthetic fungicides, to extend their shelf life. A long list of studies reported the effectiveness of the natural biopolymer chitosan in control of postharvest diseases of fresh fruit. However, these findings remain controversial, with many mixed claims in the literature. In this work, we used random-effects meta-analysis to investigate the effects of 1% chitosan on (i) postharvest decay incidence; (ii) mycelium growth of fungal pathogens *Botrytis cinerea*, *Penicillium* spp., *Colletotrichum* spp. and *Alternaria* spp.; and (iii) phenylalanine ammonia-lyase, chitinase and β -1,3-glucanase activities. Chitosan significantly reduced postharvest disease incidence (mean difference [MD], -30.22; $P < 0.00001$) and *in-vitro* mycelium growth (MD, -54.32; $P < 0.00001$). For host defence responses, there were significantly increased activities of β -1,3-glucanase (MD, 115.06; $P = 0.003$) and chitinase (MD, 75.95; $P < 0.0002$). This systematic review contributes to confirm the multiple mechanisms of mechanisms of action of chitosan, which has unique properties in the natural compound panorama. Chitosan thus represents a model plant protection biopolymer for sustainable control of postharvest decay of fresh fruit.

Keywords: defence related enzymes; fungal pathogens; natural antifungal compounds; plant protection; sustainable control of plant pathogens

1 INTRODUCTION

Postharvest fungal diseases can limit the storage period and shelf life, and thus market life, of fruit and vegetables, which results in serious economic losses worldwide (Oerke & Dehne, 2004; Romanazzi, Smilanick, Feliziani, & Droby, 2016; Palou & Smilanick, 2020). The global average loss due to the food postharvest reported by Food and Agriculture Organization, was estimated in North America, Europe and Oceania about 29%, compared to an average of about 38% in industrialized Asia, Africa, Latin America and South East Asia (Parfitt, Barthel, & Macnaughton, 2010; Food and Agriculture Organization of the United Nations, 2011; Sawicka, 2019).

The main fungal diseases (and their associated fungal pathogen) include: gray mold (*Botrytis cinerea* Pers.); Rhizopus rot (*Rhizopus stolonifer* Ehrenb.); anthracnose (*Colletotrichum* spp.); green mold (*Penicillium digitatum* Pers.); blue mold (*Penicillium italicum* Wehmer on citrus fruit, *P. expansum* Link on other fruit); and *Alternaria* rot (*Alternaria* spp.). The control of the causal fungal pathogens is therefore critical to extend the shelf-life of these fresh products (Prusky, 2011; Arah, Amaglo, Kumah, & Ofori, 2015). Despite the efficacy of synthetic fungicides in the control of postharvest decay, public concerns about chemical and toxic residues in food (Belden, McMurry, Smith, & Reilley, 2010; Mebdoua, 2018; Gonçalves et al., 2019; Liu, Yamdeu, Gong, & Orfila, 2020) and the increase in drug-resistant strains of many pathogens (Zuccolo et al., 2019) indicate the need for development of new strategies. Over the last few decades, there has been an increasing interest in the study of postharvest control methods that make use of natural resources (Palou, Smilanick & Droby, 2008; Talibi, Boubaker, Boudyach, & Ait Ben Aoumar, 2014; Souza, Yuk, Khoo, & Zhou, 2015; Guimarães, Abrunhosa, Pastrana, & Cerqueira, 2018; Ebrahimzadeh & Abrinbana, 2019; Liu et al., 2019; Liu, et al., 2020). Such alternative compounds can act as resistance inducers

and/or activators of plant defence mechanisms, or they can have strong antimicrobial activities against the main postharvest fungal pathogens (Romanazzi, Feliziani, Baños, & Sivakumar, 2017; Ribes, Fuentes, Talens, & Barat, 2018). However, only a few such natural fungicides have been approved for use as control agents for postharvest diseases, due to the strict regulatory policies for food safety. Among these, chitosan is a natural biocompatible polysaccharide emerged as a promising eco-friendly alternative to synthetic fungicides (Muzzarelli, 1983; Romanazzi, Feliziani, & Sivakumar, 2018; Betchem, Johnson, & Wang, 2019). To give some background, chitosan is a common name for the polysaccharide N-acetyl-D-glucosamine (Zargar, Asghari, & Dashti, 2015). The chitosan compound is obtained by deacetylation of chitin through exposure to NaOH solutions or to the enzyme chitinase. It is a functional cationic biopolymer that is widely studied and used across the world. Chitosan have many applications included food industry (Gutiérrez, 2017; da Silva, de Souza, & Dantas Lacerda, 2019; Morin-Crini, Lichtfouse, Torri, & Crini, 2019; Kabanov, & Novinyuk, 2020), cosmetology (Aranaz et al., 2018; Kaczmarek, Struszczyk-Swita, Li, Szczesna-Antczak, & Daroch, 2019) and human medicine (Tungland & Meyer, 2002; Leung, Liu, Koon, & Fung, 2006; Kofuji et al., 2010; Zhao et al., 2018).

Concerning the agriculture applications, the chitosan was the first compound in the list of basic substances approved in the European Union for plant protection purposes (Reg. EU 66 2014/563), for both organic agriculture and integrated pest management. For several years now, chitosan has been of interest in many studies that have shown that it can be used to prolong storage of an array of fruit and vegetables worldwide, where it has been shown to have three major activities: including biofilm formation on treated surfaces (El Ghaouth, Arul, Ponnampalam, & Boulet, 1991; Valencia-Chamorro, Palou, & Del Río, 2011; Romanazzi et al., 2018); as an antimicrobial (Goy, De Britto, & Assis, 2009; Kong, Chen, Xing, & Park, 2010; Feliziani, Landi, & Romanazzi, 2015; Cheung, Ng, Wong, & Chan, 2015; Wang, Li, & Zhang,

2017; Pétriacq, López, & Luna, 2018; Duan et al., 2019); and as an elicitor of host defence mechanisms (Landi, Feliziani, & Romanazzi, 2014; Coqueiro et al., 2015; Landi et al., 2017; Colman et al., 2019; Xoca-Orozco et al., 2019; Obianom, Romanazzi, & Sivakumar, 2019). For these reasons, chitosan can be used as a biodegradable fungicide (Rebelo, Vila, & Fanguero, R., 2016; Liang et al., 2017).

However, the heterogeneity of chitosan activities and its effectiveness across a wide range of experimental conditions have led to different interpretations of its primary use/mechanism/ actions. As a result, different recommendations for chitosan treatments have been provided (Ramos-García et al., 2012; Bill, Sivakumar, Korsten, & Thompson, 2014; Xing et al., 2016; Flores et al., 2018; Betchem et al., 2019; de Souza, Lundgren, de Oliveira, Berger, & Magnani, 2019). Furthermore, based on reports of the evaluation of chitosan across similar and different fungal strains, its value for disease reduction can vary (Herrera-Romero, Ruales, & Caviedes, 2017; Hua et al., 2019; Zahid, Maqbool, Ali, Siddiqui, & Bhatti, 2019). Also, despite the many studies in the literature that have investigated a wide range of chitosan treatments and their influences, no single study has made all of the appropriate comparisons for a full evaluation. Thus, given the mixed claims in the literature, there is the need to define the overall effectiveness of chitosan, to highlight useful aspects for its future investigation.

Meta-analyses can be applied as a tool for analysis of large amounts of data across many primary studies, in which the main purpose is to integrate and interpret the findings, to provide conclusions that the individual studies alone cannot show clearly. This statistical procedure provides an integration of the data across several to many independent studies (Maestri, Pavlicevic, Montorsi, & Marmioli, 2019). The combination of the resulting outcomes can also increase the statistical power, and make it possible to detect relatively small effects (Rosenberg, Garrett, Su, & Bowden, 2004; Nelson, Gent, & Grove., 2015; Schwingshackl, Hoffmann, Iqbal,

Schwedhelm, & Boeing, 2018; Chen, Chen, Chen, & Huang, 2019; González-Domínguez et al., 2019).

The aim of the present study was to carry out a meta-analysis to quantitatively review the data across the available studies on the effectiveness of 1% chitosan, the most common concentration that has been tested in the control of postharvest decay (Romanazzi et al., 2018). Hence, the objectives were to determine the effectiveness of 1% chitosan on: (i) reduction of postharvest diseases of fresh fruit; (ii) *in-vitro* mycelium growth of the causal agents of postharvest decay; and (iii) phenylalanine ammonia-lyase (PAL), β -1,3-glucanase and chitinase activities associated with host defence mechanisms against these causal agents at 24 h post-treatment (hpt).

2. METHODS

2.1 Search strategy and study selection

A systematic literature search from 2007 to 2019 was performed using the databases of Scopus and Web of Science and the following terms: ‘chitosan’ and ‘fruit’. Studies that used chitosan mixed with other compounds were not considered. The selection of studies was conducted according to the Preferred Reporting Items for Systematic Reviews and Meta-analyses (PRISMA) guidelines (Moher, Liberati, Tetzlaff, Altman, & PRISMA Group, 2009).

Article selection for the meta-analysis used the following inclusion criteria: 1% chitosan; disease incidence; *in-vitro* mycelium growth according to specific postharvest fungi; and activity of the enzymes involved in plant defence mechanisms. The eligibility of the articles was assessed, with the exclusion of the studies with different chitosan concentrations, with no information on disease incidence, mycelium growth or defence enzymes, and with no known fungal species.

In more detail, three categories were included for the studies related to: (i) disease incidence published from 2010 to 2019, caused by gray mold, *Rhizopus* rot, anthracnose, green/blue mold and/or *Alternaria* rot, considered as subgroups; (ii) *in-vitro* mycelium growth published from 2007 to 2019 for the decay causing fungal pathogens *B. cinerea*, *Penicillium* spp., *Colletotrichum* spp. and *Alternaria* spp., considered as subgroups; (iii) enzyme activities associated with host defence mechanisms analysed at 24 hpt published from 2009 to 2018, for PAL, chitinase and β -1,3-glucanase, considered as subgroups. All of the studies included at least two treatments, as an untreated control and the 1% chitosan treatment. The fruit varieties, the 1% chitosan application and the detection timing varied across these studies. In some studies, the treatment application times and rates were reported. In such cases, only the treatments applied at the same time as the standard treatment were considered in the meta-analysis. The risk of bias and test for asymmetry for the funnel plots were used to evaluate the publication bias. Cochran's I^2 indices, Tau^2 and χ^2 tests were used to estimate the statistical heterogeneity of the studies (Tufanaru, Munn, Stephenson, & Aromataris, 2015). If the heterogeneity was significant ($I^2 > 75\%$; and/or $P < 0.05$), a random effects model was applied to all of the subgroups included in the postharvest decay disease incidence, the decay causing fungi mycelium growth, and the defence enzyme activity categories.

2.2 Data extraction

Data were recorded from the same days of chitosan treatments in each study. All of the studies that were related to the effects of chitosan towards disease incidence were calculated as percentage effects. The studies on the effects on mycelium growth resulted on three different measurement units (percentage, mm, cm), and again these were converted to percentages. To unify the different measurement units used across the studies of the defence enzyme activities, the values were converted into percentage of the mean (% mean) with respect to the normal

control ([treatment mean/ normal control mean] $\times$ 100) (Viswanatha, Shylaj, & Moolemath, 2017). If the standard deviations (SDs) or standard errors (SEs) were not reported, the data were transformed according to the P values (Weir et al., 2018). Data were extracted from the Figures presented in the papers using Plot Digitiser software (Kadic, Vucic, Dosenovic, Sapunar, & Puljak, 2016). The change scores with the corresponding standard deviations were used, as based on the guidelines of the Cochrane handbook (<https://www.cochranelibrary.com/cdsr/doi/10.1002/14651858.CD012276/epdf/full>).

2.3 Data analysis

All of these meta-analyses were conducted using the Review Manager (RevMan) software, version 5.3. (Copenhagen: The Nordic Cochrane Centre, The Cochrane Collaboration, 2014; <http://tech.cochrane.org/revman>). The data type was selected as continuous. The statistical method was considered as inverse variance. Weighted means, effect sizes, 95% confidence intervals (CIs), which included 0, were calculated. In all of these analyses, P-value <0.05 was considered statistically significant. Differences among the groups were defined when the 95% CIs overlapped a vertical line. If the 95% CIs did not overlap, it can be suggested that the differences were significant (Yang, Scott, Mao, Tang, & Farmer, 2014; Dardiotis et al., 2018). The studies are presented as Forrest plots in the order of the statistical power.

3 RESULTS OF THE REVIEW

3.1 Chitosan-microbe interactions

The antimicrobial activity of chitosan is a complex process that depends significantly from intrinsic properties and environmental factors (Yilmaz Atay, 2019) as well as the type of bacteria, fungi or virus involved (Chirkov, 2002; Kong, et al., 2010; Hosseinnnejad, & Jafari, 2016). The precise mechanism of chitosan antimicrobial activity is still not completely

understood. Several studies have suggested that the antimicrobial action is mainly due to the polycationic structure of the chitosan. Several studies have suggested that the antimicrobial action is mainly due to the polycationic structure of the chitosan. This activity is carried out in a pH range among 5.6 and 6 (Romanazzi, Gabler, Margosan, Mackey, & Smilanick, 2009) that is below the pKa of chitosan. The chitosan, positively charged, reacts with negatively charged microbial cell membranes (Rabea, Badawy, Stevens, Smagghe, & Steurbaut, 2003; Goy et al., 2009; Kong et al., 2010). This bond alters the permeability of the membrane which is followed by an inhibition of DNA replication and subsequently cell death (Nagy et al., 2011; Divya, Vijayan, George, & Jisha, 2017). A chelating action was also observed. The chitosan molecule binds to the metallic elements present in the trace causing the inhibit of toxins production and microbial growth (Cuero, Osuji, & Washington, 1991; Chung, Wang, Chen, & Li, 2003). The effect of chitosan on fungal pathogens was to inhibits the radial growth, spore germination, and the elongation of the germ tube as well as the production of virulence factors (Palma-Guerrero, Jansson, Salinas, & Lopez-Llorca, 2008; Badawy, & Rabea, 2011).

3.2 Chitosan-plant interactions

The chitosan acts as a powerful elicitor able to inducing a defense response against pathogens in plant tissues by activating both, a local (Zuppini et al., 2003; Iriti, & Varoni, 2015) and systemic plant defense (Benhamou, Lafontaine, & Nicole, 1994; Xing, Zhu, Peng, & Qin, 2015) with the involvement several molecules related to defense mechanisms as pathogenesis-related (PR) proteins (Lopez-Moya et al., 2017; Corsi, Forni, Riccioni, & Linthorst, 2017), Reactive Oxygen Species (ROS) (Singh et al., 2019) and secondary metabolites with active roles in defense as lignin, callose, phytoalexins, PAL, peroxidases and chitinase (Ma, Yang, Yan, Kennedy, & Meng, 2013; Landi et al., 2014; Malerba, & Cerana, 2016). However, the chitosan elicitation activity depends on the reactivity of the host tissues (Romanazzi et al., 2016) as well

as from the acetylation and degree polymerization of chitosan (Cord-Landwehr, Melcher, Kolkenbrock, & Moerschbacher, 2016; Li, Xing, Liu, & Li, 2016). Until now the chitosan binding receptors are undefined (Iriti & Faoro 2009; Hidangmayum, Dwivedi, Katiyar, & Hemantaranjanm, 2019). Some researches proposed that chitosan could also interact with chromatin and directly affect gene expression (Hadwiger & Polashock, 2013; Katiyar, Hemantaranjan, Bharti, & Nishant Bhanu, 2014). However, chitosan molecular signals are transduced by messengers such as ROS or phytohormones able to induce physiological and defense response by host (Yin, Li, Zhao, Du, & Ma, 2006; Hidangmayum et al., 2019).

An effect often observed on plants tissue after chitosan treatment was the inhibition of light-induced stomatal opening (Lee et al., 1999; Iriti et al., 2009). On this regard, the transcriptome analysis performed on sweet orange (Coqueiro et al., 2015) and strawberry (Landi et al., 2017) after chitosan treatments underline early impact of compound on the light photosynthetic process affecting imbalance/balance of ROS/redox signaling (Landi et al., 2017). These entire signaling molecules contribute to the adaptive mechanism in chitosan treated plants in response to stress.

3.3 Description of included studies

A flow chart of the screening of the studies identified for the effectiveness of 1% chitosan is shown in Figure 1, with a total of 56 articles finally available for the meta-analysis according to the search criteria. These covered 117 studies, of which 49 were related to disease incidence (total cases, 8,543 [for each of control and chitosan treatment]) (Figure 2), 41 to *in-vitro* mycelium growth (total cases, 1,072) (Figure 3), and 27 to changes in defence-mechanism-related enzymes (total cases, 1,332) (Figure 4). Some of the relevant details of the articles that were included in this meta-analysis are given in Table 1. All of the selected articles were included in the assessment for risk of bias. Also, blinding of outcome assessment in these

studies (i.e., performance bias) was not necessary, so it was not included in the analysis for risk of bias. The domains considered for risk of bias were chosen based on each study that reported data and scientific information. All of the studies provided specific indication that the basic characteristics of the control and treatment groups were balanced and were treated under similar environmental conditions. None of these studies included misleading samples. As a result, the selection, detection, attrition and reporting were free of bias, and the publications were defined as at low risk of bias. The funnel plots constructed from the data for disease incidence, mycelium growth and defence enzyme activities did not reveal any significant asymmetry (Figure 5).

3.4 Effects of 1% chitosan on disease incidence

Based on this meta-analysis, the overall data demonstrated the significant effectiveness of 1% chitosan over the control treatment for reduction of disease incidence (studies, 49; total cases, 8,5473) (mean difference [MD], -30.22; 95% confidence intervals [CI], -36.48 to -23.96; I^2 , 90.0%; $P < 0.00001$) (Figure 2). The subgroup analysis here (Figure 2) showed that 1% chitosan was significantly effective for reduction of disease incidence against: gray mold (studies, 12; total cases, 1,473), (Shao, Tu, Tu, & Tu, 2012; Feliziani, Santini, Landi, & Romanazzi, 2013; Gao, Zhu, & Zhang, 2013; Romanazzi, Feliziani, Santini, & Landi, 2013; Feliziani et al., 2015; Kanetis, Exarchou, Charalambous, & Goulas, 2017; Zheng, et al., 2017; Gramisci, Lutez, Lopes, & Sangorrína, 2018; Hajji, Younes, Affes, Boufi, & Nasri, 2018) (MD, -23.97; 95% CI, -32.25 to -15.68; I^2 , 77.0%; $P < 0.00001$), as highly effective in 9 of these studies, (Shao et al., 2012; Gao et al., 2013; Romanazzi et al., 2013; Feliziani et al., 2015; Kanetis et al., 2017; Zheng, et al., 2017; Gramisci et al., 2018; Hajji et al., 2018); blue/green molds caused by *Penicillium* spp. (studies, 16; total cases, 1,968) (Xing, Xu, Che, Li, & Li, 2011; Shao et al., 2012; Cháfer, Sánchez-González, González-Martínez & Chiralt, 2012; Feliziani et al., 2013;

Romanazzi et al., 2013; Wang, Wu, Qin, & Meng, 2014; Lu et al., 2014; Shao et al., 2015; El Guilli, Hamza, Clément, Ibriz, & Ait Barka, 2016; Zheng, et al., 2017; Gramisci et al., 2018; Kharchoufi, et al., 2018; Liu, Sun, Xiu, Huang, & Zhou, 2018; Shi, Wang, Lu, & Deng, 2018) (MD, -30.85; 95% CI, -41.91 to -19.79; I^2 , 90.0%; $P < 0.00001$), as highly effective in 9 of these studies (Xing et al., 2011; Romanazzi et al., 2013; Lu, et al., 2014; Shao et al., 2015; El Guilli et al., 2016; Zheng, et al., 2017; Liu et al., 2018; Shi et al., 2018); *Rhizopus* rot (studies, 5; total cases, 1,740) (Cia, Benato, Pascholati, & Garcia, 2010; Ramos-García et al., 2012; Romanazzi et al., 2013; Xing et al., 2015) (MD, -28.80; 95% CI, -46.13 to -11.47; I^2 , 87.0%; $P = 0.001$), as effective in 3 of these studies (Cia et al., 2010; Ramos-García et al., 2012; Romanazzi et al., 2013); and anthracnose (11 studies; total cases, 2,134) (Maqbool, Ali, Ramachandran, Smith, & Alderson, 2010; Zahid, Ali, Manickam, Siddiqui, & Maqbool, 2012; Bill et al., 2014; Edirisinghe, Ali, Maqbool, & Alderson, 2014; Ali, Noh, & Mustafa, 2015; Gutiérrez-Martínez, Bautista-Banos, Berúmen-Varela, Ramos-Guerrero, & Hernández-Ibanez, 2017; Obianom et al., 2019) (MD, -46.64; 95% CI, -61.54 to -31.73; I^2 , 92.0%; $P < 0.00001$), as effective in all of these studies. For *Alternaria* rot, 1% chitosan was not significantly effective (studies, 5; total cases, 1,228) (Meng, Yang, Kennedy, & Tian, 2010; Yan et al., 2011; López-Mora, Gutiérrez-Martínez, Bautista-Baños, Jiménez-García, & Zavaleta-Mancera, 2013; Feliziani et al., 2015; Guo, Xing, Yu, Zhao, & Zhu, 2017) (MD, -8.50; 95% CI, -15.75 to -1.25; I^2 , 27.0%; $P = 0.24$), although in 1 of these studies (Guo et al., 2017) its effect reached significance.

3.5 Effects of 1% chitosan on in-vitro mycelium growth

The overall data here showed the significant effectiveness of 1% chitosan over the control treatment against *in-vitro* mycelium growth of these fungal pathogens that are involved in postharvest diseases (studies, 41; total cases, 1,072) (MD, -54.32; 95% CI, -64.35 to -44.28;

I², 95.0%; P <0.00001) (Figure 3). The subgroup analysis here (Figure 3) showed that 1% chitosan was significantly effective against *in-vitro* mycelium growth for: *B. cinerea* (studies, 5; total cases, 37) (Kanetis et al., 2017; Xu et al., 2007; Feliziani et al., 2013; Munhuweyi et al., 2017; Flores et al., 2018). (MD, -49.38; 95% CI, -72.98 to -25.79; I², 94.0%; P <0.0001), as medium high effects for all of these studies; *Penicillium* spp. (studies, 9; total cases, 65) (Xing et al., 2011; Abdel-Kader, El-Mougy & Lashin, 2011; Nisia, Noreña, & Brandelli, 2012; Wang et al., 2014; Waewthongrak, Pisuchpen, & Leelasuphakul, 2015; Shao et al., 2015; Munhuweyi et al., 2017; Madanipour, et al., 2019) (MD, -73.00; 95% CI, -89.71 to -56.30; I², 92.0%; P <0.00001), as the highest effects seen, and for all of these studies; *Colletotrichum* spp. (studies, 24; total cases, 955) (Jitareerat, Paumchai, Kanlayanarat, & Sangchote, 2007; Rahman, Mahmud, Kadir, Abdul Rahman, & Begum, 2008; Munoz, Moret, & Garces, 2009; Maqbool et al., 2010; Zahid et al., 2012; Mohamed, Clementine, Didier, Gérard, & Noëlle, 2013; Ali et al., 2014; Bill et al., 2014; Edirisinghe et al., 2014; Ali et al., 2015; Varela, Coronado Partida, Ochoa Jiménez, López, & Martínez, 2015; Gutiérrez-Martínez et al., 2017; de Oliveira, Berger, de Araújo, Camara, & de Souza, 2017; Ramos-Guerrero, González-Estrada, Hanako-Rosas, & Bautista-Banños, 2018; Xoca-Orozco, Aguilera-Aguirre, López-García, Gutiérrez-Martínez, & Chacón-López, 2018) (MD, -48.18; 95% CI, -62.83 to -33.53; I², 96.0%; P <0.00001), as the lowest effects seen based on the point estimate, with the highest effects for 16 of these studies (Jitareerat, et al., 2007; Rahman, et al., 2008; Maqbool et al., 2010; Zahid et al., 2012; Bill et al., 2014; Ali et al., 2014; Varela et al., 2015; de Oliveira et al., 2017; Ramos-Guerrero et al., 2018; Xoca-Orozco et al., 2018); and *Alternaria* spp. (3 studies; total cases, 15) (Yan et al., 2011; Feliziani et al., 2013; López-Mora et al., 2013) (MD, -55.20; 95% CI, -80.50 to -29.90; I², 90.0%; P <0.0001), as significant for all of these studies.

3.6 Effects of 1% chitosan on enzyme activities associated with host defence

The overall data for the effects of 1% chitosan on the activities of the enzymes associated with host plant defence at 24 hpt showed significantly increased activity over the control treatment (studies, 27; total cases, 1,332) (MD, 74.58; 95% CI, 41.15 to 108.01; I^2 , 99.0%; $P < 0.0001$) (Figure 4). For the details of the subgroup analysis here (Figure 4), in the treated fruit, 1% chitosan did not induce any significant difference compared to the control at 24 hpt for the PAL activity (studies, 9; total cases 575) (Zahid et al., 2012; Landi et al., 2014; Bill et al., 2014; Shao et al., 2015; Waewthongrak et al., 2015; Song et al., 2016; Jongsri, Rojsitthisak, Wangsomboondee, & Seraypheapa, 2017; Shen & Yang, 2017; Silva et al., 2018) (MD, 37.06; 95% CI, -17.28 to 91.40; I^2 , 99.0%; $P = 0.18$). However, 5 of these studies (Landi et al., 2014; Bill et al., 2014; Shao et al., 2015; Waewthongrak et al., 2015; Shen & Yang, 2017) showed significant increases in PAL activity. Furthermore, significant increases were seen overall for chitinase activity (10 studies; total cases, 491) (Hewajuliage, Sultanbawa, Wijeratnam, & Wijesundara, 2009; Feliziani et al., 2013; Bill et al., 2014; Landi et al., 2014; Ali et al., 2014; Shao et al., 2015; Jongsri, et al., 2017; Shen, & Yang, 2017) (MD, 75.95; 95% CI, 36.18 to 115.73; I^2 , 99.0%; $P = 0.0002$), as 8 of these with significance increases (Hewajuliage, et al., 2009; Feliziani et al., 2013; Landi et al., 2014; Bill et al., 2014; Ali et al., 2014; Jongsri, et al., 2017; Shen, & Yang, 2017), and overall for β -1,3-glucanase activity (8 studies; total cases 266) (Hewajuliage, et al., 2009; Wang & Gao, 2013; Landi et al., 2014; Bill et al., 2014; Ali et al., 2014; Shao et al., 2015; Jongsri, et al., 2017; Shen, & Yang, 2017) (MD, 115.06; 95% CI, 38.24 to 191.88; I^2 , 100.0%; $P = 0.003$), as 5 of these with significance increases (Hewajuliage, et al., 2009; Wang & Gao, 2013; Landi et al., 2014; Bill et al., 2014; Ali et al., 2014).

4 DISCUSSION

This study brings together and summarises the results from the literature of the effects of 1% chitosan on postharvest diseases and pathogens, according to disease incidence, *in-vitro*

mycelium growth, and induction of host defence responses through monitoring of the most commonly analysed enzymes linked to defence mechanisms. This meta-analysis emphasises the primary role of 1% chitosan against the main diseases and pathogens associated with postharvest decay (Romanazzi et al., 2018; Betchem et al., 2019). These pooled estimates highlighted that 1% chitosan is effective against the main postharvest diseases caused by several fungal pathogens that infect different plant species. Although some of these data show high heterogeneity, they also show low risk of bias and high validity for each study, with no substantial baseline differences seen between the control and treatment groups. Indeed, the funnel plots as a method to assess the potential role of publication bias (Harbord, Egger, & Sterne, 2006) indicate that no bias was detected across the studies included. Therefore, these values of $I^2 > 90\%$ indicate real differences in these studies.

Our study underlines the transversal effectiveness of chitosan in postharvest disease management. Here, the subgroup analysis of *in-vitro* mycelium growth emphasises that the most powerful growth reduction was for *Penicillium* spp., followed by *Alternaria* spp. and *B. cinerea*, while lower effectiveness was seen against *Colletotrichum* spp..

These data also show that chitosan has differential effects across these fungal species, potentially through the control of fungal development and lytic enzyme activation by chitosan (El Gueddari, Rauchhaus, Moerschbacher & Deising, 2002; Geoghegan & Gurr, 2016; Geoghegan, Steinberg, & Gurr, 2017; Ramos-Guerrero et al., 2018; Ramos-Guerrero, González-Estrada, Romanazzi, Landi, & Gutiérrez-Martínez, 2020). There are direct links between the cell wall and cell membranes, as the synthesis of key cell-wall components (e.g., glucans, chitin) occurs at the plasma membrane, with the associated synthase enzyme complexes (Maddi, & Free, 2010). The chitin is localized in the membrane proximal portion of the cell wall and is incorporated into the wall matrix by being cross-linked to the glucans (Patel & Free, 2019). Previous studies have investigated the role of plasma membrane in the sensitivity

of fungi to chitosan showing that the plasma membrane of chitosan-sensitive fungi is more fluid and richer in polyunsaturated free fatty acids than in chitosan-resistant fungi (Palma-Guerrero et al., 2009 and 2010). The authors evidenced that chitosan binds to negatively charged phospholipids. This alter plasma membrane fluidity to inducing the membrane permeabilization, which was greatest in membranes containing elevated content polyunsaturated lipids.

While this meta-analysis highlights the different reactions between the fungal species and chitosan effectiveness, it also underlines the key role of plant species in this complex relation that significantly affects the outcome of chitosan-pathogen interaction.

For this reason, the fungal pathogens can react differently to chitosan in terms of disease incidence and in *in-vitro* tests. Indeed, the meta-analysis summarized studies related to disease incidence, show significantly reducing postharvest disease incidence, although the results linked to singular disease show the highest effectiveness of chitosan against anthracnose, while it is less effective against blue/green mold, Rhizopus rot, gray mold, and particularly Alternaria rot. Therefore, it is not excluded that the involvement of mainly different fruits species on anthracnose incidence, as banana, papaya, dragon, bell pepper, soursop and avocado, not tested for the other diseases, the chitosan, could be elicited a different defence response.

This study also confirms that disease incidence is the result of a combination of the chitosan effects on film-forming, plant defence eliciting, and its antimicrobial properties (Romanazzi et al., 2018). In this context, chitosan can be considered to be a modulator of plant defences (Lopez-Moya, Suarez-Fernandez, & Lopez-Lorca, 2019). Chitosan application to plants fits into the delicate relationship between the host and pathogenic fungi and involves the primary cell-wall defence mechanisms. A link between pathogenicity and the enzymes that synthesise the fungal cell wall has been demonstrated in numerous studies (Arana et al., 2009; Leviansky et al., 2010; Lenardon, Munro, & Gow, 2010; Oliveira-Garcia, & Deising, 2013;

Geoghegan et al., 2017; Patel & Free, 2019), and depolymerisation of the cell walls of plant pathogenic fungi following the infection, evading plant immune recognition, has been reported (Geoghegan et al., 2017). It has been reported that the strategy of some fungal pathogens to evade plant immunity is to convert chitin into chitosan (Lopez-Moya, et al., 2019). Thus, both chitosan and chitin will have key roles in the control of plant immunity.

According to the concepts of systemic acquired resistance (Pieters et al., 1998; Durrant & Dong, 2004) and induced systemic resistance (Heil & Bostock, 2002; Timmermann, González, & Ruz, 2020), chitosan can induce resistance in the plants to control postharvest fungal pathogens of their fruit and as vegetables (Nandeeshkumar et al., 2008; Jia, Meng, Zeng, Wang, & Yin, 2016; Jia, Zeng, Wang, Zhang, & Yin, 2018). On this basis, the meta-analysis data related to the eliciting of the host defence enzymes by chitosan through activation of induced resistance can help us to understand this aspect (Mandal, Kar, Mukherjee, & Acharya, 2013; Walters, Ratsep, & Havis, 2013).

Although a meta-analysis of publicly available data, related to transcriptome investigations of plants defense priming, evidenced a common set of conserved transcriptional changes on plants upon stress conditions, (Bacelli, Benny, Caruso, & Martinelli, 2020), the detailed role of the chitosan in the induction of defence mechanisms has been shown for sweet oranges (Coqueiro et al., 2015) and strawberries (Landi et al., 2017). The most common approaches related to the study of enzyme activities (Wang & Gao, 2013; Ali et al., 2014; Pasquariello et al., 2015; Shao et al., 2015; Adiletta, Zampella, Coletta, & Petriccione, 2019) and the expression of individual genes (Ma et. al., 2013; Landi et al., 2014; Petriccione et al., 2017; Fooladi vanda, Shabani, & Razavizadeh, 2019; Chun & Chandrasekaran, 2019) have been investigated, both of which are associated with reactive oxygen species, specific PR proteins, cell-wall enzymes and secondary metabolites. Usually, these individual studies have

shown wide variability associated with host fruit species, application methods and times of treatment.

In the present study, we analysed the most studied of the plant defence enzymes, PAL, which is associated with the phenylpropanoid pathway (Dixon, Laphorn, & Edwards, 2002; Yadav et al., 2020), and chitinase and β -1,3-glucanase, which are linked to cell-wall hydrolysis (Gupta et al., 2015; Pusztahelyi, 2018), at the main analysis time point of 24 hpt. These data do not show any significant effects of chitosan on PAL activity at 24 hpt, while high increases in the activities of chitinase and β -1,3-glucanase were detected, independent of the host species. These findings are in agreement with the plant immunity mechanisms that indicate that chitinase and β -1,3-glucanase release the glucan oligomers from the chitin of the fungal cell walls to trigger the plant immune responses (Jones & Dang, 2006; Fesel & Zuccaro, 2016; Lopez-Moya et al., 2019;), although the induction of these defence mechanisms can vary greatly according to the time of treatment. The present study suggests that the analysis of the chitinase and β -1,3-glucanase activities at 24 hpt represents a marker for verification of induction of the plant defences by chitosan, while activation of PAL has generally been reported to occur at later times (Landi et al., 2014; Bill et al., 2014).

5 CONCLUSIONS

The present work established the first comprehensive investigation of chitosan effectiveness on postharvest pathogens using meta-analysis approach. This study provides knowledge based on three robust findings, as the effects of 1% chitosan on disease incidence, mycelium growth of decay-causing fungi, and the activities of two important defence enzymes in particular, chitinase and β -1,3-glucanase. This investigation shown the chitosan have antifungal properties against different phytopathogens highlight the versatile properties of this natural biopolymer.

It was demonstrated there are enough data about the effectiveness of chitosan in the control of postharvest diseases, also inducing resistance on fruit to postharvest pathogens.

The outcomes of this study aim to contribute to a better understanding concerning the role of chitosan in the control of postharvest decay of fresh fruit, that will be relevant for the conceptualization and measurement of future studies. Collectively, these data confirm the multiple mechanisms of action of chitosan, which has unique properties in the panorama of activities of natural compounds that define it as a model plant-protection agent for sustainable control of postharvest decay of fruit and vegetables.

References

- Abdel-Kader, M., El-Mougy, N., & Lashin, S. (2011). Evaluation of grapefruit coating with chemical preservatives as control measure against postharvest decay. *Phytopathologia*, 59, 25-38. http://www.up.poznan.pl/~ptfit1/pdf/P59/P59_03.pdf
- Adiletta, G., Zampella, L., Coletta, C., & Petriccione, M. (2019). Chitosan coating to preserve the qualitative traits and improve antioxidant system in fresh figs (*Ficus carica* L.). *Agriculture*, 9(4), 84. <https://doi.org/10.3390/agriculture9040084>
- Ali, A., Zahid, N., Manickam, S., Siddiqui, Y., Alderson, P. G., & Maqbool, M. (2014). Induction of lignin and pathogenesis related proteins in dragon fruit plants in response to submicron chitosan dispersions. *Crop Protection*, 63, 83-88. <https://doi.org/10.1016/j.cropro.2014.05.009>
- Ali, A., Noh, N. M., & Mustafa, M. A. (2015). Antimicrobial activity of chitosan enriched with lemongrass oil against anthracnose of bell pepper. *Food Packaging and Shelf Life*, 3, 56-61. <https://doi.org/10.1016/j.fpsl.2014.10.003>
- Arah, I. K., Amaglo, H., Kumah, E. K., & Ofori, H. (2015). Preharvest and postharvest factors affecting the quality and shelf life of harvested tomatoes. A mini review. *International Journal of Agronomy*, 2015, 6. <https://doi.org/10.1155/2015/478041>
- Arana, D. M., Prieto, D., Román, E., Nombela, C., Alonso-Monge, R., & Pla, J. (2009). The role of the cell wall in fungal pathogenesis. *Microbial Biotechnology*, 2(3), 308–320. <https://doi.org/10.1111/j.1751-7915.2008.00070.x>
- Aranaz, I., Acosta, N., Civera, C., Elorza, B., Mingo, J., Castro, C., ... Heras-Caballero, A. (2018) Cosmetics and cosmeceutical applications of chitin, chitosan and their derivatives. *Polymers*, 10, 213. <https://doi.org/10.3390/polym10020213>.

Baccelli, I., Benny, J., Caruso, T., & Martinelli, F. (2020). The priming fingerprint on the plant transcriptome investigated through meta-analysis of RNA-seq data. *European Journal of Plant Pathology*, 156(3), 779-797. <https://doi.org/10.1007/s10658-019-01928-3>

Badawy, M. E. I., & Rabea, E. I. (2011). A biopolymer chitosan and its derivatives as promising antimicrobial agents against plant pathogens and their applications in crop protection. *International Journal of Carbohydrate Chemistry*, ID 460381, <https://doi.org/10.1155/2011/460381>

Belden, J., McMurry, S., Smith, L., & Reilley, P. (2010). Acute toxicity of fungicide formulations to amphibians at environmentally relevant concentrations. *Environmental Toxicology and Chemistry*, 29(11), 2477-2780. <https://doi.org/10.1002/etc.297>

Benhamou, N., Lafontaine, P. J., & Nicole, M. (1994). Seed treatment with chitosan induces systemic resistance to *Fusarium* crown and root rot in tomato plants. *Phytopathology*, 84(12), 1432-1444. <https://doi.org/10.1094/phyto-84-1432>

Betchem, G., Johnson, N. A. N., & Wang, Y. (2019). The application of chitosan in the control of postharvest diseases: a review. *Journal of Plant Diseases and Protection*, 126, 495-597. <https://doi.org/10.1007/s41348-019-00248-2>

Bill, M., Sivakumar, D., Korsten, L., & Thompson, K. (2014). The efficacy of combined application of edible coatings and thyme oil in inducing resistance components in avocado (*Persea americana* Mill.) against anthracnose during postharvest storage. *Crop Protection*, 64, 159-167. <https://doi.org/10.1016/j.cropro.2014.06.015>

Cháfer, M., Sánchez-González, L., González-Martínez, C., & Chiralt A. (2012). Fungal decay and shelf life of oranges coated with chitosan and bergamot, thyme, and tea tree essential oils. *Journal of Food Science*, 77, 182-187. <https://doi.org/10.1111/j.1750-3841.2012.02827.x>

Chen, C., Chen, H. Y. H., Chen, X., & Huang Z. (2019). Meta-analysis shows positive effects of plant diversity on microbial biomass and respiration. *Nature Communications*, 10, 1332. <https://doi.org/10.1038/s41467-019-09258-y>

Cheung, R. C. F., Ng, T. B., Wong, J. H., & Chan W. Y. (2015). Chitosan: an update on potential biomedical and pharmaceutical applications. *Marine Drugs*, 13(8), 5156-5186. <https://doi.org/10.3390/md13085156>

Chirkov, S. N. (2002). The antiviral activity of chitosan (review). *Applied Biochemistry and Microbiology*, 38, 1–8. <https://doi.org/10.1023/A:1013206517442>

Chun, S. C., & Chandrasekaran, M. (2019). Chitosan and chitosan nanoparticles induced expression of pathogenesis-related proteins genes enhances biotic stress tolerance in tomato. *International Journal of Biological Macromolecules*, 125, 948-954. <https://doi.org/10.1016/j.ijbiomac.2018.12.167>

Chung, Y. C., Wang, H. L., Chen, Y. M., & Li, S. L. (2003). Effect of abiotic factors on the antibacterial activity of chitosan against waterborne pathogens. *Bioresource Technology* 88(3), 179–184. [https://doi.org/10.1016/S0960-8524\(03\)00002-6](https://doi.org/10.1016/S0960-8524(03)00002-6)

Cia, P., Benato, E. A., Pascholati, S. F., & Garcia, E. O. (2010). Chitosan on the postharvest control of soft rot in ‘Rama Forte’ persimmon. *Bragantia*, 69, 745-752. <https://doi.org/10.1590/S0006-87052010000300028>

Colman, S. L., Salcedo, M. F., Mansilla, A. Y., Iglesias M. J., Fiol, D. F., Saldaña, S. M., ... Casalongué, C. A. (2019). Chitosan microparticles improve tomato seedling biomass and modulate hormonal, redox and defense pathways. *Plant Physiology and Biochemistry*, 143, 203-211. <https://doi.org/10.1016/j.plaphy.2019.09.002>

Coqueiro, D. S., de Souza, A. A., Takita, M. A., Rodrigues, C. M., Kishi, L. T., & Machado, M. A. (2015). Transcriptional profile of sweet orange in response to chitosan and salicylic acid. *BMC Genomics*, 16, 288. <https://doi.org/10.1186/s12864-015-1440-5>

Cord-Landwehr S., Melcher R. L. J., Kolkenbrock S., & Moerschbacher B. M. (2016). A chitin deacetylase from the endophytic fungus *Pestalotiopsis* sp. efficiently inactivates the elicitor activity of chitin oligomers in rice cells. *Scientific Reports*, 6,38018. <https://doi.org/10.1038/srep38018>

Corsi, B., Forni, C., Riccioni, L. & Linthorst, J. M. H. (2017). Enhancement of *PR1* and *PR5* gene expressions by chitosan treatment in kiwifruit plants inoculated with *Pseudomonas syringae* pv. *actinidiae*. *European Journal of Plant Pathology*, 148, 163–179. <https://doi.org/10.1007/s10658-016-1080-x>

Cuero, R. G., Osuji, G., & Washington, A. (1991) N-carboxymethylchitosan inhibition of aflatoxin production: Role of zinc. *Biotechnology Letters*, 13,441–444. <https://doi.org/10.1007/BF01030998>

da Silva, S. B., de Souza, D., & Dantas Lacerda, L. (2019). Food applications of chitosan and its derivatives. In L. A. M. van den Broek, & C. G. Boeriu (Eds.), *Chitin and chitosan: properties and applications* (1st ed., pp. 315–347). John Wiley & Sons Ltd. <https://doi.org/10.1002/9781119450467.ch13>

Dardiotis, E. Tsouris, Z., Mentis, A. F. A., Siokas, V., Michalopoulou, A., Sokratous, M., ... Kountouras, J. (2018). H. pylori and Parkinson's disease: meta-analyses including clinical severity. *Clinical Neurology and Neurosurgery*, 175, 16-24. <https://doi.org/10.1016/j.clineuro.2018.09.039>

de Oliveira, K. A. R., Berger, L. R. R., de Araújo, S. A., Camara, M. P. S., & de Souza, E. L. (2017). Synergistic mixtures of chitosan and *Mentha piperita* L. essential oil to inhibit *Colletotrichum* species and anthracnose development in mango cultivar Tommy Atkins. *Food Microbiology*, 66, 96-103. <https://doi.org/10.1016/j.fm.2017.04.012>

de Souza, E. L., Lundgren, G A., de Oliveira, K. Á. R., Berger, L. R. R., & Magnani, M. (2019). An Analysis of the Published Literature on the Effects of Edible Coatings Formed

by Polysaccharides and Essential Oils on Postharvest Microbial Control and Overall Quality of Fruit. *Comprehensive Reviews in Food Science and Food Safety*, 18(6), 1947,1967. <https://doi.org/10.1111/1541-4337.12498>.

Divya, K., Vijayan, S., George, T.K., & Jisha, M. S. (2017). Antimicrobial properties of chitosan nanoparticles: Mode of action and factors affecting activity. *Fibers and Polymers*, 18, 221–230. <https://doi.org/10.1007/s12221-017-6690-1>

Dixon, D. P., Laphorn, A., & Edwards, R. (2002). Plant glutathione transferases. *Genome Biology*, 3, reviews3004.1. <https://doi.org/10.1186/gb-2002-3-3-reviews3004>.

Duan, C., Meng, X., Meng, J., Khan, I. H., Dai, L., Khan, A., ... Ni, Y. (2019). Chitosan as a preservative for fruits and vegetables: a review on chemistry and antimicrobial properties. *Journal of Bioresources and Bioproducts*, 4(1), 11-21. <https://doi.org/10.21967/jbb.v4i1.189>.

Durrant, W. E., & Dong, X. (2004). Systemic Acquired Resistance. *Annual Review of Phytopathology*, 42,185-209. <https://doi.org/10.1146/annurev.phyto.42.040803.140421>

Ebrahimzadeh, F., & Abrinbana, M. (2019). Activity of fungicide mixtures against *Botrytis cinerea* isolates resistant to benzimidazoles, strobilurins and dicarboximides. *Annals of Applied Biology*, 174(3), 301-312. <https://doi.org/10.1111/aab.12497>

Edirisinghe, M., Ali, A., Maqbool, M., & Alderson, P. G. (2014). Chitosan controls postharvest anthracnose in bell pepper by activating defense-related enzymes. *Journal of Food Science and Technology*, 51(12), 4078-4083. <https://doi.org/10.1007/s13197-012-0907-51>

El Ghaouth, A., Arul, J., Ponnampalam, R., & Boulet, M. (1991). Use of chitosan coating to reduce water loss and maintain quality of cucumbers and bell pepper fruits. *Journal of Food Processing and Preservation*, 15(5), 359-368. <https://doi.org/10.1111/j.1745-4549.1991.tb00178.x>

El Gueddari, N. E., Rauchhaus, U., Moerschbacher, B. M., & Deising, H. B. (2002). developmentally regulated conversion of surface exposed chitin to chitosan in cell walls of plant pathogenic fungi. *New Phytologist*, 156(1), 103-112. <https://doi.org/10.1046/j.1469-8137.2002.00487.x>

El Guilli, M., Hamza, A., Clément, C., Ibriz, M., & Ait Barka, E. (2016). Effectiveness of postharvest treatment with chitosan to control citrus green mold. *Agriculture*, 6(2), 12. <https://doi.org/10.3390/agriculture6020012>.

Feliziani, E., Santini, M., Landi, L., & Romanazzi, G. (2013). Pre- and postharvest treatments with alternatives to synthetic fungicides to control postharvest decay of sweet cherry. *Postharvest Biology and Technology*, 78, 133-138. <https://doi.org/10.1016/j.postharvbio.2012.12.004>

Feliziani, E., Landi, L., & Romanazzi, G. (2015). Preharvest treatments with chitosan and other alternatives to conventional fungicides to control postharvest decay of strawberry. *Carbohydrate Polymers*, 132, 111-117. <https://doi.org/10.1016/j.carbpol.2015.05.078>

Fesel, P. H., & Zuccaro, A. (2016). β -glucan: Crucial component of the fungal cell wall and elusive MAMP in plants. *Fungal Genetics and Biology*, 90, 53-60. <https://doi.org/10.1016/j.fgb.2015.12.004>

Flores, C. Flores, C., Lopez, M., Tabary, N., Neut, C., Chai, F., ... Blanchemaina, N. (2018). Preparation and characterization of novel chitosan and cyclodextrin polymer sponges for wound dressing applications. *Carbohydrate Polymers*, 173, 535-546. <https://doi.org/10.1016/j.carbpol.2017.06.026>

Food and Agriculture Organization of the United Nations. *Global Food Losses and Food Waste: Extent, Causes and Prevention*; FAO: Rome, Italy, 2011.

Fooladi vanda, G., Shabani, L. & Razavizadeh, R. (2019). Chitosan enhances rosmarinic acid production in shoot cultures of *Melissa officinalis* L. through the induction of methyl jasmonate. *Botanical Studies*, 60, 26. <https://doi.org/10.1186/s40529-019-0274-x>

Gao, P., Zhu, Z., & Zhang, P. (2013). Effects of chitosan-glucose complex coating on postharvest quality and shelf life of table grapes. *Carbohydrate Polymers*, 95(1), 371-378. <https://doi.org/10.1016/j.carbpol.2013.03.029>

Geoghegan, I. A., & Gurr, S. J. (2016). Chitosan mediates germling adhesion in *Magnaporthe oryzae* and is required for surface sensing and germling morphogenesis. *PLOS Pathogens*, 12(6), e1005703. <https://doi.org/10.1371/journal.ppat.1005703>

Geoghegan, I., Steinberg, G., & Gurr, S. (2017). The role of the fungal cell wall in the infection of plants. *Trends in Microbiology*, 25, 957-967. <https://doi.org/10.1016/j.tim.2017.05.015>

Gonçalves, A., Gkrillas, A., Dorne J. L., Dall'Asta, C., Palumbo, R., Lima, N., ... Giorni, P. (2019). Pre- and postharvest strategies to minimize mycotoxin contamination in the rice food chain. *Comprehensive Reviews in Food Science and Food Safety*, 18(2), 441-454. <https://doi.org/10.1111/1541-4337.12420>

González-Domínguez, E., Fedele, G., Caffi, T., Delière, L., Sauris, P., Gramaje, D., ... Rossi, V. (2019). A network meta-analysis provides new insight into fungicide scheduling for the control of *Botrytis cinerea* in vineyards. *Pest Management Science*, 75(2), 324–332. <https://doi.org/10.1002/ps.5116>.

Goy, R. C., De Britto, D., & Assis, O. B. G. (2009). A review of the antimicrobial activity of chitosan. *Polímeros*, 19(3), 241–247. <https://doi.org/10.1590/S0104-14282009000300013>

Gramisci, B. R., Lutez, C., Lopes, C. A., & Sangorrína, M. P. (2018). Enhancing the efficacy of yeast biocontrol agents against postharvest pathogens through nutrient profiling

and the use of other additives. *Biological Control*, 121, 151-158.
<https://doi.org/10.1016/j.biocontrol.2018.03.001>.

Guimarães, A., Abrunhosa, L., Pastrana, L. M., & Cerqueira, M. A. (2018). Edible Films and Coatings as Carriers of Living Microorganisms: a new strategy towards biopreservation and healthier foods. *Comprehensive Reviews in Food Science and Food Safety*, 17(3), 594-614. <https://doi.org/10.1111/1541-4337.12345>

Guo, H., Xing, Z., Yu, Q., Zhao, Y., & Zhu, E. (2017). Effectiveness of preharvest application of submicron chitosan dispersions for controlling *Alternaria* rot in postharvest jujube fruit. *Journal of Phytopathology*, 165(7-8), 425-431.
<https://doi.org/10.1111/jph.12576>.

Gupta, S., Banerjee, S. K., Chatterjee, A., Sharma, A. K., Kundu, M., & Basu, J. (2015). The essential protein SepF of mycobacteria interacts with FtsZ and MurG to regulate cell growth and division. *Microbiology*, 16, 1627-1638. <https://doi.org/10.1099/mic.0.000108>

Gutiérrez, T. J. (2017). Chitosan applications for the food industry. In S. Ahmed & S. Ikram (Eds.), *Chitosan: derivatives, composites and applications* (1st ed., pp. 183–232). Wiley-Scrivener. <https://doi.org/10.1002/9781119364849.ch8>

Gutiérrez-Martínez, P., Bautista-Banos, S., Berúmen-Varela, G., Ramos-Guerrero, A., & Hernández-Ibanez, A. M. (2017). *In-vitro* response of *Colletotrichum* to chitosan. Effect on incidence and quality on tropical fruit. enzymatic expression in mango. *Acta Agronómica*, 66(2), 282-289. <http://dx.doi.org/10.15446/acag.v66n2.53770>

Hadwiger, L. A. & Polashock, J. (2013). Fungal mitochondrial DNases: effectors with the potential to activate plant defences in non-host resistance. *Phytopathology*, 103, 81–90.
<http://dx.doi.org/10.1094 / PHYTO-04-12-0085-R>

Hajji, S., Younes, I., Affes, S., Boufi, S., & Nasri, M. (2018). Optimization of the formulation of chitosan edible coatings supplemented with carotenoproteins and their use

for extending strawberries postharvest life. *Food Hydrocolloid*, 83, 375-392.
<https://doi.org/10.1016/j.foodhyd.2018.05.013>.

Harbord, R. M., Egger, M., & Sterne, J. (2006). A modified test for small-study effects in meta-analyses of controlled trials with binary endpoints. *Statistics in Medicine*, 25, 3443-3457. <https://doi.org/10.1002/sim.2380>

Heil, M., & Bostock, R. M. (2002). Induced systemic resistance (ISR) against pathogens in the context of induced plant defense. *Annals of Botany*, 89, 503-512. <https://doi.org/10.1093/aob/mcf076>

Herrera-Romero, I., Ruales, C., & Caviedes, M. (2017). Postharvest evaluation of natural coatings and antifungal agents to control *Botrytis cinerea* in *Rosa* sp. *Phytoparasitica*, 45, 9-20. <https://doi.org/10.1007/s12600-017-0565-2>

Hewajuliage, I. G. N., Sultanbawa, Y., Wijeratnam, R. W., & Wijesundara, R. L. C. (2009). Mode of action of chitosan coating on anthracnose disease control in papaya. *Phytoparasitica*, 37, 437-444. <https://doi.org/10.1007/s12600-009-0052-5>

Hidangmayum, A., Dwivedi, P., Katiyar, & D. Hemantaranjanm A. (2019). Application of chitosan on plant responses with special reference to abiotic stress. *Physiology and Molecular Biology of Plants*, 25, 313–326. <https://doi.org/10.1007/s12298-018-0633-1>

Hosseinnejad, M., & Jafari, S. M. (2016). Evaluation of different factors affecting antimicrobial properties of chitosan. *International Journal of Biological Macromolecules*, 85, 467–475. <https://doi.org/10.1016/j.ijbiomac.2016.01.022>

Hua, C., Li, Y., Wang, X., Kai, K., Su, M., Shia, W., ... Liu, Y. (2019). The effect of low and high molecular weight chitosan on the control of gray mold (*Botrytis cinerea*) on kiwi fruit and host response. *Scientia Horticulturae*. 246 (2), 700-709. <https://doi.org/10.1016/j.scienta.2018.11.038>.

Iriti, M., & Faoro, F. (2009). Chitosan as a MAMP, searching for a PRR. *Plant Signaling & Behavior*, 4, 66–68. <https://doi.org/10.4161/psb.4.1.7408>

Iriti, M., & Varoni, E. M. (2015). Chitosan-induced antiviral activity and innate immunity in plants. *Environmental Science and Pollution Research*, 22, 2935–2944. <https://doi.org/10.1007/s11356-014-3571-7>

Iriti, M., Picchi, V., Rossoni, M., Gomasasca, S., Ludwig, N., Gargano, M., & Faoro, F. (2009). Chitosan antitranspirant activity is due to abscisic acid-dependent stomatal closure. *Environmental and Experimental Botany*, 66, 493–500. <http://dx.doi.org/0.1016/j.envexpbot.2009.01.004>

Jia, X., Meng, Q., Zeng, H., Wang, W., & Yin, H. (2016). Chitosan oligosaccharide induces resistance to Tobacco mosaic virus in Arabidopsis via the salicylic acid-mediated signalling pathway. *Scientific Reports*, 6, 26144. <https://doi.org/10.1038/srep26144>

Jia, X, Zeng, H, Wang, W, Zhang, F., & Yin H. (2018). Chitosan oligosaccharide induces resistance to *Pseudomonas syringae* pv. *tomato* DC3000 in *Arabidopsis thaliana* by activating both salicylic acid- and jasmonic acid-mediated pathways. *Molecular Plant-Microbe Interaction*, 31(12), 1271-1279. <https://doi.org/10.1094/MPMI-03-18-0071-R>

Jitareerat, P., Paumchai, S., Kanlayanarat, S., & Sangchote, S. (2007). Effect of chitosan on ripen Rehmaing, enzymatic activity and disease development in mango (*Mangifera indica*) fruit. *New Zealand Journal of Crop and Horticultural Science*, 35 (2), 211-218. <https://doi.org/10.1080/01140670709510187>.

Jones, J. D. G., & Dang, J. L. (2006). The plant immune system. *Nature*, 444, 323-329. <https://doi.org/10.1038/nature05286>

Jongsri, P., Rojsitthisak, P., Wangsomboondee, P., & Seraypheapa, K. (2017). Influence of chitosan coating combined with spermidine on anthracnose disease and qualities of ‘Nam

Dok Mai' mango after harvest. *Scientia Horticulturae*, 224, 180-187.
<https://doi.org/10.1016/j.scienta.2017.06.011>

Kabanov V. L., & Novinyuk L. V. (2020). Chitosan application in food technology: a review of recent advances. *Food Systems*, 3, 10-15. <https://doi.org/10.21323/2618-9771-2020-3-1-10-15>

Kaczmarek, M. B., Struszczyk-Swita, K., Li, X., Szczesna-Antczak, M., & Daroch, M. (2019). Enzymatic modifications of chitin, chitosan, and chitooligosaccharides. *Frontiers in Bioengineering and Biotechnology*, 7, 243. <https://doi.org/10.3389/fbioe.2019.00243>

Kadic, A. J., Vucic, K., Dosenovic, S., Sapunar, D., & Puljak, L. (2016). Extracting data from Figures with software was faster, with higher interrater reliability than manual extraction. *Journal of Clinical Epidemiology*, 74, 119-123. <https://doi.org/10.1016/j.jclinepi.2016.01.002>

Kanetis, L., Exarchou, V., Charalambous, Z., & Goulas, V. (2017). Edible coating composed of chitosan and *Salvia fruticosa* Mill. extract for the control of grey mould of table grapes. *Journal of the Science of Food and Agriculture*, 97(2), 452-460. <https://doi.org/10.1002/jsfa.7745>

Katiyar, D., Hemantaranjan, A., Bharti, S., & Nishant Bhanu, A. (2014). A future perspective in crop protection: chitosan and its oligosaccharides. *Advances in Plants & Agriculture Research*, 1, 23-30. <http://dx.doi.org/10.15406/apar.2014.01.00006>

Kharchoufi, S., Parafati, L., Licciardello, F., Muratore, G., Hamdi, M., Cirvilleri, G., & Restuccia, C. (2018). Edible coatings incorporating pomegranate peel extract and biocontrol yeast to reduce *Penicillium digitatum* postharvest decay of oranges. *Food Microbiology*, 74, 107-112. <https://doi.org/10.1016/j.fm.2018.03.011>

Kofuji, K., Huang, Y., Tsubaki, K., Kokido, F., Nishikawa, K., Takashi I., & Murata, Y. (2010). Preparation and evaluation of a novel wound dressing sheet comprised of β -glucan-

chitosan complex. *Reactive and Functional Polymers*, 70, 784-789.
<https://doi.org/10.1016/j.reactfunctpolym.2010.07.014>

Kong, M., Chen, X. G., Xing, K., & Park, H. J. (2010). Antimicrobial properties of chitosan and mode of action: a state-of-the-art review. *International Journal of Food Microbiology*, 144(1), 51-63. <https://doi.org/10.1016/j.ijfoodmicro.2010.09.012>

Landi, L., De Miccolis Angelini, R. M., Pollastro, S., Feliziani, E., Faretra, F., & Romanazzi, G. (2017). Global transcriptome analysis and identification of differentially expressed genes in strawberry after preharvest application of benzothiadiazole and chitosan. *Frontiers of Plant Science*, 8, 235. <https://doi.org/10.3389/fpls.2017.00235>.

Landi, L., Feliziani, E., & Romanazzi, G. (2014). Expression of defense genes in strawberry fruit treated with different resistance inducers. *Journal of Agricultural and Food Chemistry*, 62(14), 3047-3056. <http://doi.org/10.1021/jf404423x>

Lee, S., Choi, H., Suh, S., Doo, I-S., Oh, K-Y., Choi, E. J. ... Lee., Y. (1999). Oligogalacturonic acid and chitosan reduce stomatal aperture by inducing the evolution of reactive oxygen species from guard cells of tomato and *Commelina communis*. *Plant Physiology*, 121, 147-152. <http://dx.doi.org/10.1104/pp.121.1.147>

Lenardon, M. D., Munro, C. A., & Gow, N. A. (2010). Chitin synthesis and fungal pathogenesis. *Current Opinion in Microbiology*, 13(4), 416-423. <https://doi.org/10.1016/j.mib.2010.05.002>

Leung, M. Y. K., Liu, C., Koon, J. C. M., & Fung, K. P (2006). Polysaccharide biological response modifiers. *Immunology Letters*, 105(2), 101-114. <https://doi.org/10.1016/j.imlet.2006.01.009>

Levdansky, E., Kashi, O., Sharon, H., Shadkchan, Y., & Oshero, N. (2010). The *Aspergillus fumigatus* cspA gene encoding a repeat-rich cell wall protein is important for

normal conidial cell wall architecture and interaction with host cells. *Eukaryotic Cell*, 9, 1403–1415. <https://doi.org/10.1128/EC.00126-10>.

Li, K., Xing, R., Liu, S., & Li P. (2016). Advances in preparation, analysis and biological activities of single chitooligosaccharides. *Carbohydrate Polymers*, 139, 178–190. <https://doi.org/10.1016/j.carbpol.2015.12.016>

Liang, J., Yan, H., Puligundla, P., Gao, X., Zhou, Y., & Wan, X. (2017). Applications of chitosan nanoparticles to enhance absorption and bioavailability of tea polyphenols: a review. *Food Hydrocolloids*, 69, 286–292. <https://doi.org/10.1016/j.foodhyd.2017.01.041>

Liu, H., Zhao, H., Lyu, L., Huang, Z., Fan, S., Wu, W., & Li, W. (2019). Synergistic effect of natural antifungal agents for postharvest diseases of blackberry fruits. *Journal of the Science of Food and Agriculture*, 99(7), 3343–3349. <https://doi.org/10.1002/jsfa.9551>

Liu, Y., Sun, Z, Xiu, L., Huang, J., & Zhou, F. (2018). Selective antifungal activity of chitosan and sulfonated chitosan against postharvest fungus isolated from blueberry. *Journal of Food Biochemistry*, 42(6), e12658. <https://doi.org/10.1111/jfbc.12658>

Liu, Y., Yamdeu, J. H. G., Gong, Y.Y., & Orfila, C. (2020). A review of postharvest approaches to reduce fungal and mycotoxin contamination of foods. *Comprehensive Reviews in Food Science and Food Safety*, 19(4), 1521–1560. <https://doi.org/10.1111/1541-4337.12562>

López-Mora, L. I., Gutiérrez-Martínez, P., Bautista-Baños, S., Jiménez-García, L. F., & Zavaleta-Mancera, H. A. (2013). Evaluation of antifungal activity of chitosan in *Alternaria alternata* and in the quality of ‘Tommy Atkins’ mango during storage. *Revista Chapingo. Serie Horticultura*, 19(3), 315–331. <http://dx.doi.org/10.5154/r.rchsh.2012.07.038>

Lopez-Moya, F., Escudero, N., Zavala-Gonzalez, E. A., Esteve-Bruna, D., Blázquez, M. A., Alabadí, D., & Lopez-Llorca, L. V. (2017). Induction of auxin biosynthesis and WOX5

repression mediate changes in root development in *Arabidopsis* exposed to chitosan. *Scientific Reports*, 7,16813. [https://doi.org/ 10.1038/s41598-017-16874-5](https://doi.org/10.1038/s41598-017-16874-5).

Lopez-Moya, F., Suarez-Fernandez, M., & Lopez-Lorca, L.V. (2019). Molecular mechanisms of chitosan interactions with fungi and plants. *International Journal of Molecular Sciences*, 20(2), 332. <https://doi.org/10.3390/ijms20020332>.

Lu, L., Liu, Y., Yang, J., Azat, R., Yu, T., & Zheng, X. (2014). Quaternary chitosan oligomers enhance resistance and biocontrol efficacy of *Rhodosporidium paludigenum* to green mold in satsuma orange. *Carbohydrate Polymers*, 113, 174-181. <https://doi.org/10.1016/j.carbpol.2014.06.077>

Ma, Z., Yang, L., Yan, H., Kennedy, J. F., & Meng, X. (2013). Chitosan and oligochitosan enhance the resistance of peach fruit to brown rot. *Carbohydrate Polymers*, 94(1), 272-277. <https://doi.org/10.1016/j.carbpol.2013.01.012>

Madanipour, S., Alimohammadi, M., Rezaie, S., Nabizadeh, R., Jahed Khaniki, G., Hadi, M., ... Yousefzadeh, S. (2019). Influence of postharvest application of chitosan combined with ethanolic extract of liquorice on shelf life of apple fruit. *Journal of Environmental Health Science and Engineering*, 17(1), 331-336. <https://doi.org/10.1007/s4020>

Maddi, A., & Free, S. J. (2010). 1,6-Mannosylation of N-linked oligosaccharide present on cell wall proteins is required for their incorporation into the cell wall in the filamentous fungus *Neurospora crassa*. *Eukaryotic Cell*, 9(11), 1766-1775. <http://doi.org/10.1128/EC.00134-10>

Maestri, E. Pavlicevic, M., Montorsi, M., & Marmiroli, N. (2019). Meta-Analysis for correlating structure of bioactive peptides in foods of animal origin with regard to effect and stability. *Comprehensive Reviews in Food Science and Food Safety*, 18(1), 3-3 <https://doi.org/10.1111/1541-4337.12402>

Malerba, M., & Cerana, R. (2016). Chitosan effects on plant systems. *International Journal of Molecular Sciences*, 17(7), 996. <https://doi.org/10.3390/ijms17070996>

Mandal, S., Kar, I., Mukherjee, A. K., & Acharya, P. (2013). Elicitor-induced defense responses in *Solanum lycopersicum* against *Ralstonia solanacearum*. *Science World Journal*, 2013, 1-9. <https://doi.org/10.1155/2013/561056>

Maqbool, M., Ali, A., Ramachandran, S., Smith, D., & Alderson, P. (2010). Control of postharvest anthracnose of banana using a new edible composite coating. *Crop Protection*, 29, 1136-1141. <https://doi.org/10.1016/j.cropro.2010.06.005>

Mebdoua S. (2018) Pesticide Residues in Fruits and Vegetables. In: Mérillon JM., Ramawat K. (eds) *Bioactive Molecules in Food*. Reference Series in Phytochemistry. Springer, Cham.

Meng, X., Yang, L., Kennedy, J. F., & Tian, S. (2010). Effects of chitosan and oligo chitosan on growth of two fungal pathogens and physiological properties in pear fruit. *Carbohydrate Polymers*, 81(1), 70-75. <https://doi.org/10.1016/j.carbpol.2010.01.057>

Mohamed, C., Clementine, K. A., Didier, M., Gérard, L., & Noëlle, D. C. M. (2013). Antimicrobial and physical properties of edible chitosan films enhanced by lactoperoxidase system. *Food hydrocolloids*, 30(2), 576-580. <https://doi.org/10.1016/j.foodhyd.2012.07.018>

Moher, D., Liberati, A., Tetzlaff, J., Altman, D. G., & PRISMA Group. (2009). Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Medicine* 6(7), e1000097. <https://doi.org/10.1371/journal.pmed.1000097>

Morin-Crini, N., Lichtfouse, E., Torri, G., & Crini, G. (2019). Applications of chitosan in food, pharmaceuticals, medicine, cosmetics, agriculture, textiles, pulp and paper, biotechnology, and environmental chemistry. *Environmental Chemistry Letters*, 17, 1667-1692. <https://doi.org/10.1007/s10311-019-00904-x>

Munhuweyi, K., Lennox, C.L., Meitz-Hopkins, J. C., Caleb, O. J., Sigged, G. O., & Opara, U. L. (2017). Investigating the effects of crab shell chitosan on fungal mycelial growth and postharvest quality attributes of pomegranate whole fruit and arils. *Scientia Horticulturae*, 220, 78-89. <https://doi.org/10.1016/j.scienta.2017.03.038>

Munoz, Z., Moret, A., & Garces, S. (2009). Assessment of chitosan for inhibition of *Colletotrichum* spp. on tomatoes and grapes. *Crop Protection*, 28(1), 36-40. <http://doi.org/10.1016/j.cropro.2008.08.015>

Muzzarelli, R. A. A. (1983). Chitin and its derivatives: new trends of applied research. *Carbohydrate Polymers*, 3(1), 53-75. [https://doi.org/10.1016/0144-8617\(83\)90012-7](https://doi.org/10.1016/0144-8617(83)90012-7)

Nagy, A., Harrison, A., Sabbani, S., Munson, R. S. Jr., Dutta, P. K., & Waldman, W. J. (2011). Silver nanoparticles embedded in zeolite membranes: release of silver ions and mechanism of antibacterial action. *International Journal of Nanomedicine*, 6, 1833-1852. <https://doi.org/10.2147/IJN.S24019>

Nandeeshkumar, P, Sudisha, J, Ramachandra, K. K., Prakash, H. S., Niranjana, S. R., & Shekar, S. H. (2008). Chitosan induced resistance to downy mildew in sunflower caused by *Plasmopara halstedii*. *Physiological and Molecular Plant Pathology*, 72(4-6), 188-194. <http://www.sciencedirect.com/science/journal/08855765>

Nelson, M. E., Gent, D. H., & Grove., G. (2015). Meta-analysis reveals a critical period for management of powdery mildew on hopcones. *Plant Disease*, 99(5), 632-640. <https://doi.org/10.1094/PDIS-04-14-0396-RE>

Nisia C, Noreña, C.P.Z., & Brandelli, A. (2012). Antimicrobial activity of chitosan films containing nisin, peptide P34, and natamycin. *Journal of Food*, 10, 21-26. <https://doi.org/10.1080/19476337.2010.537371>

Obianom, C., Romanazzi, G., & Sivakumar, D. (2019). Effects of chitosan treatment on avocado postharvest diseases and expression of phenylalanine ammonia-lyase, chitinase

and lipoxygenase genes. *Postharvest Biology and Technology*, 147, 214-221. <https://doi.org/10.1016/j.postharvbio.2018.10.004>

Oerke, E. C., & Dehne, H. W. (2004). Safeguarding production-losses in major crops and the role of crop protection. *Crop Protection*, 23, 275-285. <https://doi.org/10.1016/j.cropro.2003.10.001>

Oliveira-Garcia, E., & Deising, H. B. (2013). Infection structure- specific expression of β -1,3-glucan synthase is essential for pathogenicity of *Colletotrichum graminicola* and evasion of β -glucan-triggered immunity in maize. *Plant Cell*, 25(6), 2356-2378. <http://doi.org/10.1105/tpc.112.103499>

Palma-Guerrero, J., Jansson, H. B., Salinas, J., & Lopez-Llorca, L. V. (2008). Effect of chitosan on hyphal growth and spore germination of plant pathogenic and biocontrol fungi. *Journal of Applied Microbiology*, 104, 541-553. <https://doi.org/10.1111/j.1365-2672.2007.03567.x>

Palma-Guerrero, J., Huang, I. C., Jansson, H. B., Salinas, J., Lopez-Llorca, L. V., & Read N. D. (2009). Chitosan permeabilizes the plasma membrane and kills cells of *Neurospora crassa* in an energy dependent manner. *Fungal Genetics Biology*, 46(8), 585-594. <http://doi.org/10.1016/j.fgb.2009.02.010>. Epub 2009 Apr 21.

Palma-Guerrero, J., Lopez-Jimenez, J. A., Pérez-Berná, A.J., Huang, I. C., Jansson, H. B., Salinas, J., ... Lopez-Llorca, L. V. (2010). Membrane fluidity determines sensitivity of filamentous fungi to chitosan. *Molecular Microbiology*, 75(4), 1021–1032. <https://doi.org/10.1111/j.1365-2958.2009.07039.x>.

Palou, L., Smilanick J. L., & Droby, S. (2008). Alternatives to conventional fungicides for the control of citrus postharvest green and blue moulds. *Stewart Postharvest Review*, 4(2), 1-16. <https://doi.org/10.2212/spr.2008.2.2>

Palou, L., & Smilanick J. L. (2020). Postharvest Pathology of Fresh Horticultural Produce. In. Palou, L., & Smilanick, J. (Eds.) (1st ed., pp. 1–842). Boca Raton: CRC Press, <https://doi.org/10.1201/9781315209180>

Parfitt, J., Barthel, M., & Macnaughton, S. (2010). Food waste within food supply chains: Quantification and potential for change to 2050. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 365, 3065-3081. <https://doi.org/10.1098/rstb.2010.0126>

Pasquariello, M. S., Di Patre, D., Mastrobuoni, F., Zampella, L., Scortichini, M., & Petriccione, M. (2015). Influence of postharvest chitosan treatment on enzymatic browning and antioxidant enzyme activity in sweet cherry fruit. *Postharvest Biology and Technology*, 109, 45-56. <https://doi.org/10.1016/j.postharvbio.2015.06.007>

Patel, P. K. & Free, S. J. (2019). The genetics and biochemistry of cell wall structure and synthesis in *Neurospora crassa*, a model filamentous fungus. *Frontiers in Microbiology*, 10, 2294. <https://doi.org/10.3389/fmicb.2019.02294>

Pétriaccq, P., López, A. & Luna, S. (2018). Fruit decay to diseases: can induced resistance and priming help? *Plants*, 7(4), 77. <https://doi.org/10.3390/plants7040077>

Petriccione, M., Mastrobuoni, F., Zampella, L., Nobis, E., Capriolo, G., & Scortichini, M. (2017). Effect of chitosan treatment on strawberry allergen-related gene expression during ripening stages. *Journal of Food Science and Technology*, 54(5), 1340–1345. <https://doi.org/10.1007/s13197-017-2554-3>

Pieterse, C. M. J., Van Wees, S. C. M., van Pelt, J. A., Knoester, M., Laan, R., Gerrits, H., ... van Loon, L. C. (1998). A novel signaling pathway controlling induced systemic resistance in Arabidopsis. *Plant Cell*, 10(9), 1571–1580. <https://doi.org/10.1105/tpc.10.9.1571>

Prusky, D. (2011). Reduction of the incidence of postharvest quality losses, and future prospects. *Journal of Food Security*, 3(4), 463-474. <https://doi.org/10.1007/s12571-011-0147-y>

Pusztahelyi T. (2018). Chitin and chitin-related compounds in plant-fungal interactions. *Mycology*, 9(3), 189–201. <https://doi.org/10.1080/21501203.2018.1473299>

Rabea, E. I., Badawy, M. E., Stevens, C. V., Smagghe, G., & Steurbaut, W. (2003). Chitosan as antimicrobial agent: applications and mode of action. *Biomacromolecules*, 4, 1457-1465. <https://doi.org/10.1021/bm034130m>

Rahman, M. A., Mahmud, T. M. M., Kadir, J., Abdul Rahman, R., & Begum, M. M. (2008). Antimicrobial activities of chitosan and calcium chloride on *in-vitro* growth of *Colletotrichum gloeosporioides* from Papaya. *Pertanika. Journal of Tropical Agricultural Science*, 31(2), 223-232.

Ramos-García, M., Bosquez-Molina, E., Hernández-Romano, J., Zavala-Padilla, G., Terrés-Rojas, E., Alia-Tejacal, ... Bautista-Baños, S. (2012). Use of chitosan-based edible coatings in combination with other natural compounds, to control *Rhizopus stolonifer* and *Escherichia coli* DH5a in fresh tomatoes. *Crop Protection*, 38, 1-6. <https://doi.org/10.1016/j.cropro.2012.02.01>

Ramos-Guerrero, A., González-Estrada, R. R, Hanako-Rosas, G., & Bautista-Banões, S. (2018). Use of inductors in the control of *Colletotrichum gloeosporioides* and *Rhizopus stolonifer* isolated from soursop fruits: *in vitro* tests. *The Food Science and Biotechnology*, 27(3), 755-763. <https://doi.org/10.1007/s10068-018-0305-5>

Ramos-Guerrero, A., González-Estrada, A. A., Romanazzi, G., L. Landi, L. & Gutiérrez-Martínez, P. (2020). Effect of chitosan in the control of postharvest anthracnose of soursop (*Annona muricata*) fruit. *Revista Mexicana de Ingenieria Quimica*, 19(1), 99-108. <https://doi.org/10.24275/rmiq/Bio527>

Rebelo, R., Vila, N. & Fangueiro, R. (2016). Poly lactic acid fibre based biodegradable stents and their functionalization techniques, in: R. Fangueiro, S. Rana (Eds.), *Natural*

fibres: advances in science and technology towards industrial applications. to Mark.,
 RILEM Book series, pp. 331-342.

Ribes, S., Fuentes, A., Talens, P., & Barat, J. (2018). Prevention of fungal spoilage in food
 products using natural compounds: A review. *Critical Reviews in Food Science and
 Nutrition*, 58(12), 2002-2016. <https://doi.org/10.1080/10408398.2017.1295017>

Romanazzi, G., Feliziani, E., Santini, M., & Landi, L. (2013). Effectiveness of postharvest
 treatment with chitosan and other resistance inducers in the control of storage decay of
 strawberry. *Postharvest Biology and Technology*, 75, 24-27.
<https://doi.org/10.1016/j.postharvbio.2012.07.007>

Romanazzi, G., Gabler, F. M., Margosan, D., Mackey, B. E., & Smilanick, J. L. (2009).
 Effect of chitosan dissolved in different acids on its ability to control postharvest gray mold
 of table grape. *Phytopathology*. 99(9), 1028-1036. <https://doi.org/10.1094/PHYTO-99-9-1028>.

Romanazzi, G., Sanzani, S. M., Bi, Y., Tian, S., Gutierrez-Martinez, P., & Alkan, N. (2016).
 Induced resistance to control postharvest decay of fruit and vegetables. *Postharvest Biology
 and Technology* 122, 82–94. <https://doi.org/10.1016/j.postharvbio.2016.08.003>

Romanazzi, G., Smilanick, J. L., Feliziani, E., & Droby, S. (2016). Integrated management
 of postharvest gray mold on fruit crops. *Postharvest Biology and Technology*, 113, 69-76.
<https://doi.org/10.1016/j.postharvbio.2015.11.003>

Romanazzi, G., Feliziani, E., Baños, S. B., & Sivakumar, D. (2017). Shelf life extension of
 fresh fruit and vegetables by chitosan treatment. *Critical Reviews in Food Science and
 Nutrition*, 57, 579-601. <https://doi.org/10.1080/10408398.2014.900474>

Romanazzi, G., Feliziani, E., & Sivakumar, D. (2018). Chitosan, a biopolymer with triple
 action on postharvest decay of fruit and vegetables: eliciting, antimicrobial and film-

forming properties. *Frontiers in Microbiology*, 9, 2745. <https://doi.org/10.3389/fmicb.2018.02745>

Rosenberg, M. S., Garrett, K. A., Su, Z. & Bowden, R. L. (2004). Meta-analysis in plant pathology: synthesizing research results. *Phytopathology*, 94(9), 1013-1017. <https://doi.org/10.1094/PHYTO.2004.94.9.1013>

Sawicka B. (2019). Post-harvest Losses of Agricultural Produce. In: Leal Filho W., Azul A., Brandli L., Özuyar P., Wall T. (Eds.) Zero Hunger. Encyclopedia of the UN Sustainable Development Goals. Springer, Cham. https://doi.org/10.1007/978-3-319-69626-3_40-1

Schwingshackl, L., Hoffmann, G., Iqbal, K., Schwedhelm, K. & Boeing, H. (2018). Food groups and intermediate disease markers: a systematic review and network meta-analysis of randomized trials. *The American Journal of Clinical Nutrition*, 108(3), 576-586. <https://doi.org/10.1093/ajcn/nqy151>

Shao, X. F. Tu, K., Tu, S., & Tu, J. (2012). A combination of heat treatment and chitosan coating delays ripening and reduces decay in (Gala) apple fruit. *Journal of Food Quality*, 35, 83-92 <https://doi.org/10.1111/j.1745-4557.2011.00429.x>

Shao, X., Cao, B., Xu, F., Xie, S., Yu, D., & Wang, H. (2015). effect of postharvest application of chitosan combined with clove oil against citrus green mold. *Postharvest Biology and Technology*. 99, 37-43. <https://doi.org/10.1016/j.postharvbio.2014.07.014>

Shen, Y., & Yang, H. (2017). Effect of preharvest chitosan-g-salicylic acid treatment on postharvest table grape quality, shelf life, and resistance to *Botrytis cinerea* induced spoilage. *Scientia Horticulturae*. 224, 367-373. <https://doi.org/10.1016/j.scienta.2017.06.046>

981 Shi, Z., Wang, F., Lu, Y., & Deng, J. (2018). Combination of chitosan and salicylic acid to
 982 control postharvest green mold caused by *Penicillium digitatum* in grapefruit fruit. *Scientia*.
 983 *Horticulturae*, 233, 54-60. <http://dx.doi.org/10.1016/j.scienta.2018.01.039>
 984 Silva W. B, Silva G. M. C., Santana D. B., Salvador, A. R., Medeiros, D. B, Belghith, I., ...
 985 Misobutsi, G. P. (2018). Chitosan delays ripening and ROS production in guava (*Psidium*
 986 *guajava* L.) fruit. *Food Chemistry*, 242, 232-238.
 987 <https://doi.org/10.1016/j.foodchem.2017.09.052>
 988 Singh, R. K., Soares, B., Goufo, P., Castro, I., Cosme, F., Pinto-Sintra, A. L., ... Falco, V.
 989 (2019). Chitosan upregulates the genes of the ROS pathway and enhances the antioxidant
 990 potential of grape (*Vitis vinifera* L. 'Touriga Franca' and 'Tinto Cão') tissues. *Antioxidants*,
 991 8, 11. <https://doi.org/10.3390/antiox8110525>
 992 Song, H., Yuan, W., Jin, P., Wang, W., Wang, X., Yanga, L., & Zhang, Y. (2016). Effects
 993 of chitosan/ nano-silica on postharvest quality and antioxidant capacity of loquat fruit
 994 during cold storage. *Postharvest Biology and Technology*, 119, 41-
 995 48. <https://doi.org/10.1016/j.postharvbio.2016.04.015>
 996 Souza, C. D., Yuk, H. G., Khoo, G. H., & Zhou, W. (2015). Application of light-emitting
 997 diodes in food production, postharvest preservation, and microbiological food safety.
 998 *Comprehensive Reviews in Food Science and Food Safety*, 14(6), 719-740.
 999 <https://doi.org/10.1111/1541-4337.12155>
 1000 Talibi, I., Boubaker, H., Boudyach, E. H., & Ait Ben Aoumar, A. (2014). Alternative
 1001 methods for the control of postharvest citrus diseases. *Journal of Applied Microbiology*,
 1002 117(1), 1-17. <https://doi.org/10.1111/jam.12495>
 1003 Timmermann, T., González, B. & Ruz, G.A. (2020). Reconstruction of a gene regulatory
 1004 network of the induced systemic resistance defense response in Arabidopsis using boolean
 1005 networks. *BMC Bioinformatics*, 21, 142. <https://doi.org/10.1186/s12859-020-3472-3>

Tufanaru, C., Munn, Z., Stephenson, M., & Aromataris, E. (2015). Fixed or random effects meta-analysis? Common methodological issues in systematic reviews of effectiveness. *International Journal of Evidence-Based Healthcare*, 13, 196-207. <https://doi.org/10.1097/XEB.0000000000000065>

Tungland, B. C., & Meyer, D. (2002). Non-digestible oligo- and polysaccharides (dietary fiber): their physiology and role in human health and food. *Comprehensive Reviews in Food Science and Food Safety*, 1(3), 90-109. <https://doi.org/10.1111/j.1541-4337.2002.tb00009.x>

Valencia-Chamorro, S. A., Palou, L., & Del Río, M. A. (2011). Antimicrobial edible films and coatings for fresh and minimally processed fruits and vegetables: a review. *Critical Reviews in Food Science and Nutrition*, 51(9), 872-900. <https://doi.org/10.1080/10408398.2010.485705>

Varela, G. B., Coronado Partida, L. D., Ochoa Jiménez, V. A., López, C. M. A., & Martínez, G. P. (2015). Effect of chitosan on the induction of disease resistance against *Colletotrichum* spp. in mango (*Mangifera indica* L.) cv. Tommy Atkins. *Investigación Ciencia*, 66, 16-21.

Viswanatha, G. L., Shylaj, H., & Moolemath, Y. (2017). The beneficial role of naringin, a citrus bioflavonoid, against oxidative stress-induced neurobehavioral disorders and cognitive dysfunction in rodents: a systematic review and meta-analysis. *Biomedicine and Pharmacotherapy*, 94, 909-929. <https://doi.org/10.1016/j.biopha.2017.07.072>

Waewthongrak, W., Pisuchpen, S., & Leelasuphakul, W. (2015). Effect of *Bacillus subtilis* and chitosan applications on green mold (*Penicillium digitatum* Sacc.) decay in citrus fruit. *Postharvest Biology and Technology*, 99, 44-49. <https://doi.org/10.1016/j.postharvbio.2014.07.016>

Walters, D. R., Ratsep, J., & Havis, N. D. (2013). Controlling crop diseases using induced resistance: challenges for the future. *The Journal of Experimental Botany*, 64(5), 1263-1280. <https://doi.org/10.1093/jxb/ert026>

Wang, L., Wu, H., Qin, G., & Meng, X. (2014). Chitosan disrupts *Penicillium expansum* and controls postharvest blue mold of jujube Fruit. *Food Control*, 41(1), 56-62. <https://doi.org/10.1016/j.foodcont.2013.12.028>

Wang, S. Y., & Gao, H. (2013). Effect of chitosan-based edible coating on antioxidants, antioxidant enzyme system, and postharvest fruit quality of strawberries (*Fragaria × ananassa* Duch). *LWT - Food Science and Technology*, 52, 71-79. <https://doi.org/10.1016/j.lwt.2012.05.003>

Wang, Y. G, Li, B. & Zhang, X. D. (2017). Low molecular weight chitosan is an effective antifungal agent against *Botryosphaeria* sp. and preservative agent for pear (*Pyrus*) fruits. *International Journal of Biological Macromolecules*, 95, 1135-1143. <https://doi.org/10.1016/j.ijbiomac.2016.10.105>

Weir, C. J., Butcher, I., Assi, V., Lewis, S. C., Murray, G. D., Langhorne, P., & Brady, M. C. (2018). Dealing with missing standard deviation and mean values in meta-analysis of continuous outcomes: a systematic review. *BMC Medical Research Methodology*, 18, 25. <https://doi.org/10.1186/s12874-018-0483-0>

Xing, K., Zhu, X., Peng, X., & Qin S. (2015). Chitosan antimicrobial and eliciting properties for pestcontrol in agriculture: a review. *Agronomy for Sustainable Development*, 35, 569-588. <https://doi.org/10.1007/s13593-014-0252>

Xing, Y., Xu, Q., Che, Z., Li, X., & Li, W. (2011). Effects of chitosan oil coating on blue mold disease and quality attributes of jujube fruits. *Food & Function*, 2(8), 466-474. <http://dx.doi.org/10.1039/c1fo10073d>

Xing, Y., Lin, H., Cao, D., Xu, Q., Han, W., Wang, R., ... Li, X. (2015). Effect of Chitosan Coating with Cinnamon Oil on the Quality and Physiological Attributes of China Jujube Fruits. *BioMed Research International*, 2015,835151 <https://doi.org/10.1155/2015/835151>

Xing, Y., Xu, Q., Yang, S. X., Chen, C., Tang, Y., Sun, S., ... Li, X. (2016). Preservation mechanism of chitosan-based coating with cinnamon oil for fruits storage based on sensor data. *Sensors*, 16(7), 1111. <https://doi.org/10.3390/s16071111>

Xoca-Orozco, L. A., Aguilera-Aguirre, S, López-García, U.M., Gutiérrez-Martínez, P., & Chacón-López, A (2018). Effect of chitosan on the *in-vitro* control of *Colletotrichum* spp. and its influence on postharvest quality in Hass avocado fruits. *Revista Bio Ciencias*, 5, 355. <http://dx.doi.org/10.15741/revbio.05.01.13>

Xoca-Orozco L. A., Aguilera-Aguirre, S., Vega-Arreguín J., Acevedo-Hernández, G., Tovar Pérez, E., Stoll, A., ...Chacón-López, A. (2019). Activation of the phenylpropanoid biosynthesis pathway reveals a novel action mechanism of the elicitor effect of chitosan on avocado fruit epicarp. *Food Research International*, 121, 586-592. <http://dx.doi.org/10.1016/j.foodres.2018.12.023>

Xu, W.T., Huang, K.L., Guo, F., Yang, J., Liang, Z., & Luo, Y. (2007). Postharvest grapefruit seed extract and chitosan treatments of table grapes to control *Botrytis cinerea*. *Postharvest Biology and Technology*, 46, 86-94. <https://doi.org/10.1016/j.postharvbio.2007.03.019>

Yadav, V., Wang, Z., Wei, Amo, A., Ahmed, B., Yang, X., & Zhang, X. (2020). Phenylpropanoid pathway engineering: an emerging approach towards plant defense. *Pathogens*, 9(4), 312. <https://doi.org/10.3390/pathogens9040312>

Yan, J., Li, J., Zhao, H., Chen, N., Cao, J., & Jiang, W. (2011). Effects of oligo chitosan on postharvest *Alternaria* rot, storage quality, and defense responses in Chinese jujube

(*Zizyphus jujuba* Mill. cv. Dongzao) fruit. *Journal of Food Protection*, 74(5), 783-788. <https://doi.org/10.4315/0362-028X.JFP-10-480>

Yang, Z., Scott, C. A., Mao, C., Tang, J. & Farmer, A. J. (2014). Resistance exercise versus aerobic exercise for type 2 diabetes: a systematic review and meta-analysis. *Sports Medicine*, 44(4), 487-499. <https://doi.org/10.1007/s40279-013-0128-8>

Yilmaz Atay, H. (2019) Antibacterial activity of chitosan-based systems. In: Jana S., Jana S. (eds) *Functional Chitosan*. Springer, Singapore. https://doi.org/10.1007/978-981-15-0263-7_15

Yin, H., Li, S., Zhao, X., Du, Y., & Ma, X. (2006) cDNA microarray analysis of gene expression in *Brassica napus* treated with oligo chitosan elicitor. *Plant Physiology and Biochemistry*, 44(11-12), 910-916. <http://dx.doi.org/10.1016/j.plaphy.2006.10.002>

Zahid, N., Ali, A., Manickam, S., Siddiqui, Y., & Maqbool, M. (2012). Potential of chitosan-loaded nanoemulsions to control different *Colletotrichum* spp. and maintain quality of tropical fruits during cold storage. *Journal of Applied Microbiology*, 113, 925-939. <https://doi.org/10.1111/j.1365-2672.2012.05398.x>

Zahid, N., Maqbool, M., Ali, A., Siddiqui, Y. & Bhatti, Q. A. (2019). Inhibition in production of cellulolytic and pectinolytic enzymes of *Colletotrichum gloeosporioides* isolated from dragon fruit plants in response to submicron chitosan dispersions. *Scientia Horticulturae*, 243, 314-319. <https://doi.org/10.1016/j.scienta.2018.08.011>

Zargar, V., Asghari, M., & Dashti, A. (2015). A review on chitin and chitosan polymers: structure, chemistry, solubility, derivatives, and applications. *ChemBioEng Reviews*, 2(3), 204-226. <https://doi.org/10.1002/cben.201400025>

Zhao, D., Yu, S., Sun, B., Gao, S., Guo, S., & Zhao, K. (2018). Biomedical applications of chitosan and its derivative nanoparticles. *Polymers (Basel)*, 10, 462. <https://doi.org/10.3390/polym10040462>

Zheng, W., Li, L., Pan, S., Liu, M., Zhang, W., Liu, H., & Zhu, C. (2017). Controls postharvest decay and elicits defense response in kiwi fruit. *Food and Bioprocess Technology*, 11, 1937-1945. <https://doi.org/10.1007/s11947-017-1957-5>

Zuccolo, M., Kunova, A., Musso, L., Forlani, F., Pinto, A., Vistoli G., ... Dellavalle S. (2019). Dual-active antifungal agents containing strobilurin and SDHI-based pharmacophores. *Scientific Reports* 9, 11377. <https://doi.org/10.1038/s41598-019-47752-x>

Zuppini, A., Baldan, B., Million, R., Favaron, F., Navazio, & L., Mariani, P. (2003). Chitosan induces Ca^{2+} mediated programmed cell death in soybean cells. *New Phytologist* 161, 557–568. <https://doi.org/10.1046/j.1469-8137.2003.00969.x>

Acknowledgements

The authors would like to thank COST Action FA1405 "Using three-way interactions between plants, microbes and arthropods to enhance crop protection and production" for organising a meta-analysis workshop in July 2016, from led to the idea for this study.

Author Contributions

R.R. performed the literature research, analysed the data, and contributed to write the manuscript; L.L designed the analysis, analysed the data, and wrote the manuscript; G.R. designed the analysis, supervised and complemented the writing, and coordinated the study.


Conflicts of Interest: The authors declare that they have no competing interests.

TABLE 1. Main characteristics of datasets that have included 1% chitosan effects on postharvest fungal pathogens.

First author	Year	Fungal pathogen	Chitosan effects measures			Defence enzyme
			Disease incidence (fruit)	In-vitro mycelium growth	Plant defence mechanism (fruit)	
Xu	2007	<i>B. cinerea</i>		Yes		-
Jitareerat	2007	<i>Colletotrichum</i> spp.		Yes		-
Rahman	2008	<i>Colletotrichum</i> spp.		Yes		-
Hewajulige	2009	-			Papaya	Chitinase, β -1,3-glucanase
Munoz	2009	<i>Colletotrichum</i> spp.		Yes		-
Meng	2010	<i>Alternaria</i> spp.	Pear			-
Maqbool	2010	<i>Colletotrichum</i> spp	Banana			-
Cia	2010	<i>Rhizopus</i> spp.		Yes		-
Yan	2011	<i>Alternaria</i> spp.	Jujube	Yes		-
Abdel-Kader	2011	<i>Penicillium</i> spp.		Yes		-
Xing	2011	<i>Penicillium</i> spp	Jujube			-
Nisia	2012	<i>Penicillium</i> spp.		Yes		-
Ramos-Garcia	2012	<i>Rhizopus</i> spp.	Tomato			-
Shao	2012	<i>Penicillium</i> spp., <i>B. cinerea</i>	Apple			-
Cháfer	2012	<i>Penicillium</i> spp.	Orange			-
Zahid	2012	<i>Colletotrichum</i> spp.	Banana, Papaya, Dragon	Yes		-
Feliziani	2013	<i>B. cinerea</i> , <i>Alternaria</i> spp., <i>Penicillium</i> spp.		Yes	Table grape	Chitinase
Wang	2013	-			Strawberry	β -1,3-Glucanase
Mohamed	2013	<i>Colletotrichum</i> spp.		Yes		-
Gao	2013	<i>B. cinerea</i>	Table grape			-
López-Mora	2013	<i>Alternaria</i> spp.	Mango	Yes		-

Romanazzi	2013	<i>Penicillium</i> spp., <i>B. cinerea</i> , <i>Rhizopus</i> spp.	Strawberry		-
Bill	2014	<i>Colletotrichum</i> spp.		Yes	Avocado PAL, chitinase, β -1,3- glucanase
Ali	2014	<i>Colletotrichum</i> spp.		Yes	Dragon Chitinase, β -1,3- glucanase
Wang	2014	<i>Penicillium</i> spp.	Jujube	Yes	-
Lu	2014	<i>Penicillium</i> spp.	Orange		-
Landi	2014				Strawberry PAL, chitinase, β -1,3- glucanase
Edirisinghe	2014	<i>Colletotrichum</i> spp.	Bell pepper	Yes	-
Zahid	2015				Dragon PAL
Feliziani	2015	<i>B. cinerea</i>	Strawberry		-
Waewthongrak	2015	<i>Penicillium</i> spp.		Yes	Citrus PAL
Varela	2015	<i>Colletotrichum</i> spp.		Yes	-
Shao	2015	<i>Penicillium</i> spp.		Yes	Mandarine PAL, chitinase, β -1,3- glucanase
Xing	2015	<i>Rhizopus</i> spp.	Jujube		-
Ali	2015	<i>Colletotrichum</i> spp.	Bell pepper	Yes	-
Song	2016				Loquat PAL
El Guilli	2016	<i>Penicillium</i> spp.	Citrus		-
Zheng	2017	<i>B. cinerea</i>	Kiwi		-
Gutiérrez-Martínez	2017	<i>Colletotrichum</i> spp.	Mango, banana, soursop	Yes	-
Guo	2017	<i>Alternaria</i> spp.	Jujube		-
Shen	2017	-			Table grape PAL, chitinase, β -1,3- glucanase

Jongsri	2017	-		Mango	PAL, chitinase, β -1,3- glucanase
de Oliveria	2017	<i>Colletotrichum</i> spp.	Yes		-
Kanetis	2017	<i>B. cinerea</i>	Table grape	Yes	-
Munhuweyi	2017	<i>B. cinerea</i>	Yes		-
Silva	2018			Guava	PAL
Gramisci	2018	<i>B. cinerea</i> , <i>Penicillium</i> spp.	Pear		-
Hajji	2018	<i>B. cinerea</i>	Strawberry		-
Kharchoufi	2018	<i>Penicillium</i> spp.	Orange		-
Flores	2018	<i>B. cinerea</i>	Yes		-
Ramos-Guerrero	2018	<i>Colletotrichum</i> spp.	Yes		-
Liu	2018	<i>Penicillium</i> spp.	Blueberry		-
Shi	2018	<i>Penicillium</i> spp.	Grapefruit		-
Xoca-Orozco	2018	<i>Colletotrichum</i> spp.	Yes		-
Obianom	2019	<i>Colletotrichum</i> spp.	Avocado		-
Madanipour	2019	<i>Penicillium</i> spp.	Yes		-



1129

1130

1131 **Figure 1.** Flow chart exhibiting the selection process of eligible studies.

1132




Figure 2. Forest plots using the RavMan 5.3 software for random effects analysis related to the effectiveness of 1% chitosan on disease incidence. Gray mold, blue/ green mold, *Rhizopus* rot., anthracnose and *Alternaria* rot were considered as subgroups. For Feliziani 2013, Kanetis 2017, Lu 2014, Shao 2012, Ramos-Garcia 2012, Gutiérrez-Martínez 2017 and Zahid 2012, several studies were included from each article into the subgroups. IV, inverse variance; CI, confidence interval.




Figure 3. Forest plot using the RavMan 5.3 software for random effects analysis related to the effectiveness of 1% chitosan on *in-vitro* mycelium growth. *Botrytis cinerea*, *Penicillium* spp., *Colletotrichum* spp. and *Alternaria* spp. were considered as subgroups. For Kanetis 2017, Kader 2011, de Oliveria 2017, Gutiérrez-Martínez 2017 and Zahid 2012, several studies were included from each article into the subgroups. IV, inverse variance; CI, confidence interval.




Figure 4. Forest plots using the RavMan 5.3 software for random effects analysis related to the effectiveness of 1% chitosan on plant defence mechanism enzyme activities. Phenylalanine ammonia-lyase (PAL), chitinase and β -1,3-glucanase were considered as subgroups. For Feliziani 2013 several studies were included from each article into the subgroups. IV, inverse variance; CI, confidence interval.




Figure 5. Funnel plots for the detect of publication bias in the studies, for the disease incidence (a), mycelium growth (b) and defence enzyme activity (c) detected after 1% chitosan treatments, compared to the controls. SE(MD) = standard error (mean difference); MD = mean difference.