



UNIVERSITÀ POLITECNICA DELLE MARCHE
Repository ISTITUZIONALE

“Mask up to keep it up”: Preliminary evidence of the association between erectile dysfunction and COVID-19

This is the peer reviewed version of the following article:

Original

“Mask up to keep it up”: Preliminary evidence of the association between erectile dysfunction and COVID-19 / Sansone, A.; Mollaioli, D.; Ciocca, G.; Colonnello, E.; Limoncin, E.; Balercia, G.; Jannini, E. A. - In: ANDROLOGY. - ISSN 2047-2919. - 9:4(2021), pp. 1053-1059. [10.1111/andr.13003]

Availability:

This version is available at: 11566/291884 since: 2024-04-10T13:18:05Z

Publisher:

Published

DOI:10.1111/andr.13003

Terms of use:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. The use of copyrighted works requires the consent of the rights' holder (author or publisher). Works made available under a Creative Commons license or a Publisher's custom-made license can be used according to the terms and conditions contained therein. See editor's website for further information and terms and conditions.

This item was downloaded from IRIS Università Politecnica delle Marche (<https://iris.univpm.it>). When citing, please refer to the published version.

(Article begins on next page)

DR ANDREA SANSONE (Orcid ID : 0000-0002-1210-2843)

PROFESSOR EMMANUELE A JANNINI (Orcid ID : 0000-0002-5874-039X)

Article type : Original Article

“Mask up to keep it up”: preliminary evidence of the association between erectile dysfunction and COVID-19.

Authors: Andrea Sansone¹, Daniele Mollaioli¹, Giacomo Ciocca², Elena Colonnello¹, Erika Limoncin¹, Giancarlo Balercia³, and Emmanuele A. Jannini¹

Affiliations: 1- Chair of Endocrinology and Medical Sexology (ENDOSEX), Department of Systems Medicine, University of Rome Tor Vergata, Roma, Italy

2- Department of Dynamic and Clinical Psychology and Health Studies "Sapienza" University of Rome, Rome, Italy

3- Division of Endocrinology, Department of Clinical and Molecular Sciences, Polytechnic University of Marche, Ancona, Italy

Short title: The link between COVID-19 and erectile dysfunction.

Keywords: erectile dysfunction; COVID-19; endothelial dysfunction; coronavirus; SARS-CoV-2; testosterone

Corresponding Author: Prof. Emmanuele A. Jannini, eajannini@gmail.com, Chair of Endocrinology and Medical Sexology (ENDOSEX), Department of Systems Medicine, University of Rome Tor Vergata, Roma, via Montpellier 1, 00133 Rome, Italy

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/ANDR.13003](https://doi.org/10.1111/ANDR.13003)

This article is protected by copyright. All rights reserved

Abstract

Background. Erectile dysfunction (ED), as the hallmark of endothelial dysfunction, could be a short- or long-term complication of COVID-19. Additionally, being ED a clinical marker and predictor of Non-communicable Chronic Diseases, particularly cardiovascular, subjects with ED could potentially have a higher risk of contracting COVID-19.

Objectives. To investigate the prevalence of ED among subjects with a reported diagnosis of COVID-19 and to measure the association of COVID-19 and ED.

Materials and methods. We reviewed data from the Sex@COVID online survey (performed between April 7th and May 4th, 2020 in Italy) to retrieve a sample of Italian male sexually active subjects with reported SARS-CoV-2 infection. A matching sample of COVID-19 negative male sexually active subjects was also retrieved using propensity score matching in a 3:1 ratio. The survey used different standardized psychometric tools to measure effects of lockdown and social distancing on the intrapsychic, relational, and sexual health of Italian subjects.

Results. 100 subjects were included in the analysis (25 COVID positive; 75 COVID negative). Prevalence of ED, measured with the Sexual Health Inventory for Men, was significantly higher in the COVID+ group (28% vs 9.33%; $p = 0.027$). Logistic regression models confirmed a significant effect of COVID-19 on the development of ED, independently of other variables affecting erectile function, such as psychological status, age and BMI [OR 5.66 ,95% CI: 1.50 – 24.01]. Likewise, subjects with ED were more likely to have COVID-19, once corrected for age and BMI [OR 5.27, 95% CI: 1.49 – 20.09].

Discussion and conclusion. On top of well-described pathophysiological mechanisms, there is preliminary evidence in a real-life population of ED as a risk factor of developing COVID-19 and possibly occurring as a consequence of COVID-19. Universal vaccination against the Covid-19 and the personal protective equipment could possibly have the added benefit of preventing sexual dysfunctions.

Introduction

COVID-19, the coronavirus disease caused by the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), features several clinical phenotypes, ranging from mild to severe forms: the “cytokine storm”^{1,2} may lead to the development of microvascular thrombotic and inflammatory processes, which in turn promote progression to possibly lethal pulmonary complications. Some evidence has also suggested that even “silent”, asymptomatic forms of COVID-19 could have subclinical microvascular involvement^{3,4}, and that long-term cardiovascular sequelae could be expected in COVID-19 patients^{5,6}. Endothelial dysfunction has been considered as the potential trigger for the onset of more severe forms, as well as the link between different comorbidities associated with COVID-19⁷: indeed, COVID-19 is by all means an endothelial disease, in which systemic manifestations of the disease can potentially be due to tissue ischemia resulting from alterations in endothelial thrombotic/fibrinolytic balance⁸. Additionally, endothelial cells express many of the co-factors used by the SARS-CoV-2 to invade host cells⁹.

Erectile dysfunction (ED) has been often considered a hallmark of endothelial dysfunction^{10,11}, and as such a potential association between ED and COVID-19 has also been postulated¹².

Another interesting take on the association between ED and COVID-19 comes from the shared risk factors for the two conditions. Indeed, severity and prevalence of both ED and COVID-19 are higher among men, and even more among subjects suffering from hypertension, obesity, diabetes, and history of cardiovascular disease¹³⁻¹⁶. Broadly speaking, ED is often considered a clinical marker of a “dysfunctional” phenotype, which often features cardiovascular events at an early age. This would possibly suggest that subjects with ED, due to the underlying conditions which impair erectile response, could also be more susceptible to contracting COVID-19.

The present study is, to our knowledge, the first one investigating the prevalence of ED and the possible association between ED and COVID-19 from real-life data in a large survey.

Materials and methods

Subjects

In order to retrieve an adequate sample for the present study, we reviewed data from the Sex@COVID study¹⁷, a previous research project by our group. The Sex@COVID study was an

anonymous web-based questionnaire investigating the psychological, relational, and sexual health of Italian subjects between April 7th and May 4th 2020.

Overall, 6,821 subjects aged 18 years or older (females, 4,177; males: 2,644; mean age 32.83 ± 11.24 years) living in Italy, stratified according to marital status and sexual activity during lockdown, participated in the Sex@COVID study. All subjects provided informed consent to the study, which has been approved by our local Ethical Committee.

985 sexually active men were therefore identified in the Sex@COVID cohort, among which 25 (2.54%) reported being tested as positive for COVID-19.

Measures

Anxiety and depression were respectively measured by the GAD-7¹⁸ (Generalized Anxiety Disorder scale) and the PHQ-9¹⁹ (Patient Health Questionnaire), two questionnaires validated for clinical use, which have already been used to assess psychological outcomes of COVID-19²⁰. For each test, scores ≥10 are considered suggestive of general anxiety disorder and depressive disorder, respectively. Erectile function was measured by the IIEF-5, or Sexual Health Inventory for Men (SHIM)^{21,22}, a shortened, five-item version of the International Index of Erectile Function²³ often used in the clinical setting: scores 21 or below are considered suggestive of ED, whereas scores 22-25 are considered normal.

Statistical analysis

Statistical analysis was performed with the statistical software R (version 3.6.2); statistical significance was set at $p < 0.05$. Propensity score matching was used in order to retrieve two matching samples, based on age, body mass index (BMI), GAD-7 and PHQ-9 scores, using a 1:3 ratio to improve reliability of the results. Assessment of normality was performed using the Shapiro-Wilk test of normality. Wilcoxon rank sum test with continuity correction and Fisher's exact test (one-tailed) were used to assess differences in the numerical and categorical variables between study groups. Chi-squared goodness of fit test was used to measure differences in the prevalence of ED between study groups. Logistic regression models were fitted to assess to which extent age, GAD-7 and PHQ-9 scores, BMI and history of COVID-19 affected erectile function and to measure the effect of age, BMI, and ED on the susceptibility to COVID-19.

Power analysis

Sample size was calculated based on a 28% prevalence of ED among the COVID+ group, using a 0.8 effect size (accounting for a large effect size), with 1:3 ratio, $\alpha = 0.05$ and power 0.95. According to these calculations, a total sample size of 87 (22 COVID+, 65 COVID-) was needed. By including 25 COVID+ and 75 COVID- subjects, the post-hoc analysis yielded a 0.97 statistical power.

Results

According to the suggested propensity score matching approach detailed above, 100 subjects from the 985 sexually active men belonging to the Sex@COVID cohort were retrieved: 25 subjects reported having contracted COVID-19, and a matching sample (based on age, BMI, GAD-7 and PHQ-9 scores) of 75 subjects was recruited among those having no history of SARS-CoV-2 infection prior to filling the Sex@COVID survey. Descriptive data of the study population are reported in Table 1. According to propensity score matching, no statistically significant difference was found for age, GAD-7 and PHQ-9 scores and BMI between the two groups. Prevalence of ED was higher in the COVID+ group (7/25, 28%) than in the COVID- group (7/75, 9.33%; p-value 0.0274) (Figure 1).

Logistic regression models confirmed the association of COVID-19 with ED of COVID-19 on the development of ED (Table 2): while age, BMI, and psychological health scores failed to reach statistical significance, history of COVID-19 was highly significant, resulting in a 5.66 odds ratio [95% confidence interval: 1.50 - 24.01] of having ED.

Since ED could be a valid clinical marker of several underlying and unaccounted conditions, such as diabetes and hypertension, we also measured in the same sample the likelihood of having a self-reported history of COVID-19 following a diagnosis of ED. Logistic regression models adjusted for age and BMI (Table 3) showed a significant association between ED and COVID-19, with a 5.27 odds ratio [95% CI: 1.49 – 20.09].

Discussion

The large majority of the studies investigating the effects of COVID-19 on male sexual and reproductive health have focused on fertility and its preservation²⁴⁻³⁰, and to the present date

there seems to be no definite evidence of the presence of SARS-CoV-2 in seminal fluid^{31,32}; however, there is reason to suspect that sexual quality of life and function might also be impaired as a consequence of COVID-19^{12,33,34}. While of course this is of relative importance to patients in intensive care units, the possible long-term effects on erectile function might be an additional cause of worry in COVID-19 patients. In the present study, to our best knowledge, we investigated for the first time the possible association between erectile function and COVID-19 in a real-life setting. Results of our study agree with the pathophysiological mechanisms linking ED, endothelial dysfunction and COVID-19. By performing propensity score matching, we removed the possible bias resulting from age and BMI, factors which contribute to both increased prevalence of ED^{13,14} and increased susceptibility to COVID-19^{15,16}. Therefore, our results are highly suggestive of the role of the infection in the development of the sexual dysfunction and of the possible clinical relevance of COVID-19 as an additional risk factor for the development of ED. Additionally, considering the bidirectional interaction between sexual activity and psychological well-being¹⁷, the removal of the possible influence of anxiety and depression confirmed that the increased prevalence of ED here found is not only a consequence of the psychological burden of lockdown, but also prominently due to other, *bona fide* organic factors, among which endothelial dysfunction is the most likely culprit. On the other hand, there is another plausible explanation: ED is a well-recognized surrogate marker of systemic health¹¹, and therefore ED patients could already carry several underlying and unexplored risk factors, such as dyslipidemia, diabetes and hypertension, which could increase the likelihood of contracting COVID-19. Hence, we also investigated whether subjects with ED, owing to the worse systemic health¹¹, could possibly be more at risk of developing COVID-19. Based on our preliminary results, ED and COVID-19 seem to be strongly associated, with COVID-19 increasing the chances of developing ED and ED being a marker of increased susceptibility to infection. While more adequately tailored studies are needed, we believe that our study highlights the possible relevance for sexual medicine and andrology of COVID-19 spectrum of disease and its short and long-term health consequences. There is a solid pathogenetic background for the mechanisms through which COVID-19 could affect erectile function¹². Many machineries through which SARS-CoV-2 invades the host cells have been identified: the virus uses angiotensin-converting enzyme 2 (ACE2) as an entry point to the cells, and primes the spike protein, which facilitates viral entry into target cells, by employing

the transmembrane protease, serine 2 TMPRSS2³⁵. The SARS-CoV-2 therefore shows some similarities with another coronavirus^{36,37}, namely SARS-CoV, the virus strain responsible for the SARS outbreak occurring in 2003-2004³⁷⁻³⁹. Since adult Leydig cells express ACE-2, testicular involvement due to COVID-19 has been hypothesized⁴⁰ and confirmed in autopsy reports⁴¹. Also, at least one case of acute orchitis following SARS-CoV-2 infection has been reported⁴². The testicular injury reported in these studies could also be a consequence of alterations in the coagulative status, resulting in development of ischemia at a microvascular level⁴³. Independently of the etiology, testicular damage can possibly lead to the development of a form of hypergonadotropic hypogonadism⁴⁴⁻⁴⁸: as testosterone modulates endothelial function⁴⁹, the possible effects of COVID-19 on erection could also be indirectly due to impaired testosterone secretion from the affected testis, besides the known direct effects of testosterone in male sexual response⁵⁰. Additionally, as higher testosterone levels are also associated with lower levels of pro-inflammatory cytokines such as tumor necrosis factor alpha (TNF- α), interleukins (IL-6 and IL-1 β) and higher levels of anti-inflammatory cytokines (such as IL-10)⁵¹, the “immunothrombotic” mechanism described for COVID-19⁵² and involving the same cytokines^{5,53} could be promoted by the hypogonadal state encountered in affected patients. The immunothrombosis could potentially affect the penile vessels, triggering endothelial dysfunction – therefore impairing vascular function and promoting progression to more severe forms of sexual dysfunction^{54,55}. Other cardiovascular complications of COVID-19, such as cardiomyopathy and myocarditis^{56,57}, could also be involved in the pathogenesis of ED even after the end of the acute phase, potentially becoming long-term cardiovascular sequelae^{5,6}. Other potential factors could contribute to impaired erectile function in COVID-19 patients, such as pulmonary fibrosis^{58,59} causing hypoxia in the penile vascular bed^{60,61}, or anosmia and ageusia, both manifestations of COVID-19^{62,63} with possible negative effects on sexual health⁶⁴⁻⁶⁷. While these factors could possibly have a minor influence when considered individually, the likely presence of most, if not all, of them at the same time could easily allow the progression from a subclinical to an overt form of sexual dysfunction^{68,69}. Considering the subclinical, yet non-negligible, cardiopulmonary damage reported in asymptomatic subjects⁴, we speculate that the same progression from subclinical to an overt sexual dysfunction could also occur in subjects with

“silent” forms of COVID-19 in the presence of additional risk factors – being the proverbial straw that breaks the camel’s back.

Based on the presented evidence and on similarities to previous coronavirus diseases, ED could therefore be both a short-term and a long-term complication of COVID-19. The potential association between ED and COVID-19 might increase awareness of the importance of personal protective equipment, such as masks, and social distancing in a harm reduction perspective for long-term consequences⁷⁰. The use of masks as devices for the prevention of sexual dysfunctions is perhaps a bit stretched, but at present we believe that this could also possibly be an additional strategy to promote the use of personal protective equipment – as people have already posted on Twitter, using the hashtag “#MaskUpToKeepItUp”. Additionally, subjects with a sudden onset or worsening of ED might also consider precautionary quarantine or nasopharyngeal swab, as COVID-19 might act as a potential initiating trigger for the onset of erectile impairment, or an aggravating factor for its progression to more severe forms.

Likewise, subjects with ED should consider their erectile impairment as a sign of possible underlying conditions which could increase the likelihood of suffering from COVID-19. Therefore, the importance of investigating the possible causes of ED becomes of paramount importance in times like these: identifying and treating potential comorbidities would have potentially beneficial effects on erectile function, while at the same time reducing the risk of contracting SARS-CoV-2 or developing more severe forms of COVID-19.

The present study has several limitations, including its retrospective nature, the recall bias associated with the use of online questionnaires, and the inclusion of diagnosis of COVID-19 based on the response to the survey, rather than on nasopharyngeal swabs. Additionally, as the survey did not investigate the potential comorbidities (such as diabetes, endocrine disorders and hypertension ^{13,50,54,71-73}), treatments (such as anti-depressants ^{54,71}) and lifestyles (such as smoking ^{14,54,74}) affecting sexual health, these findings only provide preliminary evidence of the association between COVID-19 and erectile dysfunction. However, at the time of the first Italian lockdown here explored, swabs were rarely available, and an online survey was the most reliable way to obtain information on a large cohort of patients without breaking restrictions. While our study could not include such data, based on the study design, more studies adequately tailored to investigate the endocrine function, most importantly testosterone, and penile vascular

dynamics in patients with a history of COVID-19 are needed to provide definite evidence; however, at present our findings, no matter how limited, are highly suggestive of a potential long-term risk for male sexual function following COVID-19.

Acknowledgments

None.

Funding Information

This work was carried out within the framework of an Italian Ministry of University project and was partly supported by the PRIN 2017S9KTNE_002 grant by the Italian Ministry of Education, University and Research.

Conflicts of Interest

The Authors declare no competing interests for the present study.

Author's contributions

Conceptualization, AS, DM and EAJ; data curation, AS and DM; formal analysis, AS and DM; investigation, AS, DM, GC, EL, EC; methodology, AS and DM.; project administration and supervision, EAJ; validation, EAJ; visualization, AS; writing—original draft, AS and DM; writing—review and editing, GC, EL, EC, GB and EAJ. We attest that all authors contributed significantly to the creation of this manuscript. We confirm that the order of authors listed in the manuscript has been approved by all named authors. All authors have read and agreed to the published version of the manuscript.

References

1. Tay MZ, Poh CM, Renia L, MacAry PA, Ng LFP. The trinity of COVID-19: immunity, inflammation and intervention. *Nat Rev Immunol.* Jun 2020;20(6):363-374. <https://doi.org/10.1038/s41577-020-0311-8>.
2. Jose RJ, Manuel A. COVID-19 cytokine storm: the interplay between inflammation and coagulation. *The Lancet Respiratory Medicine.* 2020;8(6):e46-e47. [https://doi.org/10.1016/s2213-2600\(20\)30216-2](https://doi.org/10.1016/s2213-2600(20)30216-2).

3. Oran DP, Topol EJ. Prevalence of Asymptomatic SARS-CoV-2 Infection : A Narrative Review. *Ann Intern Med.* Sep 1 2020;173(5):362-367. <https://doi.org10.7326/M20-3012>.
4. Long QX, Tang XJ, Shi QL, et al. Clinical and immunological assessment of asymptomatic SARS-CoV-2 infections. *Nat Med.* Aug 2020;26(8):1200-1204. <https://doi.org10.1038/s41591-020-0965-6>.
5. Liu PP, Blet A, Smyth D, Li H. The Science Underlying COVID-19: Implications for the Cardiovascular System. *Circulation.* Jul 7 2020;142(1):68-78. <https://doi.org10.1161/CIRCULATIONAHA.120.047549>.
6. Vittori A, Lerman J, Cascella M, et al. COVID-19 Pandemic Acute Respiratory Distress Syndrome Survivors: Pain After the Storm? *Anesth Analg.* Jul 2020;131(1):117-119. <https://doi.org10.1213/ANE.00000000000004914>.
7. Bernard I, Limonta D, Mahal LK, Hobman TC. Endothelium Infection and Dysregulation by SARS-CoV-2: Evidence and Caveats in COVID-19. *Viruses.* Dec 26 2020;13(1). <https://doi.org10.3390/v13010029>.
8. Libby P, Luscher T. COVID-19 is, in the end, an endothelial disease. *Eur Heart J.* Sep 1 2020;41(32):3038-3044. <https://doi.org10.1093/eurheartj/ehaa623>.
9. Pons S, Fodil S, Azoulay E, Zafrani L. The vascular endothelium: the cornerstone of organ dysfunction in severe SARS-CoV-2 infection. *Crit Care.* Jun 16 2020;24(1):353. <https://doi.org10.1186/s13054-020-03062-7>.
10. Guay AT. ED2: erectile dysfunction = endothelial dysfunction. *Endocrinol Metab Clin North Am.* Jun 2007;36(2):453-63. <https://doi.org10.1016/j.ecl.2007.03.007>.
11. Jannini EA. SM = SM: The Interface of Systems Medicine and Sexual Medicine for Facing Non-Communicable Diseases in a Gender-Dependent Manner. *Sex Med Rev.* Jul 2017;5(3):349-364. <https://doi.org10.1016/j.sxmr.2017.04.002>.
12. Sansone A, Mollaioli D, Ciocca G, et al. Addressing male sexual and reproductive health in the wake of COVID-19 outbreak. *J Endocrinol Invest.* Jul 13 2020. <https://doi.org10.1007/s40618-020-01350-1>.
13. Romanelli F, Sansone A, Lenzi A. Erectile dysfunction in aging male. *Acta Biomed.* 2010;81 Suppl 1:89-94.

14. Mollaioli D, Ciocca G, Limoncin E, et al. Lifestyles and sexuality in men and women: the gender perspective in sexual medicine. *Reprod Biol Endocrinol*. Feb 17 2020;18(1):10. <https://doi.org10.1186/s12958-019-0557-9>.
15. Mahase E. Covid-19: Why are age and obesity risk factors for serious disease? *BMJ*. Oct 26 2020;371:m4130. <https://doi.org10.1136/bmj.m4130>.
16. Caci G, Albini A, Malerba M, Noonan DM, Pochetti P, Polosa R. COVID-19 and Obesity: Dangerous Liaisons. *J Clin Med*. Aug 4 2020;9(8). <https://doi.org10.3390/jcm9082511>.
17. Mollaioli D, Sansone A, Ciocca G, et al. Benefits of Sexual Activity on Psychological, Relational, and Sexual Health During the COVID-19 Breakout. *J Sex Med*. Jan 2021;18(1):35-49. <https://doi.org10.1016/j.jsxm.2020.10.008>.
18. Lowe B, Decker O, Muller S, et al. Validation and standardization of the Generalized Anxiety Disorder Screener (GAD-7) in the general population. *Med Care*. Mar 2008;46(3):266-74. <https://doi.org10.1097/MLR.0b013e318160d093>.
19. Kroenke K, Spitzer RL, Williams JB. The PHQ-9: validity of a brief depression severity measure. *J Gen Intern Med*. Sep 2001;16(9):606-13. <https://doi.org10.1046/j.1525-1497.2001.016009606.x>.
20. Lai J, Ma S, Wang Y, et al. Factors Associated With Mental Health Outcomes Among Health Care Workers Exposed to Coronavirus Disease 2019. *JAMA Netw Open*. Mar 2 2020;3(3):e203976. <https://doi.org10.1001/jamanetworkopen.2020.3976>.
21. Corona G, Jannini EA, Maggi M. Inventories for male and female sexual dysfunctions. *Int J Impot Res*. May-Jun 2006;18(3):236-50. <https://doi.org10.1038/sj.ijir.3901410>.
22. Rosen RC, Cappelleri JC, Smith MD, Lipsky J, Pena BM. Development and evaluation of an abridged, 5-item version of the International Index of Erectile Function (IIEF-5) as a diagnostic tool for erectile dysfunction. *Int J Impot Res*. Dec 1999;11(6):319-26. <https://doi.org10.1038/sj.ijir.3900472>.
23. Rosen RC, Riley A, Wagner G, Osterloh IH, Kirkpatrick J, Mishra A. The international index of erectile function (IIEF): a multidimensional scale for assessment of erectile dysfunction. *Urology*. Jun 1997;49(6):822-30. [https://doi.org10.1016/s0090-4295\(97\)00238-0](https://doi.org10.1016/s0090-4295(97)00238-0).

24. Song C, Wang Y, Li W, et al. Absence of 2019 novel coronavirus in semen and testes of COVID-19 patients. *Biol Reprod.* Jun 23 2020;103(1):4-6. <https://doi.org/10.1093/biolre/ioaa050>.
25. Pan F, Xiao X, Guo J, et al. No evidence of severe acute respiratory syndrome-coronavirus 2 in semen of males recovering from coronavirus disease 2019. *Fertil Steril.* Jun 2020;113(6):1135-1139. <https://doi.org/10.1016/j.fertnstert.2020.04.024>.
26. Paoli D, Pallotti F, Colangelo S, et al. Study of SARS-CoV-2 in semen and urine samples of a volunteer with positive naso-pharyngeal swab. *J Endocrinol Invest.* Dec 2020;43(12):1819-1822. <https://doi.org/10.1007/s40618-020-01261-1>.
27. Holtmann N, Edimiris P, Andree M, et al. Assessment of SARS-CoV-2 in human semen-a cohort study. *Fertil Steril.* Aug 2020;114(2):233-238. <https://doi.org/10.1016/j.fertnstert.2020.05.028>.
28. Ma L, Xie W, Li D, et al. Evaluation of sex-related hormones and semen characteristics in reproductive-aged male COVID-19 patients. *J Med Virol.* Jul 4 2020. <https://doi.org/10.1002/jmv.26259>.
29. Guo L, Zhao S, Li W, et al. Absence of SARS-CoV-2 in semen of a COVID-19 patient cohort. *Andrology.* Jun 29 2020. <https://doi.org/10.1111/andr.12848>.
30. Li D, Jin M, Bao P, Zhao W, Zhang S. Clinical Characteristics and Results of Semen Tests Among Men With Coronavirus Disease 2019. *JAMA Netw Open.* May 1 2020;3(5):e208292. <https://doi.org/10.1001/jamanetworkopen.2020.8292>.
31. Corona G, Baldi E, Isidori AM, et al. SARS-CoV-2 infection, male fertility and sperm cryopreservation: a position statement of the Italian Society of Andrology and Sexual Medicine (SIAMS) (Societa Italiana di Andrologia e Medicina della Sessualita). *J Endocrinol Invest.* Aug 2020;43(8):1153-1157. <https://doi.org/10.1007/s40618-020-01290-w>.
32. Paoli D, Pallotti F, Turriziani O, et al. SARS-CoV-2 presence in seminal fluid: Myth or reality. *Andrology.* May 26 2020. <https://doi.org/10.1111/andr.12825>.
33. Li G, Tang D, Song B, et al. Impact of the COVID-19 Pandemic on Partner Relationships and Sexual and Reproductive Health: Cross-Sectional, Online Survey Study. *J Med Internet Res.* Aug 6 2020;22(8):e20961. <https://doi.org/10.2196/20961>.

34. Panzeri M, Ferrucci R, Cozza A, Fontanesi L. Changes in Sexuality and Quality of Couple Relationship During the COVID-19 Lockdown. *Front Psychol.* 2020;11:565823. <https://doi.org/10.3389/fpsyg.2020.565823>.
35. Zhang H, Penninger JM, Li Y, Zhong N, Slutsky AS. Angiotensin-converting enzyme 2 (ACE2) as a SARS-CoV-2 receptor: molecular mechanisms and potential therapeutic target. *Intensive Care Med.* Apr 2020;46(4):586-590. <https://doi.org/10.1007/s00134-020-05985-9>.
36. Hoffmann M, Kleine-Weber H, Schroeder S, et al. SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked by a Clinically Proven Protease Inhibitor. *Cell.* Apr 16 2020;181(2):271-280 e8. <https://doi.org/10.1016/j.cell.2020.02.052>.
37. Rabaan AA, Al-Ahmed SH, Haque S, et al. SARS-CoV-2, SARS-CoV, and MERS-COV: A comparative overview. *Infez Med.* Ahead Of Print Jun 1 2020;28(2):174-184.
38. Chan KS, Zheng JP, Mok YW, et al. SARS: prognosis, outcome and sequelae. *Respirology.* Nov 2003;8 Suppl(s1):S36-40. <https://doi.org/10.1046/j.1440-1843.2003.00522.x>.
39. Gralinski LE, Bankhead A, 3rd, Jeng S, et al. Mechanisms of severe acute respiratory syndrome coronavirus-induced acute lung injury. *mBio.* Aug 6 2013;4(4). <https://doi.org/10.1128/mBio.00271-13>.
40. Douglas GC, O'Bryan MK, Hedger MP, et al. The novel angiotensin-converting enzyme (ACE) homolog, ACE2, is selectively expressed by adult Leydig cells of the testis. *Endocrinology.* Oct 2004;145(10):4703-11. <https://doi.org/10.1210/en.2004-0443>.
41. Yang M, Chen S, Huang B, et al. Pathological Findings in the Testes of COVID-19 Patients: Clinical Implications. *Eur Urol Focus.* Sep 15 2020;6(5):1124-1129. <https://doi.org/10.1016/j.euf.2020.05.009>.
42. La Marca A, Busani S, Donno V, Guaraldi G, Ligabue G, Girardis M. Testicular pain as an unusual presentation of COVID-19: a brief review of SARS-CoV-2 and the testis. *Reprod Biomed Online.* Nov 2020;41(5):903-906. <https://doi.org/10.1016/j.rbmo.2020.07.017>.
43. Flaifel A, Guzzetta M, Occidental M, et al. Testicular Changes Associated With Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2). *Arch Pathol Lab Med.* Jan 1 2021;145(1):8-9. <https://doi.org/10.5858/arpa.2020-0487-LE>.

44. Rastrelli G, Di Stasi V, Inglese F, et al. Low testosterone levels predict clinical adverse outcomes in SARS-CoV-2 pneumonia patients. *Andrology*. May 20 2020. <https://doi.org10.1111/andr.12821>.
45. Ma L, Xie W, Li D, et al. Effect of SARS-CoV-2 infection upon male gonadal function: A single center-based study. *medRxiv*. 2020:2020.03.21.20037267. <https://doi.org10.1101/2020.03.21.20037267>.
46. Pal R, Banerjee M. COVID-19 and the endocrine system: exploring the unexplored. *J Endocrinol Invest*. Jul 2020;43(7):1027-1031. <https://doi.org10.1007/s40618-020-01276-8>.
47. Okcelik S. COVID-19 pneumonia causes lower testosterone levels. *Andrologia*. Nov 19 2020:e13909. <https://doi.org10.1111/and.13909>.
48. Kadihasanoglu M, Aktas S, Yardimci E, Aral H, Kadioglu A. SARS-CoV-2 Pneumonia Affects Male Reproductive Hormone Levels: A Prospective, Cohort Study. *J Sex Med*. Feb 2021;18(2):256-264. <https://doi.org10.1016/j.jsxm.2020.11.007>.
49. Isidori AM, Buvat J, Corona G, et al. A critical analysis of the role of testosterone in erectile function: from pathophysiology to treatment-a systematic review. *Eur Urol*. Jan 2014;65(1):99-112. <https://doi.org10.1016/j.eururo.2013.08.048>.
50. Sansone A, Romanelli F, Gianfrilli D, Lenzi A. Endocrine evaluation of erectile dysfunction. *Endocrine*. Aug 2014;46(3):423-30. <https://doi.org10.1007/s12020-014-0254-6>.
51. Mohamad NV, Wong SK, Wan Hasan WN, et al. The relationship between circulating testosterone and inflammatory cytokines in men. *Aging Male*. Jun 2019;22(2):129-140. <https://doi.org10.1080/13685538.2018.1482487>.
52. Nakazawa D, Ishizu A. Immunothrombosis in severe COVID-19. *EBioMedicine*. Sep 2020;59:102942. <https://doi.org10.1016/j.ebiom.2020.102942>.
53. Varga Z, Flammer AJ, Steiger P, et al. Endothelial cell infection and endotheliitis in COVID-19. *Lancet*. May 2 2020;395(10234):1417-1418. [https://doi.org10.1016/S0140-6736\(20\)30937-5](https://doi.org10.1016/S0140-6736(20)30937-5).
54. Yafi FA, Jenkins L, Albersen M, et al. Erectile dysfunction. *Nat Rev Dis Primers*. Feb 4 2016;2:16003. <https://doi.org10.1038/nrdp.2016.3>.

55. Maiorino MI, Bellastella G, Giugliano D, Esposito K. From inflammation to sexual dysfunctions: a journey through diabetes, obesity, and metabolic syndrome. *J Endocrinol Invest*. Nov 2018;41(11):1249-1258. <https://doi.org/10.1007/s40618-018-0872-6>.
56. Khalil A, Naneishvili T, Prasad N, Glancy J. SARS-CoV-2 myocarditis: what physicians need to know. *BMJ Evid Based Med*. Jan 5 2021. <https://doi.org/10.1136/bmjebm-2020-111527>.
57. Magadum A, Kishore R. Cardiovascular Manifestations of COVID-19 Infection. *Cells*. Nov 19 2020;9(11). <https://doi.org/10.3390/cells9112508>.
58. Spagnolo P, Balestro E, Aliberti S, et al. Pulmonary fibrosis secondary to COVID-19: a call to arms? *Lancet Respir Med*. Aug 2020;8(8):750-752. [https://doi.org/10.1016/S2213-2600\(20\)30222-8](https://doi.org/10.1016/S2213-2600(20)30222-8).
59. George PM, Wells AU, Jenkins RG. Pulmonary fibrosis and COVID-19: the potential role for antifibrotic therapy. *Lancet Respir Med*. Aug 2020;8(8):807-815. [https://doi.org/10.1016/S2213-2600\(20\)30225-3](https://doi.org/10.1016/S2213-2600(20)30225-3).
60. Graney BA, Wamboldt FS, Baird S, et al. Looking ahead and behind at supplemental oxygen: A qualitative study of patients with pulmonary fibrosis. *Heart Lung*. Sep - Oct 2017;46(5):387-393. <https://doi.org/10.1016/j.hrtlng.2017.07.001>.
61. Cavezzi A, Troiani E, Corrao S. COVID-19: hemoglobin, iron, and hypoxia beyond inflammation. A narrative review. *Clin Pract*. May 19 2020;10(2):1271. <https://doi.org/10.4081/cp.2020.1271>.
62. Eshraghi AA, Mirsaeidi M, Davies C, Telischi FF, Chaudhari N, Mittal R. Potential Mechanisms for COVID-19 Induced Anosmia and Dysgeusia. *Front Physiol*. 2020;11:1039. <https://doi.org/10.3389/fphys.2020.01039>.
63. Han AY, Mukdad L, Long JL, Lopez IA. Anosmia in COVID-19: Mechanisms and Significance. *Chem Senses*. Jun 17 2020. <https://doi.org/10.1093/chemse/bjaa040>.
64. Ottaviano G, Zuccarello D, Frasson G, et al. Olfactory sensitivity and sexual desire in young adult and elderly men: an introductory investigation. *Am J Rhinol Allergy*. May-Jun 2013;27(3):157-61. <https://doi.org/10.2500/ajra.2013.27.3879>.
65. Siegel JK, Kung SY, Wroblewski KE, Kern DW, McClintock MK, Pinto JM. Olfaction Is Associated With Sexual Motivation and Satisfaction in Older Men and Women. *J Sex Med*. Jan 7 2021. <https://doi.org/10.1016/j.jsxm.2020.12.002>.

66. Schafer L, Mehler L, Hahner A, Walliczek U, Hummel T, Croy I. Sexual desire after olfactory loss: Quantitative and qualitative reports of patients with smell disorders. *Physiol Behav.* Mar 15 2019;201:64-69. <https://doi.org/10.1016/j.physbeh.2018.12.020>.
67. Bendas J, Hummel T, Croy I. Olfactory Function Relates to Sexual Experience in Adults. *Arch Sex Behav.* Jul 2018;47(5):1333-1339. <https://doi.org/10.1007/s10508-018-1203-x>.
68. Jannini EA, Lenzi A, Isidori A, Fabbri A. Subclinical erectile dysfunction: proposal for a novel taxonomic category in sexual medicine. *J Sex Med.* Sep 2006;3(5):787-794. <https://doi.org/10.1111/j.1743-6109.2006.00287.x>.
69. Colonnello E, Ciocca G, Limoncin E, Sansone A, Jannini EA. Redefining a sexual medicine paradigm: subclinical premature ejaculation as a new taxonomic entity. *Nat Rev Urol.* Jan 13 2021. <https://doi.org/10.1038/s41585-020-00417-1>.
70. Sansone A, Limoncin E, Colonnello E, et al. Harm Reduction in Sexual Medicine. *Sex Med Rev.* accepted.
71. Shamloul R, Ghanem H. Erectile dysfunction. *Lancet.* Jan 12 2013;381(9861):153-65. [https://doi.org/10.1016/S0140-6736\(12\)60520-0](https://doi.org/10.1016/S0140-6736(12)60520-0).
72. Foresta C, Caretta N, Corona G, et al. Clinical and metabolic evaluation of subjects with erectile dysfunction: a review with a proposal flowchart. *Int J Androl.* Jun 2009;32(3):198-211. <https://doi.org/10.1111/j.1365-2605.2008.00932.x>.
73. Mulhall JP, Giraldi A, Hackett G, et al. The 2018 Revision to the Process of Care Model for Evaluation of Erectile Dysfunction. *J Sex Med.* Sep 2018;15(9):1280-1292. <https://doi.org/10.1016/j.jsxm.2018.06.005>.
74. Corona G, Sansone A, Pallotti F, et al. People smoke for nicotine, but lose sexual and reproductive health for tar: a narrative review on the effect of cigarette smoking on male sexuality and reproduction. *J Endocrinol Invest.* Apr 22 2020. <https://doi.org/10.1007/s40618-020-01257-x>.

Table 1. Characteristics of the study population. All data expressed as median [Interquartile range] except prevalence of erectile dysfunction, expressed as n (%). BMI: body mass index; PHQ-9: Patient Health Questionnaire; GAD-7: Generalized Anxiety Disorder scale. °Wilcoxon rank sum test with continuity correction; §Fisher's exact test (one-tailed)

	COVID+ (n=25)	COVID- (n=75)	p-value
Age (years)	39.00 [29.00, 45.00]	42.00 [32.50, 49.00]	0.142°
BMI (kg/m²)	22.65 [20.83, 23.74]	22.74 [20.98, 24.53]	0.266°
GAD-7 Score	4.00 [2.00, 6.00]	4.00 [2.00, 5.00]	0.741°
PHQ-9 Score	5.00 [3.00, 6.00]	4.00 [2.00, 5.00]	0.873°
Erectile dysfunction	7 (28%)	7 (9.33%)	0.027§


Table 2. Results of logistic regression models for erectile dysfunction (ED) according to age, body mass index (BMI), psychological health and self-referred history of COVID-19. PHQ-9: Patient Health Questionnaire; GAD-7: Generalized Anxiety Disorder scale.

	Estimate	Std. Error	p-value	Odds ratio [95% CI]
Intercept	-9.607	3.216	0.003	0.00 [0.00 - 0.02]
Age (years)	0.062	0.027	0.023	1.06 [1.01 - 1.13]
BMI	0.143	0.106	0.178	1.15 [0.93 - 1.43]
GAD-7 Score	0.114	0.193	0.556	1.12 [0.77 - 1.66]
PHQ-9 Score	0.148	0.143	0.301	1.16 [0.88 - 1.55]
Previous COVID-19	1.734	0.694	0.013	5.66 [1.50 - 24.01]

Table 3. Results of logistic regression models for COVID according to age, body mass index (BMI) and erectile dysfunction (ED).

	Estimate	Std. Error	p-value	Odds ratio [95% CI]
Intercept	1.910	2.194	0.384	6.75 [0.11 - 661.99]
Age (years)	-0.027	0.021	0.197	0.97 [0.93 - 1.01]
BMI	-0.096	0.089	0.281	0.91 [0.76 - 1.07]
Presence of ED	1.662	0.652	0.011	5.27 [1.49 - 20.09]

Figure 1. Prevalence of erectile dysfunction in the COVID+ and COVID- groups.



andr_13003_f1.tiff