



UNIVERSITÀ POLITECNICA DELLE MARCHE  
Repository ISTITUZIONALE

One is not enough: Monitoring microplastic ingestion by fish needs a multispecies approach

This is the peer reviewed version of the following article:

*Original*

One is not enough: Monitoring microplastic ingestion by fish needs a multispecies approach / Valente, Tommaso; Pelamatti, Tania; Avio, Carlo Giacomo; Camedda, Andrea; Costantini, Maria Letizia; de Lucia, Giuseppe Andrea; Jacomini, Carlo; Piermarini, Raffaella; Regoli, Francesco; Sbrana, Alice; Ventura, Daniele; Silvestri, Cecilia; Matiddi, Marco. - In: MARINE POLLUTION BULLETIN. - ISSN 0025-326X. - STAMPA. - 184:(2022). [10.1016/j.marpolbul.2022.114133]

*Availability:*

This version is available at: 11566/314553 since: 2024-04-11T14:38:18Z

*Publisher:*

*Published*

DOI:10.1016/j.marpolbul.2022.114133

*Terms of use:*

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. The use of copyrighted works requires the consent of the rights' holder (author or publisher). Works made available under a Creative Commons license or a Publisher's custom-made license can be used according to the terms and conditions contained therein. See editor's website for further information and terms and conditions.

This item was downloaded from IRIS Università Politecnica delle Marche (<https://iris.univpm.it>). When citing, please refer to the published version.

(Article begins on next page)

***One is not enough: Monitoring microplastic ingestion by fish needs a multispecies approach.***

*Tommaso Valente<sup>a,b,\*</sup>, Tania Pelamatti<sup>b</sup>, Carlo Giacomo Avio<sup>c</sup>, Andrea Camedda<sup>d</sup>, Maria Letizia Costantini<sup>a</sup>, Giuseppe Andrea de Lucia<sup>d</sup>, Carlo Jacomini<sup>b</sup>, Raffaella Piermarini<sup>b</sup>, Francesco Regoli<sup>c</sup>, Alice Sbrana<sup>b,e</sup>, Daniele Ventura<sup>a</sup>, Cecilia Silvestri<sup>b</sup>, Marco Matiddi<sup>b</sup>.*

<sup>a</sup>'La Sapienza' University of Rome, Department of Environmental Biology, Piazzale Aldo Moro 5, 00185, Rome, RM, Italy.

<sup>b</sup>ISPRA, Italian National Institute for Environmental Protection and Research, Via di Castel Romano 100, 00128, Rome, RM, Italy.

<sup>c</sup>Marche Polytechnic University, Department of Life and Environmental Sciences (DiSVA), Via Brecce Bianche snc, 60131, Ancona, AN, Italy.

<sup>d</sup>IAS-CNR, Institute of Anthropic Impact and Sustainability in Marine Environment, National Research Council Oristano Section, Località Sa Mardini, 09170, Torregrande, OR, Italy.

<sup>e</sup>PhD Program in Evolutionary Biology and Ecology, Department of Biology, University of Rome 'Tor Vergata', Via della Ricerca Scientifica snc, 00133, Rome, RM, Italy.

**\*Corresponding author: Tommaso Valente**

University of Rome 'La Sapienza'

Department of Environmental Biology

5 P.le Aldo Moro

Rome, RM, ITALY 00185

E-mail: [tommaso.valente@uniroma1.it](mailto:tommaso.valente@uniroma1.it)

Tel.: +39 0649914773

Fax: +39 0649914773

24    **Abstract**

25    The development of monitoring programmes based on bioindicators is crucial for assessing the impact of  
26    microplastic ingestion on marine organisms. This study presents results from an Italian pilot action aimed at  
27    investigating the suitability of a monitoring strategy based on a multispecies approach. The benthic-feeder *Mullus*  
28    *barbatus*, the demersal species *Merluccius merluccius*, and the pelagic-feeder species of the genus *Scomber* were  
29    used to assess the environmental contamination by microplastics in three different marine areas, namely Ancona  
30    (Adriatic Sea), Anzio (Tyrrhenian Sea), and Oristano (Western Sardinia). Microplastic ingestion frequencies were  
31    higher in samples from Anzio (26.7%) and Ancona (25.0%) than Oristano (14.4%), suggesting a relationship  
32    between microplastic bioavailability and the proximity to urban settlements and river flows. Furthermore,  
33    microplastic ingestion was affected by the feeding habits of the examined species. The detected differences  
34    reinforce the hypothesis that a multispecies approach is needed to evaluate microplastic ingestion by marine  
35    animals.

36    **Keywords:** marine litter; micro-litter; bioindicator; feeding habits; Mediterranean Sea; MSFD

## 1. Introduction

Since the 1950s, the establishment of a consumption-based human society determined the release of significant quantities of waste into the environment (Zalasiewicz et al., 2016). Most are made of plastic (Woodall et al., 2014), a widespread set of synthetic polymers with low biodegradability (Palmisano and Pettigrew, 1992). Despite the low toxicity associated with most plastic materials (Worm et al., 2017), plastic pollution is one of the major environmental threats of current times (Horton et al., 2017). Indeed, plastics are often loaded with hazardous chemicals, such as additives, dyes, and flame retardants (Deanin, 1975; Lithner et al., 2011; Fries et al., 2013). Furthermore, the environmental persistence of plastics results in a progressive fragmentation process, impacting items exposed to ultraviolet radiation, chemical oxidation, mechanical abrasion, and biological agents (Liu et al., 2020; Turner et al., 2020). As a result, all the environmental compartments are today contaminated by the presence of microplastics (MPs), generally defined as tiny pieces of plastic smaller than 5 mm in size (Arthur et al., 2009).

Oceans and seas represent the main sink for MPs (Hale et al., 2020). In the aquatic media, MPs can sorb many kinds of organic and inorganic pollutants, including toxic and carcinogenic substances (Pelamatti et al., 2021; Rai et al., 2022). At the same time, MPs overlap the size range of prey of a wide variety of marine animals that can ingest these small particles either accidentally, or intentionally – by confusing plastic particles with natural or potential preys, as well as due to secondary ingestion (*i.e.*, items already ingested by prey) (Anbumani and Kakkar, 2018; Fossi et al., 2018; Prinz and Korez, 2020). Therefore, MPs are hazardous contaminants that can act as vectors of chemicals through the marine food webs, posing threats to ecosystem functioning (Carbery et al., 2018). Several laboratory experiments linked plastic ingestion to negative physiological effects, such as energy depletion, starvation, increased immune response, and decreased fecundity (von Moos et al., 2012; Wright et al., 2013; Rios-Fuster et al., 2021). Field studies also reported physical impacts, including blockage of the digestive tract, abrasions, and difficulties in breathing (Sharma and Chatterjee, 2017).

Several legal and policy frameworks address the “marine litter” topic, such as the Global Partnership on Marine Litter (GPML, <https://www.gpmarinelitter.org/>), the Honolulu Strategy, and the G7 Countries Agenda (Löhr et al., 2017). Even within the UN Sustainable Development Agenda, 4 of 17 Sustainable Development Goals (SDGs) pose targets concerning the reduction and mitigation of plastic pollution in marine environments by 2030 (<https://sdgs.un.org/>). In this context, the UN Environmental Assembly (UNEA-5.2 <https://www.unep.org/environmentassembly/>) aims to forge an international agreement for a global transition to a circular economy. At the Mediterranean level, the Regional Plan on Marine Litter Management sets legally binding targets by 2025 to deal with marine litter from both land- and sea-based sources (UN Environment/Mediterranean Action Plan, <https://www.unep.org/unepmap/>). Similarly, the 2008/56/EC Marine Strategy Framework Directive (MSFD) fixes the aim of achieving the Good Environmental Status (GES, “*the environmental status of marine waters where these provide ecologically diverse and dynamic oceans and seas which are clean, healthy and productive*”) by defining the criteria D10C3 as “*The amount of litter and micro-litter ingested by marine animals is at a level that does not adversely affect the health of the species concerned*” (Commission Decision 2017/848/EU).

To drive and confirm the strength of the programs of measures implemented by legislative frameworks, the UNEP/MAP recognized the importance of developing appropriate monitoring strategies based on bioindicators for assessing the occurrence and the impacts of MP ingestion on marine organisms (Galgani, 2017). The MSFD

Technical Group on Marine Litter (TG-ML) established essential criteria for the selection of bioindicators of marine litter ingestion, such as sample availability, regular litter consumption, and sufficient knowledge of the biology of the involved species, including habitat, trophic level, feeding behavior, spatial distribution, commercial importance, and conservation status (Fossi et al., 2018). The seabird *Fulmarus glacialis* in the Northern European seas and the loggerhead sea turtle *Caretta caretta* in the Mediterranean basin are the target species for monitoring macro-litter (> 5 mm) ingestion (van Franeker et al., 2011; Matiddi et al., 2017; 2019). In contrast, a micro-litter monitoring strategy is not yet defined, though fish species are good candidates for assessing this impact (Bray et al., 2019).

Several recent studies show that MP ingestion occurs in many fish species, including bony fishes and elasmobranchs of commercial importance that showed to be regular litter consumers (Wang et al., 2020). Previous studies successfully proposed monitoring programs based on the assessment of MP ingestion by examining a single target species (e.g., Tsangaris et al., 2020). However, it is known that the feeding habits of different fish species may determine differences in MP ingestion rates (Miller et al., 2020; Justino et al., 2021). Moreover, the distribution of MPs in the marine environment varies according to their different shape, size, and chemical composition (Palazzo et al., 2021). Several environmental factors (such as waves, tides, and currents) at different geographical scales contribute to determining different accumulation pathways for different MP types (Li et al., 2020). In this view, it is likely that more than one fish species should be selected for describing MP contamination of the marine environment. In particular, the target species should have different feeding behaviors and habitat uses to investigate different marine compartments within the examined area (Matiddi et al., 2021).

This study presents results from an Italian pilot action, which aims to investigate the adequacy of a monitoring strategy based on the assessment of MP ingestion by fish species with different feeding habits. This activity was planned by a national consortium involving the Italian National Institute for Environmental Protection and Research (ISPRA), the National Research Council – Institute of Anthropic Impacts and Sustainability in Marine Environment (CNR-IAS), and the Marche Polytechnic University – Department of Life and Environmental Sciences (UNIVPM-DiSVA). Three marine areas characterized by different MP contamination sources and circulation patterns were investigated to highlight site-specific variations in MPs ingested by different fish species. The selected species are the benthic-feeder *Mullus barbatus*, the demersal species *Merluccius merluccius*, and the pelagic-feeder species of the genus *Scomber*. These species were designated because of their different feeding habits and considering commercial importance, their availability in the selected areas, and the well-documented occurrence of MP ingestion (Giani et al., 2019; Avio et al., 2020; Bianchi et al., 2020; Capillo et al., 2020). The red mullet *M. barbatus* is a benthivorous species that feeds on the sea bottom by swallowing sediment with the prey and expelling the sediment through the gills (Labropoulou and Eleftheriou, 1997). Therefore, it could be regarded as an indicator of MP ingestion within the strictly benthic compartment. Differently, the European hake *M. merluccius* is a demersal top predator that frequently moves from the sea floor to mid- and surface-waters, where it typically feeds upon fast-moving preys, such as cephalopods and fish (Carpentieri et al., 2005). Finally, the Atlantic mackerel *S. scombrus* and the Atlantic chub mackerel *S. colias* are pelagic species preferring zooplankton and small pelagic fish (Lopes et al., 2020), and therefore might represent the pelagic compartment.


The main aim of this study is to provide further information for improving the future development of marine micro-litter monitoring systems by: i) investigating possible differences in MP ingestion by different fish species within

the same area; ii) highlighting the diversity of MPs ingested by the same species in three different marine areas; iii) verifying the ecological relevance of a multi-species monitoring approach.

## 2. Materials and methods

### 2.1 Study areas and sampling activities

Fish were sampled in August-October 2019 at 3 sampling sites within 3 different Italian coastal areas: Ancona in the Adriatic Sea, Anzio in the Tyrrhenian Sea, and Oristano on the Western side of Sardinia (Figure 1). The marine area off the city of Ancona is characterized by low depths and it is affected by the inputs of the Po River and other relevant runoffs. The Po River is the main Italian river, which flows into the Adriatic Sea  $\approx 150$  km north of the sampling area. The effect of the West Adriatic current (flowing toward the South) together with the intense urbanization of the coast and the presence of commercial seaports make this area a prospective site of plastic accumulation within the Mediterranean Sea (UNEP/MAP, 2012; Liubartseva et al., 2016; Giani et al., 2019). Similarly, the Tyrrhenian Sea area off Anzio coast is affected by the discharge of the Tiber River (distance from the mouth  $\approx 50$  km), the second largest river catchment in Italy, and shows a high coastal urbanization. On the other hand, the basin is characterized by a wider bathymetric range, a complex bottom topography, and a highly variable circulation that isolates this area from other Mediterranean sub-regions during the warm season (Inghilesi et al., 2012), determining the potential accumulation of waste from local inputs. Finally, the Gulf of Oristano is not closely affected by large rivers and important urban settlements. Therefore, due to the distance of the island of Sardinia from the mainland, the plastic contamination in this area is mainly due to the transport conveyed by the prevailing winds, which blow from West/North-West, and by the inflow of Modified Atlantic Water (MAW) (Camedda et al., 2021; Palazzo et al., 2021).



**Figure 1.** Map of sampling areas: Ancona (Adriatic Sea), Anzio (Tyrrhenian Sea), and Oristano (Western Sardinia).

In each of the three areas, fish were caught by local fisheries using trawling nets for commercial fishery, with a hemp-coloured cod-end and a 40 mm square mesh. During each fishing trip (geographical centroids: Ancona – 43°41'45"N, 013°33'32"E; Anzio – 41°18'31"N, 012°28'43"E; Oristano – 39°30'19"N, 008°20'29"E), the nets were towed for 3 hours at an average speed of 3 knots (estimated swept area  $\approx 0.8 \text{ km}^2$ ). A total of 264 specimens were collected: 84 from Ancona (24 *M. merluccius*, 30 *M. barbatus*, and 30 *S. scombrus*), 90 from Anzio (30 *M. merluccius*, 30 *M. barbatus*, and 30 *S. colias*), and 90 from Oristano (30 *M. merluccius*, 30 *M. barbatus*, and 30 *S. colias*). Table 1 reports information on the mean values ( $\pm$  se) of total length, total weight, gastrointestinal weight, and relative condition factor (Kn) recorded for each species within each sampling area.

	<i>M. merluccius</i>	<i>M. barbatus</i>	<i>Scomber</i> spp.
<b>a) Total length [cm]</b>			
Ancona	30.0 $\pm$ 1.5	16.5 $\pm$ 1.2	22.2 $\pm$ 0.9
Anzio	28.9 $\pm$ 1.3	15.5 $\pm$ 0.7	27.2 $\pm$ 1.3
Oristano	23.0 $\pm$ 1.6	13.2 $\pm$ 0.7	26.8 $\pm$ 2.0
All Locations	27.4 $\pm$ 3.1	15.1 $\pm$ 1.6	25.4 $\pm$ 2.7
<b>b) Total weight [g]</b>			
Ancona	198.9 $\pm$ 26.1	57.1 $\pm$ 13.4	93.3 $\pm$ 12.3
Anzio	172.5 $\pm$ 23.3	36.6 $\pm$ 4.8	157.6 $\pm$ 24.7
Oristano	114.2 $\pm$ 22.0	32.9 $\pm$ 5.6	209.2 $\pm$ 49.7
All Locations	159.3 $\pm$ 42.4	42.2 $\pm$ 13.8	153.4 $\pm$ 57.7
<b>c) GI weight [g]</b>			
Ancona	7.1 $\pm$ 3.5	1.6 $\pm$ 1.0	6.5 $\pm$ 2.6
Anzio	6.3 $\pm$ 5.3	1.4 $\pm$ 0.4	7.2 $\pm$ 1.9
Oristano	5.9 $\pm$ 3.3	2.0 $\pm$ 0.7	12.8 $\pm$ 6.8
All Locations	6.4 $\pm$ 4.1	1.7 $\pm$ 0.8	8.8 $\pm$ 5.2
<b>d) Relative condition factor (Kn)</b>			
Ancona	1.05 $\pm$ 0.10	1.11 $\pm$ 0.09	1.14 $\pm$ 0.19
Anzio	1.01 $\pm$ 0.06	0.88 $\pm$ 0.06	0.85 $\pm$ 0.04
Oristano	1.23 $\pm$ 0.12	1.29 $\pm$ 0.13	1.17 $\pm$ 0.07
All Locations	1.10 $\pm$ 0.13	1.09 $\pm$ 0.19	1.05 $\pm$ 0.19

**Table 1.** Morpho-anatomical data (mean  $\pm$  sd) of fish samples (*Merluccius merluccius*, *Mullus barbatus*, and *Scomber* spp.) collected in August-October 2019 within three Italian marine areas (Ancona, Anzio, and Oristano): **a)** total length [cm]; **b)** total weight [g]; **c)** gastrointestinal (GI) weight [g]; **d)** relative condition factor (Kn). Kn was computed according to the expression  $\text{Kn} = \text{TW} \cdot (a \cdot \text{TL}^b)^{-1}$ , where  $a$  and  $b$  are the log-transformed length-weight relationship parameters, TW is the total wet weight expressed in grams, and TL is the total length expressed in cm.

After the collection onboard or at landing, all the individuals were immediately stored at -20 °C until further analyses. Specimens showing diseases, evidence of net feeding, or regurgitation were discarded.

## 2.2 Lab analyses

Lab analyses were performed in the laboratories of ISPRA (Anzio), CNR-IAS (Oristano), and DiSVA (Ancona) following the guidelines provided in Matiddi et al. (2021). Following this protocol, the MPs in this study were defined as “All sorts of small particles of plastic, less than 5 mm in size in two of the three dimension or diameter that pass through a 5 mm mesh screen but are retained by a lower one, according to the chosen size class”, fixing the lower size limit to 100  $\mu\text{m}$  due to operational limits.

Total length and total wet weight of each individual were recorded after thawing at room temperature. Then, fish were dissected to extract the entire gastrointestinal tract (GI), which was weighed to the nearest 0.1 g. Each GI was placed into individual glass jars and filled with 15%  $\text{H}_2\text{O}_2$  to digest the biogenic component. Extraction of micro-items larger than 100  $\mu\text{m}$  was performed using a vacuum pump for filtering the digestates on glass microfiber (Whatman GF/D<sup>TM</sup>; pore size: 2.7  $\mu\text{m}$ ) or cellulose nitrate (Sartorius 11301; pore size: 8  $\mu\text{m}$ ) membranes. MP identification was made using a dissecting microscope. Shape type (filament, film, fragment,

granule, or pellet), colour, and size class (class 1: 100 µm – 330 µm; class 2: 330 µm – 1 mm; class 3: 1 mm – 5 mm) was recorded for each collected item. Polymeric characterization was determined using Fourier Transformed Infra-Red Micro-Spectrometry (µFT-IR). Fibres from textiles were not considered. The discrimination between fibres and filaments was based on an accurate shape analysis of each thread-like item. Filaments were defined as rod-like particles with regular diameter, while microfibers were characterized as items with a ribbon-like shape, not regular diameter, and frayed ends. Polymeric characterization was performed for all the questionable items.

#### 2.2.1 Quality assurance and quality control

All the precautions suggested by Matiddi et al. (2021) were adopted during every analytical step to reduce secondary contamination. Dissecting procedures were performed in a clean room where airflow and staff presence were limited. Workbenches, instruments, and tools were cleaned with ethanol or rinsed with ultrapure water before every use. Only glass and metal labware was used whenever applicable and the wearing of synthetic clothes was avoided. All dissecting tools, containers, and sieves were additionally cleaned with compressed air. Aluminium foils were used to cover labware and samples exposed to airborne contamination. Covered Petri dishes were used to store and preserve membranes.

A contamination control was treated in parallel to each batch of 10 samples. Blank controls only revealed the presence of microfibers made of natural polymers (cellulosic-made or wool; mean  $\pm$  se =  $2.61 \pm 0.79$  items  $\cdot$  control<sup>-1</sup>). Since natural microfibers were not considered in this study, no results adjustment techniques were adopted.

#### 2.3 Data and statistical analyses

Relative fish condition factor (Kn) was computed for each individual. Kn calculation was performed according to the expression  $Kn = TW \cdot (a \cdot TL^b)^{-1}$ , where  $a$  and  $b$  are the log-transformed length-weight relationship parameters, TW is the total wet weight expressed in grams, and TL is the total length expressed in cm. Frequency of Occurrence (FO) was computed as no. of individuals with ingested MPs  $\cdot$  no. of individuals examined<sup>-1</sup>. The abundance of ingested MPs was expressed as the average ( $\pm$  standard error, se) no. of ingested MPs.

Differences in FO were tested using Generalized Linear Models (GLMs) assuming a binomial error structure and a logit link function. Differences in MP abundance were analyzed using a negative-binomial GLM. Relationships between biological parameters (namely total length, total weight, GI weight, and Kn) and the occurrence of ingested MPs were investigated through binomial Generalized Linear Mixed Models (GLMMs), setting sampling area as random effect. Correspondence analysis was performed to highlight differences in the types of ingested MPs by species and sampling area. Association between variables was tested through chi-square statistics. The significance level was set at  $p < 0.05$ .

Statistical analyses were performed with R4.1.0 (R Core Team, 2021) using packages lme4 (Bates et al., 2015), FactoMineR (Le et al., 2008), factoextra (Kassambara and Mundt, 2020), ggplot2 (Wickham, 2016), and gplots (Warnes et al., 2020).

### 3. Results

In all three marine areas, MP ingestion occurred in all the examined species, with an overall frequency of occurrence of 22.0%. FO was higher in samples from Anzio (26.7%) and Ancona (25.0%) than Oristano (14.4%).




Comparing species, the highest FO was recorded in *Scomber* spp. (33.3%), followed by *M. merluccius* (20.2%), and *M. barbatus* (12.2%). The 58 individuals with ingested MPs contained a total of 73 items (mean  $\pm$  se =  $1.26 \pm 0.09$  items  $\cdot$  individual<sup>-1</sup>). The max numbers of MPs extracted from an individual were: 2 for *M. barbatus*, 2 for *M. merluccius*, and 5 for *Scomber* spp. FOs and MP abundances (average  $\pm$  se) for species by sampling area are available in Table 2.

	<i>M. merluccius</i>	<i>M. barbatus</i>	<i>Scomber</i> spp.	All species
<b>a) FO (%)</b>				
Ancona	33.3	16.7	26.7	25.0
Anzio	20.0	10.0	50.0	26.7
Oristano	10.0	10.0	23.3	14.4
All Locations	20.2	12.2	33.3	22.0
<b>b) MP abundance (considering all the examined individuals)</b>				
Ancona	0.38 $\pm$ 0.12	0.17 $\pm$ 0.07	0.30 $\pm$ 0.10	0.27 $\pm$ 0.05
Anzio	0.23 $\pm$ 0.09	0.13 $\pm$ 0.08	0.73 $\pm$ 0.20	0.37 $\pm$ 0.08
Oristano	0.13 $\pm$ 0.08	0.13 $\pm$ 0.08	0.30 $\pm$ 0.11	0.19 $\pm$ 0.05
All Locations	0.24 $\pm$ 0.06	0.14 $\pm$ 0.04	0.44 $\pm$ 0.08	0.28 $\pm$ 0.04
<b>c) MP abundance (considering only individuals with ingested MP)</b>				
Ancona	1.12 $\pm$ 0.12	1.00 $\pm$ 0.00	1.12 $\pm$ 0.12	1.10 $\pm$ 0.07
Anzio	1.17 $\pm$ 0.17	1.33 $\pm$ 0.33	1.47 $\pm$ 0.29	1.38 $\pm$ 0.19
Oristano	1.33 $\pm$ 0.33	1.33 $\pm$ 0.33	1.29 $\pm$ 0.19	1.31 $\pm$ 0.13
All Locations	1.18 $\pm$ 0.12	1.18 $\pm$ 0.12	1.33 $\pm$ 0.15	1.16 $\pm$ 0.09

**Table 2.** Microplastic ingestion events recorded in fish samples (*Merluccius merluccius*, *Mullus barbatus*, and *Scomber* spp.) collected within three Italian marine areas (Ancona, Anzio, and Oristano) in August-October 2019: **a)** Frequency of occurrence (FO = no. of individuals with ingested MPs  $\cdot$  no. of total individuals examined<sup>-1</sup>); **b)** average  $\pm$  se no. of ingested MPs considering all the individuals examined; **c)** average  $\pm$  se no. of ingested MPs considering only the individuals with ingested MPs.

Comparing marine areas, the highest abundance of filaments was extracted in samples from Anzio (42.4% of the total number of MPs found). Fragments were always among the most represented shape categories (39.4% at Anzio, 47.8% at Ancona, 29.4% at Oristano). Pellets and granules were isolated only in samples from Oristano. Considering colours, most of the MPs were black (23.29%) or blue (19.19%). Further colours frequently found were green (16.44%) and white (16.44%), followed by transparent (10.96%) and red (9.69%). Other colours were rarer (4.11%). Following the proposed size classification of MPs, most items ranged between 100 and 330  $\mu$ m (56.16%). Items in the other two size classes (330  $\mu$ m - 1 mm, and 1 mm - 5 mm) had equal frequencies (21.92% both). A detailed summary of MP types found ingested by species and sampling area is reported in Figure 2.



**Figure 2.** Heatmap representing the diversity of microplastic types in the gastrointestinal tract of fish samples (*Merluccius merluccius*, *Mullus barbatus*, and *Scomber spp.*) collected in three Italian marine areas (Ancona, Anzio, and Oristano) in August-October 2019.


The polymeric composition of 48 out of the 73 recovered MPs was determined through spectroscopic analyses (65.7%). The most represented polymer was polyethylene (PE=33.33%), followed by polypropylene (PP=14.58%), polyester (PEST=12.50%), polyacrylate (PAK=8.33%), polyethylene terephthalate (PET=6.25%), and nylon (PA=6.25%). Other particles found were made of copolymers (PE/PP and Ethylene/Metacrilate=16.67%), or other plastic types such as EDPM rubber and MATER-BI bioplastic. Polymer frequencies by species and sampling areas are available in Table 3.

	Polymer							
	PE	PP	PEST	PAK	PET	PA	COP	OTH
<b>a) Sampling area</b>								
Ancona	25.0	-	20.8	0.8	-	12.5	16.7	16.7
Anzio	40.0	25.0	-	5.0	15.0	5.0	10.0	-
Oristano	40.0	40.0	20.0	-	-	-	-	-
<b>b) Species</b>								
<i>M. merluccius</i>	44.4	-	-	11.1	-	22.2	22.2	-
<i>M. barbatus</i>	-	-	57.1	-	28.6	-	-	14.3
<i>Scomber spp.</i>	37.5	21.9	6.25	6.25	3.13	3.13	12.5	9.4

**Table 3.** Percentages of polymers found ingested in fish samples collected in August-October 2019 by: **a)** sampling areas; **b)** fish species. Abbreviations: polyethylene, PE; polypropylene, PP; polyester, PEST; polyacrylate, PAK; polyethylene terephthalate, PET; nylon, PA; copolymers, COP; other, OTH.

Statistical analysis of absence/presence data showed a two-group partitioning of the samples, the former including samples with FOs equal to 10.0% (i.e., *M. barbatus* from Anzio and Oristano, and *M. merluccius* from Oristano),


and the latter including all the other samples with FOs equal to or higher than 16.7% (Figure 3; Table 1S). No significant differences among samples were detected considering the number of ingested MPs (all p-values > 0.05).



**Figure 3.** Barplot of the frequency of microplastic ingestion (FO) detected in fish samples (*Merluccius merluccius*, *Mullus barbatus*, and *Scomber* spp.) collected within three Italian marine areas (Ancona, black; Anzio, dark grey; and Oristano, light grey) in August-October 2019. Bars with \* indicate significant lower values according to a Generalized Linear Model (GLM) computed by assuming a binomial error structure and a logit link function.

Relationships between biological parameters (total length, total weight, GI weight, and Kn) and the occurrence of ingested MPs were investigated through GLMMs. Only a slightly significant correlation for the effect of GI weight on frequency of ingestion by *M. merluccius* was found (p-value = 0.031; Table 2S).

Correspondence analysis showed a significant association between the combined variable “species-sampling area” and the shape ( $X^2 = 46.771$ ;  $df = 24$ ; p-value <0.01) and size ( $X^2 = 42.604$ ;  $df = 16$ ; p-value <0.001) of ingested MPs. Figure 4 showed how the types of MPs ingested by the examined species were different according to both their feeding habits and the sampling location. Figure 4a highlighted different proportions of MP shape-types ingested by *M. merluccius* in the three sampling areas, while the between-sites differences for the other two species seemed less relevant. Figure 4b showed a high prevalence of small MPs (size class 1: 100  $\mu\text{m}$  - 330  $\mu\text{m}$ ) in *M. barbatus*.



**Figure 4.** Biplots from correspondence analysis describing the diversity of microplastics ingested by different fish species (*Merluccius merluccius*, *Mullus barbatus* and *Scomber* spp.) sampled within three Italian marine areas (Ancona, Anzio, and Oristano) in August-October 2019: **a)** shape categories (i.e., filament, film, fragment, and other); **b)** size classes (size class 1: 100  $\mu$ m – 330  $\mu$ m; class 2: 330  $\mu$ m – 1 mm; class 3: 1 mm – 5 mm). Abbreviations: B, *M. barbatus*; M, *M. merluccius*; S, *Scomber* spp.; 1, Ancona; 2, Anzio; 3, Oristano. Percentages of explained variance are labelled on the relevant axis.

#### 4. Discussion

The impact of plastic pollution on marine ecosystems is one of the main environmental issues of emerging concern (Thushari and Senevirathna, 2020). Although monitoring programs of environmental matrices are currently underway (e.g., in the case of seawater or beach litter within the European MSFD), the fulfilment of new tools for assessing MP bioavailability is crucial for understanding the threat posed by marine litter to marine life. As a result of snapshot in time, the present study shows that MP ingestion occurs in all the examined species within the three sampling areas, confirming that MP ingestion by marine fish is widespread (Savoca et al., 2021). The overall FO (22.0%) is relatively low compared to other studies from European seas (median FO = 42.0%; Wootton et al., 2021), but we must notice that our results are not fully comparable with other studies due to the exclusion of fibres, which must be considered in future specific investigations aimed at describing their diversity within different marine compartments (Athey and Erdle, 2022). Nevertheless, the detected differences among species and sampling areas provide interesting insights for developing a monitoring strategy based on fish species as bioindicators of MP impact on biota.

##### 4.1 Frequency of occurrence

Coastal anthropogenic pressures are recognized as key predictors of spatial variations in MP ingestion rates (Sbrana et al., 2020; Tsangaris et al., 2020). In the present study, FO is slightly lower in samples from Oristano (14.4%) than Ancona and Anzio (25.0% and 26.7%, respectively). This difference could be explained considering that Anzio and Ancona marine areas have more densely populated coasts (with urban settlements hosting more than 100,000 inhabitants compared to the 30,000 inhabitants living at Oristano; ISTAT, 2018), are crossed by important shipping lanes, and are affected by the flows of the two main Italian rivers (the Tiber River and the Po River, respectively) (Inghilesi et al., 2012; Liubartseva et al., 2016). The relationship between the proximity to the contamination sources and MP bioavailability seems to suggest that the marine environment is continuously subject to diffuse contamination through new MP inputs. Then, the definition of the assessment areas and the

sampling strategy should be carefully planned to obtain reliable estimates of temporal variations of MP loads from land-based activities (Bai et al., 2022).

However, it is important to notice that the differences among the three locations emerge only considering all the species together. Indeed, *M. barbatus* from Anzio and Oristano have the same FO. Likewise, no significant FO differences distinguished the *Scomber* spp. samples collected in the three marine areas. A similar picture of the overall conditions (*i.e.*, Anzio  $\approx$  Ancona  $>$  Oristano) is obtained from the analysis of *M. merluccius*, but only the integration with the data from the other two species returns a comprehensive description of the contamination status (see Subsection 4.4). Our results show that different species can show different MP ingestion frequencies within the same location. Nevertheless, each single species has different FOs within different locations. This evidence suggests that the environmental contamination by MPs cannot be described using a one-time assessment based on the examination of only one species.

As requested by the MSFD for EU Member States, a reliable strategy for monitoring the impact of litter ingestion on marine organisms should provide an ecologically relevant assessment. Therefore, study areas must be selected according to the several factors that may affect MP distribution, such as the presence of contamination sources (*e.g.*, industrial, fluvial, urban) and oceanographic features (*i.e.*, wind stress, water depth, waves, tides, and currents). Moreover, the assessment of MP ingestion should be based on the analysis of a reasonable number of species that can be regarded as indicators of MP ingestion in different marine compartments (Avio et al., 2020). On the other hand, the number of individuals analyzed per single species must be sufficient to allow reliable statistical analysis (Matiddi et al., 2021). A monitoring program should then optimize the work effort (in terms of sampling activities, lab analysis, and staff employment) to obtain a proper characterization of MP ingestion events occurring in a determined area. This study suggests that the analysis of about 30 individuals of at least three different fish species (for a minimum of  $\approx 90$  samples from each sampling area) might provide a fair quantity of information. However, more extensive targeted studies will need to determine the efforts required to obtain reliable data on spatial and temporal variability, as well as in terms of statistical power. For instance, the number of individuals needed may depend on the degree of contamination of the assessment area (Matiddi et al., 2021). Furthermore, the number of target species that should be examined to obtain an ecologically relevant assessment could be larger and variable according to the complexity of the resident trophic web (Avio et al., 2020).

## 4.2 Abundance of ingested MPs

The number of MPs ingested by individuals is low, with a maximum value of 5 recorded in a *S. colias* specimen from Anzio. This result is in line with previous studies showing that in regions with a high incidence of plastic contamination MPs are ingested in relatively low quantities by many specimens of different species (*e.g.*, Moore et al., 2001; Lusher et al., 2013; Foekema et al., 2013; Valente et al., 2019). The low number of MPs per individual associated with high FOs suggests that the excretion rates of MPs are greater than ingestion rates (Sbrana et al., 2020). As a result, the residence time of MPs in the gastrointestinal tract of fish is relatively short, and it does not lead to their bioaccumulation over time or biomagnification along food chains. However, this could not be true either in severely contaminated areas – where MP encounter rates exceed the rates of depuration, or for MPs with smaller sizes (the operational lower size limit in this study is 100  $\mu\text{m}$ ) (Avio et al., 2017; Covernton et al., 2021; Miller et al., 2021).

In the perspective of a monitoring program, the absence of bioaccumulation and biomagnification processes ensures that fish can be considered as appropriate bioindicators of MP ingestion at local scale, since it is unlikely that they retain MPs ingested far from sampling locations (Sbrana et al., 2020). On the other hand, it is difficult to highlight species-specific, spatial, and temporal differences in MP ingestion through MPs abundance or MP concentration values (Covernton et al., 2021). Therefore, FO appears to be the most reliable index for evaluating the incidence of MP ingestion for monitoring purposes and for studies comparison (Avio et al., 2020). Nevertheless, it should be considered that the retention time of MPs varies according to the range of specific biological traits and the diet composition of the examined species (Valente et al., 2019). This could affect the estimation of MP ingestion probabilities through FO values and future studies will need to understand how biological traits of different fish species may alter the retention time of ingested MPs. Moreover, these considerations are valid for the size range of MPs that are commonly documented in biota (*i.e.*, >100  $\mu\text{m}$ ; Covernton et al., 2021) and further considerations would be made for MPs of smaller sizes.

#### 4.3 Relationships between biological parameters and MP ingestion

Previous studies describe some relationships between the biological parameters of fish and MP ingestion. Alomar and Deudero (2017) and Valente et al. (2019) highlight a positive correlation between GI fullness of elasmobranch species and the number ingested MPs, suggesting that the retention time of MPs and food items could be the same. In this study, we observed a significant correlation between the GI weight and the frequency of MP ingestion by *M. merluccius*. This could be due to the strict predatory habit of this species (Carpentieri et al., 2005), which is potentially exposed to secondary ingestion through contaminated prey. Indeed, the ingestion of many preys (in turn exposed to MP ingestion) could lead to an increase in the probability of MP secondary ingestion and affect the diversity of ingested MP types. However, we have no tools to unequivocally confirm this speculation, which should be considered in future studies that should also provide new methodological approaches allowing MP detection and diet analysis simultaneously.

Further research will also be needed to consider the potential influence of fish conditions on MP ingestion. In this study, we did not find significant direct correlation between the occurrence of ingested MPs and the relative condition factor of individuals. Although MP ingestion seems not to affect individual body conditions in these species (Compa et al., 2018; Sbrana et al., 2020), considering *M. merluccius* and *Scomber* spp., we noted a correspondence between Kn and FO at the site-specific level. In particular, *M. merluccius* from Anzio (FO = 20.0%; Kn =  $1.01 \pm 0.06$ ) and Ancona (FO = 33.3%; Kn =  $1.05 \pm 0.10$ ) have lower values of Kn and significant higher values of FO compared to *M. merluccius* from Oristano (FO = 10.0%; Kn =  $1.23 \pm 0.12$ ). Similarly, *Scomber* sp. from Anzio (FO = 50.0%; Kn =  $0.86 \pm 0.05$ ) have very low Kn values and a very high FO, while the relative condition factor and FOs computed for samples from Ancona (FO = 26.7%; Kn =  $1.14 \pm 0.19$ ) and Oristano (FO = 13.3%; Kn =  $1.18 \pm 0.07$ ) are similar. Considering the degree of coastal anthropogenic pressure impacting Ancona and Anzio marine areas (see Subsections 2.1 and 4.1), MP ingestion seems to reflect the overall environmental conditions of the marine compartments exploited by the examined species.

Since a capillary data collection is crucial to increase the knowledge about the possible predictors and effects of MP ingestion by fish, the recording of data on biological parameters (*e.g.*, sex, maturity, and weight of liver and gonads) and the evaluation of fish conditions should be strongly recommended during monitoring activities.

#### 4.4 MP typologies

The results of this study show that the types of MPs ingested by the examined species are different both according to the feeding habits and the sampling areas. Comparing species, *M. barbatus* tends to ingest smaller MPs, mostly filaments made of high-density polymers such as PEST, PET, and EDPM that, due to their physical characteristics, are more susceptible to wind mixing, vertical transport, and therefore sinking to the sea bottom (Woodall et al., 2014; Bergmann et al., 2017; Courtené-Jones et al., 2017; 2018). Therefore, it is not surprising to find these types of MPs in the GI of benthic feeder species like *M. barbatus*, which search for food in the sediment (Labropoulou and Eleftheriou, 1997). In contrast, the analysis of MPs ingested by *M. merluccius* and *Scomber* spp. reveals a predominance of polymers with lower densities (such as PE, PP, and Ethylene/Metacrilate copolymers) and a wider variety of shape-types and sizes. Overall, *Scomber* spp. showed the highest variability in ingested MP types in each sampling area. On the other hand, the between-sites differences were more relevant in *M. merluccius*. This could be likely related to the feeding habits of these species. Lopes et al. (2020) found that MP ingestion by *Scomber* spp. is significantly related to its diet composition, suggesting that a size-selection mechanism might drive MP ingestion in these species. Differently, the predatory behavior of *M. merluccius* might result in a higher incidence of secondary ingestion (see Subsection 4.3). Indeed, *M. merluccius* can be exposed to different MP types depending on the selectivity of its prey, which can vary among areas and life stages (Mahe et al., 2007).

All these results reinforce the hypothesis that a multispecies approach is needed to assess MP ingestion. There are many site-specific factors governing the distribution of MPs, including the presence of different contamination sources (e.g., industrial, fluvial, urban) and oceanographic features (i.e., wind stress, water depth, waves, tides, and currents). Furthermore, the bioavailability of different MP types could be affected by species-specific biological and ecological traits such as feeding habits, which vary considerably between different distribution areas. Further studies will need to clarify the relative importance of the several mechanisms affecting the complex interaction patterns between MPs and marine organisms. Particular attention must be paid to the selection of target species for monitoring programs. In fact, only an exhaustive description of the MP ingestion events occurring in an area will allow the collection of relevant and comparable data through time and space.

## 5. Conclusions

The development of monitoring programs is essential to reveal the effectiveness of measures that could be applied by policy to reduce plastic pollution, such as the ban of single-use plastic and the implementation of a wide educational plan to increase the awareness of new generations. Through a snapshot in time, this study provides a set of suggestions that can improve the future development of monitoring programs of MP ingestion, such as the ones required by the Marine Strategy Framework Directive for the European Seas. Our results confirm that fish species are suitable bioindicators of MP ingestion at a local scale. The relationship between MP ingestion frequency and the proximity to MP contamination sources (such as urban settlements, river flows, and shipping lanes) suggests that the marine environment is continuously subject to new MP inputs. Therefore, monitoring programs should investigate marine areas at different distances from land-based and sea-based human activities, including both highly anthropized sites and theoretically pristine areas. Furthermore, this study shows that different fish species can show different MP ingestion frequencies within the same marine area and even that biological and ecological traits of different species may affect the bioavailability of different MP types. Therefore, an ecologically relevant assessment of MP ingestion cannot be based the examination of only one species. A multispecies approach should be developed considering that the choice of target species can vary according to the complexity of the

399 resident trophic web. Further research is needed to select appropriate target species within different marine areas  
400 and to develop a common strategy to obtain suitable and comparable data through time and space.



## **Funding**

This study was supported by the project INDICIT II, funded by the European Commission – DG Environment [Grant N. 11.0661/2016/748064/SUB/ENV.C2]. TV was supported by a PhD fellowship granted by ‘La Sapienza’ University of Rome. AS was supported by a research fellowship granted by University of Rome ‘Tor Vergata’, Department of Biology.

## **CRedit authorship contribution statement**

**Tommaso Valente:** Conceptualization, Methodology, Investigation, Data curation, Formal analysis, Writing – Original Draft; **Tania Pelamatti:** Investigation, Writing – Review and Editing; **Carlo Giacomo Avio:** Methodology, Investigation, Data curation; Writing – Review and Editing; **Andrea Camedda:** Data curation, Writing – Review and Editing; **Maria Letizia Costantini:** Supervision, Writing – Review and Editing; **Giuseppe Andrea de Lucia:** Supervision, Writing – Review and Editing; **Carlo Jacomini:** Writing – Review and Editing; **Raffaella Piermarini:** Resources, Project administration; **Francesco Regoli:** Supervision, Writing – Review and Editing; **Alice Sbrana:** Investigation, Writing – Review and Editing; **Daniele Ventura:** Supervision, Writing – Review and Editing; **Cecilia Silvestri:** Supervision, Project administration, Resources, Writing – Review and Editing; **Marco Matiddi:** Conceptualization, Supervision, Project administration, Resources, Writing – Review and Editing.

## **Declaration of competing interest**

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

## **Acknowledgements**

This study was planned within the framework of the INDICIT II European Project (<https://indicit-europa.eu>) – Activity 4: Implementation of the indicator “Micro-debris ingested by fish and sea turtles” at the OSPAR and Barcelona RSCs and MSFD Areas. We are grateful to the two anonymous reviewers for their helpful comments and suggestions. We are grateful to the two anonymous reviewers for their helpful comments and suggestions. Special thanks to the local fisheries for their contribution during the sampling activities.

## **Data statement**

Data will be made available on request.

## References

- Alomar, C., Deudero, S., 2017. Evidence of microplastic ingestion in the shark *Galeus melastomus* Rafinesque, 1810 in the continental shelf off the western Mediterranean Sea. *Environ. Pollut.* 223, 223-229. <https://doi.org/10.1016/j.envpol.2017.01.015>
- Anbumani, S., Kakkar, P., 2018. Ecotoxicological effects of microplastics on biota: a review. *Environ. Sci. Pollut. R.* 25(15), 1473–1496. <https://doi.org/10.1007/s11356-018-1999-x>
- Arthur, C., Baker, J., Bamford, H., 2009. Proceedings of the international research workshop on the occurrence, effects, and fate of microplastic marine debris. NOAA marine debris program. Technical memorandum NOS-OR&R-30. <https://marinedebris.noaa.gov/proceedings-second-research-workshop-microplastic-marine-debris> (Accessed 28 March 2022)
- Athey, S.N., Erdle, L.M., 2022. Are we understanding anthropogenic microfiber pollution? A critical review of occurrence, methods, and reporting. *Environ. Toxicol. Chem.* 41(4), 822-837. <https://doi.org/10.1002/etc.5173>
- Avio, C.G., Cardelli, L.R., Gorbi, S., Pellegrini, D., Regoli, F., 2017. Microplastics pollution after the removal of the Costa Concordia wreck: first evidences from a biomonitoring case study. *Environ. Pollut.* 227, 207-214. <https://doi.org/10.1016/j.envpol.2017.04.066>
- Avio, C.G., Pittura, L., d'Errico, G., Abel, S., Amorello, S., Marino, G., Gorbi, S., Regoli, F., 2020. Distribution and characterization of microplastic particles and textile microfibers in Adriatic food webs: General insights for biomonitoring strategies. *Environ. Pollut.* 258, 113766. <https://doi.org/10.1016/j.envpol.2019.113766>
- Bai, M., Lin, Y., Hurley, R.R., Zhu, L., Li, D., 2022. Controlling factors of microplastic riverine flux and implications for reliable monitoring strategies. *Environ. Sci. Technol.* 56(1), 48-61. <https://doi.org/10.1021/acs.est.1c04957>
- Bates, D., Maechler, M., Bolker, B., Walker, S., 2015. Fitting Linear Mixed-Effects Models using lme4. *J. Stat. Softw.* 67(1), 1-48. <https://doi.org/10.18637/jss.v067.i01>
- Bergmann, M., Wirzberger, V., Krumpfen, T., Lorenz, C., Primpke, S., Tekman, M.B., Gerdts, G., 2017. High quantities of microplastic in Arctic deep-sea sediments from the HAUSGARTEN observatory. *Environ. Sci. Technol.* 51(19), 11000-11010. <https://doi.org/10.1021/acs.est.7b03331>
- Bianchi, J., Valente, T., Scacco, U., Cimmaruta, R., Sbrana, A., Silvestri, C., Matiddi, M., 2020. Food preference determines the best suitable digestion protocol for analysing microplastic ingestion by fish. *Mar. Pollut. Bull.* 154, 111050. <https://doi.org/10.1016/j.marpolbul.2020.111050>
- Bray, L., Digka, N., Tsangaris, C., Camedda, A., Gambaiani, D., de Lucia, G.A., Matiddi, M., Miaud, C., Palazzo, L., Pérez-del-Olmo, A., Raga, J.A., Silvestri, C., Kaberi, H., 2019. Determining suitable fish to monitor plastic ingestion trends in the Mediterranean Sea. *Environ. Pollut.* 247, 1071-1077. <https://doi.org/10.1016/j.envpol.2019.01.100>
- Camedda, A., Coppa, S., Palazzo, L., Marra, S., Massaro, G., Serrentino, F., Vencato, S., Brundu, R., de Lucia, G.A., 2021. Characterization and assessment of micro and macroscopic litter in Sardinian beaches (Western Mediterranean Sea). *Water Air Soil Poll.* 232(2), 1-14. <https://doi.org/10.1007/s11270-021-04993-9>
- Capillo, G., Savoca, S., Panarello, G., Mancuso, M., Branca, C., Romano, V., D'Angelo, G., Bottari, T., Spanò, N., 2020. Quali-quantitative analysis of plastics and synthetic microfibers found in demersal species from Southern Tyrrhenian Sea (Central Mediterranean). *Mar. Pollut. Bull.* 150, 110596. <https://doi.org/10.1016/j.marpolbul.2019.110596>
- Carbery, M., O'Connor, W., Palanisami, T., 2018. Trophic transfer of microplastics and mixed contaminants in the marine food web and implications for human health. *Environ. Int.* 115, 400-409. <https://doi.org/10.1016/j.envint.2018.03.007>

471 Carpentieri, P., Colloca, F., Cardinale, M., Belluscio, A., Ardizzone, G.D., 2005. Feeding habits of European hake  
472 (*Merluccius merluccius*) in the central Mediterranean Sea. Fish B-NOAA 103(2), 411-416.  
473 <https://aquadocs.org/handle/1834/26234>

474 Compa, M., Ventero, A., Iglesias, M., Deudero, S., 2018. Ingestion of microplastics and natural fibres in *Sardina*  
475 *pilchardus* (Walbaum, 1792) and *Engraulis encrasicolus* (Linnaeus, 1758) along the Spanish Mediterranean coast.  
476 Mar. Pollut. Bull. 128, 89-96. <https://doi.org/10.1016/j.marpolbul.2018.01.009>

477 Courtene-Jones, W., Quinn, B., Ewins, C., Gary, S.F., Narayanaswamy, B.E., 2019. Consistent microplastic  
478 ingestion by deep-sea invertebrates over the last four decades (1976-2015), a study from the North East Atlantic.  
479 Environ. Pollut. 244, 503-512. <https://doi.org/10.1016/j.envpol.2018.10.090>

480 Courtene-Jones, W., Quinn, B., Gary, S.F., Mogg, A.O.M., Narayanaswamy, B.E., 2017. Microplastic pollution  
481 identified in deep-sea water and ingested by benthic invertebrates in the Rockall Trough, North Atlantic Ocean.  
482 Environ. Pollut. 231, 271-280. <https://doi.org/10.1016/j.envpol.2017.08.026>

483 Covernton, G.A., Davies, H.L., Cox, K.D., El-Sabaawi, R., Juanes, F., Dudas, S.E., Dower, J.F., 2021. A Bayesian  
484 analysis of the factors determining microplastics ingestion in fishes. J. Haz. Mat. 413, 125405.  
485 <https://doi.org/10.1016/j.jhazmat.2021.125405>

486 Deanin, R.D., 1975. Additives in plastics. Environ. Health Persp. 11, 35-39. <https://doi.org/10.1289/ehp.751135>

487 Foekema, E.M., De Gruijter, C., Mergia, M.T., van Franeker, J.A., Murk, A.J., Koelmans, A.A., 2013. Plastic in  
488 North Sea fish. Environ. Sci. Technol. 47, 8818-8824. <https://doi.org/10.1021/es4000931b>

489 Fossi, M.C., Pedà, C., Compa, M., Tsangaris, C., Alomar, C., Claro, F., Ioakeimidis, C., Galgani, F., Hema, T.,  
490 Deudero, S., Romeo, T., Battaglia, P., Andaloro, F., Caliani, I., Casini, S., Panti, C., Baini, M., 2018. Bioindicators  
491 for monitoring marine litter ingestion and its impacts on Mediterranean biodiversity. Environ. Pollut. 237, 1023-  
492 1040. <https://doi.org/10.1016/j.envpol.2017.11.019>

493 Fries, E., Dekiff, J.H., Willmeyer, J., Nuelle, M.-T., Ebert, M., Remy, D., 2013. Identification of polymer types  
494 and additives in marine microplastic particles using pyrolysis-GC/MS and scanning electron microscopy. Environ.  
495 Sci. Process. Impacts 15, 1949-1956. <https://doi.org/10.1039/c3em00214d>

496 Galgani, F., 2017. Specially Protected Areas Protocol Regional Activity Centre (Barcelona Convention), Defining  
497 the Most Representative Species for IMPA Common Indicator 24. SPA/RAC, Tunis.

498 Giani, D., Baini, M., Galli, M., Casini, S., Fossi, M.C., 2019. Microplastic occurrence in edible fish species (*Mullus*  
499 *barbatus* and *Merluccius merluccius*) collected in three different geographical sub-areas of the Mediterranean Sea.  
500 Mar. Pollut. Bull. 140, 129-137. <https://doi.org/10.1016/j.marpolbul.2019.01.005>

501 Hale, R.C., Seeley, M.E., La Guardia, M.J., Mai, L., Zeng, E.Y., 2020. A global perspective on microplastics. J.  
502 Geophys. Res.-Oceans 125(1), e2018JC014719. <https://doi.org/10.1029/2018JC014719>

503 Horton, A.A., Walton, A., Spurgeon, D.J., Lahive, E., Svendsen, C., 2017. Microplastics in freshwater and  
504 terrestrial environments: Evaluating the current understanding to identify the knowledge gaps and future research  
505 priorities. Sci. Total. Environ. 586, 127-141. <https://doi.org/10.1016/j.scitotenv.2017.01.190>

506 Inghilesi, R., Ottolenghi, L., Orasi, A., Pizzi, C., Bignami, F., Santoleri, R., 2012. Fate of river Tiber discharge  
507 investigated through numerical simulation and satellite monitoring. Ocean Sci. 8, 773-786.  
508 <https://doi.org/10.5194/os-8-773-2012>

509 ISTAT, 2018. Principali Statistiche Geografiche Sui Comuni. World Wide Web electronic publication, version  
510 (12/18).

511 Justino, A.K.S., Lenoble, V., Pelage, L., Ferreira, G.V.B., Passarone, R., Frédou, T., Frédou, F.L., 2021.  
512 Microplastic contamination in tropical fishes: An assessment of different feeding habits. Reg. Stu. Mar. Sci. 45,  
513 101857. <https://doi.org/10.1016/j.rsma.2021.101857>

514 Kassambara, A., Mundt, F., 2020. factoextra: Extract and visualize the results of multivariate data analyses. R  
515 packages version 1.0.7. <https://CRAN.R-project.org/package=factoextra>

516 Labropoulou, M., Eleftheriou, A., 2005. The foraging ecology of two pairs of congeneric demersal fish species:  
517 Importance of morphological characteristics in prey selection. J. Fish Biol. 50(2), 324-340.  
518 <https://doi.org/10.1111/j.1095-8649.1997.tb01361.x>

519 Le, S., Josse, J., Husson, F., 2008. FactoMineR: An R package for multivariate analysis. J. Stat. Softw. 25(1), 1-  
520 18. <https://doi.org/10.18637/jss.v025.i01>

521 Li, Y., Zhang, H., Tang, C., 2020. A review of possible pathways of marine microplastics transport in the ocean.  
522 Anthropocene Coasts 3(1), 6-13. <https://doi.org/10.1139/anc-2018-0030>

523 Lithner, D., Larsson, Å., Dave, G., 2011. Environmental and health hazard ranking and assessment of plastic  
524 polymers based on chemical composition. Sci. Total Environ. 409(18), 3309-3324.  
525 <https://doi.org/10.1016/j.scitotenv.2011.04.038>

526 Liu, P., Zhan, X., Wu, X., Li, J., Wang, H., Gao, S., 2020. Effect of weathering on environmental behavior of  
527 microplastics: Properties, sorption and potential risk. Chemosphere 242, 125193.  
528 <https://doi.org/10.1016/j.chemosphere.2019.125193>

529 Liubartseva, S., Coppini, G., Lecci, R., Creti, S., 2016. Regional approach to modeling the transport of floating  
530 debris in the Adriatic Sea. Mar. Pollut. Bull. 103, 115-127. <http://dx.doi.org/10.1016/j.marpolbul.2015.12.031>

531 Löhr, A., Savelli, H., Beunen, R., Kalz, M., Ragas, A., Van Belleghem, F., 2017. Solutions for global marine litter  
532 pollution. Curr. Opin. Env. Sust. 28, 90-99. <https://doi.org/10.1016/j.cosust.2017.08.009>

533 Lopes, C., Raimundo, J., Caetano, M., Garrido, S., 2020. Microplastic ingestion and diet composition of  
534 planktivorous fish. Limnol. Oceanogr.-Let. 5, 103-112. <https://doi.org/10.1002/lol2.10144>

535 Lusher, A.L., McHugh, M., Thompson, R.C., 2013. Occurrence of microplastic in the gastrointestinal tract of  
536 pelagic and demersal fish from the English Channel. Mar. Pollut. Bull. 67, 94-99.  
537 <https://doi.org/10.1016/j.marpolbul.2012.11.028>

538 Mahe, K., Amara, R., Bryckaert, T., Kacher, M., Brylinski, J.M., 2007. Ontogenetic and spatial variation in the  
539 diet of hake (*Merluccius merluccius*) in the Bay of Biscay and the Celtic Sea. ICES J. Mar. Sci. 64(6), 1210-1219.  
540 <https://doi.org/10.1093/icesjms/fsm100>

541 Matiddi, M., Hochscheid, S., Camedda, A., Baini, M., Cocumelli, C., Serena, F., Tomassetti, P., Travaglini, A.,  
542 Marra, S., Campani, T., Scholl, F., Mancusi, C., Amato, E., Briguglio, P., Maffucci, F., Fossi, M.C., Bentivegna,  
543 F., de Lucia, G.A., 2017. Loggerhead sea turtles (*Caretta caretta*): A target species for monitoring litter ingested  
544 by marine organisms in the Mediterranean Sea. Environ. Pollut. 230, 199-209.  
545 <https://doi.org/10.1016/j.envpol.2017.06.054>

546 Matiddi, M., de Lucia, G.A., Silvestri, C., Darmon, G., Tomás, J., Pham, C.K., Camedda, A., Vandeperre, F.,  
547 Claro, F., Kaska, Y., Kaberi, H., Revuelta, O., Piermarini, R., Daffina, R., Pisapia, M., Genta, D., Sözbilen, D.,  
548 Bradai, M.N., Rodríguez, Y., Gambaiani, D., Tsangaris, C., Chaieb, O., Moussier, J., Loza, A.L., Miaud, C., 2019.  
549 Data collection on marine litter ingestion in sea turtles and thresholds for Good Environmental Status. J. Vis. Exp.  
550 147, e59466. <https://doi.org/10.3791/59466>

551 Matiddi, M., Pham, C.K., Anastasopoulou, A., Andresmaa, E., Avio, C.G., Bianchi, J., Chaieb, O., Palazzo, L.,  
552 Darmon, G., de Lucia, G.A., Deudero, S., Sozbilen, D., Eriksson, J., Fischer, E., Gómez, M., Herrera, A., Hattia,  
553 E., Kaberi, H., Kaska, Y., Kühn, S., Lips, I., Miaud, C., Gambaiani, D., Nelms, S., Piermarini, R., Regoli, F.,  
554 Sbrana, A., Setälä, O., Settiti, S., Soederberg, L., Tomás, J., Tsangaris, C., Vale, M., Valente, T., Silvestri, C.,  
555 2021. Monitoring micro-litter ingestion in marine fish: A harmonized protocol for MSFD and RSCs areas.  
556 INDICIT II Project. [https://indicit-europa.eu/cms/wp-content/uploads/2021/06/Monitoring-microlitter-ingestion-](https://indicit-europa.eu/cms/wp-content/uploads/2021/06/Monitoring-microlitter-ingestion-in-marine-fish-1.pdf)  
557 [in-marine-fish-1.pdf](https://indicit-europa.eu/cms/wp-content/uploads/2021/06/Monitoring-microlitter-ingestion-in-marine-fish-1.pdf) (Accessed 30 March 2022).

558 Miller, M.E., Hamann, M., Kroon, F.J., 2020. Bioaccumulation and biomagnification of microplastics in marine  
559 organisms: A review and meta-analysis of current data. Plos One 15(10), e0240792.  
560 <https://doi.org/10.1371/journal.pone.0240792>

561 Miller, E., Sedlak, M., Lin, D., Box, C., Holleman, C., Rochman, C.M., Sutton, R., 2021. Recommended best  
562 practices for collecting, analyzing, and reporting microplastics in environmental media: Lessons learned from  
563 comprehensive monitoring of San Francisco Bay. J. Hazard. Mater. 409, 124770.  
564 <https://doi.org/10.1016/j.jhazmat.2020.124770>

565 Moore, C.J., Moore, S.L., Leecaster, M.K., Weisberg, S.B., 2001. A comparison of plastic and plankton in the  
566 North Pacific central Gyre. Mar. Pollut. Bull. 42, 1297-1300. [https://doi.org/10.1016/S0025-326X\(01\)00114-X](https://doi.org/10.1016/S0025-326X(01)00114-X)

567 Palazzo, L., Coppa, S., Camedda, A., Cocca, M., De Falco, F., Vianello, A., Massaro, G., de Lucia, G.A., 2021. A  
568 novel approach based on multiple fish species and water column compartments in assessing vertical microlitter  
569 distribution and composition. Environ. Pollut. 272, 116419. <https://doi.org/10.1016/j.envpol.2020.116419>

570 Palmisano, A.C., Pettigrew, C.A., 1992. Biodegradability of Plastics. BioScience 42(9), 680-685.  
571 <https://doi.org/10.2307/1312174>

572 Pelamatti, T., Cardelli, L.R., Rios-Mendoza, L.M., 2021. The Role of Microplastics in Bioaccumulation of  
573 Pollutants. In: Rocha-Santos, T., Costa, M., Mouneyrac, C., (Eds) Handbook of Microplastics in the Environment.  
574 Springer, Cham. <https://doi.org/10.1007/978-3-030-10618-8>

575 Prinz, N., Korez, Š., 2020. Understanding How Microplastics Affect Marine Biota on the Cellular Level Is  
576 Important for Assessing Ecosystem Function: A Review. Pp. 101-120 in: Jungblut S, Liebich V, Bode-Dalby M  
577 (Eds) YOUMARES 9 The Oceans: Our Research, Our Future Proceedings of the 2018 conference for YOUNg  
578 MARine REsearcher in Oldenburg, Germany. Springer Open  
579 <https://library.oapen.org/bitstream/handle/20.500.12657/22870/1007291.pdf>

580 R Core Team, 2021. R: A language and environment for statistical computing. R Foundation for Statistical  
581 Computing, Vienna, Austria. <https://www.R-project.org/>

582 Rai, P.K., Sonne, C., Brown, R.J.C., Younis, S.A., Kim, K.-H., 2022. Adsorption of environmental contaminants  
583 on micro- and nano-scale plastic polymers and the influence of weathering processes on their adsorptive attributes.  
584 J. Haz. Mat. 427, 127903. <https://doi.org/10.1016/j.jhazmat.2021.127903>

585 Rios-Fuster, B., Arechavala-Lopez, P., García-Marcos, K., Alomar, C., Compá, M., Álvarez, E., Julià, M.M.,  
586 Martí, A.S., Sureda, A., Deudero, S., 2021. Experimental evidence of physiological and behavioral effects of  
587 microplastic ingestion in *Sparus aurata*. Aquat. Toxicol. 231, 105737.  
588 <https://doi.org/10.1016/j.aquatox.2020.105737>

589 Savoca, M.S., McInturf, A.G., Hazen, E.L., 2021. Plastic ingestion by marine fish is widespread and increasing.  
590 Glob. Change Biol. 27(10), 2188-2199. <https://doi.org/10.1111/gcb.15533>

591 Sbrana, A., Valente, T., Scacco, U., Bianchi, J., Silvestri, C., Palazzo, L., de Lucia, G.A., Valerani, C., Ardizzone,  
592 G., Matiddi, M., 2020. Spatial variability and influence of biological parameters on microplastic ingestion by  
593 *Boops boops* (L.) along the Italian coasts (Western Mediterranean Sea). Environ. Pollut. 263, 114429.  
594 <https://doi.org/10.1016/j.envpol.2020.114429>

595 Sharma, S., Chatterjee, S., 2017. Microplastic pollution, a threat to marine ecosystems and human health: A short  
596 review. Environ. Sci. Pollut. R. 24, 21530-21547. <https://doi.org/10.1007/s11356-017-9910-8>

597 Thushari, G.G.N., Senevirathna, J.D.M., 2020. Plastic pollution in the marine environment. Heliyon 6(8), e04709.  
598 <https://doi.org/10.1016/j.heliyon.2020.e04709>

599 Tsangaris, C., Digka, N., Valente, T., Aguilar, A., Borrell, A., de Lucia, G.A., Gambaiani, D., Garcia-Garin, O.,  
600 Kaberi, H., Martin, J., Mauriño, E., Miaud, C., Palazzo, L., Pérez-del-Olmo, A., Raga, J.A., Sbrana, A., Silvestri,  
601 C., Skylaki, E., Vighi, M., Wongdontree, P., Matiddi, M., 2020. Using *Boops boops* (Osteichthyes) to assess



602 microplastic ingestion in the Mediterranean Sea. Mar. Pollut. Bull. 158, 111397.  
603 <https://doi.org/10.1016/j.marpolbul.2020.111397>

604 Turner, A., Arnold, R., Williams, T., 2020. Weathering and persistence of plastic in the marine environment:  
605 Lessons from LEGO. Environ. Pollut. 262, 114299. <https://doi.org/10.1016/j.envpol.2020.114299>

606 UNEP/MAP, 2012. State of the Mediterranean Marine and Coastal Environment. United Nations Environment  
607 Program - Mediterranean Action Plan, Barcelona Convention, Athens, 2012.

608 Valente, T., Sbrana, A., Scacco, U., Jacomini, C., Bianchi, J., Palazzo, L., de Lucia, G.A., Silvestri, C., Matiddi,  
609 M., 2019. Exploring microplastic ingestion by three deep-water elasmobranch species: A case study from the  
610 Tyrrhenian Sea. Environ. Pollut. 253, 342-350. <https://doi.org/10.1016/j.envpol.2019.07.001>

611 van Franeker, J.A., Blaize, C., Danielsen, J., Fairclough, K., Gollan, J., Guse, N., Hansen, P.-L., Heubeck, M.,  
612 Jensen, J.-K., Le Guillou, G., Olsen, B., Olsen, K.-O., Pedersen, J., Stienen, E.W.M., Turner, D.M., 2011.  
613 Monitoring plastic ingestion by the northern fulmar *Fulmarus glacialis* in the North Sea. Environ. Pollut. 159,  
614 2609-2615. <https://doi.org/10.1016/j.envpol.2011.06.008>

615 von Moos, N., Burkhardt-Holm, P., Köhler, A., 2012. Uptake and effects of microplastics on cells and tissue of  
616 the blue mussel *Mytilus edulis* L. after an experimental exposure. Environ. Sci. Technol. 46(20), 11327-11335.  
617 <https://doi.org/10.1021/es302332w>

618 Wang, W., Ge, J., Yu, X., 2020. Bioavailability and toxicity of microplastics to fish species: A review. Ecotox.  
619 Environ. Safe. 189, 109913. <https://doi.org/10.1016/j.ecoenv.2019.109913>

620 Warnes, G.R., Bolker, B., Bonebakker, L., Gentleman, R., Huber, W., Liaw, A., Lumley, T., Maechler, M.,  
621 Magnusson, A., Moeller, S., Schwartz, M., Venables, B., 2020. gplots: Various R Programming Tools for Plotting  
622 Data. R package version 3.1.1. <https://CRAN.R-project.org/package=gplots>

623 Wickham, H., 2016. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag, New York, 2016.

624 Woodall, L.C., Sanchez-Vidal, A., Canals, M., Paterson, G.L.J., Coppock, R., Sleight, V., Calafat, A., Rogers,  
625 A.D., Narayanaswamy, B.E., Thompson, R.C., 2014. The deep sea is a major sink for microplastic debris. R. Soc.  
626 Open Sci. 1, 140317. <https://doi.org/10.1098/rsos.140317>

627 Wootton, N., Reis-Santos, P., Gillanders, B.M., 2021. Microplastic in fish – A global synthesis. Rev. Fish Biol.  
628 Fisheries 31, 753-771. <https://doi.org/10.1007/s11160-021-09684-6>

629 Worm, B., Lotze, H.K., Jubinville, I., Wilcox, C., Jambeck, J., 2017. Plastic as a persistent marine pollutant. Annu.  
630 Rev. Env. Resour. 42, 1-26. <https://doi.org/10.1146/annurev-environ-102016-060700>

631 Wright, S.L., Rowe, D., Thompson, R.C., Galloway, T.S., 2013. Microplastic ingestion decreases energy reserves  
632 in marine worms. Curr. Biol. 23, R1031-1033. <https://doi.org/10.1016/j.cub.2013.10.068>

633 Zalasiewicz, J., Waters, C.N., Ivar do Sul, J.A., Corcoran, P.L., Barnosky, A.D., Cearreta, A., Edgeworth, M.,  
634 Galuskza, A., Jeandel, C., Leinfelder, R., McNeill, J.R., Steffen, W., Summerhayes, C., Waprich, M., Williams,  
635 M., Wolfe, A.P., Yonan, Y., 2016. The geological cycle of plastics and their use as a stratigraphic indicator of the  
636 Anthropocene. Anthr 13, 4-17. <https://doi.org/10.1016/j.ancene.2016.01.002>