



UNIVERSITÀ POLITECNICA DELLE MARCHE  
Repository ISTITUZIONALE

Microcirculation as a guide for therapy: do not condemn an innocent without a fair trial

This is the peer reviewed version of the following article:

*Original*

Microcirculation as a guide for therapy: do not condemn an innocent without a fair trial / Damiani, E., Scorcella, C., Carsetti, A., Donati, A., Adrario, E.. - In: INTENSIVE CARE MEDICINE. - ISSN 0342-4642. - 49:10(2023), pp. 1270-1271. [10.1007/s00134-023-07192-8]

*Availability:*

This version is available at: 11566/325655 since: 2024-11-18T10:08:10Z

*Publisher:*

*Published*

DOI:10.1007/s00134-023-07192-8

*Terms of use:*

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. The use of copyrighted works requires the consent of the rights' holder (author or publisher). Works made available under a Creative Commons license or a Publisher's custom-made license can be used according to the terms and conditions contained therein. See editor's website for further information and terms and conditions.

This item was downloaded from IRIS Università Politecnica delle Marche (<https://iris.univpm.it>). When citing, please refer to the published version.

*Publisher copyright:*

Springer (article) - Postprint/Author's accepted Manuscript

This version of the article has been accepted for publication, after peer review (when applicable) and is subject to Springer Nature's AM terms of use <https://www.springernature.com/gp/open-research/policies/accepted-manuscript-terms>, but is not the Version of Record and does not reflect post-acceptance improvements, or any corrections. The Version of Record is available online at: 10.1007/s00134-023-07192-8.

(Article begins on next page)

## CORRESPONDENCE

### **Microcirculation as a guide for therapy: do not condemn an innocent without a fair trial**

Elisa Damiani<sup>1,2\*</sup>, Claudia Scorcella<sup>2</sup>, Andrea Carsetti<sup>1,2</sup>, Abele Donati<sup>1,2</sup> and Erica Adrario<sup>1</sup>

We read with interest the article by Bruno et al. [1] in a recent issue of Intensive Care Medicine. We wish to congratulate the authors for performing this study, which is the largest randomized-controlled trial investigating the impact of integrating microcirculatory monitoring in the management of patients with shock. Disappointingly, this study showed no reduction in 30-day mortality. However, several considerations need to be taken into account for a proper interpretation of the results.

First, microvascular monitoring could improve survival only if associated with an effective treatment targeting tissue perfusion. In this study, a treatment protocol with a clear strategy for microvascular resuscitation in the different types of shock was missing. Recommendations were made on fluid and vasopressor dosing according to the microcirculation. However, physicians were free to deviate from the protocol. Indeed, a mismatch between the announced and performed treatment changes was found in 41 out of 69 patients in the interventional group (59.4%) [1]. The reason for not including inotropes/vasodilators is also unclear since >50% of the inclusions were patients with cardiogenic shock [2].

Second, the use of 30-day mortality as the primary outcome appears inappropriate as it may be biased by several confounders. Changes in organ function or parameters of tissue oxygenation/perfusion would be more adequate.

In fact, microvascular parameters were similar after 24 h in the two study arms: this is in line with the lack of difference in survival. Moreover, life-sustaining therapy was limited in almost half of the patients on day two [1]: this raises questions about the appropriateness of the patient selection criteria. In addition, the use of 19% difference in mortality for sample size calculation appears to be arbitrary and represents an overestimation of the expected effect. Third, the inclusion of different types of

shock is questionable. Microcirculatory impairment has different triggers: pump failure in cardiogenic shock and myocardial stunning after cardiac surgery, vasoplegia, and microcirculatory shunting in septic shock or after cardio-pulmonary bypass. The microvascular response to treatments, such as fluids or vasopressors, may be different as well [3]. Fourth, video quality was unacceptable in 28% of videos. This is not a negligible percentage, and low-quality videos may produce spurious microcirculatory data [4]. Video-quality was assessed a posteriori. For a reliable microcirculatory assessment, low-quality videos should be excluded from the analysis [5], especially if using an automated software that cannot distinguish pressure artifacts from real flow alterations. Importantly, the AVA 4.3C software has not been validated against manual offline analysis. Therefore, the reliability of the microcirculatory data collected remains uncertain. In conclusion, we agree with the authors when they say: “[...] the results are probably more a sign for the failure of the selected interventions than the failure of microcirculatory monitoring”. This study sheds light on the fact that we lack well-established effective treatment algorithms for the optimization of microvascular perfusion. Future efforts should be directed to identify effective resuscitation protocols based on the integration of microcirculatory monitoring with the standard hemodynamic parameters.

#### **Author details:**

1 Department of Biomedical Sciences and Public Health, Università Politecnica delle Marche, via Tronto 10/a, 60126 Ancona, Italy.

2 Clinica di Anestesia e Rianimazione Generale, Respiratoria e del Trauma Maggiore, Azienda Ospedaliero Universitaria delle Marche, Ancona, Italy. Declarations Conflicts of interest None.

#### **References:**

1. Bruno RR, Wollborn J, Fengler K et al (2023) Direct assessment of microcirculation in shock: a randomized-controlled multicenter study. *Intensive Care Med.* <https://doi.org/10.1007/s00134-023-07098-5>
2. Merdji H, Levy B, Jung C, Ince C, Siegemund M, Meziani F (2023) Microcirculatory dysfunction in cardiogenic shock. *Ann Intensive Care* 13:38
3. Guven G, Hilthy MP, Ince C (2020) Microcirculation: physiology, pathophysiology and clinical application. *Blood Purif* 49:143–150

4. Damiani E, Ince C, Scorcella C, Domizi R, Carsetti A, Mininno N, Pierantozzi S, Adrario E, Romano R, Pelaia P, Donati A (2017) Impact of microcirculatory video quality on the evaluation of sublingual microcirculation in critically ill patients. *J Clin Monit Comput* 31:981–988
5. Ince C, Boerma EC, Cecconi M et al (2018) Second consensus on the assessment of sublingual microcirculation in critically ill patients: results from a task force of the European society of intensive care medicine. *Intensive Care Med* 44:281–299