



UNIVERSITÀ POLITECNICA DELLE MARCHE
Repository ISTITUZIONALE

Modulatory effect of Andean blackberry polyphenols on genes related to antioxidant and inflammatory responses, the NLRP3 inflammasome, and autophagy

This is the peer reviewed version of the following article:

Original

Modulatory effect of Andean blackberry polyphenols on genes related to antioxidant and inflammatory responses, the NLRP3 inflammasome, and autophagy / Castejón-Vega, Beatriz; Kyriakidis, Nikolaos C.; Alcócer-Gómez, Elizabet; Giampieri, Francesca; González-Paramás, Ana M.; Cordero, Mario D.; Alvarez-Suarez, José M.. - In: JOURNAL OF BERRY RESEARCH. - ISSN 1878-5093. - 14:1(2024), pp. 41-59. [10.3233/jbr-230058]

Availability:

This version is available at: 11566/327467 since: 2024-03-07T15:58:13Z

Publisher:

Published

DOI:10.3233/jbr-230058

Terms of use:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. The use of copyrighted works requires the consent of the rights' holder (author or publisher). Works made available under a Creative Commons license or a Publisher's custom-made license can be used according to the terms and conditions contained therein. See editor's website for further information and terms and conditions.

This item was downloaded from IRIS Università Politecnica delle Marche (<https://iris.univpm.it>). When citing, please refer to the published version.

(Article begins on next page)

Modulatory effect of Andean blackberry polyphenols on genes related to antioxidant and inflammatory responses, the NLRP3 inflammasome and autophagy

Beatriz Castejón-Vega^a, Nikolaos C. Kyriakidis^b, Elizabet Alcócer-Gómez^c, Francesca Giampieri^{d,e}, Ana M. González-Paramás^f, Mario D. Cordero^{g, h*}, and José M. Alvarez-Suarez^{i,j*}

^a Institute of Molecular, Cell and Systems Biology, University of Glasgow, Glasgow G12 8QQ, United Kingdom

^b Cancer Research Group, Faculty of Medicine, Universidad de Las Américas, Quito, Ecuador

^c Departamento de Psicología Experimental, Facultad de Psicología, Universidad de Sevilla, Seville, 41009, Spain

^d Research Group on Food, Nutritional Biochemistry and Health, Universidad Europea del Atlántico, 39011 Santander, Cantabria, Spain

^e Department of Clinical Sciences, Polytechnic University of Marche, Ancona, 60131, Italy

^f Grupo de Investigación en Polifenoles, Universidad de Salamanca, Campus Miguel de Unamuno, Salamanca, 37008, Spain

^g Instituto de Investigación e Innovación en Ciencias Biomédicas de Cádiz (INiBICA), Cadiz, 11009, Spain.

^h CIBER de Enfermedades Respiratorias (CIBERES), Instituto de Salud Carlos III, Madrid, 28220, Spain

ⁱ Laboratorio de Bioexploración. Universidad San Francisco de Quito USFQ, Quito, 170901, Ecuador

^j Laboratorio de Investigación de Ingeniería en Alimentos (LabInAli). Colegio de Ciencias e Ingeniería, Universidad San Francisco de Quito USFQ, Quito, 170901, Ecuador

Corresponding authords: José M. Alvarez-Suarez (jalvarez@usfq.edu.ec) and Mario D. Cordero (mario.cordero@inibica.es)

Abstract

BACKGROUND: The Andean blackberry (*Rubus glaucus* Benth) is one of Ecuador's iconic berries that grows in its Andean region, of which a high anthocyanin content has been described.

OBJECTIVE: The aim of the present study was to determine the chemical composition and anti-inflammatory potential of the Andean blackberry from Ecuador, with emphasis on its effects on NLRP3 inflammasome activation and autophagy processes. **RESULTS:** Andean blackberry extracts were rich in hydroxycinnamates (coumaric acid and derivatives), in addition to quercetin and kaempferol as main flavonols. Cyanidin and its glycosides were identified as the main anthocyanins present. Andean blackberry extracts efficiently reduced oxidative stress markers in the lipopolysaccharide-stimulated RAW 264.7 cells. The extract also caused a moderate decrease in the expression of the pro-inflammatory and antioxidant genes NFκB1, TNF, IL-1, IL-6 and NOS2 expression, while it significantly increased the mRNA levels of both SOD1 and NFE2L2 genes. Andean blackberry extracts significantly decreased the activation of the NLRP3 inflammasome complex, as well as p62 levels and the LC3I/LC3II ratio increased, suggesting a direct action of Andean blackberry compounds on the inflammatory response and restoration of the autophagy process. **CONCLUSION:** These results suggest the potential anti-inflammatory effect of Andean blackberry, through their ability to regulate genes related to the inflammatory and antioxidant response, as well as modulate the activation of the NLRP3 inflammasome complex and autophagy processes.

Keywords: Anthocyanins, Autophagy, Blackberries, Inflammation, NLRP3 inflammasome

1. Introduction

Current epidemiological evidence increasingly highlights the relationship between nutritional patterns based on frequent consumption of fruits and vegetables with a low incidence of noncommunicable diseases, such as diabetes, cardiovascular and neurodegenerative diseases and cancer [1]. All these pathologies have a common pathophysiological feature in their development and progression based on high production of pro-inflammatory mediators and are characterized by excessive production of reactive oxygen species (ROS) with the consequent oxidative damage on macromolecules and cellular structures [2]. Thus, the inhibition of the excessive inflammatory response and oxidative damage has been proposed as an alternative to minimize their damage. It is in this context that a diet rich in fruits and vegetables, with its important contribution of antioxidants and anti-inflammatory compounds, plays a fundamental role based mainly on the ability of these compounds not only to block the excessive circulating free radicals but also on their biological effects that go beyond the simple antioxidant character [3]. Indeed, in the last decades an increasing body of articles have described the anti-inflammatory and immunoregulatory properties that polyphenols have on different pathways related to the antioxidant response and, therefore, their contribution to protection against oxidative damage [4,5]. Within the polyphenols described with anti-inflammatory activity, anthocyanins (ACNs) play a prominent role. Anthocyanins are natural phenolic pigments with broad and outstanding biological activity. They have potent antioxidant and anti-inflammatory activity, properties that have been closely related to other of their biological effects such as their antidiabetic and anticancer activity, and their role in cardiovascular and neuroprotective prevention [6,7]. Anthocyanins are widespread in the plant world, providing the characteristic red/blue colour to various fruits, vegetables, flowers and leaves, and their content is influenced by various factors, such as species, cultivars, climatic conditions and level of maturity [8], and others [9]. Among its natural sources, berries stand out (strawberries, blueberries, blackberries, blackcurrant, redcurrant,

and raspberries) for being a rich source of anthocyanins, with levels ranging from about 100 to about 700 mg/100 g of fresh fruits [9].

The Andean blackberry (*Rubus glaucus* Benth) is one of Ecuador's iconic berries that grows wild or cultivated, isolated, dispersed or in groups with other plants in the Andean highlands of Ecuador. They are traditionally collected and commercialized by the indigenous people of these regions [10] and they have become a popular fruit among the Ecuadorian population in general consumed in the form of jams, wine, and boiled drinks such as the famous "*Colada morada*" (purple strained) that is consumed in the day of the dead celebration [11]. Although several studies have highlighted the chemical composition and biological effects of blackberries from other geographical regions [12–15], few studies have been developed in the case of the Andean blackberry. Only a few investigations have described its chemical composition, evidencing its high anthocyanin content [16–18], as well as its biological properties, such as its ability to protect against oxidative damage [19]. Thus, more studies are necessary to define its chemical composition, as well as the biological effects and mechanisms that it uses to generate them, to gradually determine its true health potential and therefore justifying its consumption. With this background, in the present study we firstly determined the polyphenolic composition of the Andean blackberry fruit. In order to assess its anti-inflammatory effects, a murine macrophage RAW 264.7 cells were used as an *in vitro* validation model both in the presence or absence of the endotoxin lipopolysaccharide (LPS). The levels of the principal biomarkers related to oxidative damage and inflammatory markers were estimated together with the activity of the main antioxidant enzymes. Finally, the expression of different genes related to antioxidant and inflammatory response was also determined with the objective of discovering the possible molecular pathways involved in the anti-inflammatory action of Andean blackberry. Finally, given the importance of the NLRP3 inflammasome complex as an important mediator of inflammatory processes, we studied the effects of Andean blackberry compounds on its activation by using an *in vitro* model of THP-1 human monocytes cells. At the same time, given the repairing effect of

the autophagy process and therefore its importance in cellular hemostasis, we also studied the effect of the polyphenols of this fruit on the activation of this process using this cellular model.

2. Material and methods

2.1. Chemical and reagents

All chemicals and solvents were of analytical grade. Solvents, lipopolysaccharides (LPS) from *Escherichia coli* O55:B5, RPMI and 1640 Medium, fetal bovine serum, penicillin-streptomycin, RIPA Buffer, hydrogen peroxide solution (H₂O₂), reduced glutathione (GSH), β -Nicotinamide adenine dinucleotide 2'-phosphate reduced tetrasodium salt hydrate (NADPH), glutathione peroxidase from bovine erythrocytes (GPx), L-glutathione reduced (GSH), glutathione S-transferase from equine liver (GST), 2-Thiobarbituric acid, malondialdehyde tetrabutylammonium salt (MDA), 2,4-Dinitrophenylhydrazine (DNPH), guanidine hydrochloride, streptomycin, Tris-HCl, Triton[™] X-100, phenylmethylsulfonyl fluoride, leupeptin and Bradford reagent were purchased from Sigma-Aldrich Co. (St. Louis, MO, USA), while the 2'-7'-dichlorofluorescein diacetate (DCFH) kit was purchased from Merck, Darmstadt, Germany. TRIzol[™] reagent, Turbo[™] DNase and SuperScript III Platinum SYBR Green One-Step RT-qPCR kit were purchased from Invitrogen Carlsbad, CA, USA) while specific monoclonal antibodies were obtained from Santa Cruz Biotechnology (Dallas, TX, USA). A cocktail of protease inhibitors (cOmplete Protease Inhibitor Cocktail) was purchased from Boehringer Mannheim (Indianapolis, IN). The Immun-Star horseradish peroxidase (HRP) substrate kit was obtained from Bio-Rad Laboratories Inc. (Hercules, CA, USA).

2.2. Plant materials and sample preparation

Ripe Andean blackberries (*Rubus glaucus* Benth) fruits were harvested during the berries crop season (October – December, 2018) from plants growing in the highlands of the province of Tungurahua, located in the inter-Andean region of Ecuador at an altitude between 3000 and 3200 meters above sea level. Fruits were collected on two different occasions at random from different plants based on a similar degree of ripeness and without blemishes or any signs of damage or

decay. Two batches of samples (~1kg) were collected on each occasion and the specimens were identified using the reference vouchers for specimens deposited in the herbarium of the Jardín Botánico de Quito, Ecuador. During collection time, the fruits were kept at 4°C protected from light and transported to the laboratory the same day they were harvested. Afterwards, samples were cleaned, removing remains of leaves, stems and other residues, washed with distilled water and allowed to dry on absorbent paper at room temperature and protected from light. Finally, fruits were frozen at -20 °C overnight and then freeze-dried, ground to a fine powder, and stored at -20°C until analysis.

For chemical composition determination and *in vitro* assays, a polyphenol-rich extract was obtained as previously reported [20]. Briefly, the fine powder of fruits (10g) was added to 100 mL of the extraction solution consisting of methanol acidified with 0.1% HCl and Milli-Q water (80%, v/v) and stirred overnight at 3–5°C. Then, the solution was centrifuged for 15 minutes at 4000g at 4°C and filtered through a 2.5 µm pore size Whatman® cellulose filter paper (Sigma-Aldrich Co., St. Louis, MO, USA). The solid residues were washed with the extraction solution as many times as necessary until no red colour was observed in the extracted solution. Finally, the resulting extraction solutions were combined, water was added, and the final solution was concentrated in a vacuum rotary evaporator (~30°C) until obtaining a dry crude extract.

2.3. HPLC-DAD/ESI-MSⁿ characterization of phenolic acids, flavonols and anthocyanins

The Andean blackberry crude extract (ABCext) was dissolved in 0.5 mL of 0.1% trifluoroacetic acid/acetonitrile (70:30, v/v), filtered through a 0.22 µm disposable LC filter disk for analysis by HPLC and analyzed using double online detection by diode array spectrophotometry and mass spectrometry (MS). The HPLC system consisted in a Hewlett-Packard 1200 chromatograph (Hewlett-Packard 1200, Agilent Technologies, Santa Clara, CA, USA) equipped with a binary pump and a diode array detector (DAD) coupled with an HP ChemStation (rev. A.05.04) data processing station coupled with an MS detector API 3200 Qtrap (Applied Biosystems, Darmstadt, Germany) that was controlled by the Analyst 5.1 software. As stationary phase, an AQUA®

(Phenomenex, Torrance, CA, USA) reverse phase C18 column (5 μ m, 150 mm \times 4.6 mm) thermostated at 35°C was used. The solvents were: (A) 0.1% trifluoroacetic acid and (B) acetonitrile. The elution gradient established was: isocratic 10% B for 3 min, 10–15% B in 12 min, isocratic 15% B for 5 min, 15–18% B over 5 min, 18–30% B over 20 min, 30–35% B over 5 min, and re-equilibration of the column to initial solvent conditions. The flow rate used was 0.5 mL/min. For detection, the DAD was set using 520, 330 and 280 nm as preferred wavelengths, while the MS operated in the positive ion mode for the analysis of anthocyanins and in the negative mode for the rest of the compounds, the spectra were recorded between m/z 100-1000. Zero grade air served as the nebulizer gas (30 psi) and as turbo gas (400°C) for solvent drying (40 psi). Nitrogen served as the curtain (20 psi) and collision gas (medium). Both quadrupoles were set at unit resolution and EMS and EPI analyses were also performed. The EMS parameters were set as follows: ion spray voltage –4500V, DP -50V, EP -6V, CE -10V and cell exit potential (CXP) –3V, whereas EPI settings were: DP -50V, EP -6V, CE -30V and CES 10V. The individual phenolic compounds were tentatively identified based on their UV and mass spectra, and comparison with data reported in the literature, as well as with internal standards when available.

2.4. Evaluation of anti-inflammatory properties

2.4.1. Cell culture and treatments

Murine RAW 264.7 macrophage cell line was purchased from ATCC (ATCC-TIB71) (Manassas, VA, USA) and maintained in RPMI 1640 medium supplemented with 10% fetal bovine serum and 1% penicillin-streptomycin antibiotics (100 IU/mL penicillin and 100 mg/mL streptomycin) in a humidified atmosphere at 37°C with 5% CO₂. For all the assays cells were used between the 4th and the 6th passage. The Andean blackberry crude extract (ABCext) was resuspended in RPMI 1640 medium to achieve the final concentration of 80 μ g/mL. RAW macrophages were treated with: (i) RPMI 1640 medium only (Ctrl group), (ii) lipopolysaccharides (LPS) (*Escherichia coli* serotype 055:B5) at 1 μ g/mL for 24h (LPS group), (iii) ABCext (80 μ g/mL) for 24h (ABCext group), or (iv) LPS (1 μ g/mL) for 24h and then ABCext (80 μ g/mL) for

another 24h (LPS/ABCext group), (v) ABCext (80 µg/mL) for 24h and then LPS (1 µg/mL) for another 24h (ABCext/LPS group) and (vi) ABCext (80 µg/mL) + LPS (1 µg/mL) for 24h (ABCext+LPS group). The appropriate dose/time combination of ABCext and LPS treatments was selected according to our previously reported studies using the same type of extract [19] as well as the same cell model according to the culture conditions in our laboratory [21].

2.4.2. Measurement of intracellular ROS level

Intracellular reactive oxygen species (ROS) levels were determined using the 2'-7'-dichlorofluorescein diacetate (DCFH) kit according to the manufacturer's instructions. RAW cells previously treated under the same/above-mentioned conditions were incubated with 5 µmol/L of DCFH at 37°C for 30 min in the dark at room temperature. Next, the fluorescence intensity was read using a microplate reader (Thermo Scientific Microplate Reader, Multiskan®, Winooski, VT, USA) at 485/530 nm excitation/emission wavelength. Total protein levels were determined by the Bradford method [22] and results were expressed as arbitrary units of fluorescent intensity/µg cell proteins.

2.4.3. Determination of antioxidant enzyme activities, biomarkers of oxidative damage in lipid and proteins and nitrite production

After treatment, murine RAW cells were washed with PBS, incubated with RIPA buffer on ice for 5 min and the obtained lysate was stored at -80°C until analysis. For the analysis of antioxidant enzymes, superoxide dismutase (SOD) was analyzed based on the inhibition of the formation of NADHphenazine methosulphate-nitroblue tetrazolium (NBT) formazon by SOD [23]. Catalase (CAT) was analyzed following the decomposition of hydrogen peroxide [24]. Glutathione peroxidase (GPx) activity was determined according to the ability of GPx to remove H₂O₂ by coupling its reduction to H₂O with oxidation of reduced glutathione in the presence of NADPH [25]. Glutathione reductase (GR) activity was determined based on the capacity of GR to reduce oxidized glutathione back to reduced glutathione (GSH), in presence of NADPH [26], while glutathione transferase (GST) activity was determined by measuring the content of dinitrophenyl

thioether produced by the conjugation of 1-chloro-2,4-dinitrobenzene (CDNB) with reduced glutathione [27]. SOD and CAT activity were expressed as U/mg prot/min, GPx and GR activity were expressed as nmole NADPH oxidized/mg prot/min, while GST activity was expressed as nmol 1-chloro-2,4-dinitro benzene (CDNB)-reduced glutathione (GSH) conjugate/mg protein/min. Protein carbonyl content [28] and thiobarbituric acid-reactive substance (TBARS) [29] were used as markers of protein and lipid oxidative damage, respectively and the results were expressed as nmol/mg protein for total protein carbonyl content, and nmol/100 mg protein for TBARS levels. Nitrite oxide (NO) accumulation in cell culture media was determined by the Griess reagent system following the manufacturer's protocol and NO content was expressed as μM of $\text{NaNO}_2/10^5$ cells.

2.4.4. RT-qPCR analysis of related genes to antioxidant and inflammatory responses

RNA from RAW cells was extracted using the TRIzol™ reagent followed by DNase I treatment to eliminate genomic DNA interference (Turbo™ DNase kit) according to the manufacturer's instructions. RNA concentrations and purity were determined using a NanoDrop 2000 Spectrophotometer (Thermo Scientific, Waltham, MA, USA). Reverse transcription and subsequent quantitative real-time PCR were performed on a CFX96 Real-Time PCR Systems (Bio-Rad Laboratories Inc., Hercules, CA, USA) using the SuperScript III Platinum SYBR Green One-Step RT-qPCR kit and 100 ng per RNA sample. cDNA synthesis was carried out for 3 minutes at 50°C followed by an initial denaturation step at 95°C for 5 minutes. The cycling conditions comprised an initial denaturation step at 95°C for 5 min, followed by 40 cycles of: denaturation at 95°C for 30 s, primer annealing at 65°C for 30s and extension at 72°C for 30s, followed by a final elongation step at 40°C for 1 min. For gene expression analysis primers for 10 different genes associated with pro-inflammatory, anti-inflammatory, and oxidative stress responses designed online with the NCBI Primer-BLAST tool were used (Table 1). Beta actin (ACTB) was used as a housekeeping gene. Finally, a melting curve analysis, whereby the qPCR products were run from

65°C to 95°C and an increment of 0.5°C for 0.05s per step, was used to verify specific product formation. Relative gene expression analysis was calculated using the $\Delta\Delta C_t$ method [30].

2.5. Assessment of inflammasome and autophagy activation

2.5.1. Cell culture and treatments

THP-1 cells, a monocyte isolated from peripheral blood from an acute monocytic leukemia patient, were purchased from ATCC (ATCC® TIB-202™) (Manassas, VA, USA) and cultured in RPMI media with 10% fetal serum and penicillin (100 U/ml)/streptomycin (100 µg/ml) at 37°C under an atmosphere of 5% CO₂. To determine the effects of the Andean blackberry crude extract (ABCext) on the LPS + ATP-induced inflammasome, the cells were divided into 3 groups: (i) RPMI 1640 medium only (Ctrl group), (ii) LPS (*Escherichia coli* serotype 055:B5) at 500 ng/mL LPS for 4h followed by stimulation with ATP (5 mM) for 6h (LPS+ATP group) and LPS (500 ng/mL) for 4h followed by ATP (5 mM) and ABCext (80 µg/mL) for 6h (LPS+ATP+ABCext group). The appropriate dose/time combination of LPS and ATP treatments for inflammasome activation was previously established by researchers at the Research Laboratory, Oral Medicine Department, University of Seville, Seville, Spain [31]. Moreover, ABCext was established according to preliminary dosage assays for protein expression induction (data not shown).

2.5.2. Western blotting

Whole cellular lysate from THP-1 cells was prepared by gentle shaking with a buffer containing 0.9% NaCl, 20 mM Tris-HCl, pH 7.6, 0.1% triton X-100, 1 mM phenylmethylsulfonylfluoride and 0.01% leupeptin. The protein content was determined by the Bradford method [22]. Electrophoresis was carried out in a 10-15 % acrylamide SDS/PAGE and proteins were transferred to Immobilon membranes (Amersham Pharmacia, Piscataway, NJ, USA). Next, membranes were washed with PBS, blocked over night at 4°C and incubated with the primary antibody solution (1:1000) specific for the detection of the following proteins or multiproteic complexes: NLRP3 inflammasome, caspase 1, caspase 1 p20 unit (p20), interleukin 1 beta (IL-1β), mature form of IL-1β (p17), the autophagy markers microtubule-associated protein light chain 3 (LC3I and LC3III)

and p62. Membranes were then probed with their respective secondary antibody (1:2500). Immunolabeled proteins were detected by chemiluminescence method (Immun Star HRP substrate kit, Bio-Rad Laboratories Inc., Hercules, CA, USA). Western blot images were quantified using ImageJ software (see: <http://rsb.info.nih.gov/ij/download.html>).

2.6. Statistical analysis

Statistical analysis was carried out with SPSS v.24 software and the Jamovi built of R statistical language [32–34]. Median \pm standard deviation (SD) of three independent experiments are presented. Correlations between the different stimulation groups were processed using one-way ANOVA analysis and the Tukey post hoc test with a $p < 0.05$ was considered statistically significant, with a $p < 0.01$ being considered as highly significant.

3. Results and Discussion

3.1. Phenolic composition of Andean blackberries fruit

In this study, a hydroalcoholic extract obtained from Andean blackberries fruit was analyzed for its polyphenol composition by HPLC-DAD-ESI/MSⁿ. Figure 1A shows representative HPLC chromatograms of the polyphenol profile obtained at different λ (330 and 360 nm for phenolic acid derivatives and flavonoids and 520 nm for anthocyanins), while Fig. 1B shows the specific profile for anthocyanins and Fig. 1C, the specific profile of phenolic acids and flavonoids. On the other hand, Table 2 compiles the tentative identity of the peaks, assigned based on their UV-Vis spectra, mass spectral data of their (pseudo)molecular ions and MS² fragmentation patterns. The detected phenolic acids corresponded mainly to hydroxycinnamoyl derivatives, whereas the majority of flavonoid peaks corresponded to the group of flavonols and anthocyanins. Hydroxycinnamates here identified (coumaric acid and ferulic acid derivatives) have been previously reported in berries [4,13], similarly to quercetin and kaempferol which have been reported as the main flavonols present in this fruit [13,18,35]. Among the anthocyanins reported in berries, cyanidin and its glycosides have been identified as the main form of this group of pigments [13,35] in fact, in our study cyanidin was identified predominantly in its 3-form glycosidic and only two minor peaks of

pelargonidin derivatives were detected. In this context, the chemical composition reported here result crucial to justify part of the biological effects reported here below, as well as those previously reported [35]. Hydroxycinnamates, for example, are among the most widely distributed phenylpropanoids present in the insoluble-bound, conjugated-soluble and free forms in plants and render relevant antioxidant activity by their capacity to scavenge different free radicals (FR) such as hydroxyl radical (HO^\bullet), superoxide radical anion ($\text{O}_2^{\bullet-}$), peroxy radical (ROO^\bullet), peroxynitrite (NO_3^-), singlet oxygen ($^1\text{O}_2$) and several organic radicals [36]. Coumaric acids have been reported as an important antioxidant component in different berries [37], while in addition to its antioxidant capacity, it has also been associated with antibacterial, anticancer, antihistaminic effects and an enhancement of ATP production [38]. Quercetin and kaempferol were also associated with significant antioxidant, anti-inflammatory and antitumor effects [7,39], similar to anthocyanins that have been widely studied for their powerful antioxidant and anti-inflammatory effects [7,35,40], antitumoral capacity [7,41], and protective effects against cardiovascular diseases [5,7,42].

3.2 Effects of Andean blackberry extract on radical levels, antioxidant enzyme and oxidative biomarkers in LPS-stimulated cells

During inflammation, different macrophage subsets have a dual behavior: on one side, macrophages displaying an M1 phenotype secrete pro-inflammatory cytokines and produce elevated amounts of reactive oxygen species (ROS) and nitric oxide (NO), enhancing the inflammatory status of the immune response. On the other hand, alternatively activated M2 macrophages are involved in the inflammation resolution and tissue reconstruction, being responsible for apoptotic and necrotic cell phagocytosis and secretion of anti-inflammatory, proangiogenic and profibrotic cytokines, such as interleukin 10 (IL-10), vascular endothelial growth factors (VEGF) and transforming growth factor β (TGF- β), respectively. The excessive production of ROS and NO has been correlated with a chronic inflammation typical of some diseases, including, for example, asthma, ulcerative colitis and rheumatoid arthritis [43].

Therefore, it is very important to investigate the role that these mediators have on these processes, together with the effects of natural extracts in improving inflammation. In line with the above, this work analyzed the effect of ABCext on the intracellular production of ROS and NO in RAW 264.7 macrophages, as well as on a group of oxidative stress markers related to oxidative stress derived from the inflammatory response (Table 3).

Regarding ROS, LPS significantly ($p<0.05$) increased the intracellular % of ROS up to 2.8 times compared to control cells, while the treatment with ABCext led to a slight decrease compared to the control; in the combined treatments, we found that ABCext significantly ($p<0.05$) decreased the intracellular ROS levels, in LPS/ABCext group, but especially in the ABCext/LPS group, compared to LPS group, highlighting the protective effects of blackberry extract before the LPS damage (Table 3). The similar trend was found for NO production (Table 3), with the highest values ($p<0.05$) found in LPS group, the lowest ($p<0.05$) in ABCext group and with the pretreatment with ABCext the most effective ($p<0.05$) compared to the other combined treatments. Our results agree with previous data, which highlighted the ability of the main blackberry compounds, such as anthocyanins, to decrease basal intracellular production of inflammatory-based ROS and NO as a predisposing factor for several noncommunicable chronic degenerative diseases [6,44–46]. In fact, through different pathways, anthocyanins possess the ability to act as reducing agents in the electron-transfer reaction pathway, donating electrons to the free radicals with unpaired electrons [47]. However, recent studies have shown that the biological activities of anthocyanin also extend to other implicated pathways such as improvement of mitochondrial function [48], immunomodulatory properties [49], regulation of lipid metabolism [50], as well as mitochondrial biogenesis and functionality, modulation of intracellular ROS production and regulation of antioxidant enzymes production involved in intracellular antioxidant defense [51].

Regarding the activities of the main antioxidant enzymes, in LPS treated cells, a significant ($p<0.05$) decrease in the activities of GPx, GR, GST, SOD and CAT was observed compared to the control group (Table 3), while in ABCext treated cells similar values with the control group

were measured. Interestingly, the pretreatment with ABCext (ABCext/LPS group) exerted a more protective effect against LPS damage, compared to the LPS/ABCext and ABCext+LPS. The effects of ABCext on ROS and NO levels, as well as on antioxidant enzyme activities, followed a similar behavior to protein and lipid damage (Table 3). The highest values ($p<0.05$) of the carbonyl group and TBARS were found in LPS group, while the lowest ($p<0.05$) were measured in ABCext group and in the ABCext/LPS group, confirming the efficacy of the ACNs-rich ABCext treatment in counteracting oxidative damage, as previously reported by our group in ANCs-rich extracts from different red fruits [21,40].

3.2. Effects of Andean blackberry extract on the expression of proinflammatory and antioxidant genes in LPS-stimulated RAW 264.7 cells

Given the direct effect of various natural compounds from, the same families as those reported here on gene regulation beyond their antioxidant power [35,49,50], here we investigate the effect of the ABCext on the expression levels of genes that codify for key proinflammatory and antioxidant mediators in LPS-stimulated murine macrophages. Among the genes analyzed, we observed some interesting changes in the expression profile of 7 of these important mediators (Fig. 2). More specifically, although the nuclear factor kappa B (NF κ B) expression was relatively low, not allowing for statistically significant differences between the different stimulation groups to be revealed, we observed that treating RAW cells with ABCext alone or with LPS and ABCext simultaneously shows a tendency to lower NF κ B expression compared with LPS-stimulated cells (Fig. 2A). In the same line, pretreating murine macrophages with ABCext before LPS stimulation appeared to confer a protective effect though lower NF κ B expression did not reach statistical significance compared to LPS-stimulated cells.

On the other hand, the analysis for tumor necrosis factor (TNF) expression showed statistically significant differences between the different stimulation groups ($p<0.05$) (Fig. 2B). Additionally, we observed that mean TNF expression in both RAW cells treated first with LPS and then ABCext and cells treated first with ABCext and then LPS was almost half of the LPS-treated cells, although

these values did not reach statistical significance, while stimulating RAW cells simultaneously with LPS and ABCext did not change *TNF* expression compared with LPS stimulation alone.

Interestingly, similar expression patterns were found between the other two proinflammatory cytokines, namely IL-1 β and IL-6 (Fig.2C and D, respectively). Indeed, for both genes codifying for these cytokines, treating cells with ABCext alone lowered expression levels compared with LPS-stimulated macrophages ($p<0.001$) for each gene, respectively) as well as with constitutive expression levels. However, treating cells simultaneously with LPS and ABCext, first with LPS and then with ABCext or first ABCext and then LPS did not manage to significantly lower the expression levels of either of these genes compared with the levels induced by LPS alone stimulation (Fig. 2).

Nitric oxide synthase 2 (NOS2) expression exhibited a similar pattern whereby ABCext-stimulated cells were found to significantly reduce NOS2 mRNA levels against LPS-treated cells ($p<0.05$) and even reach lower expression levels than in unstimulated cells (Fig. 2E). However, LPS appears to be a strong promotor of NOS2 activation as either of the treatments that involved LPS and ABCext were shown to induce similar NOS2 expression levels as the LPS alone condition.

Finally, NFE2L2 and SOD1 expression patterns were largely similar (Fig. 2F and G, respectively). Treating RAW cells with LPS was shown to lower expression levels of both genes, although the differences with basal expression levels did not reach statistical significance. However, treating cells with ABCext after LPS stimulation incremented the mRNA levels of both genes reaching statistical significance for both superoxide dismutase (SOD1) and nuclear factor erythroid 2-related factor 2 (NFE2L2) ($p<0.05$). Toll-like receptor (TLR) ligands like LPS and cytokines like IFNG drive M1 polarization. The outcome of this stimulation is a significant increment in the expression of proinflammatory cytokines, such as IL-1 β , IL-12, TNF, IL-6, chemokines like IL-8 (CXCL8), MHC-II molecules and production of nitrous oxide and ROS. The principal promoter of the aforementioned mediators is NF κ B, a pivotal transcription factor that

bridges TLR signalling and the expression of the whole armamentarium of molecules employed by classically activated macrophages [52] Indeed, LPS, a master activator of TLR4, induced overexpression of NFκB, TNF, IL-1β, IL-6, and NOS2, while slightly downregulating the expression of NFE2L2 that codifies for the NRF2 protein, a known regulator of anti-inflammatory, detoxifying and antioxidant responses [53].

Treatment with ABCext, whether before, after, or concomitantly with LPS, largely reversed the LPS-induced upregulation of NFκB suggesting an important protective role for ABCext in excessive inflammation, although generally low LPS-induced NFκB expression did not allow for statistical significance to be reached. Moreover, treating murine macrophages with ABCext before or after LPS stimulation seemed to partially mitigate TNF, but not IL-1β or IL-6 upregulation (Fig. 2C and D, respectively). This finding implies that the expression of these proinflammatory cytokines is either not exclusively NFκB mediated in RAW cells, as LPS-induced NFκB expression levels suggest, or that these cells are so sensitive to LPS stimulation that even lower concentrations of this ligand are needed to allow the full anti-inflammatory effect of ABCext to be revealed. Additionally, the induction of NFE2L2 and SOD mRNA expression by ABCext in RAW cells, especially if added after LPS stimulation, revealed a possible antioxidant role of this extract in an endeavour to minimize oxidative stress adverse effects.

Overall, gene expression analysis in stimulated RAW 264.7 cells strongly suggest an important anti-inflammatory and antioxidant role of ABCext as previously has been demonstrated in ANCs-rich extracts from different fruits [15,48–50,54]. Nonetheless, the anti-inflammatory and antioxidant effects of ABCext warrant further investigation in *in vivo* models and the exact components exerting these effects need to be identified.

3.3 Effects of Andean blackberry extract on NLRP3 inflammasome and autophagy process

Given the direct effect that molecular regulation has on the inflammatory response, and therefore on the deleterious effects that it can produce, considering the results presented here, it is imperative to know how the polyphenolic constituents of Andean blackberry can influence the regulation of the response. With this aim, we set out to determine whether ABCext administration can alleviate the inflammatory response by mediating regulation of the NLRP3 inflammasome complex. For this, we used an *in vitro* model of THP-1 cells. Since NLRP3 activation requires two signals, the priming, and the activation signal, we treated the cells with LPS and ATP, respectively [55]. The NLRP3 inflammasome is the best characterized NOD-type receptors (NLR) which plays an essential role in the inflammation response. It is a multiprotein complex, also known as caspase-1 activation platform, which once triggered by pathogens, stress, or danger-signals, leads to the activation of caspase-1(CASP1) and thus to the maturation and secretion of interleukin 1 beta (IL-1 β) and interleukin 18 (IL-18) which initiate the inflammatory signalling [56,57]. NLRP3 inflammasome plays a crucial role in the inflammation response maintaining the homeostasis by repairing tissues, eliminating pathogens, and sensing and adapting stress levels. However, dysregulation of its activation leads to a wide range of diseases such as metabolic, inflammatory, cardiovascular, neurological and cancer. In addition, the ageing process is commonly accompanied by low grade sterile chronic inflammation, being NLRP3 inflammasome activated and thus leading to the accumulation of pro-inflammatory mediators. This process is already known as inflaming [58,59]. For this reason, it is increasingly necessary to identify compounds, such as those present in food, that target the inflammasome to improve our general health and well-being.

The treatment with LPS+ATP in THP-1 cells (LPS+ATP group) significantly increased ($p<0.01$) the expression levels of NLRP3 proteins (1.35 times), activated CASP1 (p20) (1.24 times) and activated IL-1 β (p17) (0.65 times) compared to the control group (Ctrl group) (Fig. 3A), indicating that the NLRP3 inflammasome was triggered and therefore the inflammatory response was on. However, after treatment with ABCext (LPS+ATP+ABCext group), the levels of these proteins showed a significant decrease ($p<0.01$) (0.89 times for NLRP3 protein; 0.73 times

for activated caspase-1 (p20) and 0.95 times for activated IL-1 β (p17))—compared to the group treated with LPS and ATP (LPS+ATP group). These values are close to the basal levels, where, in the case of activated IL-1 β (p17), the levels decreased below the basal values of the control group (Ctrl group) (Fig. 3A). In addition, our results showed that there is a downward trend in NF κ B levels after treating the cells with ABCext (Section 3.2 of the Results and Discussion), which correlated with the observed decrease in NLRP3 inflammasome and IL-1 β levels after the treatment with ABCext, since NF κ B mediates their expression after the priming signal which in our case is the LPS addition. These results are promising since they suggest a possible modulatory effect of Andean blackberry polyphenols, more probably the anthocyanins, on the NLRP3 inflammasome complex, due to its ability to decrease its activation, as well as activated CASP1 and IL-1 β protein levels, evidencing its potential anti-inflammatory effects. In fact, several works have already reported the modulating effect of polyphenols, as anthocyanins on the NLRP3 complex [60]. It was recently reported that anthocyanin extract from Purple Sweet Potato was able to restrict NLRP3 inflammasome activation in alveolar macrophages infected with *Klebsiella pneumoniae* [61]. On the other hand, the anthocyanins extracted from the petals of *Hibiscus syriacus* L. were able to inhibit NLRP3 Inflammasome in BV2 microglia cells by alleviating NF κ B and endoplasmic reticulum stress induced Ca²⁺ accumulation and mitochondrial ROS production [62]. In general, the anti-inflammatory molecular mechanisms of anthocyanins have been associated with some mechanisms, among which it is worth noting, and which are associated with the results presented here: (i) their ability to suppress the activation of NF κ B [63], and (ii) suppress LRR, NACHT and PYD domains-containing protein 3 (NLRP3) inflammasomes by activation of *NRF2* and the thioredoxin-1/thioredoxin-interacting protein (Trx1/TXNIP) inhibitory complex [64].

After NLRP3 activation, we observed a significant increase ($p<0.01$) in the levels of p62 and the LC3I/LC3II ratio in the LPS+ATP group, proteins that play a fundamental role in the autophagy process (Fig. 3B). It consists in a guided degradation of unnecessary or dysfunctional

cellular components such as misfolded proteins and dysfunctional organelles by a double membrane vesicle, or autophagosome, and then delivered to the lysosomes for their degradation [65]. High levels of p62 together with high levels of LC3I/LC3II ratio suggest an impairment in the autophagy with the accumulation of non-degraded substrates. However, after treatment with ABCext (LPS+ATP+ABCext group), the p62 levels showed a significant decrease ($p<0.01$) and the LC3I/LC3II ratio increase ($p < 0.05$), compared to the LPS+ATP group, indicating a restoration of the autophagy process (Fig. 3B). Autophagic dysfunction, as well as NLRP3 dysregulation, are associated with ageing and age-related diseases. By treating the cells with ABCext after triggering the inflammatory response, we have observed a reduction in NLRP3 protein level which leads, according to other studies [66–68], to a decrease in p62 and an increase in the LC3I/LC3II ratio, restoring the autophagy flux. In light of these results the consumption of Andean blackberry polyphenols, more probably the anthocyanins seem certainly promising to treat inflammatory-based and age-related diseases. The results presented here agree with several studies that have demonstrated that anthocyanins from different natural sources induce autophagy [69,70]. More recently it was demonstrated that anthocyanins from *Sambucus canadensis* could significantly reduce cell senescence and lens ageing by increasing autophagic and mitophagy flux and enhancing mitochondria and cell renewal through the inhibition of the PI3K/AKT/mTOR signaling pathway, which leads to attenuation of ageing [71].

4. Conclusions

According to the results presented here, it was possible to demonstrate a modulating effect of inflammation by the polyphenolic components of the Andean blackberry. This effect was not only due to its antioxidant capacity, evidenced in the improvement of markers of oxidative damage to cellular macromolecules, but also through its ability to regulate certain genes related to the antioxidant and inflammatory response. The components of the Andean blackberry showed a modulatory effect of the NLRP3 inflammasome complex, decreasing its activation, in addition to restoring the autophagy process, which suggests a direct action of Andean blackberry compounds

on the inflammatory response and restoration of the autophagy process. Our findings require further investigation, and more efforts in *in vivo* studies and intervention models evaluating the impact of Andean blackberry components on modulating inflammatory and antioxidant responses and the direct involvement of the NLRP3 complex in these effects should be done.

Conflicts of interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

This work was supported by Universidad de Las Américas (Grant Number: BIO.YGB.21.02) and Universidad San Francisco de Quito (Grant Number: 15748) and Proyecto PID2019-106167RB-I00 financed by MCIN/ AEI /10.13039/501100011033. Access to plant genetic resources was granted by means of the “Framework agreement for access to plant genetic resources: MAE-DNB-CM-2017-0072-M-001” celebrated between the Environment Ministry of Ecuador and the Universidad de Las Américas, Ecuador.

References

- [1] WHO. Increasing fruit and vegetable consumption to reduce the risk of noncommunicable diseases Biological , behavioural and contextual rationale. World Health Organization 2016:7–12.
- [2] Camps J, García-Heredia A. Introduction: Oxidation and inflammation, a molecular link between non-communicable diseases. *Advances in Experimental Medicine and Biology*. 2014;824:1–4.
- [3] Majdan M, Bobrowska-Korczak B. Active compounds in fruits and inflammation in the body. *Nutrients*. 2022;14:2496.
- [4] Del Rio D, Borges G, Crozier A. Berry flavonoids and phenolics: Bioavailability and evidence of protective effects. *British Journal of Nutrition*. 2010;104:S67–90.
- [5] Santhakumar AB, Battino M, Alvarez-Suarez JM. Dietary polyphenols: Structures, bioavailability and protective effects against atherosclerosis. *Food and Chemical Toxicology*. 2018;113:49–65.

- 522 [6] Giampieri F, Cinciosi D, Alvarez-Suarez JM, Quiles JL, Forbes-Hernández TY, Navarro-
523 Hortal MD, et al. Anthocyanins: What do we know until now? *Journal of Berry Research*.
524 2023;13:1–6.
- 525 [7] Salehi B, Sharifi-Rad J, Cappellini F, Reiner Ž, Zorzan D, Imran M, et al. The therapeutic
526 potential of anthocyanins: Current approaches based on their molecular mechanism of
527 action. *Frontiers in Pharmacology*. 2020;11.
- 528 [8] Guevara-Terán M, Padilla-Arias K, Beltrán-Novoa A, González-Paramás AM, Giampieri
529 F, Battino M, et al. Influence of altitudes and development stages on the chemical
530 composition, antioxidant, and antimicrobial capacity of the wild Andean Blueberry
531 (*Vaccinium floribundum* Kunth). *Molecules*. 2022;27:7525.
- 532 [9] Mattioli R, Francioso A, Mosca L, Silva P. Anthocyanins: A comprehensive Review of their
533 chemical properties and health effects on cardiovascular and neurodegenerative diseases.
534 *Molecules*. 2020;25:3809.
- 535 [10] Carrillo-Perdomo E, Aller A, Cruz-Quintana SM, Giampieri F, Alvarez-Suarez JM. Andean
536 berries from Ecuador: A review on botany, agronomy, chemistry and health potential.
537 *Journal of Berry Research*. 2015;5:49–69.
- 538 [11] Trujillo León J. La colada morada: antropología de la culinaria ritual ecuatoriana. *Anales*
539 *de La Universidad Central Del Ecuador* 2019;1:123–44.
- 540 [12] Azofeifa G, Quesada S, Boudard F, Morena M, Cristol JP, Pérez AM, et al. Antioxidant and
541 anti-inflammatory in vitro activities of phenolic compounds from tropical highland
542 blackberry (*rubus adenotrichos*). *Journal of Agricultural and Food Chemistry*.
543 2013;61:5798–804.
- 544 [13] Kaume L, Howard LR, Devareddy L. The Blackberry Fruit: A review on its composition
545 and chemistry, metabolism and bioavailability, and health benefits. *Journal of Agricultural*
546 *and Food Chemistry*. 2012;60:5716–27.
- 547 [14] Lavefve L, Howard LR, Carbonero F. Berry polyphenols metabolism and impact on human
548 gut microbiota and health. *Food and Function*. 2020;11:45–65.
- 549 [15] Meireles M, Marques C, Norberto S, Fernandes I, Mateus N, Rendeiro C, et al. The impact
550 of chronic blackberry intake on the neuroinflammatory status of rats fed a standard or high-
551 fat diet. *Journal of Nutritional Biochemistry* 2015;26:1166–73.
- 552 [16] Arozarena Í, Ortiz J, Hermosín-Gutiérrez I, Urretavizcaya I, Salvatierra S, Córdova I, et al.
553 Color, ellagitannins, anthocyanins, and antioxidant activity of Andean Blackberry (*Rubus*
554 *glaucus* Benth.) wines. *Journal of Agricultural and Food Chemistry*. 2012;60:7463–73.
- 555 [17] Osorio C, Hurtado N, Dawid C, Hofmann T, Heredia-Mira FJ, Morales AL. Chemical
556 characterisation of anthocyanins in tamarillo (*Solanum betaceum* Cav.) and Andes berry
557 (*Rubus glaucus* Benth.) fruits. *Food Chemistry*. 2012;132:1915–21.
- 558 [18] Vasco C, Riihinen K, Ruales J, Kamal-Eldin A. Phenolic Compounds in Rosaceae Fruits
559 from Ecuador. *Journal of Agricultural and Food Chemistry*. 2009;57:1204–12.
- 560 [19] Alarcón-Barrera KS, Armijos-Montesinos DS, García-Tenesaca M, Iturralde G, Jaramilo-
561 Vivanco T, Granda-Albuja MG, et al. Wild Andean blackberry (*Rubus glaucus* Benth) and
562 Andean blueberry (*Vaccinium floribundum* Kunth) from the highlands of Ecuador:

563 Nutritional composition and protective effect on human dermal fibroblasts against cytotoxic
564 oxidative damage. *Journal of Berry Research*. 2018;8:223–36.

565 [20] Giampieri F, Alvarez-Suarez JM, Cordero MD, Gasparrini M, Forbes-Hernandez TY, Afrin
566 S, et al. Strawberry consumption improves aging-associated impairments, mitochondrial
567 biogenesis and functionality through the AMP-activated protein kinase signaling cascade.
568 *Food Chemistry*. 2017;234:464–71.

569 [21] Alvarez-Suarez JM, Carrillo-Perdomo E, Aller A, Giampieri F, Gasparrini M, González-
570 Pérez L, et al. Anti-inflammatory effect of Capuli cherry against LPS-induced cytotoxic
571 damage in RAW 264.7 macrophages. *Food and Chemical Toxicology*. 2017;102:46–52.

572 [22] Bradford MM. A rapid and sensitive method for the quantitation of microgram quantities
573 of protein utilizing the principle of protein-dye binding. *Analytical Biochemistry*.
574 1976;72:248–54.

575 [23] Kakkar P, Das B, Viswanathan PN. A modified spectrophotometric assay of superoxide
576 dismutase. *Indian Journal of Biochemistry and Biophysics*. 1984;21:130–2.

577 [24] Aebi H. Catalase in vitro. *Methods in Enzymology*. 1984;105:121–6.

578 [25] Sies H, Koch OR, Martino E, Boveris A. Increased biliary glutathione disulfide release in
579 chronically ethanol-treated rats. *FEBS Letters*. 1979;103:287–90.

580 [26] Carlberg I, Mannervik B. Glutathione reductase. *Methods in Enzymology*. 1985;113:484–
581 90.

582 [27] Habig WH, Pabst MJ, Jakoby WB. Glutathione S-Transferases. *Journal of Biological*
583 *Chemistry* 1974;249:7130–9.

584 [28] Levine RL, Garland D, Oliver CN, Amici A, Climent I, Lenz A-G, et al. [49] Determination
585 of carbonyl content in oxidatively modified proteins. *Methods in Enzymology*.
586 1990;186:464–78.

587 [29] Ohkawa H, Ohishi N, Yagi K. Assay for lipid peroxides in animal tissues by thiobarbituric
588 acid reaction. *Analytical Biochemistry*. 1979;95:351–8.

589 [30] Livak KJ, Schmittgen TD. Analysis of relative gene expression data using Real-Time
590 Quantitative PCR and the 2- $\Delta\Delta$ CT method. *Methods*. 2001;25:402–8.

591 [31] González-Dominguez A, Montañez R, Castejón-Vega B, Nuñez-Vasco J, Lendines-
592 Cordero D, Wang C, et al. Inhibition of the NLRP3 inflammasome improves lifespan in
593 animal murine model of Hutchinson–Gilford Progeria. *EMBO Molecular Medicine*.
594 2021;13.

595 [32] Jamovi Project. The Jamovi Project. 2022 2022:Version 2.3.

596 [33] R Core Team. R: A Language and environment for statistical computing 2021:Version 4.1.

597 [34] Fox J, Wesiberg S, Price B, Adler D. Companion to Applied Regression [R package car
598 version 3.0-12] 2021.

599 [35] Van de Velde F, Esposito D, Grace MH, Pirovani ME, Lila MA. Anti-inflammatory and
600 wound healing properties of polyphenolic extracts from strawberry and blackberry fruits.
601 *Food Research International*. 2019;121:453–62.

- 602 [36] Shahidi F, Chandrasekara A. Hydroxycinnamates and their in vitro and in vivo antioxidant
603 activities. *Phytochemistry Reviews*. 2010;9:147–70.
- 604 [37] Sova M, Saso L. Natural Sources, Pharmacokinetics, Biological Activities and Health
605 Benefits of Hydroxycinnamic Acids and Their Metabolites. *Nutrients*. 2020;12:2190.
- 606 [38] Pei K, Ou J, Huang J, Ou S. *p* -Coumaric acid and its conjugates: dietary sources,
607 pharmacokinetic properties and biological activities. *Journal of the Science of Food and*
608 *Agriculture*. 2016;96:2952–62.
- 609 [39] Christensen LP, Christensen KB. The Role of direct and indirect polyphenolic antioxidants
610 in protection against oxidative stress. *Polyphenols in human health and disease*, Elsevier;
611 2014, p. 289–309.
- 612 [40] Gasparini M, Forbes-Hernandez TY, Giampieri F, Afrin S, Alvarez-Suarez JM, Mazzoni
613 L, et al. Anti-inflammatory effect of strawberry extract against LPS-induced stress in RAW
614 264.7 macrophages. *Food and Chemical Toxicology*. 2017;102:1–10.
- 615 [41] Amatori S, Mazzoni L, Alvarez-Suarez JM, Giampieri F, Gasparini M, Forbes-Hernandez
616 TY, et al. Polyphenol-rich strawberry extract (PRSE) shows in vitro and in vivo biological
617 activity against invasive breast cancer cells. *Scientific Reports*. 2016;6: 30917.
- 618 [42] Alvarez-Suarez JM, Giampieri F, Tulipani S, Casoli T, Di Stefano G, González-Paramás
619 AM, et al. One-month strawberry-rich anthocyanin supplementation ameliorates
620 cardiovascular risk, oxidative stress markers and platelet activation in humans. *Journal of*
621 *Nutritional Biochemistry*. 2014;25:289–94.
- 622 [43] Leyane TS, Jere SW, Houreld NN. Oxidative stress in ageing and chronic degenerative
623 pathologies: Molecular mechanisms involved in counteracting oxidative stress and chronic
624 inflammation. *International Journal of Molecular Science*. 2022;23:7273.
- 625 [44] Calvano A, Izuora K, Oh EC, Ebersole JL, Lyons TJ, Basu A. Dietary berries, insulin
626 resistance and type 2 diabetes: an overview of human feeding trials. *Food and Function*.
627 2019;10:6227–43.
- 628 [45] Speer H, D’Cunha NM, Alexopoulos NI, McKune AJ, Naumovski N. Anthocyanins and
629 human health—A focus on oxidative stress, inflammation and disease. *Antioxidants*.
630 2020;9:366.
- 631 [46] Röhrl C, Steinbauer S, Bauer R, Roitinger E, Otteneder K, Wallner M, et al. Aqueous
632 extracts of lingonberry and blackberry leaves identified by high-content screening
633 beneficially act on cholesterol metabolism. *Food and Function*. 2021;12:10432–42.
- 634 [47] Alvarez-Suarez JM, Cuadrado C, Redondo IB, Giampieri F, González-Paramás AM,
635 Santos-Buelga C. Novel approaches in anthocyanin research - Plant fortification and
636 bioavailability issues. *Trends in Food Science and Technology*. 2021;117:92–105.
- 637 [48] Solverson P, Albaugh GP, Harrison DJ, Luthria DL, Baer DJ, Novotny JA. High-dose
638 administration of purified cyanidin-3-glucose or a blackberry extract causes improved
639 mitochondrial function but reduced content in 3T3-L1 adipocytes. *Food Frontiers*.
640 2022;3:276–84.

- 641 [49] Cenk E, Schmutz C, Pahlke G, Oertel A, Kollarova J, Mock HP, et al. Immunomodulatory
642 properties of blackberry anthocyanins in thp-1 derived macrophages. *International Journal*
643 *of Molecular Science*. 2021;22.
- 644 [50] Pemmari T, Hämäläinen M, Ryyti R, Peltola R, Moilanen E. Cloudberry (*Rubus*
645 *chamaemorus* L.) supplementation attenuates the development of metabolic inflammation
646 in a high-fat diet mouse model of obesity. *Nutrients*. 2022;14.
- 647 [51] Giampieri F, Alvarez-Suarez JM, Battino M. Strawberry and human health: Effects beyond
648 antioxidant activity. *Journal of Agricultural and Food Chemistry*. 2014;62:3867–76.
- 649 [52] Liu T, Zhang L, Joo D, Sun S-C. NF- κ B signaling in inflammation. *Signal Transduction*
650 *and Targeted Therapy*. 2017;2:17023.
- 651 [53] He F, Ru X, Wen T. NRF2, a transcription factor for stress response and beyond.
652 *International Journal of Molecular Science*. 2020;21:4777.
- 653 [54] Ma Z, Du B, Li J, Yang Y, Zhu F. An insight into anti-Inflammatory activities and
654 inflammation related diseases of anthocyanins: A Review of both in vivo and in vitro
655 investigations. *International Journal of Molecular Science*. 2021;22:11076.
- 656 [55] Kelley N, Jeltema D, Duan Y, He Y. The NLRP3 inflammasome: An overview of
657 mechanisms of activation and regulation. *International Journal of Molecular Science*.
658 2019;20:3328.
- 659 [56] Alcocer-Gómez E, Castejón-Vega B, López-Sánchez M, Cordero MD. Inflammasomes in
660 clinical practice: A brief introduction, In: Cordero, M., Alcocer-Gómez, E. (eds) 641
661 *Inflammasomes: Clinical and Therapeutic Implications*. *Experientia Supplementum*, vol
662 642 108. Springer, Cham. 2018, p. 1–8.
- 663 [57] Navarro-Pando JM, Alcocer-Gómez E, Castejón-Vega B, Navarro-Villarán E, Condés-
664 Hervás M, Mundi-Roldan M, et al. Inhibition of the NLRP3 inflammasome prevents
665 ovarian aging. *Science Advances*. 2021;7.
- 666 [58] Chung HY, Kim DH, Lee EK, Chung KW, Chung S, Lee B, et al. Redefining chronic
667 inflammation in aging and age-related diseases: Proposal of the senoinflammation concept.
668 *Aging and Disease*. 2019;10:367.
- 669 [59] Cordero MD, Williams MR, Ryffel B. AMP-Activated Protein Kinase regulation of the
670 NLRP3 inflammasome during aging. *Trends in Endocrinology & Metabolism* 2018;29:8–
671 17.
- 672 [60] Castejon-Vega B, Giampieri F, Alvarez-Suarez JM. Nutraceutical compounds targeting
673 inflammasomes in human diseases. *International Journal of Molecular Sciences*. 654
674 2020;21:4829.
- 675 [61] Dong G, Xu N, Wang M, Zhao Y, Jiang F, Bu H, et al. Anthocyanin Extract from Purple
676 Sweet Potato Exacerbate Mitophagy to Ameliorate Pyroptosis in *Klebsiella pneumoniae*
677 Infection. *International Journal of Molecular Sciences*. 2021;22:11422.
- 678 [62] Molagoda IMN, Lee KT, Choi YH, Jayasingha JACC, Kim G-Y. Anthocyanins from
679 *Hibiscus syriacus* L. inhibit NLRP3 inflammasome in BV2 microglia cells by alleviating
680 NF- κ B- and ER stress-Induced Ca²⁺ accumulation and mitochondrial ROS production.
681 *Oxidative Medicine and Cellular Longevity*. 2021;2021:1–17.

- [63] Poulouse SM, Fisher DR, Larson J, Bielinski DF, Rimando AM, Carey AN, et al. Anthocyanin-rich Açai (*Euterpe oleracea* Mart.) fruit pulp fractions attenuate inflammatory stress signaling in mouse brain BV-2 microglial cells. *Journal of Agricultural and Food Chemistry*. 2012;60:1084–93.
- [64] Cui H-X, Chen J-H, Li J-W, Cheng F-R, Yuan K. Protection of anthocyanin from *Myrica rubra* against cerebral ischemia-reperfusion injury via modulation of the TLR4/NF- κ B and NLRP3 pathways. *Molecules*. 2018;23:1788.
- [65] Parzych KR, Klionsky DJ. An Overview of Autophagy: Morphology, Mechanism, and Regulation. *Antioxidants & Redox Signaling*. 2014;20:460–73.
- [66] Cañadas-Lozano D, Marín-Aguilar F, Castejón-Vega B, Ryffel B, Navarro-Pando JM, Ruiz-Cabello J, et al. Blockade of the NLRP3 inflammasome improves metabolic health and lifespan in obese mice. *Geroscience*. 2020;42:715–25.
- [67] Marín-Aguilar F, Lechuga-Vieco A V., Alcocer-Gómez E, Castejón-Vega B, Lucas J, Garrido C, et al. NLRP3 inflammasome suppression improves longevity and prevents cardiac aging in male mice. *Aging Cell*. 2020;19.
- [68] Marín-Aguilar F, Castejón-Vega B, Alcocer-Gómez E, Lendines-Cordero D, Cooper MA, de la Cruz P, et al. NLRP3 inflammasome inhibition by MCC950 in aged mice improves health via enhanced autophagy and PPAR α activity. *The Journals of Gerontology: Series A*. 2020;75:1457–64.
- [69] Li J, Zhao R, Zhao H, Chen G, Jiang Y, Lyu X, et al. Reduction of aging-induced oxidative stress and activation of autophagy by Bilberry anthocyanin supplementation via the AMPK–mTOR signaling pathway in aged female rats. *Journal of Agricultural and Food Chemistry*. 2019;67:7832–43.
- [70] Choe Y-J, Ha TJ, Ko K-W, Lee S-Y, Shin SJ, Kim H-S. Anthocyanins in the black soybean (*Glycine max* L.) protect U2OS cells from apoptosis by inducing autophagy via the activation of adenosyl monophosphate-dependent protein kinase. *Oncology Reports*. 2012;28:2049–56.
- [71] Hu X, Yang Y, Tang S, Chen Q, Zhang M, Ma J, et al. Anti-Aging Effects of Anthocyanin Extracts of *Sambucus canadensis* Caused by Targeting Mitochondrial-Induced Oxidative Stress. *International Journal of Molecular Sciences*. 2023;24:1528.

FIGURE CAPTION

Figure. 1. HPLC chromatograms of phenolic acid derivatives and flavonoid profile in the crude extract of Andean blackberry (ABCext). A, shows representative HPLC chromatograms of the polyphenol profile for phenolic acid derivatives and flavonoids recorded at 330 and 360 nm and for anthocyanins at 520 nm. B, shows the specific profile of anthocyanins (at 520 nm) and C, the specific profile of phenolic acids and flavonoids (at 330 nm).

719

720 **Figure. 2.** Expression of genes related to the inflammatory response and antioxidant response in
721 RAW 264.7 murine macrophages treated with different combinations of ABCext and/or LPS. Data
722 are expressed as mean values \pm SD. $p<0.05$, $p<0.01$, significant differences between the
723 experimental groups.

724

725 **Figure. 3.** Expression levels of the proteins studied. Data are expressed as mean values \pm SD.
726 $*p<0.05$, $**p<0.01$, significant differences between the experimental groups.