



UNIVERSITÀ POLITECNICA DELLE MARCHE
Repository ISTITUZIONALE

Organochlorines and Polycyclic Aromatic Hydrocarbons as fingerprint of exposure pathways from marine sediments to biota

This is the peer reviewed version of the following article:

Original

Organochlorines and Polycyclic Aromatic Hydrocarbons as fingerprint of exposure pathways from marine sediments to biota / Traina, A., Ausili, A., Bonsignore, M., Fattorini, D., Gherardi, S., Gorbi, S., Quinci, E., Romano, E., Salvagio Manta, D., Tranchida, G., Regoli, F., Sprovieri, Mario.. - In: MARINE POLLUTION BULLETIN. - ISSN 0025-326X. - 170:(2021). [10.1016/j.marpolbul.2021.112676]

Availability:

This version is available at: 11566/328698 since: 2024-04-11T13:48:40Z

Publisher:

Published

DOI:10.1016/j.marpolbul.2021.112676

Terms of use:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. The use of copyrighted works requires the consent of the rights' holder (author or publisher). Works made available under a Creative Commons license or a Publisher's custom-made license can be used according to the terms and conditions contained therein. See editor's website for further information and terms and conditions.

This item was downloaded from IRIS Università Politecnica delle Marche (<https://iris.univpm.it>). When citing, please refer to the published version.

(Article begins on next page)

Organochlorines and Polycyclic Aromatic Hydrocarbons as fingerprint of exposure pathways from marine sediments to biota

Anna Traina^a, Antonella Ausili^b, Maria Bonsignore^{c*}, Daniele Fattorini^d, Serena Gherardi^e, Stefania Gorbi^d, Enza Quinci^c, Elena Romano^b, Daniela Salvagio Manta^a, Giorgio Tranchida^c, Francesco Regoli^d, Mario Sprovieri^c

^a Institute of Anthropic Impacts and Sustainability in marine environment - National Research Council of Italy (IAS-CNR), Lungomare Cristoforo Colombo 452, 90149 Palermo, Italy

^b Institute for Environmental Protection and Research (ISPRA), Via V. Brancati 60, 00144 Rome, Italy

^c Institute of Anthropic Impacts and Sustainability in marine environment - National Research Council of Italy (IAS-CNR), via del Mare 3, 91021 Torretta Granitola (Trapani), Italy

^d Department of Life and Environmental Sciences (DiSVA), Polytechnic University of Marche, Via Breccie Bianche, Monte Dago, 60131 Ancona, Italy

^e National Research Council of Italy-Institute of Marine Science (ISMAR-CNR), Calata Porta di Massa, 80133 Naples, Italy

* Corresponding author: maria.bonsignorei@ias.cnr.it

Abstract

To elucidate the dynamics of a suite of organochlorine contaminants (PCBs, HCB), PAHs and Hg and verify the potential of these pollutants as reliable fingerprints of sources, an ensemble of marine sediments and organisms (finfish, shellfish species and *Mytilus galloprovincialis*) were analysed from the contaminated Augusta Bay (Southern Italy). The Hg and HCB concentration in the sediments exceeded the EQS of the Directive 2000/60/EU. Similarly, Σ PCB and selected PAHs were above the threshold limit set by regulation. The marine organisms showed Hg concentrations above CE 1881/2006. Contaminants in transplanted mussel evidenced an increased accumulation overtime and different distribution patterns between sampling sites. Analysis of the homolog composition of PCB congeners revealed comparable patterns between sediments and marine organisms and offered the opportunity to define a robust fingerprint for tracing contaminants transfer from the abiotic to the biotic compartments. These results were confirmed by the Fluoranthene/Pyrene, Hg and HCB distribution modes.

Keywords: Organochlorines, PAHs, Biota, Sediment, Source fingerprint.

1. Introduction

Understanding the transfer of contaminants from marine sediments to biota represents a critical aspect for environmental risk assessment, modelling and specific remediation actions (Gobas and Arnot, 2010). Distribution patterns of pollutants, isotope measurements, compositional analysis and investigation of variable proportions of single congeners of specific organic pollutants were shown as valuable tools for tracing sources of contamination to the trophic chains (e.g., Sahu et al., 2009; Baskaran, 2011; Rodenburg and Leidos, 2017; Habibullah-Al-Mamun et al., 2019; Bonsignore et al., 2020).

42 Although several actions have been taken to reduce or even eliminate input of contaminants in the
43 environment (UNEP 2001, 2017, 2019), compounds such as Persistent Organic Pollutants (POPs)
44 and some trace metals, including Hg, are accumulated in different environmental matrices thus
45 representing a global environmental threat for marine organisms and human health (García-Flor et
46 al., 2009; Bellas et al., 2011; Sanchez-Avila et al., 2012; Berrojalbiz et al., 2014; Olenycz et al.,
47 2015; Batang et al., 2016; Jepson et al., 2016; Beiras, 2018). Due to their hydrophobic behaviour in
48 aquatic ecosystems, most of the pollutants such as e.g. Polychlorinated biphenyls -[PCBs],
49 Hexachlorobenzene [HCB] and Polycyclic Aromatic Hydrocarbons [PAHs], readily bind to
50 particles (suspended in seawater and/or into the sediments) and are strongly associated with organic
51 phase. Once in the sediments they persist for very long time (Minh et al., 2007) with high potential
52 to be incorporated into the food web via benthic–pelagic coupling (Thomann et al., 1992). As an
53 example of extremely stable compounds, Hexachlorobenzene (HCB) was produced for industrial
54 and agricultural applications worldwide (Courtney, 1979; Bailey, 2001; Meijer et al., 2003), and
55 although its production was stopped before 2000 in most countries, it is still abundant in the
56 environment due to its long-half life in water and sediments (Barber et al., 2005). PCBs were
57 produced in the past century, up to the 1970s-1980s, and largely used for different industrial
58 applications (e.g. as dielectric fluid in electrical transformers and capacitors, active constituent of
59 pesticides) (ATSDR, 2000; Crinnion, 2011). Among 209 congeners, the Stockholm Convention
60 reports as mandatory to measure at least six indicator congeners non-dioxin-like (NDL) PCBs 28,
61 52, 101, 138, 153 and 180 to estimate environmental PCBs contamination (JECFA, 2018), while
62 ICES (Webster et al., 2013) recommends the measure of 7 PCBs congeners (including also
63 congener 118) which represent a large percentage of the total congeners measured in environmental
64 samples and human fluids (EFSA, 2010; IARC, 2016). Likewise, Hg is a trace metal efficiently
65 transferred and biomagnified along food web reaching high concentrations in the upper trophic
66 levels and thus representing a risk to human health (e.g., Signa et al., 2017).

67 Finally, PAHs represent another group of chemicals of priority concern which include the largest
68 known class of carcinogens and chemical mutagens (Keith and Telliard, 1979). PAHs tend to
69 adsorb rapidly on particles (Neff, 1979; Landrum and Robbins, 1990) and since their solubility
70 decreases with increasing molecular weight, the bioaccumulation of these chemicals from
71 sediments to marine organisms is generally higher for those with lower molecular weight (Porte and
72 Albaigés, 1993; Djomo et al., 1996).

73 Over the last decade, a number of scientific contributions explored the distribution patterns of
74 classes of contaminants (e.g., PAHs, PCBs, PCDD/Fs, Hg and HCB) to trace transfer dynamic from
75 the sediments to the biotic compartment, and to characterize sources of variability in contaminant
76 bioavailability to aquatic biota (e.g. Selck et al., 2012; McLeod et al., 2015).

77 Fish and other key species have been used as bioindicators to investigate the presence and toxic
78 effects of chemical pollutants (Ueno et al., 2003). It is well-known that biotransformation pathways
79 significantly modulate absorption, distribution and excretion of organic xenobiotics in fish while
80 filter feeders, such as bivalve, tend to accumulate these pollutants in their tissues from both water
81 column and sediments thus providing additional information on the environmental state of pollution
82 (Regoli et al., 1998; Wiberg et al., 2002; Olenycz et al., 2015; Farrington et al., 2016; Beyer et al.,
83 2017).

84 However, PCBs congeners distribution in the biota generally remains similar to the un-weathered
85 PCBs mixtures in the sediments (mainly when those with higher chlorine content are considered),
86 thus allowing the identification of these sources with a satisfactory degree of confidence. Some
87 exercises of PCBs fingerprinting in biota have been performed in the Hanford Site in Washington
88 State (Rodenburg et al., 2015) and Portland Harbor in Oregon (Rodenburg et al., 2019) offering
89 robust information on the transfer of contaminants from the abiotic to the biotic compartments.

90 In this work, we present an unprecedented dataset of Hg, HCB, PCBs and 16 priority PAHs
91 measured on surface sediments, mussels, and several species of marine organisms from the highly
92 contaminated marine area of Augusta Bay (southern Italy). Patterns of PCBs congeners, combined

to spatial distribution of PAHs, HCB and Hg in sediments and biota were elaborated to evaluate the fingerprint association between sediments (sources) and biota, and their opportunity for high-resolution tracing of priority contaminants sources.

2. Study area

Augusta Bay is a $\sim 25 \text{ km}^2$ area located in the eastern Sicilian coast (Fig. 1) which hosts one of the most important harbours of the Mediterranean Sea and, since 1950s, one of the largest petrochemical complexes in Europe, characterized by oil refineries and chlor-alkali plant. Uncontrolled industrial discharges led to significant contamination of sediments by metals and organics, mainly Hg, PCBs, PAHs and HCB (ICRAM, 2008; Bellucci et al., 2012; Croudace et al., 2015), and all the area was included in the National Remediation Plan by the Italian Ministry of Environment. Several investigations have highlighted a strong contamination of the marine environment with i) abnormal levels of Hg, PCBs and HCB in sediments (ICRAM, 2005, 2008, Bellucci et al., 2012; Sprovieri et al., 2011; Orecchio and Polizzotto, 2013), ii) active fluxes of Hg from the sediment to the water column (Salvagio Manta et al., 2016; Denaro et al., 2020) and iii) Hg evasion processes to the atmosphere (Sprovieri et al., 2011; Bagnato et al., 2013). Serious impacts on the ecosystem have been documented by ecotoxicological investigations which revealed genotoxic damages and high Hg contents, exceeding the regulatory limits for food consumption in the tissues of fish and mussels (Ausili et al., 2008; ENVIRON, 2008; ICRAM, 2008; Tomasello et al., 2012; Bonsignore et al., 2013; Signa et al., 2017), thus advising for possible human health implications due to consumption of seafood from this area (Ausili et al., 2008; Bonsignore et al., 2013; Di Bella et al., 2020).

The role of polluted sediments as the main carrier of Hg to the ecosystem and local fish consumers has been further revealed through the exploration of Hg isotopes signature in sediments and fish (Bonsignore et al., 2015).

119 3. Sampling activity

120

121 3.1 Sediments

122 The sediment sampling, carried out in October 2017 during an oceanographic cruise on board R/V
123 “Luigi Sanzo”, provided sediment cores from 4 stations (A3, A7, A9, A11) inside the bay, from the
124 northern to the southern area (Fig. 1), using an oceanic box-corer. Sediment cores were immediately
125 stored at –20 °C until the analyses.

126

127 3.2 Marine organisms

128 In November 2013, wild mussels (*Mytilus galloprovincialis*) with size range 3.6-6 cm, were
129 manually collected from industrial wharf and rapidly transported to laboratory for the tissue’s
130 dissection. In the same period, a total of 62 specimens of finfish (*Sphyraena sphyraena*, *Mullus*
131 *barbatus*, *Pagellus* spp., *Sparus aurata*, *Serranus cabrilla*, *Diplodus* spp.) were obtained through
132 local fisherman (Fig. 1).

133 In September 2017, mussels (*Mytilus galloprovincialis*) with size range of 4-6 cm were obtained
134 from a commercial farm. A pool of individuals (about 2 kg) was used as control (*M.*
135 *galloprovincialis* CTRL), while other two pools (about 10 kg) were transplanted in the northern (*M.*
136 *galloprovincialis* N) and the southern (*M. galloprovincialis* S) areas of Augusta Bay, caged in nylon
137 net bags, secured to a rope and maintained at approximately 3 m water depth. After a period of 5
138 weeks, a subgroup of mussels was collected from each site (N1 and S1), and an additional sampling
139 was performed 7 months later (N2 and S2).

140 In October 2017, a total of 96 specimens of finfish (*Sphyraena sphyraena*, *Trigla lucerna*, *Mullus*
141 *barbatus*, *Pagellus* spp., *Diplodus* spp.) and shellfish (*Parapaeneus kerathurus* and *Sepia* spp.)
142 were obtained through local fishermen (Fig. 1).

143

144 4. Analytical methods

145 4.1 Samples preparation

146 In laboratory, the sediment cores were defrosted, extruded and immediately sliced into 1 cm
147 intervals. The samples were then dried at 35°C and homogenized by an agate mortar. For the goals
148 of this study only the most superficial levels (between 0-10 cm) were analysed.

149 The total length (mm) and the weight (g) for each organism were measured and muscles and soft
150 tissues were dissected by stainless steel scissors. Individuals of each species with comparable size
151 were pooled, homogenized and stored at -80°C (Table 1). The tissues samples were then freeze-
152 dried and powdered by an agate mortar prior the analyses.

153

154 4.2 Mercury determination in sediments and marine organisms

155 The concentration of Hg in organisms and sediments was determined by a direct mercury analyzer
156 (milestone-DMA-80 atomic absorption spectrophotometer), according to analytical procedures
157 reported in EPA 7473 method (2007). About 0.050 g of dried tissue and ~0.010 g of dry sediment
158 were loaded in nickel boats and transferred into the DMA-80 system. The Certified Reference
159 Materials-TORT-2 Lobster hepatopancreas and (PACS-2 Marine sediment, NRC-CNRC) were used
160 to assess accuracy (estimated % error =3%) and precision (routinely better than 4%; RSD%, n = 3).
161 About 20% of the total number of samples was duplicated to estimate reproducibility (better than
162 7%). Acid-cleaned laboratory materials were used in order to minimize contamination risks during
163 sample preparation and analyses procedures. Analyses were performed at the biogeochemical
164 laboratory of the Institute of Anthropic impacts and Sustainability in marine environment (IAS-
165 CNR) of Capo Granitola (Trapani, Italy). Results were expressed as on wet weight basis.

166 Results relative to tissues were converted from dried to wet-weight ($\mu\text{g g}^{-1}$) applying a conversion
167 factor previously calculated using the following formula: $C_w = C_d \times (100 - \%H_2O / 100)$ where C_d and
168 C_w are the concentration expressed relatively to dry and wet mass respectively. $\%H_2O$ is the
169 percentage of humidity (ranging around 80% for almost species) calculated after the freeze-drying
170 process (Di Bella et al., 2020 and references therein).

171

172 4.3 PCBs, HCB and PAHs determination in sediments and marine organisms

173 The sediment and biota samples were freeze dried and homogenized, then an aliquot (about 1 and 2
174 g of tissues and sediments, respectively) was placed in an ASE 200 steel extraction cell, spiked with
175 surrogate standards (o2si smart solution ® custom deuterated PAH mix; Wellington Laboratories
176 Inc PCB mix; CDN Isotope Inc HCB standard solution) for recovery monitoring and extracted by
177 pressurized fluid extraction using a moderately polar solvent mixture. The resulting extract was
178 cleaned up according to the matrix (sediment or biota) and the analytes (organochlorines or PAHs)
179 by a combination of column chromatography on silica (PAH) and florisil (organochlorine)
180 adsorbents and a shaking of the extract with a NaOH solution (PAH in biota) or with concentrated
181 sulphuric acid (organochlorine). For the determination of organochlorine compounds in sediment
182 samples the extract was also shaken with activated copper powder for sulphur removal. The cleaned
183 up concentrated extract was spiked with injection internal standard and analysed by GC/MS/MS for
184 organochlorine and GC/MS for PAHs.

185 PAH contents in sediments were compared to values established by the Italian regulation
186 (Legislative Decree 172/2015). Benzo(a)pyrene and Fluorantene were monitored in biota,
187 specifically in molluscs as suggested by Legislative Decree 172/2015.

188 Reference materials for sediments (SRM 1941b-NIST, organics in marine sediments) and biota
189 (NIST 2974a, organics in freeze-dried mussel's tissue - *Mytilus edulis*) were analysed to estimate
190 the accuracy (recoveries for each analyte of PAHs, PCBs and HCB ranged between 94% and
191 107%). The reproducibility, estimated on tripled samples, was better than $\pm 10\%$ and analytical
192 precision was routinely better than 4% (RSD%, n = 3).

193 For the PCB class of contaminants, we referred to the following notation: Σ PCBs (sum of all the
194 measured congeners), Σ PCBs¹ (sum of PCB congeners according to Italian Legislative Decree
195 172/2015 for marine sediments), Σ NDL-PCBs (PCB 28, 52, 101, 153, 138 and 180) and Σ 7PCBs
196 (Σ 6PCBs + PCB 118).

197

198 *4.4 Statistical analysis*

199 All the statistical analyses and graphics were performed using the statistical software R 3.6.3 (R
200 Development Core Team, 2020). Box-whisker plots of measured Hg, HCB and PCBs were used to
201 provide synthetic and direct comparisons for these parameters among the cores. For statistical
202 analysis, all values below the limit of detection (LOD) were set-up as $\frac{1}{2}$ LOD.

203 A hierarchical cluster analysis (HCA) with Euclidean distance and Ward's grouping method was
204 applied to the ratio 'individual PCB congener/ Σ PCB' in order to create homogeneous clusters of
205 samples (sediments, fishes and mussels) based on similarity of PCBs composition. The elbow
206 method was used to determine the most statistically reliable and representative number of clusters,
207 minimising the total intra-cluster variation.

208

209 **5. Results and Discussion**

210

211 *5.1 Contaminants in sediments*

212 *5.1.1. Mercury*

213 Mercury concentrations measured in the A3, A7, A9 and A11 cores ranged from 3.07 to 12.2 $\mu\text{g g}^{-1}$
214 (Tab. 2; Tab. S1) widely exceeding (generally more than one order of magnitude) the
215 Environmental Quality Standard (EQS), defined in accordance with Directive 2000/60/EU as
216 criteria for the achieving of the Good Chemical and Ecological Status of water bodies, and
217 established by the Italian regulation (Legislative Decree 172/2015; 0.3 mg kg^{-1}). The highest values
218 ($\text{mean} \pm 1\sigma = 11.4 \pm 0.57 \mu\text{g g}^{-1}$) were found in A9 core collected from the south-western sector of the
219 bay, in front of the past chlor-alkali plant (Tab. 2; Fig 2a). The lowest concentration ($\text{mean} \pm 1\sigma =$
220 $3.64 \pm 0.27 \mu\text{g g}^{-1}$) was recorded in the A3 core sampled in the northern area (Tab. 2, Fig. 2a).
221 Finally, A7 and A11 cores, collected from the central and southern part of the bay, showed
222 comparable levels of Hg ($\text{mean} \pm 1\sigma = 6.71 \pm 0.80 \mu\text{g g}^{-1}$ and $6.54 \pm 1.28 \mu\text{g g}^{-1}$, respectively).

223

224 5.1.2. Organochlorine compounds (HCB and PCBs) and PAHs

225 The analysis of organochlorine compounds in sediments showed very high concentrations of HCB
226 in all the analysed cores, ranging from 1.98 to 330 ng g⁻¹ (Tab. 2; Tab. S1). The mean values
227 registered from each core, again considering only the superficial 10 cm of sediments, were
228 considerably above the EQS for marine sediments (0.4 ng g⁻¹), with highest values measured in the
229 A9 core (mean±1σ = 172 ± 87.6 ng g⁻¹) (Tab. 2; Fig. 2b).

230 Also ΣPCBs and Σ7PCB registered higher values in A9 core (41.9 ± 5.60 ng g⁻¹; Tab. 2; Tab. S1;
231 Fig. 2c,d). The ΣPCBs also exceeded EQS value (8 ng g⁻¹) for sediments in all cores (Tab. 2).

232 The average concentrations of the individual PAHs were, in some cases, above the EQS indicated in
233 Tab 1/A of the Legislative Decree 172/2015 Specifically, the Anthracene concentrations (range =
234 23.4-286 ng g⁻¹) were above the EQS (24 ng g⁻¹) in all the analyzed samples; most of the A3 levels
235 showed concentrations of Benzo-a-Pyrene (mean±1σ = 64.7 ± 20.1 ng g⁻¹; EQS = 30 ng g⁻¹),
236 Benzo-b-Fluoranthene (mean±1σ = 54.6 ± 20.4 ng g⁻¹; EQS = 40 ng g⁻¹) and Benzo-k-Fluoranthene
237 (mean±1σ = 33.6 ± 24.3 ng g⁻¹; EQS = 20 ng g⁻¹) above the EQS values; furthermore, Indenopyrene
238 concentrations higher than EQS (70 ng g⁻¹) were recorded in the upper part (between 0-6 cm) of the
239 A3 core (Tab. 2; Tab. S2).

240

241 5.2 Contaminants in marine organisms

242 5.2.1. Mercury

243 The lowest Hg mean values referred to *Sepia* spp. collected in 2017 (0.21±0.08 µg g⁻¹) and *S.*
244 *aurata* sampled in 2013 (0.32±0.09 µg g⁻¹). In most cases, such as *S. sphyræna*, *T. lucerna*, *M.*
245 *barbatus*, *Diplodus* spp. and *P. kerathurus* (sampled in 2017) and *S. sphyræna*, *S. cabrilla* and
246 *Diplodus* spp. (sampled in 2013), the Hg levels exceed the threshold limits set by EU regulation for
247 contaminants in seafood (CE 1881/2006) (Tab. 3). Excluding the effect of age and length, the
248 contents of Hg found in organisms resulted in most cases higher than values previously measured in

the study area. Specifically, the concentrations measured in *Diplodus* spp. ($1.49 \pm 1.77 \mu\text{g g}^{-1}$ in 2017) and *Pagellus* spp. ($0.86 \pm 0.44 \mu\text{g g}^{-1}$ in 2017 and $0.53 \pm 0.12 \mu\text{g g}^{-1}$ in 2013) were higher than those reported by Bonsignore et al. (2013), Ausili et al. (2003) and ICRAM (2005, 2008), for the same species (between $0.56\text{--}0.90 \mu\text{g g}^{-1}$ and $0.36\text{--}0.41 \mu\text{g g}^{-1}$, respectively). In red mullet (*M. barbatus*), the concentrations relative to 2013 ($0.44 \pm 0.20 \mu\text{g g}^{-1}$) were in the same range of measurements ($0.46\text{--}0.82 \mu\text{g g}^{-1}$) of the previous investigations (Bonsignore et al., 2013; ICRAM, 2005, 2008) while those relative to 2017 (2.11 and $1.71 \mu\text{g g}^{-1}$) resulted significantly higher and in the same range of those reported by ICRAM in 2003 ($1.49\text{--}1.92 \mu\text{g g}^{-1}$). Finally, the Hg concentration in *T. lucerna* ($0.66 \mu\text{g g}^{-1}$) resulted slightly higher than that reported by ICRAM in 2003 ($0.57 \mu\text{g g}^{-1}$).

The bioavailability of Hg in the study area has been confirmed by analyses on both native and caged mussels with concentrations always higher than the EQS ($0.02 \mu\text{g g}^{-1}$) (Tab.3). Wild specimens collected in industrial wharf exhibited concentration of approximately about $0.2 \mu\text{g g}^{-1}$, ~20-fold higher than those measured in control organisms and comparable to those observed by Ausili et al. (2008) (Tab. 3). Bioaccumulation of Hg demonstrated to be very rapid in caged mussels showing, after only 5 weeks, comparable values to those of native organisms; after 7 months the concentrations increased by about 5-fold (Tab. 3). In particular, in the northern area, Hg in mussels 5 weeks after transplant (N1: $0.02 \pm 0.002 \mu\text{g g}^{-1}$) appeared double than in the control sample ($0.01 \mu\text{g g}^{-1}$), while after 7 months, the Hg increase was nearly one order of magnitude higher (N2: $0.12 \pm 0.01 \mu\text{g g}^{-1}$) (Tab. 3). Worthy to note, this value is reasonably in agreement with the results recently reported for mussels in the Augusta Bay ($0.17 \pm 0.07 \mu\text{g g}^{-1}$; Caricato et al., 2019). In the southern area, when comparing the concentration in pre-deployed mussels, the levels of Hg were ~23 times higher in those collected after 5 weeks (S1: $0.23 \pm 0.02 \mu\text{g g}^{-1}$) and up to 89 times higher after 7 months (S2: $0.89 \mu\text{g g}^{-1}$). The latter value exceeds the threshold limits set up by EU regulation for contaminants in seafood ($0.5 \mu\text{g g}^{-1}$; Reg. CE 1881/2006) and appeared much higher than those reported for many other Mediterranean sites (range: $0.01\text{--}0.20 \mu\text{g g}^{-1}$; Amodio-Cocchieri

et al., 2003; Ipolyi et al., 2004; Licata et al., 2004; Cardellicchio et al., 2010; Kljaković-Gašpić et al., 2010; Spada et al., 2011, 2012, 2013; Caricato et al., 2019).

277

5.2.2. HCB, PCBs and PAHs

Concentrations of HCB, PCBs and PAHs measured in organisms are reported in Table 3. Values of HCB were <LOD (0.05 ng g⁻¹) in all species of finfish and shellfish sampled in 2017, except for *S. sphyraena* (1.40 ng g⁻¹) and *M. barbatus* (0.24 and 0.65 ng g⁻¹); on the other hand, in 2013 they ranged between 0.09 ±0.02 (*Pagellus* spp.) and 1.36±0.31 ng g⁻¹ (*Mullus* spp.). All the concentrations are below the EQS of 10 ng g⁻¹.

HCB together with PCBs are ubiquitous contaminants in Mediterranean marine coastal areas (Solé et al., 2000). HCB can be stored in sediments and accumulated in benthic organisms. Being a lipophilic compound, it has a greater affinity for tissues with high lipid content, thus explaining the low concentrations measured in almost all the muscle samples of fish analysed in this study. Other authors have reported data related to HCB in fishes, but a direct comparison appears problematic due to the high variability of investigated organs/tissues (skin, gills or mussels' tissues) and the different used reference system (wet, dry or lipid weight) (Domingo and Bocio, 2007). Nonetheless, when comparable, our results are within the range of concentration reported from other studies for the Mediterranean Sea (Rodríguez-Hernández et al., 2016; Junquè et al., 2018).

PCBs were detected in all the analysed species. The mean values of Σ PCBs ranged from 28.7 ng g⁻¹ (*Diplodus* spp.) to 241.7 ng g⁻¹ (*S. sphyraena*). The Σ NDL-PCBs showed variable concentrations among the analysed species, with a mean value above the limit of 75 ng g⁻¹ (European Commission, 2011) measured in *S. sphyraena* (168 ng g⁻¹) and *M. barbatus* (88.2 ng g⁻¹) collected in 2017 and *Mullus* spp. sampled in 2013 (96.2 ng g⁻¹). All the analysed species sampled in 2013 and 2017, exceeded Σ 7PCBs threshold limit (10 ng g⁻¹) with the lowest and highest value measured in *T. lucerna* (19.0 ng g⁻¹) and *S. sphyraena* (180.9 ng g⁻¹), respectively (Tab. 3); this value represents the

300 limit above which effects on marine organisms might be expected according to the Ecotoxicological
 301 Assessment Criteria (EACs) of the OSPAR Convention (Campillo et al., 2017). A direct
 302 comparison of the $\Sigma 7$ PCBs with a previous study (Ferrante et al., 2007) on various edible species
 303 sampled in the Gulf of Naples (Southern Tyrrhenian Sea) showed systematically higher
 304 concentrations in samples from Augusta Bay.

305 The highly-chlorinated congeners CB153 and CB138 (hexa-chlorinated) and CB180 (epita-
 306 chlorinated) exhibited the higher abundance in biota (Tab. 3) and dominated the PCB group in all
 307 the analysed samples, thus mirroring results from other studies (Miao et al., 2000; Solé et al., 2000;
 308 Green and Knutzen, 2003; Castro-Jimenez et al., 2008; Storelli et al., 2009; Scarpato et al., 2010;
 309 Xia et al., 2012; Suarez et al., 2013; Herceg-Romanic et al., 2014; Mohebbi- Nozar et al., 2014;
 310 Kampire et al., 2015; Batang et al., 2016). The major occurrence of these congeners is related to the
 311 high degree of chlorination on the aromatic rings which results in a lower degradation rate by
 312 organisms and a potential bioaccumulation (Jönsson et al., 2003; Storelli et al., 2009). Specific
 313 differences in PCBs content could reflect the dissimilar behaviour and feeding patterns, trophic
 314 levels, physiological status of organisms or metabolic detoxification capacity (Ashley et al., 2003)
 315 typical of the studied species. In particular, *S. sphyraena*, which is an active predatory fish,
 316 occupying a high level in the trophic chain (Premolatha and Manojkumar, 1990), showed the
 317 highest concentration of PCBs with respect to other species. Also, the *Mullus* species exhibited
 318 higher values of Σ PCBs most likely related to their ecological features (bottom-feeder) and
 319 consequently greater probability of exposure to PCBs from sediments.

320 PAHs were measured only in organisms sampled in 2017, revealing differences among the analysed
 321 species both as total content and individual PAHs (Tab. 3). Finfish and shellfish showed a major
 322 content of low molecular weight (LMW) PAHs, while high molecular weight (HMW) PAHs were
 323 all <LOD (0.8 ng g⁻¹). The Σ PAHs ranged between 9.22 ng g⁻¹ (*S. sphyraena*) and 22.0 ng g⁻¹
 324 (*Sepia* spp.). The Benzo-a-pyrene (BaP), used as reference of PAHs presence in seafood (Reg. EC

1881/2006 as amended), was <LOD in all finfish and shellfish except in mussels. The latter exceed threshold limit of Benzo-a-pyrene established for molluscs (5 ng g⁻¹; Legislative Decree 172/2015). The OCs pollutants in transplanted mussels showed higher values than those measured in the control samples, suggesting significant bioaccumulation (Tab. 3). HCB detected in transplanted mussel samples revealed differences between the two sites, with always higher HCB values in samples from the southern sites over both the sampling periods (S1 and S2), with levels up to 7 times higher (1.91 and 3.15 ng g⁻¹, respectively) than those measured in the northern site (0.44 and 0.41 ng g⁻¹) (Tab. 3). Recorded values, compared to those from other Mediterranean areas (Ferrante et al., 2007), resulted higher and suggested that mussels from Augusta Bay were exposed to high levels of this pollutant.

The Σ PCBs in S1 mussels (35.2 ng g⁻¹) was >3 times than values measured in N1 organisms (Σ PCBs= 11.66 ng g⁻¹). A higher content of Σ PCBs, compared to N2 (27.1 ng g⁻¹), was confirmed in S2 mussels (39.3 ng g⁻¹). Furthermore, Σ 7PCBs resulted above the OSPAR limit in all samples except in N1 mussels. The analysis of PCB congeners showed that the penta (101, 118), hexa (138-153) and epta-chlorinated PCBs (180) were generally more abundant than the less chlorinated forms (28 and 52) (Tab. 3) reflecting previous data reported by Ferrante et al. (2007), Beiras et al. (2012) and Campillo et al. (2017).

Also, PAHs concentrations were higher in transplanted mussels (Σ PAHs ranged between 34.2 and 76.4 ng g⁻¹) than in the control sample (Σ PAHs= 25.0 ng g⁻¹) with B(a)P showed values above the EQS (5 ng g⁻¹) (Tab. 3). In particular, S1 and S2 organisms exhibited a marked increase of B(a)P (11.3 and 11.02 ng g⁻¹, respectively) in relation to the initial value of control samples (4.3 ng g⁻¹). All samples showed values above the limit of 30 ng g⁻¹ for Fluoranthene (Tab. 3).

347

348 **Discussion and conclusive remarks**

349 Recent studies revealed a systematic correlation between PCBs in biota and in nearly un-weathered

350 PCB mixtures in sediments (Rodenburg et al., 2015, 2019). Several investigations (e.g., Wiegel
351 and Wu, 2000; Bedard et al., 2005; Zanaroli et al., 2015; Praveckova et al., 2016) evidenced that
352 dehalogenation processes affect PCBs under strictly anaerobic conditions and low redox potential
353 and thus changing the congener patterns from highly to low chlorinated congeners.

354 The PCBs composition in surface sediments appears dominated by the higher chlorinated penta,
355 hexa and hepta CBs, accounting for >60% of Σ PCBs in all samples (Fig. 3). As previously
356 mentioned, differences in Σ PCBs content measured in the various species, primarily depend on a
357 number of different ecological features (Ashley et al., 2003; Arnot and Gobas, 2006; Martinez-Silva
358 et al., 2018) and might reflect a non-homogeneous proportion of the various congeners.

359 Nonetheless, the compositional patterns of the PCB congeners in the studied fishes appear
360 reasonably comparable (Fig. 3), and their distribution mode in finfish and shellfish maintains a
361 systematic order of concentration: hexa->penta->hepta->tetra->tri-CBs, contributing for 58%, 24%,
362 21%, 4.5% and 0.3% (Fig. 3). Particularly, the major contribution of penta and hexa-CBs is in
363 excellent agreement with data reported for biota by other authors (Naso et al., 2005; Ferrante et al.,
364 2007; Howell et al., 2008; Pan et al., 2016; Habibullah-Al-Mamun et al., 2019). The mussels
365 showed a similar profile although with a minor percentage of epta-CBs accounting for 5.1-8.5% of
366 Σ PCBs (Fig. 3).

367 The composition pattern of the 7 PCB congeners (28, 52, 101, 118, 138, 153 and 180) in all samples
368 was explored by cluster analysis (Fig. 4), using the elbow method to identify the optimal number of
369 groups primarily driving the high variability in the studied data population. Except for *Diplodus*
370 spp., all the finfish and shellfish samples appear strictly grouped in the cluster together with
371 sediment samples from the A9 core, while all the mussel samples and the other sediment samples
372 (core A3, A7 and A11) grouped in two separate clusters (Fig. 4).

373 The statistically significant relationship between sediments from the A9 core and finfish and
374 shellfish suggests a first-order, although robust, insight on the intimate link between the abiotic and

375 biotic compartments and reasonably supports in the study area, the use of the PCB congeners as
376 potential source fingerprint for these contaminants. Indeed, sediments from the A9 core showed
377 highly comparable percentages of high-chlorinated congeners (CB 153, 138 and 180) whereas
378 mussels (native and transplanted) and sediments from the A3, A7 and A11 cores evidence different
379 PCB patterns. Although the period of permanence of the organisms in the area of the A9 core could
380 not be precisely assessed, our results suggest that this zone would represent a primary
381 contamination ‘hot spot’ area for Augusta Bay. Differently, the other sediment cores, characterized
382 by different and relatively lower PCBs concentration, might represent secondary sources of these
383 contaminants to the biotic compartment.

384 Analogue pieces of evidence, supporting transfer of contaminants from sediment to biota, again
385 emerge from the analysis of the distribution patterns of specific PAHs, which in finfish and
386 shellfish are commonly dominated by LMW hydrocarbons, in particular Fluoranthene and Pyrene.
387 Indeed, these two molecules have a large range of stability and are good indicators of
388 thermodynamic vs. kinetic processes (Soclo et al., 2000). The Fluoranthene/Pyrene has been
389 frequently applied to identify and characterize sources of PAHs in the marine environment (Magi et
390 al., 2002). On the other side, the distribution pattern of PAHs in mussels has been reported to reflect
391 accumulation of bioavailable fraction from the water column (Baumard et al., 1998) with
392 preferential accumulation of LMW PAHs (Varanasi and Gmur, 1981; Broman et al., 1990). The
393 dotchart in figure 5 evidence a significant correlation between biota and sediments, with
394 comparable Fluoranthene/Pyrene ratios in finfish, shellfish and A9 sediments. Worthy to note, also
395 Fluoranthene/Pyrene measured in N2, S1 and S2 mussels which seem to mirror a potential impact
396 from the A9 sediments. This could primarily suggest an effect of that ‘hot-spot’ area in terms of
397 resuspension and re-distribution of highly contaminated particles on a relatively wider distance
398 within the study area (see Denaro et al., 2020 for a complete modelling discussion on the dynamic
399 within Augusta Bay). Thus, if PCBs seem to offer a primary fingerprint to follow sediment-to-biota

transfer of contaminants, the combined use of specific PAHs patterns convincingly supports the hypotheses of dynamic transfer of those organics in the various environmental compartments. Particularly, a clear ‘hot-spot’ effect of the A9 sediments on the investigated biotic compartment can be also documented taking into account the Hg and HCB concentration patterns mirroring coherent higher concentrations in the analysed benthic fish. Conclusively, the ensemble of achieved results suggests that combined information from a wide spectrum of contaminants provides a consistent fingerprint to trace their transfer from sediments to biotic compartments and thus supports specific remediation decisions and sediment mitigation strategies. Despite the statistical robustness, the obtained results could primarily reflect specific dynamics of local biogeochemistry and a critical role of local effects on their distribution patterns, mainly at the abiotic and biological marine interfaces. However, these promising results encourage to the use of PCB congeners and pattern of pollutants as fingerprint of exposure pathways from marine sediments to biota in other marine environmental contexts.

413

414 **Acknowledgements**

This work is part of the CISAS (MIUR-CNR) project, funded by the Italian Ministry of Education, University and Research. The authors wish to thank the colleagues involved in the scientific collaboration among the National Research Council of Italy, Carmelo Buscaino and Gaspare Buffa for their support in samples collection during the CNR oceanographic surveys and all the laboratory personnel of IAS-CNR of Capo Granitola.

420

421 **References**

Amodio-Cocchieri, R., Amoroso, S., Arnese, A., Cirillo, T., Montuori, P., Triassi, M., 2003. Pollution by mercury, arsenic, lead, chromium, cadmium, and polycyclic aromatic

424 hydrocarbons of fish and mussels from the Gulf of Naples, Italy. *Environ. Contam. Tox.* 71,
425 551-560.

426 Arnot, J.A., Gobas, F.A.P.C., 2006. A review of bioconcentration factor (BCF) and
427 bioaccumulation factor (BAF) assessments for organic chemicals in aquatic organisms.
428 *Environ. Rev.* 14, 257-297.

429 Ashley, J. T., Horwitz, R., Steinbacher, J. C., Ruppel, B., 2003. A comparison of congeneric PCB
430 patterns in American eels and striped bass from the Hudson and Delaware River Estuaries.
431 *Mar. Pollut. Bull.* 46, 1294-1308.

432 ATSDR, Agency for Toxic Substances and Disease Registry, 2000. Toxicological Profile for
433 Selected Polychlorinated Biphenyls. U.S. Department of Health and Human Services, Public
434 Health Service, Atlanta, GA.

435 Ausili, A., Gabellini, M., Cammarata, G., Fattorini, D., Benedetti, M., Pisanelli, B., Gorbi, S.,
436 Regoli, F., 2008. Ecotoxicological and human health risk in a petrochemical district of
437 southern Italy. *Mar. Environ. Res.* 66, 215-217.

438 Bagnato, E., Sprovieri, M., Barra, M., Bitetto, M., Bonsignore, M., Calabrese, S., Di Stefano, V.,
439 Oliveri, E., Parello, F., Mazzola, S., 2013. The sea-air exchange of mercury (Hg) in the
440 marine boundary layer of the Augusta basin (southern Italy): Concentrations and evasion
441 flux. *Chemosphere* 93, 2024–2032, <https://doi.org/10.1016/j.chemosphere.2013.07.025>,
442 2013.

443 Bailey, R.E., 2001. Global hexachlorobenzene emissions. *Chemosphere* 43:167–82.

444 Barber, J.L., Sweetman, A.J., van Wijk, D., Jones, K.C., 2005. Hexachlorobenzene in the global
445 environment: emissions, levels, distribution, trends and processes. *Sci. Total. Environ.* 349,
446 1–44.

447 Baskaran, M. (2011). *Handbook of Environmental Isotope Geochemistry*. ISBN 978-3-642-10636-1
448 (Printed in 2 volumes) DOI 10.1007/ 978-3-642-10637-8. Springer Heidelberg Dordrecht
449 London New York.

450 Batang, Z.B., Alikunhi, N., Gochfeld, M., Burger, J., Al-Jahdali, R., Al-Jahdali, H., Aziz, M.A., Al-
451 Jebreen, D., Al-Suwailem, A., 2016. Congener-specific levels and patterns of
452 polychlorinated biphenyls in edible fish tissue from the Central Red Sea coast of Saudi
453 Arabia. *Sci. Total Environ.* 572, 915–925.

454 Baumard, P., Budzinski, H., Michcon, Q., Garrigues, P., Burgeot, T., and Bellocq, J., 1998. Origin
455 and bioavailability of PAHs in the Mediterranean Sea from mussel and sediment records.
456 *Estuarine, Coastal and Shelf Science* 47, 77-90.

457 Bedard D.L., Pohl E.A., Bailey J.J., Murphy A., 2005. Characterization of the PCB substrate range
458 of microbial dechlorination process LP. *Environ Sci Technol.* 39, 6831–8.

459 Bellas, J., Gonz alez-Quijano, A., Vaamonde, A., Fumega, J., Soriano, J.A., Gonza lez, J.J., 2011.
460 PCBs in wild mussels (*Mytilus galloprovincialis*) from the N-NW Spanish coast: current
461 levels and long-term trends during the period 1991-2009. *Chemosphere* 85 (3), 533e541.

462 Bellucci, L.G., Giuliani, S., Romano, S., Albertazzi, S., Mugnai, C., Frignani, M., 2012. An
463 integrated approach to the assessment of pollutant delivery chronologies to impacted areas:
464 Hg in the Augusta Bay (Italy). *Environ. Sci. Technol.* 46, 2040-2046.

465 Berrojalbiz, N., Castro-Jiménez, J., Mariani, G., Wollgast, J., Hanke, G., & Dachs, J., 2014.
466 Atmospheric occurrence, transport and deposition of poly- chlorinated biphenyls and
467 hexachlorobenzene in the Mediterranean and black seas. *Atmos. Chem. and Phys.* 14, 8947–
468 8959. <https://doi.org/10.5194/acp-14-8947-2014>.

- 469 Beyer, J., Green, N., Brooks, S., Allan, I.J., Ruus, A., Gomes, T., Bråte, I.L., Schøyen, M., 2017.
470 Blue mussels (*Mytilus edulis* spp.) as sentinel organisms in coastal pollution monitoring: a
471 review. *Mar. Environ. Res.* 130, 338–365.
- 472 Bonsignore, M., Manta, D., Salvagio, Oliveri, E., Sprovieri, M., Basilone, G., Bonanno, A., Falco,
473 F., Traina, A., Mazzola, S., 2013. Mercury in fishes from Augusta Bay (southern Italy): risk
474 assessment and health implication. *Food Chem. Toxicol.* 56, 184-194.
- 475 Bonsignore, M., Salvagio Manta, D., Barsanti, M., Conte, F., Delbono, I., Horvat, M., Quinci, E.
476 M., Schirone, A., Shlyapnikov, Y., Sprovieri, M., 2020. Mercury isotope signatures in
477 sediments and marine organisms as tracers of historical industrial pollution. *Chemosphere*
478 258,127435.
- 479 Bonsignore, M., Tamburrino, S., Oliveri, E., Marchetti, A., Durante, D., Berni, A., Quinci, E. M.,
480 Sprovieri, M., 2015. Tracing mercury pathways in Augusta Bay (southern Italy) by total
481 concentration and isotope determination. *Environ. Pollut.*, 205, 178-185.
- 482 Broman, D., Näuf, C., Lundbergh, I., Zebühr, Y., 1990. An in situ study on the distribution,
483 biotransformation and flux of polycyclic aromatic hydrocarbons (PAHs) in an aquatic food
484 chain (seston-*Mytilus edulis* L. *Somateria mollissima* L.) from the baltic: An
485 ecotoxicological perspective. *Environ. Toxicol. and Chem.* 9 (4), pp. 429-442.
- 486 Campillo, J.A., Fernández, B., García, V., Benedicto, J., León, V.M., 2017. Levels and temporal
487 trends of organochlorine contaminants in mussels from Spanish Mediterranean waters.
488 *Chemosphere.* 182:584–94. doi.org/10.1016/j.chemosphere.2017.05.025.
- 489 Cardellicchio, N., Annicchiarico, C., Assennato, G., Blonda, M., Di Leo, A. et al., 2010.
490 Preliminary results of pollut- ants biomonitoring in coastal marine and transitional wa- ters
491 of Apulia Region (Southern Italy). *Fresen. Environ. Bull.*, 19 (9), 1841-1847.
- 492 Caricato, R., Giordano, M.E., Schettino, T., Maisano, M., Mauceri, A., Giannetto, A., Cappello, T.,
493 Parrino, V., Ancora, S., Caliani, I., Bianchi, N., Leonzio, C., Mancini, G., Cappello, S.,
494 Fasulo, S., Lionetto, M.G., 2019. Carbonic anhydrase integrated into a multimarker
495 approach for the detection of the stress status induced by pollution exposure in *Mytilus*
496 *galloprovincialis*: a field case study. *Sci. Tot. Environ.*, 690, 140-150.
- 497 Castro-Jiménez, J., Deviller, G., Ghiani, M., Loos, R., Mariani, G., Skejo, H., Umlauf Wollgast,
498 J.G., Laugier, T., Héas-Moisán, K., Léauté, F., Munschy, C., Tixier, C., Tronczyński, J.,
499 2008. PCDD/F and PCB multi-media ambient concentrations, congener patterns and
500 occurrence in a Mediterranean coastal lagoon (Etang de Thau, France). *Environ. Pollut.* 156,
501 123–135.
- 502 Courtney, K.D., 1979. Hexachlorobenzene (HCB): A review. *Environ. Res.*, 20 (2), pp. 225-266.
- 503 Crinnion, W.J., 2011. Polychlorinated biphenyls: persistent pollutants with immunological,
504 neurological and endocrinological consequences. *Altern. Med. Rev.*, 16, 5–13.
- 505 Croudace, I.W., Romano, E., Ausili, A., Bergamin, L., Rothwell, G., 2015. X-ray core scanners as
506 an environmental forensic tool: a case study of polluted harbor sediment (Augusta Bay,
507 Sicily). In: *Micro-XRF Studies of sediment cores*, chapter 15. Ian W. Croudace and R. Guy
508 Rothwell (Eds.), Springer.
- 509 Denaro, G., Salvagio Manta, D., Borri, A., Bonsignore, M., Valenti, D., Quinci, E., Cucco, A.,
510 Spagnolo, B., Sprovieri, M., De Gaetano, A., 2020. HR3DHG version 1: modeling the
511 spatiotemporal dynamics of mercury in the Augusta Bay (southern Italy). *Geosci. Molel*
512 *Dev.* 13, 2073-2093.
- 513 Di Bella, C., Traina, A., Giosuè, C., Carpintieri, D., Lo Dico, G.M., Bellante, A., Del Core, M.,
514 Falco, F., Gherardi, S., Uccello, M.M., Ferrantelli, V., 2020. Heavy metals and PAHs in

515 meat, milk, and seafood from Augusta area (Southern Italy): contamination levels, dietary
516 intake, and human exposure assessment. *Front. Public Health* 8:273.
517 doi.org/10.3389/fpubh.2020.00273.

518 Djomo, J. E., Garrigues, P., & Narbonne, J. F., 1996. Uptake and depuration of polycyclic aromatic
519 hydrocarbons from sediment by the zebrafish (*Bracydanio rerio*). *Environ. Toxicol. Chem.*
520 15, 1177-1181.

521 Domingo, J.L., Bocio, A., 2007. Levels of PCDD/PCDFs and PCBs in edible marine species and
522 human intake: a literature review. *Environ. Int.* 33, 397–405.

523 EFSA (European Food Safety Authority), 2010. Results of the monitoring of non- dioxin-like PCBs
524 in food and feed. *EFSA J.* 8, 35.

525 ENVIRON, 2008. Appendix A Sediment Investigation Activities Performed by ENVIRON
526 Summer and Fall. Augusta Bay, Sicily, Italy.

527 European Commission, 2011. Commission Regulation (EU) No 1259/2011 of 2 December 2011
528 amending Regulation (EC) No 1881/2006 as regards maximum levels for dioxins, dioxin-
529 like PCBs and non-dioxin-like PCBs in foodstuffs. In: *Official Journal of the European*
530 *Union, L*, 320/18-23.

531 Farrington, J.W., Tripp, B.W., Tanabe, S., Subramanian, A., Sericano, J.L., Wade, T.L., Knap,
532 A.H., Edward, D., 2016. Goldberg’s proposal of “the mussel watch”: reflections after 40
533 years *Mar. Pollut. Bull.* 110 (1), 501-510.

534 Ferrante, M.C., Cirillo, T., Naso, B., Clausi, M.T., Lucisano, A., Amodio Cocchieri, R., 2007.
535 Polychlorinated biphenyls and organochlorine pesticides in seafood from the Gulf of Naples
536 (Italy). *J. Food Protect.* 70 (3), 706–715.

537 García-Flor, N., Dachs, J., Bayona, J.M., Albaige s, J., 2009. Surface waters are a source of
538 polychlorinated biphenyls to the coastal atmosphere of the North-Western Mediterranean
539 Sea. *Chemosphere* 75, 1144e1152.

540 Gobas, F.A.P.C., Arnot, J.A., 2010. Food web bioaccumulation model for polychlorinated
541 biphenyls in san Francisco Bay, California, USA. *Environ. Toxicol. Chem.* 29, 1385e1395.

542 Green, N.W., Knutzen, J., 2003. Organohalogens and metals in marine fish and mussels and some
543 relationships to biological variables at reference localities in Norway. *Mar. Pollut. Bull.* 46,
544 362–374.

545 Habibullah-Al-Mamun, M., Ahmed, M.K., Islam, M.S., Hossain, A., Tokumura, M., Masunaga, S.,
546 2019. Polychlorinated biphenyls (PCBs) in commonly consumed sea- food from the coastal
547 area of Bangladesh: occurrence, distribution, and human health implications. *Environ. Sci.*
548 *Pollut. Res.* 26, 1355–1369.

549 Herceg-Romanic, S., Kljakovic-Gaspic, Z., Klincic, D., Ujevic, I., 2014. Distribution of persistent
550 organic pollutants (POPs) in cultured mussels from the Croatian coast of the Adriatic Sea.
551 *Chemosphere* 114, 69-75.

552 Howell, N.L., Suarez, M.P., Rifai, H.S., Koenig, L., 2008. Concentrations of polychlorinated
553 biphenyls (PCBs) in water, sediment, and aquatic biota in the Houston Ship Channel, Texas.
554 *Chemosphere*, 70 593-606.

555 IARC, International Agency for Research on Cancer, 2016. IARC Monographs on Polychlorinated
556 Biphenyls and Polybrominated Biphenyls. Vol. 107. Lyon (France).

557 Ausili, A., Cammarata, G., Gabellini, M., Regoli, F., 2003. Relazione consulenza tecnica-incarico
558 del 10-04-03. PPN 8388-01-RG-SR, 75pp.

ICRAM, 2005. Elaborazione e valutazione dei risultati della caratterizzazione ambientale della Rada di Augusta-aree prioritarie ai fini della messa in sicurezza di emergenza # BoI-Pr-SI-GP-Rada di Augusta- 01.04. Technical report, 92p.

ICRAM, 2008. Progetto preliminare di bonifica della Rada di Augusta inclusa nel sito di bonifica di interesse nazionale di Priolo. Fase I e II. Elaborazione definitiva # BoI-Pr-SI-PR-Rada di Augusta-03.22. Technical report, 253 pp.

Ipolyi, I., Massanisso, P., Sposato, S., Fodor, P., Morabito, R., 2004. Concentration levels of total and methylmercury in mussel samples collected along the coasts of Sardinia Island (Italy). *Anal. Chim. Acta* 505 (1), 145-151.

JECFA, 2018. Evaluations of the Joint FAO/WHO Expert Committee on Food Additives.

Jepson, P.D., Deaville, R., Barber, J.L., Aguilar, A., Borrell, A., Murphy, S., Barry, J., Brownlow, A., Barnett, J., Berrow, S., Cunningham, A.A., Davison, N.J., Ten Doeschate, M., Esteban, R., Ferreira, M., Foote, A.D., Genov, T., Gimenez, J., Loveridge, J., Llavona, A., Martin, V., Maxwell, D.L., Papachlimitzou, A., Penrose, R., Perkins, M.W., Smith, B., de Stephanis, R., Tregenza, N., Verborgh, P., Fernandez, A., Law, R.J., 2016. PCB pollution continues to impact populations of orcas and other dolphins in European waters. *Sci. Rep.* 6. <http://dx.doi.org/10.1038/srep18573>.

Jönsson, A., Gustafsson, Ö., Axelman, J., Sundberg, H., 2003. Global accounting of PCBs in the continental shelf sediments. *Environ. Sci. Technol.* 37, 245-255.

Junque, E., Gari, M., Llull, R.M., Grimalt, J.O., 2018. Drivers of the accumulation of mercury and organochlorine pollutants in Mediterranean lean fish and dietary significance. *Sci. Total Environ.* 634, 170–180.

Kampire, E., Rubidge, G., Adams, B., 2015. Distribution of polychlorinated biphenyl residues in sediments and blue mussels (*Mytilus galloprovincialis*) from Port Elizabeth harbour, South Africa. *Mar. Pollut. Bull.*, 91, 173–179.

Keith, L.H., Telliard, W.A., 1979. Priority pollutants. A perspective review, *Environ.Sci. Technol.* 13, 416–423. ^[1]_{SEP}

Kljaković-Gašpić, Z., Herceg-Romanić, S., Kožul, D., Veža, J., 2010. Biomonitoring of organochlorine compounds and trace metals along the eastern Adriatic coast (Croatia) using *Mytilus galloprovincialis*. *Mar. Pollut. Bull.* 60, 1879–1889.

Landrum, P.F. and Robbins, J.A., 1990. Bioavailability of sediment-associated contaminants to benthic invertebrates. In: R. Baudo, J.P. Giesy, and H. Munteau (eds) *Sediments: Chemistry and Toxicity of In-Place Pollutants*, pp. 237-263. Lewis Publishers, Ann Arbor.

Licata, P., Trombetta, D., Cristani, M., Naccari, C., Martino, D., Calò, M., Naccari, F., 2004. Heavy metals in liver and muscle of Bluefin tuna (*Thunnus thunnus*) caught in the strait of Messina (Sicily, Italy). *Environ. Monit. Assess.*, 30, 1–10.

Martinez-Silva, M.A., Audet, C., Winkler, G., et al., 2018. Prey quality impact on the feeding behavior and lipid composition of winter flounder (*Pseudopleuronectes americanus*) larvae[J]. *Aquac. Fish.* 3 (4), 145-155.

McLeod, A.M., Arnot, J.A., Borga, K., Selck, H., Kashian, D.R., Krause, A., Paterson, G., Haffner, G.D., Drouillard, K.G., 2015. Quantifying uncertainty in the trophic magnification factor related to spatial movements of organisms in a food web. *Integr. Environ. Assess. Manag.* 11(2), 306–318.

Meijer S.N. Meijer, W.A. Ockenden, A. Sweetman, K. Breivik, J.O. Grimalt, K.C., 2003. Jones Global distribution and budget of PCBs and HCB in background surface soils: implications or sources and environmental processes. *Environ. Sci. & Technol.* 37, pp. 667-672.

Miao, X.S., Swenson, C. Woodward, L.A., Li, Q.X., 2000. Distribution of polychlorinated

605 biphenyls in marine species from French Frigate Shoals, North Pacific Ocean. *Sci Total*
606 *Environ.* 257(1), 17–28.

607 Minh, N.H., Minh, T.B., Kajiwara, N., Kunisue, T., Iwata, H., Viet, P.H., Tu, N.P.C., Tuyen, B.C.,
608 Tanabe, S., 2007. Pollution sources and occurrences of selected persistent organic pollutants
609 (POPs) in sediments of the Mekong River delta, South Vietnam. *Chemosphere* 67 (9), 1794–
610 1801.

611 Mohebbi-Nozar, S.L., Ismail, W.R., Zakaria, M.P., Mortazavi, M.S., Zahed, M.A., Jahanlu, A.,
612 2014. Health risk of PCBs and DDTs in seafood from southern Iran. *Hum. Ecol. Risk*
613 *Assess.* 20(5), 1164–1176.

614 Naso, B., Perrone, D., Ferrante, M.C., Bilancion, M., Lucisano, A., 2005. Persistent organic
615 pollutants in edible marine species from the Gulf of Naples, Southern Italy. *Sci. Total*
616 *Environ.* 343, 83–95.

617 Neff, J.M., 1979. Polycyclic aromatic hydrocarbons in the aquatic environment, sources, fates and
618 biological effects. Applied Science Publishers Ltd., Essex, England. Olenycz, M.,
619 Sokołowski, A., Niewin ska, A., Wołowicz, M., Namie snik, J., Hummel, H., Jansen, J.,
620 2015. Comparison of PCBs and PAHs levels in European coastal waters using mussels from
621 the *Mytilus edulis* complex as biomonitors. *Oceanologia*, 57 (2), 196–211.

622 Orecchio, S., Polizzotto G., 2013. Fractionation of mercury in sediments during draining of Augusta
623 (Italy) coastal area by modified Tessier method. *Microchem. J.* 110, 452–457.

624 Pan, H., Geng, J., Qin, Y., Tou, F., Zhou, J., Liu, M., Yang, Y., 2016. PCBs and OCPs in fish along
625 coastal fisheries in China: distribution and health risk assessment. *Mar. Pollut. Bull.* 111 (1–
626 2), 483–487.

627 Porte, C., Albaigés, J., 1993. Bioaccumulation patterns of hydrocarbons and polychlorinated
628 biphenyls in bivalves, crustaceans, and fishes. *Arch. Environ. Contam. Toxicol.* 26, 273–
629 281.

630 Praveckova M., Brennerova M.V., Holliger C., De Alencastro F., Rossi P., 2016. Indirect Evidence
631 Link PCB Dehalogenation with Geobacteraceae in Anaerobic Sediment-Free Microcosms.
632 *Front. Microbiol.* 7, 933. doi: 10.3389/fmicb.2016.00933

633 Premolatha, P., Manojkumar, P.P., 1990. Some biological aspects of two species of barracudas from
634 the South West coast of India. *Indian J. Fish.* 37 (4), 289–295.

635 R Core Team, 2020. R: A language and environment for statistical computing. R Foundation for
636 Statistical Computing, Vienna, Austria. <https://www.R-project.org/>.

637 Regoli, F., Winston, G.W., Mastrangelo, V., Bompadre, S., 1998. Total oxyradical scavenging
638 capacity in mussel *Mytilus* sp. as a new index of biological resistance to oxidative stress.
639 *Chemosphere* 37, 2773–2783.

640 Rodenburg, L. A., and Delistraty, D. A., 2019. Alterations in fingerprints of polychlorinated
641 biphenyls in benthic biota at the Portland Harbor Superfund Site (Oregon, USA) suggest
642 metabolism. *Chemosphere* 223, 74–82. doi: 10.1016/j.chemosphere.2019.02.039

643 Rodenburg, L.A., Delistraty, D., Meng, Q., 2015. Polychlorinated biphenyl congener patterns in
644 fish near the Hanford site (Washington state, USA). *Environ. Sci. Technol.* 49, 2767–2775.

645 Rodenburg, L.A., Leidos, 2017. Green-duwamish River Watershed PCB Congener Study: Phase 2
646 Source Evaluation. Seattle, WA. Rodríguez-Hernández, A., Camacho, M., Henríquez-
647 Hernández, L.A., Boada, L.D., Ruiz-Suárez, N., Valerón, P.F., Almeida, M.G., Zaccaroni,
648 A., Zumbado, M., Luzardo, O.P., 2016. Assessment of human health hazards associated
649 with the dietary exposure to organic and inorganic contaminants through the consumption of

650 fishery products in Spain. *Sci. Total Environ.* 557-558, 808–818. Sahu, S.K., Ajmal, P.Y.,
651 Pandit, G.G., Puranik, V.D., 2009. Vertical distribution of polychlorinated biphenyl
652 congeners in sediment core from Thane Creek area of Mumbai, India. *J. Hazard. Mater.* 164
653 (2–3), 1573–1579.

654 Salvagio Manta, D., Bonsignore, M., Oliveri, E., Barra, M., Tranchida, G., Giaramita, L., Mazzola,
655 S., Srovieri, M., 2016. Fluxes and the mass balance of mercury in Augusta Bay (Sicily,
656 southern Italy). *Estuar. Coast. Shelf S.* 181, 134-143.

657 Sanchez-Avila, J., Tauler, R., Lacorte, S., 2012. Organic micropollutants in coastal waters from
658 NW Mediterranean Sea: sources distribution and potential risk. *Environ. Int.* 46, 50-62.

659 Scarpato, A., Romanelli, G., Galgani, F., Andral, B., Amici, M., Giordano, P., Caixach, J., Calvo,
660 M., Campillo, J.A., Albadalejo, J.B., Cento, A., Benbrahim, S., Sammari, C., Deudero, S.,
661 Boulahdid, M., Giovanardi, F., 2010. Western Mediterranean coastal waters - monitoring
662 PCBs and pesticides accumulation in *Mytilus galloprovincialis* by active mussel watching:
663 the Mytilos Project. *J. Environ. Monit.* 12 (4), 924-935. Selck, H., Drouillard, K., Eisenreich,
664 K., Koelmans, A.A., Palmqvist, A., Ruus, A., Salvito, D., Schultz, I., Stewart, R., Weisbrod,
665 A. et al., 2012. Explaining differences between bioaccumulation measurements in laboratory
666 and field data through use of a probabilistic modeling approach. *Integr. Environ. Assess.*
667 *Manag.* 8 (1), 42–63.

668 Signa, G., Mazzola, A., Doriana Tramati, C., Vizzini, S., 2017. Diet and habitat use influence Hg
669 and Cd transfer to fish and consequent biomagnification in a highly contaminated area:
670 Augusta Bay (Mediterranean Sea). *Environ. Pollut.*, 230, 394-404. Soclo H.H., Garrigues P.,
671 Ewald M., 2000. Origin of polycyclic aromatic hydrocarbons (PAHs) in coastal marine
672 sediments: case studies in Cotonou (Benin) and Aquitaine (France) areas. *Mar. Pollut. Bull.*
673 40, 387-396. Solé, M., Porte, C., Barcelo, D., Albaige s, J., 2000. Bivalves residue analysis
674 for the assessment of coastal pollution in the Ebro Delta (NW Mediterranean). *Mar. Pollut.*
675 *Bull.* 40 (9), 746-753.

676 Spada, L., Annicchiarico, C., Cardellicchio, N., Giandomenico, S., Di Leo, A., 2011. Mercury and
677 methylmercury concentrations in Mediterranean seafood and surface sediments, intake
678 evaluation and risk for consumers. *Int. J. Hyg. Environ. Health* 21, 418-426.

679 Spada, L., Annicchiarico, C., Cardellicchio, N., Giandomenico, S., Di Leo, A., 2013. Heavy metals
680 monitoring in mussels *Mytilus galloprovincialis* from the Apulian coasts (southern Italy).
681 *Mediterr. Mar. Sci.* 14, 99-108.

682 Spada, L., Annicchiarico, C., Cardellicchio, N., Giandomenico, S., Di Leo A., 2012. Mercury and
683 methylmercury concentrations in Mediterranean seafood and surface sediments, intake
684 evaluation and risk for consumers. *Int. J. Hyg. Environ. Health* 215 (3), 418-426.

685 Sprovieri M., Oliveri E., Di Leonardo R., Romano E., Ausili A., Gabellini M., Saggiomo V., Barra
686 Tranchida M., Bellanca A., Neri R., Budillon F., Mazzola S., 2011. The key role played by
687 the Augusta basin (southern Italy) in the mercury contamination of the Mediterranean Sea. *J.*
688 *Environ. Monitor.* 13, 1753-1760. Storelli, M.M., Losada, S., Marcotrigiano, G.O., Roosens,
689 L., Barone, G., Neels, H., Covaci, A., 2009. Polychlorinated biphenyl and organochlorine
690 pesticide contamination signatures in deep-sea fish from the Mediterranean Sea. *Environ.*
691 *Res.* 109 (7), 851–856.

692 Suarez, P., Ruiz, Y., Alinso, A., San Juan, F., 2013. Organochlorine compounds in mussels cultured
693 in the Ría of Vigo: accumulation and origin. *Chemosphere* 90, 7-19. Thomann, R.V.,
694 Connolly, J.P., Parkerton, T.F., 1992. An equilibrium model of organic chemical

695 accumulation in aquatic food webs with sediment interaction. Environ. Toxicol. Chem. 11
696 (5), 615–629.

697 Tomasello, B., Copat, C., Pulvirenti, V., Ferrito, V., Ferrante, M., Renis, M., Sciacca, S., Tigano,
698 C., 2012. Biogeochemical and bioaccumulation approaches for investigating marine
699 pollution using Mediterranean rainbow wrasse, *Coris julis* (Linnaeus 1978). Ecotoxicol.
700 Environ. Safe. 86, 168-175. Ueno, D., Takahashi, S., Tanaka, H., Subramanian, A.N.,
701 Fillmann, G., Nakata, H., Lam, P.K.S., Zheng, J., Muchtar, M. Prudente, M., Chung, K.H.,
702 Tanabe, S., 2003. Global pollution monitoring of PCBs and organochlorine pesticides using
703 skipjack tuna as a bioindicator. Arch. Environ. Contam. Toxicol. 45, 378–389.

704 UNEP, 2001. Stockholm Convention on Persistent Organic Pollutants (POPs), United Nations
705 Environment Programme UNEP/POPs/CONF/4

706 UNEP, 2017. Stockholm Convention on Persistent Organic Pollutants (POPs), United Nations
707 Environment Programme UNEP/POPS/POPRC.13/7

708 UNEP, 2019.. Stockholm Convention on Persistent Organic Pollutions (POPs), United Nation
709 Environment Programme (Accessed in 02/15/2020).

710 USEPA Method 7473, 2007. Mercury in Solids and Solutions by Thermal Decomposition,
711 Amalgamation, and Atomic Absorption Spectrophotometry, Part of Test Methods for
712 Evaluating Solid Waste, Physical/Chemical Methods. Varanasi, U., Gmur, D. J., 1981.
713 Hydrocarbons and metabolites in English sole (*Parophrys vetulus*) exposed simultaneously
714 to [3H]-benzo[a]pyrene and [14C]naphthalene in oil-contaminated sediment. Aquat.
715 Toxicol. 1, 49–67. Webster, L., Roose, P., Bersuder, P., Vorkamp, K., Kotterman, M.,
716 Haarich, M., 2013. Deter- mination of Polychlorinated Biphenyls (PCBs) in Sediment and
717 Biota. ICES Techniques in Marine Environmental Sciences No. 53, 18 pp.

718 Wiberg, P.L., Courtney, K., Harris, C.K., 2002. Desorption of p,p'-DDE from sediment during
719 resuspension events on the Palos Verdes Shelf, California: a modeling approach. Cont. Shelf
720 Res. 22, 1005–1023.

721 Wiegel, J., Wu, Q., 2000. Microbial reductive dehalogenation of polychlorinated biphenyls, FEMS
722 Microbiol Ecol. 32, 1–15.

723 Xia, C., Lam, J.C., Wu, X., Xie, Z., Lam, P.K., 2012. Polychlorinated biphenyls (PCBs) in marine
724 fishes from China: levels, distribution and risk assessment. Chemosphere 89 (8), 944–949.

725 Zanaroli, G., Negroni, A., Häggblom, M.M., Fava, F., 2015. Microbial dehalogenation of
726 organohalides in marine and estuarine environments. Current Opinion in Biotechnology 33,
727 287-295.

728

729

730 **Figure Captions**

731

732 **Figure 1:** Map and details of the sampling in Augusta Bay.

733

734 **Figure 2:** Box plot of the concentrations of Hg (a), HCB (b), Σ PCBs (c) and Σ PCBs NDL (d) in
735 sediment cores.

736

737 **Figure 3:** Relative contribution of PCB homologs (% composition) in sediments, seafood and
738 mussel samples.

739

740 **Figure 4:** Heatmap generated from hierarchical clustering analysis. The dendrograms of sample
741 clustering (on the left) and of congener clustering (on the top) were added. The colour bars inside
742 the graph indicate the different proportion of congeners (X axis) for each sample (Y axis). The rows
743 were splitted based on number of identified clusters.

744

745 **Figure 5:** Dotchart of Fluo/Py in sediments and marine organisms

746

747

	Specie	Category	Habitat	Total lenght range (med ±st.dv) cm	n. total individuals	n. pool
2013	<i>Mytilus galloprovincialis</i>	Molluscs	benthic	3.6-6 (4.60±0.58)	200	20
	<i>Sphyraena sphyraena</i>	Fish	pelagic	46-44 (45±1.41)	2	2
	<i>Mullus barbatus</i>	Fish	benthic	17.5-21 (18.7±1.14)	30	6
	<i>Pagellus spp.</i>	Fish	demersal	15-26 (20±3.25)	15	5
	<i>Sparus aurata</i>	Fish	demersal	21-24 (21.6±1.21)	9	3
	<i>Serranus cabrilla</i>	Fish	demersal	20	1	1
	<i>Diplodus spp.</i>	Fish	demersal	14.1-16.3 (15.1±1.1)	3	2
2017	<i>Sphyraena sphyraena</i>	Fish	pelagic	37.4	1	1
	<i>Trigla lucerna</i>	Fish	benthic	22.3-22.5	2	1
	<i>Mullus barbatus</i>	Fish	benthic	17.7-20.5 (18.8±1.1)	6	2
	<i>Pagellus spp.</i>	Fish	demersal	15.3-25.4 (18.9±2.7)	37	8
	<i>Diplodus spp.</i>	Fish	demersal	13.5-20 (16.8 ±3.2)	4	3
	<i>Penaeus kerathurus</i>	Crustacean	benthic	4.5-6.5 (5.2±0.3)	39	4
	<i>Sepia spp.</i>	Molluscs	bento-nectonic	9.9-15.5 (12.5±2.0)	7	6
	<i>Mytilus galloprovincialis N</i>	Molluscs	benthic	4.4-5.9 (5.20±1.06)	392	13
	<i>Mytilus galloprovincialis S</i>	Molluscs	benthic	5.0-5.4 (5.2±0.3)	284	10

750 **Table 1.** Characteristics, number of specimens and pool of marine organisms caught in the study area.
751

CORE	Unit	A3	A7	A9	A 11
		mean ±st.dv	mean ±st.dv	mean ±st.dv	mean ±st.dv
Hg	µg g ⁻¹	3.64±0.27	6.71±0.80	11.4 ±0.57	6.54±1.28
HCB	ng g ⁻¹	8.53 ± 2.06	25.3± 8.03	172.6 ± 87.6	22.8 ± 14.45
PCB 28	ng g ⁻¹	10.36 ± 6.66	2.26 ± 0.82	0.40 ± 0.12	9.65 ± 12.6
PCB 52	ng g ⁻¹	0.42 ± 0.53	<LOD	1.35 ± 0.30	0.56 ± 0.94
PCB 81	ng g ⁻¹	<LOD	<LOD	<LOD	<LOD
PCB 77	ng g ⁻¹	<LOD	<LOD	0.72±0.13	<LOD
PCB 101	ng g ⁻¹	<LOD	<LOD	5.22 ± 1.40	<LOD
PCB-118	ng g ⁻¹	6.90 ± 4.66	0.97 ± 0.77	3.79 ± 0.96	8.74 ± 16.9
PCB 114	ng g ⁻¹	<LOD	<LOD	<LOD	<LOD
PCB 123	ng g ⁻¹	<LOD	<LOD	11.2 ± 34.6	<LOD
PCB 153	ng g ⁻¹	10.4 ± 5.77	3.43±1.95	10.4 ± 1.07	16.2 ± 13.3
PCB 105	ng g ⁻¹	1.18 ± 1.88	<LOD	1.26 ± 0.80	1.24 ± 2.84
PCB 138	ng g ⁻¹	7.59 ± 7.11	2.83 ± 2.56	9.72 ± 1.53	17.3 ± 20.0
PCB 126	ng g ⁻¹	<LOD	<LOD	1.06 ± 0.24	<LOD
PCB 128+157	ng g ⁻¹	<LOD	0.16±0.20	0.62 ± 0.22	1.10 ± 1.21
PCB 156	ng g ⁻¹	<LOD	0.19 ± 0.29	0.58 ± 0.34	0.80 ± 1.65
PCB 167	ng g ⁻¹	<LOD	<LOD	1.04 ± 0.08	0.84 ± 1.75
PCB 180	ng g ⁻¹	0.81 ± 1.17	1.53 ± 1.58	7.89 ±1.45	4.30 ± 3.26
PCB 169	ng g ⁻¹	<LOD	<LOD	<LOD	<LOD
PCB 170	ng g ⁻¹	<LOD	0.34 ± 0.54	3.85 ± 0.36	2.35±2.03
PCB 189	ng g ⁻¹	<LOD	<LOD	0.25 ± 0.09	<LOD
Σ PCBs	ng g ⁻¹	38.7 ± 26.2	12.6 ± 6.7	59.5 ± 35.6	38.2 ± 19.9
Σ PCBs ¹	ng g ⁻¹	29.6 ± 26.6	11.8 ±6.50	41.9 ± 5.60	34.3 ± 16.2
Σ 7PCB	ng g ⁻¹	36.6 ± 24.4	11.2 ± 6.39	38.7 ± 5.20	32.2 ± 13.6
Phenanthrene	ng g ⁻¹	37.6 ± 8.52	26.4 ± 6.56	67.8 ± 23.3	27.7 ± 11.1
Anthracene	ng g ⁻¹	180 ± 68.9	186.8 ± 29.6	56.3 ± 14.6	91.7 ± 27.6
Fluoranthene	ng g ⁻¹	61.8 ± 32.2	35.3 ± 18.1	45.2 ± 24.7	27.5 ± 12.8
Pyrene	ng g ⁻¹	33.1 ± 13.7	24.5 ± 8.11	61.2 ± 22.3	25.4 ± 10.2
Benzo-a-Anthracene	ng g ⁻¹	34.2 ± 11.1	17.0 ± 6.18	26.4 ± 10.6	13.5 ± 3.94
Chrysene	ng g ⁻¹	38.5 ± 10.2	20.6 ± 6.72	48.2 ± 13.1	17.8 ± 4.99
Benzo-b-Fluoranthene	ng g ⁻¹	54.6 ± 20.4	32.5 ± 7.45	30.5 ± 8.51	14.4 ± 4.24
Benzo-k-Fluoranthene	ng g ⁻¹	33.6 ± 24.3	9.26 ± 2.54	10.3 ± 4.59	7.09 ± 2.03

Benzo-a-Pyrene	ng g ⁻¹	64.6 ± 20.1	29.3 ± 8.53	23.5 ± 6.94	20.5 ± 6.42
Indeno-123-cd-Pyrene	ng g ⁻¹	70.1 ± 25.5	19.5 ± 6.19	9.88 ± 5.95	13.1 ± 2.30
Dibenzo-ah-Anthracene	ng g ⁻¹	12.9 ± 6.30	1.87 ± 1.36	7.90 ± 4.29	2.91 ± 1.94
Benzo-ghi-Perylene	ng g ⁻¹	50.6 ± 21.7	17.6 ± 7.72	11.15 ± 5.30	15.7 ± 4.06
Σ PAHs	ng g ⁻¹	671.87 ± 125.97	420 ± 72.73	398 ± 103	277 ± 64.8

Values exceeding environmental quality standards (Dlgs. 172/2015) are indicated in bold

Σ PCBs¹: sum of PCBs congeners indicated by Dlgs.172/2015

Σ 7PCB: sum of CB 28+52+101+118+138+153+180

Table 2. Hg, HCB, PCBs and PAHs concentrations (d.w.) in sediment collected in the sampling site.

Parameters	Unit	Sampling 2013										Sampling 2017												
		<i>S. sphyraena</i>	<i>M. barbatus</i>	<i>Pagellus</i> spp.	<i>S. aurata</i>	<i>S. cabrilla</i>	<i>Diplodus</i> spp.	<i>M. galloprovincialis</i> wild	<i>S. sphyraena</i>	<i>T. lucerna</i>	<i>M. barbatus</i>	<i>Pagellus</i> spp.	<i>Diplodus</i> spp.	<i>P. kerathurus</i>	<i>Sepia</i> spp.	<i>M. gallorprovincialis</i>								
		med ±st.dv										med ±st.dv										CTRL	N1	S1
Hg	µg g ⁻¹	0.52 ±0.04 ^a	0.44 ± 0.20	0.53 ± 0.12	0.32 ± 0.09	0.77 ^a	0.503±0.137 ^a	0.2 ± 0.04 ^a	0.78 ^a	0.66 ^a	2.11;1.71 ^a	0.86±0.44	1.49±1.77 ^a	0.59±0.02 ^a	0.21 ± 0.08	0.01	0.02 ± 0.002 ^b	0.23 ± 0.02 ^b	0.12 ± 0.01 ^b	0.89 ^{a,b}				
HCB	ng g ⁻¹	n.d	1.36 ± 0.31	0.09 ± 0.02	0.60 ± 0.43	n.d	n.d	0.13 ± 0.00	1.40	<LOD	0.24;0.65	<LOD	<LOD	<LOD	<LOD	0.35	0.44	1.91	0.41	3.15				
PCB 28	ng g ⁻¹	n.d	0.36 ± 0.06	0.06 ± 0.01	0.16±0.09	n.d	n.d	0.15±0.00	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD				
PCB 52	ng g ⁻¹	n.d	6.31 ± 2.27	2.77 ± 0.63	4.16±2.32	n.d	n.d	4.95±1.54	2.03	<LOD	1.29;2.42	0.48 ± 0.50	<LOD	<LOD	<LOD	<LOD	<LOD	0.72	<LOD	<LOD				
PCB 81	ng g ⁻¹	n.d	0.59 ± 0.10	0.18 ± 0.08	0.21±0.11	n.d	n.d	0.26±0.01	<LOD	<LOD	1.69;<LOD	<LOD	<LOD	0.41 ± 0.34	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD				
PCB 77	ng g ⁻¹	n.d	<LOD	<LOD	<LOD	n.d	n.d	0.05±0.01	2.01	<LOD	1.0;1.49	0.69 ± 0.61	0.40	0.64 ± 0.45	0.65 ± 0.55	<LOD	<LOD	<LOD	<LOD	<LOD				
PCB 101	ng g ⁻¹	n.d	8.29 ± 4.97	3.89±1.46	4.66±2.39	n.d	n.d	5.82±1.22	12.7	1.17	3.45;5.46	3.17 ± 2.93	0.42	1.54 ± 0.48	0.93 ± 1.05	0.39	1.36	3.91	2.26	2.70				
PCB-118	ng g ⁻¹	n.d	9.91 ± 1.95	3.29 ± 1.30	5.03±2.41	n.d	n.d	2.87±0.44	12.7	<LOD	6.72;7.26	1.96 ±2.42	2.37	1.70 ± 1.79	<LOD	0.47	1.09	2.84	3.39	3.20				
PCB 114	ng g ⁻¹	n.d	0.08 ± 0.06	<LOD	0.05±0.03	n.d	n.d	<LOD	<LOD	<LOD	7.56;<LOD	2.46 ±2.86	2.59	2.91 ± 1.35	0.38 ± 0.55	<LOD	0.59	2.50	2.88	2.76				
PCB 123	ng g ⁻¹	n.d	<LOD	<LOD	<LOD	n.d	n.d	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD				
PCB 153	ng g ⁻¹	n.d	36.68 ± 2.93	13.5±4.40	18.5±8.28	n.d	n.d	10.6±1.37	62.8	7.81	40.30	16.6 ± 12.3	0.74	10.3 ± 0.87	10.58 ± 8.81	0.93	2.58	9.64	8.44	13.66				
PCB 105	ng g ⁻¹	n.d	1.78 ± 0.66	0.67±0.28	1.09±0.67	n.d	n.d	0.83±0.01	2.61	<LOD	1.01;1.32	0.52 ± 0.31	0.66	0.68 ± 0.38	<LOD	<LOD	0.41	0.56	0.66	0.49				
PCB 138	ng g ⁻¹	n.d	27.29 ± 2.51	10.6±3.64	13.9±6.06	n.d	n.d	8.46±0.97	52.4	5.80	35.34;3.19	15.7 ± 9.8	9.22	10.5 ± 0.99	1.81 ± 1.93	0.78	2.87	9.53	5.51	11.26				
PCB 126	ng g ⁻¹	n.d	0.36 ± 0.04	0.13±0.07	0.16±0.09	n.d	n.d	0.14±0.01	39.6	6.68	38.84;22.20	8.70 ± 9.35	<LOD	5.02 ± 5.87	4.53 ± 6.83	<LOD	<LOD	0.42	<LOD	0.58				
PCB 128+157+167	ng g ⁻¹	n.d	4.32	1.68	2.28	n.d	n.d	1.75	1.70	0.69	2.29	1.79	0.49	0.77	0.77	0.32	0.45	1.35	1.11	0.70				
PCB 156	ng g ⁻¹	n.d	1.42 ± 0.14	0.67±0.26	0.73±0.32	n.d	n.d	0.44±0.02	1.64	0.72	4.02;3.01	1.51 ± 1.35	1.00	1.12 ± 0.37	<LOD	<LOD	<LOD	0.35	0.32	0.44				
PCB 180	ng g ⁻¹	n.d	17.31 ± 2.69	8.39±1.92	6.84±3.12	n.d	n.d	1.45±0.06	38.1	3.76	25.24;19.49	10.9 ± 7.17	8.24	5.22 ± 0.62	8.42 ± 6.61	<LOD	0.63	1.98	1.05	1.63				
PCB 169	ng g ⁻¹	n.d	<LOD	<LOD	<LOD	n.d	n.d	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD				
PCB 170	ng g ⁻¹	n.d	6.77 ± 0.52	3.55±0.82	3.31±1.47	n.d	n.d	0.42±0.03	12.44	1.83	0.43;9.32	3.07 ± 2.28	1.46	1.82±1.12	0.44 ± 0.36	<LOD	0.16	0.52	0.31	0.73				
PCB 189	ng g ⁻¹	n.d	0.32 ± 0.04	0.14±0.04	0.15±0.08	n.d	n.d	0.07±0.01	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD				
Σ PCB _s	ng g ⁻¹	n.d	121.8 ± 1.47	49.5±13.9	1.3±22.2	n.d	n.d	38.4±5.27	241.7	30.1	175.1;113.8	68.5 ± 37.8	28.70	43.6 ± 6.93	30.1± 24.0	4.98	11.7	35.2	27.1	39.3				
Σ NDL-PCB	ng g ⁻¹	n.d	96.2 ± 9.81	39.2±10.6	48.3±16.2	n.d	n.d	31.5±5.0	168.2 ^a	18.9	110.5;66.3 ^a	47.0 ± 31.2	18.90	27.9 ± 2.32	22.1 ± 17.5	2.59	7.87	25.94	17.58	29.57				
Σ 7PCB	ng g ⁻¹	n.d	106 ± 11.4	42.5±11.9	53.3±18.5	n.d	n.d	34.4±5.5	180.90	19.0	117.2; 73.6	48.9 ± 32.7	21.30	29.7 ± 2.87	22.4 ± 17.7	3.06	8.96	28.8	20.9	32.8				
Phenanthrene	ng g ⁻¹	n.d	n.d	n.d	n.d	n.d	n.d	n.d	1.52	3.97	7.50;8.56	2.64 ± 1.07	1.33	3.61 ± 1.54	2.15±1.69	4.60	3.74	2.91	2.77	1.63				
Anthracene	ng g ⁻¹	n.d	n.d	n.d	n.d	n.d	n.d	n.d	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	6.57	11.83				
Fluoranthene	ng g ⁻¹	n.d	n.d	n.d	n.d	n.d	n.d	n.d	<LOD	1.54	2.85;1.40	2.53 ± 2.90	3.16	0.81 ± 0.48	4.32± 3.26	3.02	13.73	4.07	7.62	8.96				
Pyrene	ng g ⁻¹	n.d	n.d	n.d	n.d	n.d	n.d	n.d	3.70	5.34	7.49;5.18	9.03 ± 11.36	12.73	3.60 ± 1.05	11.97 ± 12.30	6.80	7.42	5.85	10.09	26.99				
Benzo-a-Anthracene	ng g ⁻¹	n.d	n.d	n.d	n.d	n.d	n.d	n.d	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	1.68	0.65	3.61	1.09				
Chrysene	ng g ⁻¹	n.d	n.d	n.d	n.d	n.d	n.d	n.d	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	3.08	1.19	5.09	2.74				
Benzo-b-Fluoranthene	ng g ⁻¹	n.d	n.d	n.d	n.d	n.d	n.d	n.d	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	2.11	1.24	5.21	2.93				
Benzo-k-Fluoranthene	ng g ⁻¹	n.d	n.d	n.d	n.d	n.d	n.d	n.d	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	1.94	0.69				
Benzo-a-Pyrene	ng g ⁻¹	n.d	n.d	n.d	n.d	n.d	n.d	n.d	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	4.31	8.50 ^{a,b}	11.3 ^{a,b}	6.57 ^{a,b}	11.0 ^{a,b}			
Indeno-123-cd-Pyrene	ng g ⁻¹	n.d	n.d	n.d	n.d	n.d	n.d	n.d	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	3.42	15.72	5.35	8.49	7.16			
Dibenzo-ah-Anthracene	ng g ⁻¹	n.d	n.d	n.d	n.d	n.d	n.d	n.d	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD				
Benzo-ghi-Perylene	ng g ⁻¹	n.d	n.d	n.d	n.d	n.d	n.d	n.d	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	1.20	0.97				
Σ PAHs	ng g ⁻¹	n.d	n.d	n.d	n.d	n.d	n.d	n.d	9.22	14.4	21.4;18.7	18.0 ± 13.8	21.70	11.6 ± 3.00	22.0 ± 16.9	25.0	57.6	34.2	59.6	76.4				

Values exceeding reference limits are expressed in bold; ^(a) Reg CE 1881/2006 as amended; ^(b) Dlgs.172/2015; Σ 7PCB sum of CB 28+52+101+118+138+153+180 (Webster et al., 2013).

Table 3. Concentrations (w.w.) of Hg, HCB, PCBs and PAHs in mare organisms from the Augusta Bay.

Figures




Figure 1: Map and details of the sampling in Augusta Bay.




Figure 2: Box plot of the concentrations of Hg (a), HCB (b), ΣPCBs (c) and ΣPCBs NDL (d) in sediment cores.




Figure 3: Relative contribution of PCB homologs (% composition) in sediments, seafood and mussel samples.




Figure 4: Heatmap generated from hierarchical clustering analysis. The dendrograms of sample clustering (on the left) and of congener clustering (on the top) were added. The colour bars inside the graph indicate the different proportion of congeners (X axis) for each sample (Y axis). The rows were splitted based on number of identified clusters.




Figure 5: Dotchart of Fluo/Py in sediments and marine organisms

Supplementary materials

Table S1 Hg, HCB and PCBs concentrations (d.w.) in 0-10 cm levels of sediment collected in the sampling site.

Table S1 Hg, HCB and PCBs concentrations (d.w.) in 0-10 cm levels of sediment collected in the sampling site.

Core	level	Hg	HCB	PCB 28	PCB 52	PCB 81	PCB 77	PCB 101	PCB 118	PCB 114	PCB 123	PCB 153	PCB 105	PCB 138	PCB 126	PCB 128+157	PCB 156	PCB 167	PCB 180	PCB 169	PCB 170	PCB 189	Σ PCBs	Σ PCB _s ¹	Σ 7PC _B
A 3	cm	μg g ⁻¹	ng g ⁻¹																						
	0-1	3.73	10.6	5.85	0.31	<LOD	<LOD	<LOD	3.77	<LOD	<LOD	6.34	<LOD	4.42	<LOD	<LOD	<LOD	<LOD	1.66	<LOD	<LOD	<LOD	23.3	22.9	22.4
	1-2	3.07	11.2	4.98	<LOD	<LOD	<LOD	<LOD	3.36	<LOD	<LOD	5.91	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	1.21	<LOD	<LOD	<LOD	16.6	16.1	15.7
	2-3	3.73	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
	3-4	3.81	6.29	14.7	<LOD	<LOD	<LOD	<LOD	8.23	<LOD	<LOD	14.7	<LOD	7.09	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	45.9	45.4	45.0
	4-5	3.71	9.06	4.55	<LOD	<LOD	<LOD	<LOD	2.74	<LOD	<LOD	5.90	<LOD	3.90	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	18.2	17.8	17.3
	5-6	3.96	8.42	16.0	0.41	<LOD	<LOD	<LOD	12.2	<LOD	<LOD	15.41	4.2	14.4	<LOD	<LOD	<LOD	<LOD	3.25	<LOD	<LOD	<LOD	66.8	62.2	61.8
	6-7	3.56	8.98	4.92	0.78	<LOD	<LOD	<LOD	3.59	<LOD	<LOD	5.37	0.65	3.92	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	20.2	19.2	18.7
	7-8	3.77	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
	8-9	3.81	8.72	9.41	<LOD	<LOD	<LOD	<LOD	6.05	<LOD	<LOD	9.17	<LOD	4.92	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	30.7	30.2	29.8
	9-10	3.31	4.95	22.4	1.58	<LOD	<LOD	0.24	15.3	<LOD	<LOD	20.7	4.21	21.96	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	87.7	82.8	82.3
mean		3.64	8.53	10.36	0.42	<LOD	<LOD	<LOD	6.90	<LOD	<LOD	<LOD	1.18	7.59	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	38.7	29.6	36.6
st.dev		0.27 ^a	2.06 ^a	6.66	0.53				4.66			5.77	1.88	7.11					1.17				26.2	26.6 _a	24.4 _b

A 7	0-1	6.89	23.0	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
	1-2	6.98	ND	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
	2-3	6.99	16.0	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
	3-4	6.93	11.6	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.

	4-5	6.65	32.1	1.62	<LOD	<LOD	<LOD	<LOD	1.02	<LOD	<LOD	0.83	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	4.67	4.22	3.77
	5-6	6.88	28.8	1.74	<LOD	<LOD	<LOD	<LOD	1.24	<LOD	<LOD	4.46	<LOD	5.00	<LOD	<LOD	0.78	0.43	2.66	<LOD	1.43	<LOD	18.6	16.4	15.3
	6-7	6.92	32.8	2.01	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	4.15	<LOD	3.56	<LOD	0.56	<LOD	<LOD	2.04	<LOD	<LOD	<LOD	13.4	12.9	12.0
	7-8	5.26	22.1	3.79	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	2.47	<LOD	0.47	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	7.92	7.47	7.02
	8-9	5.47	26.0	1.82	0.21	<LOD	<LOD	<LOD	1.43	<LOD	<LOD	2.32	<LOD	1.52	<LOD	<LOD	<LOD	<LOD	0.44	<LOD	0.29	<LOD	8.93	8.27	7.82
	9-10	8.08	35.8	<LOD	<LOD	<LOD	<LOD	<LOD	2.00	<LOD	<LOD	6.35	<LOD	6.38	<LOD	<LOD	<LOD	<LOD	<LOD	3.90	<LOD	<LOD	<LOD	22.2	21.8
mean		6.71	25.3	2.26	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	0.19	<LOD	1.53	<LOD	<LOD	<LOD	<LOD	11.8	11.2
st.dv		0.80 ^a	8.03 ^a	0.82					0.77			1.95					0.29		1.58					6.50 _a	6.39 _b

A 9	0-1	10.7	65.9	0.32	1.01	0.09	0.55	3.88	2.81	<LOD	0.15	8.93	0.59	8.96	0.83	0.49	0.56	0.89	6.44	<LOD	3.46	0.26	40.4	35.0	32.4
	1-2	11.6	134	0.38	2.01	<LOD	0.89	8.40	5.90	0.03	0.50	12.7	1.45	13.5	0.94	0.96	1.31	0.92	7.54	0.13	4.00	0.32	62.0	54.8	50.5
	2-3	10.7	331	0.55	1.37	<LOD	0.78	5.74	2.95	<LOD	110	11.0	1.19	10.3	1.25	0.66	0.76	1.04	7.32	0.17	4.16	0.39	160	42.9	39.2
	3-4	12.2	140	0.29	1.17	<LOD	0.52	4.29	3.08	<LOD	0.44	9.63	0.76	9.03	0.93	0.50	0.76	1.09	6.69	0.22	3.44	0.26	43.3	37.2	34.2
	4-5	11.3	328	0.34	1.12	<LOD	0.58	4.42	3.33	0.01	0.26	9.96	0.62	9.34	0.84	0.54	0.44	1.10	7.99	<LOD	4.24	0.25	45.5	39.1	36.5
	5-6	11.3	142	0.48	1.25	<LOD	0.82	4.72	3.65	<LOD	0.15	10.13	0.53	9.22	1.24	0.48	0.73	1.08	11.62	<LOD	4.44	0.13	50.9	44.5	41.1
	6-7	12.0	155	0.34	1.25	<LOD	0.82	4.21	3.47	0.02	<LOD	9.55	1.05	8.11	0.99	0.43	0.48	1.02	7.84	<LOD	3.76	0.32	43.9	37.7	34.8
	7-8	11.4	104	0.36	1.27	<LOD	0.79	5.23	4.19	<LOD	<LOD	10.8	2.97	9.68	1.18	0.44	0.40	1.09	8.42	<LOD	3.70	0.12	50.8	42.9	39.9
	8-9	12.1	162	0.31	1.72	<LOD	0.61	6.74	4.86	<LOD	<LOD	11.0	1.18	10.62	0.83	0.67	0.29	0.99	6.99	<LOD	3.90	0.14	51.2	44.8	42.3
	9-10	10.8	164	0.64	1.30	<LOD	0.82	4.52	3.70	0.03	<LOD	9.96	2.28	8.45	1.55	1.07	0.08	1.14	7.99	<LOD	3.40	0.32	47.5	40.2	36.6
mean		11.4	172.6	0.40	1.35	<LOD	<LOD	5.22	3.79	<LOD	11.2	10.4	1.26	9.72	1.06	0.62	0.58	1.04	7.89	<LOD	<LOD	0.25	59.5	41.9	38.7
st.dv		0.57 ^a	87.6 ^a	0.12	0.30		0.13	1.40	0.96		34.6	1.07	0.80	1.53	0.24	0.22	0.34	0.08	1.45		0.36	0.09	35.6	5.60 _a	5.20 _b

A 11	0-1	6.94	17.4	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
	1-2	7.18	27.9	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
	2-3	7.04	35.5	5.05	<LOD	<LOD	<LOD	<LOD	5.08	<LOD	<LOD	14.3	<LOD	19.1	<LOD	3.0	4.2	4.4	8.3	<LOD	5.4	<LOD	69.5	59.4	52.0
	3-4	7.74	54.1	4.30	<LOD	<LOD	<LOD	<LOD	2.09	<LOD	<LOD	6.44	<LOD	2.1	<LOD	<LOD	<LOD	<LOD	1.1	<LOD	<LOD	<LOD	17.1	16.7	16.2
	4-5	6.97	28.2	3.10	<LOD	<LOD	<LOD	<LOD	2.01	<LOD	<LOD	9.07	<LOD	4.3	<LOD	0.4	0.3	0.4	5.7	<LOD	1.2	<LOD	27.1	25.2	24.2
	5-6	6.63	16.0	5.18	0.64	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	12.3	<LOD	8.7	<LOD	1.0	<LOD	<LOD	3.8	<LOD	2.3	<LOD	34.8	32.1	30.8

	6-7	6.69	17.4	5.03	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	<LOD	12.2	<LOD	13.4	<LOD	<LOD	<LOD	<LOD	6.9	<LOD	4.1	<LOD	42.7	38.2	37.8
	7-8	7.45	18.0	35.3	2.42	<LOD	<LOD	<LOD	43.11	<LOD	<LOD	42.8	7.04	56.1	<LOD	2.2	<LOD	<LOD	<LOD	<LOD	1.1	<LOD	n.d.	n.d.	n.d.
	8-9	5.33	11.6	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
	9-10	3.40	1.98	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
mean		6.54	22.8	9.65	0.56	<LOD	<LOD	<LOD	8.74	<LOD	<LOD	16.2	1.24	17.3	<LOD	1.10	0.80	0.84	4.30	<LOD	2.35	<LOD	38.2	34.3	32.2
st.dv		1.28 ^a	14.45 ^a	12.6	0.94				16.9			13.3	2.84	20.0		1.21	1.65	1.75	3.26		2.03		19.9	16.2 ^a	13.6 _b

Table S2: PAHs concentrations (d.w.) in 0-10 cm levels of sediment collected in the sampling site.

Core	Layer	PHENANTHRENE	ANTHRACENE	FLUORANTHENE	PYRENE	BENZO[a]ANTHRACENE	CHRYSENE	BENZO[b]FLUORANTHENE	BENZO[k]FLUORANTHENE	BENZO[a]PYRENE	INDENO[123-cd]PYRENE	DIBENZO[a,h]ANTHRACENE	BENZO[ghi]PERYLENE	ΣPAHs
A3	cm	<i>ng g⁻¹</i>												
	0-1	53.5	136	15.9	15.0	20.9	30.9	63.0	91.9	77.1	112.2	29.6	96.6	742
	1-2	28.9	87.0	50.7	37.8	26.7	28.0	52.5	20.9	53.8	76.5	10.3	56.9	530
	2-3	49.9	118	128.9	65.8	61.8	64.1	84.6	37.8	111.4	104.0	13.7	72.4	913
	3-4	33.6	158	49.5	27.6	27.9	32.2	60.0	24.6	55.2	68.1	8.3	37.2	583
	4-5	39.3	230	63.4	29.8	31.5	35.6	48.9	23.3	54.2	74.0	14.6	49.5	694
	5-6	35.4	264	52.3	26.4	32.1	37.1	87.2	24.7	68.9	83.1	14.5	55.5	782
	6-7	32.9	286	44.6	36.0	34.5	40.7	30.0	13.5	37.4	49.6	9.4	21.7	636
	7-8	26.8	111	54.9	24.7	29.9	33.1	55.5	59.3	52.2	44.3	9.6	32.1	533
	8-9	35.6	184	52.7	26.9	35.4	38.5	29.2	16.6	62.6	32.0	8.9	36.1	558
	9-10	40.4	227	105.5	41.2	41.5	44.4	34.9	23.1	73.7	56.8	10.5	48.5	747
mean		37.6	180	61.8	33.1	34.2	38.5	54.6	33.6	64.6	70.1	12.9	50.6	671.87
st.dv		8.52	68.9	32.2	13.7	11.1	10.2	20.4	24.3	20.12	25.5	6.30	21.7	125.97

A7	0-1	41.8	186	81.1	34.3	32.7	37.8	49.8	14.8	50.2	34.4	4.77	30.9	599
	1-2	28.7	112	45.1	38.3	18.9	22.5	34.0	10.8	32.2	23.0	3.62	24.4	394
	2-3	28.0	184	37.5	25.8	18.6	23.1	33.0	7.2	29.1	20.4	1.85	25.5	434
	3-4	25.6	184	33.9	22.8	15.3	17.8	33.0	7.7	27.8	19.0	1.62	15.9	404

	4-5	23.9	201	29.3	21.5	15.8	18.9	35.4	9.0	27.4	21.6	1.41	16.7	422
	5-6	23.0	218	28.4	20.9	13.4	16.5	30.0	9.9	24.8	16.0	1.01	11.2	413
	6-7	28.2	211	32.2	25.7	17.7	19.9	32.8	8.7	29.6	18.1	1.72	20.1	446
	7-8	18.3	177	16.5	13.6	10.5	14.4	23.1	6.6	18.2	13.1	0.00	8.1	319
	8-9	19.2	209	18.7	12.7	11.6	14.7	23.0	6.7	21.9	14.1	1.54	7.2	360
	9-10	27.8	186	30.7	28.8	15.6	20.4	31.4	11.1	31.3	14.7	1.17	16.1	415
mean		26.4	186.8	35.3	24.5	17.0	20.6	32.5	9.26	29.3	19.5	1.87	17.6	420
st.dv		6.56	29.6	18.1	8.11	6.18	6.72	7.45	2.54	8.53	6.19	1.36	7.72	72.73

A 9	0-1	54.6	70.4	31.6	37.1	18.9	37.7	24.3	9.28	19.1	6.20	6.92	19.8	336
	1-2	46.1	76.2	32.6	40.7	18.8	37.0	25.2	6.61	18.0	6.27	7.53	13.7	329
	2-3	59.3	47.5	37.2	61.1	25.2	58.2	31.3	7.20	24.3	9.05	6.91	10.2	378
	3-4	60.0	65.7	34.2	50.6	20.3	40.1	24.6	7.74	20.0	9.54	6.22	9.12	348
	4-5	68.6	53.0	57.9	71.8	36.7	59.5	36.2	12.00	30.9	16.67	13.37	20.7	477
	5-6	130	23.4	112	118	53.0	78.0	51.9	22.43	39.7	23.91	3.75	9.90	666
	6-7	64.4	61.0	35.0	55.7	23.2	41.2	26.6	9.59	24.6	4.39	2.94	5.93	355
	7-8	62.0	56.3	34.3	60.2	22.6	41.7	27.4	9.59	19.1	7.03	16.34	6.91	364
	8-9	76.0	59.1	41.9	61.4	23.5	46.9	32.2	10.68	21.4	8.89	10.46	6.50	399
	9-10	57.4	50.3	34.9	55.7	21.7	41.6	25.0	7.61	18.1	6.89	4.57	8.64	332
mean		67.8	56.3	45.2	61.2	26.4	48.2	30.5	10.3	23.5	9.88	7.90	11.15	398
st.dv		67.8 ± 23.3	14.6	24.7	22.3	10.6	13.1	8.51	4.59	6.94	5.95	4.29	5.30	103

A 11	0-1	25.2	78.2	19.6	22.2	12.8	17.1	18.0	6.10	21.7	13.3	3.55	22.2	260
	1-2	28.3	93.7	18.3	24.7	11.6	15.9	14.4	6.56	20.5	14.7	4.20	19.3	272
	2-3	52.6	84.0	41.8	32.9	15.5	20.6	16.9	9.14	25.1	14.7	5.79	17.6	337
	3-4	28.9	65.6	29.4	41.2	15.6	19.5	14.5	7.80	21.8	13.9	3.71	17.7	280
	4-5	32.3	112.9	25.2	21.9	13.7	17.7	11.9	5.27	17.4	14.0	4.89	14.7	292
	5-6	21.2	96.3	27.2	21.3	14.5	18.8	15.0	10.3	22.5	14.2	3.06	16.4	281
	6-7	33.8	85.8	54.8	40.4	20.8	27.5	21.6	9.26	32.2	14.9	0.00	15.1	356
	7-8	27.4	158.1	30.2	25.5	14.7	19.9	15.5	7.10	21.9	13.6	1.89	15.5	351

	8-9	12.3	61.9	16.4	13.8	9.0	11.7	7.53	5.54	13.5	8.57	0.00	7.71	168
	9-10	15.8	80.2	12.1	10.1	6.34	9.24	8.54	3.82	8.51	9.15	2.01	11.3	177
mean		27.7	91.7	27.5	25.4	13.5	17.8	14.4	7.09	20.5	13.1	2.91	15.7	277
st.dv		11.1	27.6	12.8	10.2	3.94	4.99	4.24	2.03	206.42	2.30	1.94	4.06	64.8

