



UNIVERSITÀ POLITECNICA DELLE MARCHE
Repository ISTITUZIONALE

Effects of BPA on zebrafish gonads: Focus on the endocannabinoid system

This is the peer reviewed version of the following article:

Original

Effects of BPA on zebrafish gonads: Focus on the endocannabinoid system / FORNER PIQUER, Isabel; Beato, S.; Piscitelli, F.; Santangeli, S.; Di Marzo, V.; Habibi, H. R.; Maradonna, F.; Carnevali, O.. - In: ENVIRONMENTAL POLLUTION. - ISSN 0269-7491. - 264:(2020). [10.1016/j.envpol.2020.114710]

Availability:

This version is available at: 11566/283156 since: 2024-03-27T14:34:08Z

Publisher:

Published

DOI:10.1016/j.envpol.2020.114710

Terms of use:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. The use of copyrighted works requires the consent of the rights' holder (author or publisher). Works made available under a Creative Commons license or a Publisher's custom-made license can be used according to the terms and conditions contained therein. See editor's website for further information and terms and conditions.

This item was downloaded from IRIS Università Politecnica delle Marche (<https://iris.univpm.it>). When citing, please refer to the published version.

note finali coverage

(Article begins on next page)

27 July 2024

Manuscript Details

Manuscript number	ENVPOL_2020_788_R2
Title	Effects of BPA on zebrafish gonads: focus on the endocannabinoid system
Article type	Research Paper

Abstract

Bisphenol A (BPA), a monomer used for polycarbonate manufacture, has been widely reported as an endocrine-disrupting chemical (EDC). Among other alterations, BPA induces reproductive dysfunctions. Changes in the endocannabinoid system (ECS) has been recently shown to be associated with reproductive disorders. ECS is a lipid-based signaling system (cannabinoid receptors, endocannabinoids and enzymatic machinery) involved in several physiological functions. The main goal of the present study was to assess the effects of two environmental concentrations of BPA (10 and 20 µg/L) on the ECS in 1-year zebrafish gonads. In the males, BPA increased the gonadosomatic index (GSI) and altered testicular levels of endocannabinoids as well as reduced the testicular area occupied by spermatogonia. In the male liver, exposure to 20 µg/L BPA significantly increased vitellogenin (vtg) transcript levels. In female zebrafish, BPA altered ovarian endocannabinoid levels, elevated hepatic vtg mRNA levels as well as increased the percentage of vitellogenic oocytes in the ovaries. In conclusion, exposure to two environmentally relevant concentrations of BPA altered ECS and consequently, gonadal function in both male and female zebrafish.

Keywords	Danio rerio; endocannabinoid; Endocrine disruptors; reproduction; vitellogenin
Corresponding Author	Oliana Carnevali
Corresponding Author's Institution	Università Politecnica delle Marche
Order of Authors	Isabel Forner-Piquer, Silvia Beato, Fabiana Piscitelli, Stefania Santangeli, Vincenzo Di Marzo, Hamid Habibi, Francesca Maradonna, Oliana Carnevali
Suggested reviewers	Renata Moreira, Cristoforo Silvestri, Fabiana Lo Nostro, Vanesa Robles, Gary Hardiman

Submission Files Included in this PDF

File Name [File Type]

Lett_ed.pdf [Cover Letter]

rebuttal_R2.docx [Response to Reviewers]

Highlights.docx [Highlights]

graphical abstract.jpg [Graphical Abstract]

BPA_R2.docx [Manuscript File]

declaration-of-competing-interests.pdf [Conflict of Interest]

author statement.docx [Author Statement]

S1.docx [Supplementary Material]

Submission Files Not Included in this PDF

File Name [File Type]

S2.xlsx [Supplementary Material]

To view all the submission files, including those not included in the PDF, click on the manuscript title on your EVISE Homepage, then click 'Download zip file'.

Research Data Related to this Submission

There are no linked research data sets for this submission. The following reason is given:
Data will be made available on request



January 30th 2020

Dear Editor,

I enclose an original article entitled “**Effects of BPA on zebrafish gonads: focus on the endocannabinoid system**” co-authored by Isabel Forner-Piquer, Silvia Beato, Fabiana Piscitelli, Stefania Santangeli, Vincenzo di Marzo, Hamid R. Habibi, Francesca Maradonna and myself, Oliana Carnevali, for consideration as an article in Environmental Pollution.

In the present manuscript, we reported the results obtained after 21 days exposure to two different concentrations of Bisphenol A (BPA), 10 and 20 µg/L (based on BPA environmental concentrations), on the endocannabinoid system in both ovary and testis. The endocannabinoid system was used as a novel target due to its role on reproductive physiology.

Thus, this study can be considered as continuum of the recent effort on investigating the effects of Bisphenol A. Indeed, our results reported that BPA induced alterations on the levels of endocannabinoids and transcriptomic changes in genes coding for the endocannabinoid system and reproductive markers. In addition, the histological analyses on gonads evidenced alterations of gametogenesis in both ovary and testis.

Finally, we declare no conflict of interest associated with this publication, and we state that none of the material contained in the present manuscript has been published or is under consideration in any other place.

We believe that our findings will be of interest to the readers of your journal.

Thank you for considering this article and please, let us know if there is further information we can provide.

Sincerely,

Prof Oliana Carnevali

Department of Life and Environmental sciences
Polytechnic University of Marche

Reviewer 4

Minor points:

1) GSI should be defined in the main text the first time it is used (not just the abstract).


GSI has been defined in line 106 (materials and methods) and 176 (results).

2) The authors are expressing their data in deltaCt in table 3 and 4 and figure 3, and this must be stated. Note that these are not 'arbitrary units' as indicated in the methods. The authors must explicitly state this and indicate where this is different from the fold change analysis $2^{-\Delta\Delta Ct}$. Therefore, I suggest that the legend for table 3 and 4 and figure 3 be changed from "Data are expressed as means \pm SEM" to "Data are expressed as mean dCt \pm SEM".

As the reviewer suggested, the image legends have been modified.

1 **HIGHLIGHTS**

- 2 1. BPA altered the endocannabinoid system in zebrafish gonads.
- 3 2. BPA (20 µg/L) decreased the percentage of the area occupied by spermatogonia in the testis.
- 4 3. BPA (20 µg/L) up-regulated the hepatic vtg expression in male and females.
- 5 4. BPA increased male GSI.



Effects of BPA on zebrafish gonads: focus on the endocannabinoid system

Isabel Forner-Piquer¹, Silvia Beato¹, Fabiana Piscitelli¹, Stefania Santangeli¹, Vincenzo di Marzo^{2,3},
Hamid R. Habibi⁴, Francesca Maradonna¹ and Oliana Carnevali^{1,5*}

¹Dipartimento Scienze della Vita e dell'Ambiente, Università Politecnica delle Marche, Via Brecce Bianche, 60131 Ancona, Italy

²Endocannabinoid Research Group, Istituto di Chimica Biomolecolare, Consiglio Nazionale delle Ricerche, Via Campi Flegrei, 80078 Pozzuoli, Italy

³Canada Excellence Research Chair on the Microbiome-Endocannabinoidome Axis in Metabolic Health, Université Laval, Quebec City, Canada

⁴Department of Biological Sciences, University of Calgary, Calgary, Alberta, T2N 1N4, Canada

⁵INBB - Consorzio Interuniversitario di Biosistemi e Biostrutture, 00136 Roma, Italy.

*Corresponding author: Prof. Oliana Carnevali, Dipartimento Scienze della Vita e dell'Ambiente, Università Politecnica delle Marche, Via Brecce Bianche, 60131 Ancona, Italy

e-mail: o.carnevali@univpm.it // ORCID: <https://orcid.org/0000-0001-5994-0572>

HIGHLIGHTS

1. BPA altered the endocannabinoid system in zebrafish gonads.
2. BPA (20 µg/L) decreased the percentage of the area occupied by spermatogonia.
3. BPA (20 µg/L) up-regulated the hepatic vitellogenin (*vtg*) expression.
4. BPA increased male gonadosomatic index (GSI).

ABSTRACT

Bisphenol A (BPA), a monomer used for polycarbonate manufacture, has been widely reported as an endocrine-disrupting chemical (EDC). Among other alterations, BPA induces reproductive dysfunctions. Changes in the endocannabinoid system (ECS) has been recently shown to be associated with reproductive disorders. ECS is a lipid-based signalling system (cannabinoid receptors, endocannabinoids and enzymatic machinery) involved in several physiological functions. The main goal of the present study was to assess the effects of two environmental concentrations of BPA (10 and 20 µg/L) on the ECS in 1-year zebrafish gonads. In the males, BPA increased the gonadosomatic

35 index (GSI) and altered testicular levels of endocannabinoids as well as reduced the testicular area
36 occupied by spermatogonia. In the male liver, exposure to 20 µg/L BPA significantly increased
37 vitellogenin (*vtg*) transcript levels. In female zebrafish, BPA altered ovarian endocannabinoid levels,
38 elevated hepatic *vtg* mRNA levels as well as increased the percentage of vitellogenic oocytes in the
39 ovaries. In conclusion, exposure to two environmentally relevant concentrations of BPA altered ECS
40 and consequently, gonadal function in both male and female zebrafish.

41


42 **KEY WORDS:** *Danio rerio*; endocannabinoid; endocrine disruptors; reproduction; vitellogenin.

43

44 **Capsule:** 21-day BPA exposure altered the zebrafish endocannabinoid system and changed the
45 gonad morphology.

46

47 GRAPHICAL ABSTRACT



48

49 **ABBREVIATIONS:** AEA: anandamide; 2-AG: 2-arachidonoylglycerol; OEA: *N*-oleyl ethanolamine; PEA:
50 *N*-palmityl ethanolamine; CB1/cnr1: endocannabinoid receptor type 1; CB2/cnr2: endocannabinoid
51 receptor type 2; TRPV1: transient receptor potential vanilloid type I ion channel; GPR55: G protein-
52 coupled receptor 55; NAPE-PLD: *N*-acyl phosphatidylethanolamine phospholipase D, FAAH: Fatty acid
53 amide hydrolase, DAGLα: Diacylglycerol lipase alpha, MGLL: monoglyceride lipase, ABDH4:
54 abhydrolase domain containing 4, VTG: vitellogenin.

55

56 1. INTRODUCTION

57 Since 1940, BPA has been used for the manufacture of polycarbonate products and epoxy resins in
58 the plastic industry because of its capacity to increase heat resistance and elasticity [1]. Due to its
59 extensive use, BPA can be globally found in the aquatic environment and therefore, in humans and
60 wildlife [2]. BPA is known to enter the aquatic ecosystem by diffusion from wastewater treatment
61 plants and landfill sites [3]. Overall, BPA concentrations in the aquatic environment are variable,
62 depending on the location, reaching on average 21 µg/L in surface waters [4,5]. BPA can interact with
63 nuclear estrogen receptors (ER) alpha and its membrane-bound form (mER) [6,7] , ER beta [8] and
64 the G protein-coupled receptor 30 (GPR30) [9]. Additionally, BPA exerts estrogenic activity at low
65 concentrations [10] and can interfere with androgen mediated pathways [11–14]. Consequently, BPA
66 has been linked with alterations in male and female reproduction and epigenetic markers [15–18],
67 gonadotrophic cell hypertrophy and hyperplasia [19], follicular atresia [20], transcriptomic profile of
68 genes coding for reproductive signals [15,19,21–23], sperm physiology and motility [21,24,25],
69 hormone levels [23,26,27], sexual differentiation [28], abnormal neuronal development [11],
70 morphological development [29], and metabolic dysfunction [30].

71

72 It is now well established that the ECS is involved in different reproductive processes [31–33]. The
73 ECS is a lipid signaling system based on lipid mediators, the endocannabinoids (Anandamide: AEA, 2-
74 arachidonoylglycerol: 2-AG). It can be expanded to include endocannabinoid-like mediators such as
75 *N*-oleoyl ethanolamine (OEA) and *N*-palmitoyl ethanolamine (PEA). AEA and 2-AG exert their activity
76 through G protein-coupled receptors, i.e. the cannabinoid receptor type I (CB1 or *CNR1*) and type II
77 (CB2 or *CNR2*), as well as other molecular targets. The endocannabinoids are synthesized on demand,
78 mainly through the *N*-acyl phosphatidylethanolamine-specific phospholipase D (NAPE-PLD, for AEA)
79 and Diacylglycerol Lipase Alpha (DAGLα, for 2-AG), from membrane phospholipid precursors after
80 reuptake into the cytosol, facilitated by intracellular degrading enzymes such as fatty acid amide
81 hydrolase (FAAH) and monoglyceride lipase (MGLL). NAPE-PLD and FAAH also regulate OEA and PEA
82 tissue concentrations.

83 In humans, higher levels of AEA impaired sperm viability and motility [34], and in mice, lower levels
84 of 2-AG promoted epididymal sperm cell start-up [35]. Inactivation of CB1 caused ineffective histone
85 displacement, reduced chromatin condensation and DNA damage in mouse sperm [36]. In females,
86 a correct tone of AEA was reported to be necessary during human oocyte maturation [37,38];
87 however, higher levels of AEA may also decrease the levels of progesterone, which is associated with

88 implantation failure and/or abnormal development of the mammalian fetus [39]. Indeed, a crosslink
89 between endocannabinoids and sex steroids is reported elsewhere [40].
90 Finally, recent studies demonstrated the ability of BPA to deregulate the ECS in different species
91 [22,26,41–43]. Thus, we here examined the adverse effects of BPA on the gonadal ECS in adult
92 zebrafish, following 21-day exposure to two different environmental concentrations (10 and 20 µg
93 BPA/L).

94

95 **2. MATERIALS AND METHODS**

96 **2.1. BPA treatment**

97 BPA exposure was conducted in one-year zebrafish (AB strain) following previous publications
98 [4,5,15,22,44]. Two nominal concentrations of BPA were tested: 10 and 20 µg/L. Absolute ethanol
99 (100% EtOH) was used as a solvent carrier with a final concentration of 10 µL EtOH / L water (0.001%
100 v/v), which was well below the activity threshold (Hutchinson et al., 2006). After three weeks, fish
101 were euthanized with MS-222 (3-aminobenzoic acid ethyl ester) (Sigma-Aldrich) according to
102 University of Calgary Animal Care protocol (AC15-0183). Ovaries and testes were immediately frozen
103 in dry ice and stored at -80 °C until use for subsequent RNA extraction and endocannabinoid assays.
104 For histological procedures, the gonads were fixed in Bouin solution overnight, washed and stored
105 within EtOH 70%.

106 **Gonadosomaic Index** (GSI) was calculated: [(gonad weight/ total zebrafish weight) * 100]

107

108 **2.2 Quantitative real-time PCR (RT-qPCR).**

109 RNA was extracted from the gonads after gently homogenisation with TRIzol reagent (Invitrogen),
110 followed by chloroform disaggregation into a light aqueous layer in the upper part containing RNAs
111 and a lower one with proteins and DNAs. Then, the upper layer was transferred to a new Eppendorf
112 and the RNA was precipitated with isopropanol and washed twice with absolute ethanol. Finally, the
113 samples were treated with DNase following [15]. Quantification of the RNA was determined by
114 spectrophotometry using a Nanophotometer TM P-Class (Implen). Quality of the mRNA (integrity,
115 presence of genomic-DNA) was assessed by electrophoresis in 1% agarose gel.

116 Reverse transcription was conducted from 1 µg mRNA with High-Capacity cDNA Reverse
117 Transcription Kit (Applied Biosystems) following the manufacturer guidelines. The cDNA obtained
118 with the retrotranscription (MyCycler Thermal Cycler System, Bio-Rad) was considered as the stock

119 (1:1), this stock was diluted with milliQ water (1:10) to obtain the working concentrations. All the
120 cDNAs were kept at - 20 °C.

121 For the RT-qPCRs, SYBR Green method was used with an iQ5 multicolor Real-Time PCR detection
122 system (Bio-Rad) with single transparent strips. All replicates (n = 5) were run in duplicate. For each
123 reaction, the mix contained: 1 µL of cDNA (1:10) + 5 µL iQ SYBR Green Supermix (Bio-Rad) + 3.8 µL
124 milliQ water + 0.1 µL forward primer + 0.1 µL reverse primer. Final concentration of primers was 10
125 pmol/µL. The thermal cycling was as follow: 95 °C for 3 minutes; 45 cycles of denaturation
126 (10 seconds at 95 °C) followed by 20 seconds for annealing at 60 °C for *faah*, *abhd4*, *vtg*; 59 °C for
127 *dagla*, *trpv1*; 57 °C for *napepld*; 55 °C for *cnr1*; 53 °C for *gpr55* and *mgll*; and finally, 20 seconds for
128 elongation at 72 °C.

129 Two *no template control* (NTC) were added in each run to guarantee absence of contamination.
130 Annealing temperatures (Ta) for each primer were optimized with temperature gradient assays. Primer
131 specificities were assessed with the absence of primer-dimer formation and dissociation curves.
132 Additionally, for each pair of primers, the efficiencies were evaluated with a mix of cDNA (CTRL group)
133 at different concentrations (1:1, 1:10, 1:100, 1:1000).

134 The reference genes were *rplp0* (ribosomal protein large P0) and *18s* (ribosomal subunit 18s) for
135 ovary, and *rplp0* and *rpl13a* (ribosomal protein L13a) for testes and liver for being their expression
136 stable and abundant along all the experimental groups. List of primers is stated in Table 1 and
137 additional information can be found in Table S1 (Supplemental material).

138 RT-qPCR results have been analysed with the C_t (cycle threshold) values of both target and reference
139 genes with the Delta Delta C_t method [45,46] using the spreadsheet provided by Bio-Rad (Gene
140 Expression analysis for iCycler iQ Rela-time PCR detection system version 2.0).

141 **Table 1.** Primer list.

GENE	ABB	FORWARD (5' - 3')	REVERSE (5' - 3')	Accession number
Cannabinoid receptor 1	<i>cnr1</i>	TCTGTGGGAAGCCTGTTTC	ACCGAGTTGAGCCGTTTG	NM_212820.2
Transient receptor potential cation channel, subfamily V, member 1	<i>trpv1</i>	TGATCGTCGCTGGTGCTT	GACTGGGCTCTCTGAACG	NM_001126399.1
G protein-coupled receptor 55	<i>gpr55</i>	AACTGAAGGTGTGGATAC	ACGCCATAATGTTTAACG	XM_005163567.3
N-acyl phosphatidylethanolamine phospholipase D	<i>napepld</i>	CTCAAGGACATGCACTCA	GAGCACAATCTTCAAGACAAT	NM_001080613.2
Monoglyceride lipase	<i>mgll</i>	AAGTGAAGGTGAGAGGAT	AATGTCCAACGATGAAGA	NM_200297.2

Fatty acid amide hydrolase	<i>faah</i>	CATTCTCGGACCAGCCTTCA	CCTTCACTGCCTGTTGGAAGA	NM_001109825.1
Diacylglycerol lipase alpha	<i>dagla</i>	GAGGGTTTTCCGTCGTAC	TGTTCCCTCCAGCAATGATCC	XM_692781.8
Abhydrolase domain containing 4	<i>abhd4</i>	GAAGAGCAGTTTGTTCCTCCATA G	GACTCACTCTTTCTGGGTATT GGAT	NM_001017613.1
Vitellogenin 1	<i>vtg</i>	GCCAAAAAGCTGGGTAAACA	AGTTCCGTCTGGATTGATGG	NM_001044897.3
Ribosomal subunit 18s	<i>18s</i>	TCGGAAAACGGTGAACCTG	AAGGTCTTTGAACCCACGG	NR_195818.1
Ribosomal protein large P0	<i>rplp0</i>	CTGAACATCTCGCCCTTCTC	TAGCCGATCTGCAGACACAC	NM_131580.2
Ribosomal protein L13a	<i>rpl13a</i>	TCTGGAGGACTGTAAGAGGTATGC	AGACGCACAATCTTGAGAGCA G	NM_212784.1

142

143 **2.3 Measurement of endocannabinoids (AEA, 2-AG) and endocannabinoid-like mediators (PEA,**
144 **OEA) in testis and ovary.**

145 Endocannabinoids and endocannabinoids-like mediators were extracted, purified and quantified
146 from 5 testis and ovaries as previously described [47,48].

147

148 **2.4 Gonad histology and image analysis**

149 After being fixed overnight in Bouin, gonads were rinsed and stored in EtOH 70 % till processing.
150 Tissues were dehydrated with an increasing series of EtOH (70-100 %) and embedded in paraffin
151 overnight. Then, the paraffin blocs were sectioned with a microtome (Leica RM2125 RTS) at 4 µm
152 and stained with Eosin - Mayer's haematoxylin (Bio-Optica). Microphotographies were taken using a
153 Zeiss Axio Imager.M2 microscope coupled with a high-resolution camera Zeiss Aixocam 105 color.
154 One slide from the anterior, medium and posterior part of the testis and ovaries were chosen for the
155 analysis. Three microphotographies (40x) were taken per slide. Using *Image J* (ImageJ, NIH, USA,
156 <https://imagej.nih.gov/ij/>), the percentage of the area occupied by spermatogonia and spermatozoa
157 was evaluated from nine microphotographs per gonad [18] and 5 gonads per experimental group.
158 The area covered with spermatozoa and spermatogonia were manually delimited and evaluated
159 against the total surface of the slide and expressed as percentage. For the oocytes, using Cell Counter
160 plugging (Image J), three classes of oocytes were calculated: previtelogenic oocytes (oogonia and
161 cortical alveoli), vitelogenic oocyte and mature oocyte.

162

163 **2.5 Statistical analysis**


164 All the data was analysed by one-way analysis of variance (ANOVA) followed by Dunnett's multiple
165 comparison test. When the collected data was expressed in percentage, arcsin transformation was
166 conducted before ANOVA. When data did not meet the conditions for using ANOVA, Kruskal-Wallis
167 (non-parametric test) was applied. Superscript asterisks (*) evidenced statistical differences respect

168 the control (CTRL): * ($p < 0.05$), ** ($p < 0.01$). Data are reported as mean \pm SEM (Standard Error of
 169 the Mean). The statistical software: *GraphPad Prism 6* and *SigmaStat 3.5*. Expression heatmaps of
 170 genes were created by fold change means (BPA/CTRL) with *Genesis software v1.7.7*. When fold
 171 changes were < 1 , values were transformed to the negative inverse following: $[-1 / \text{fold change}]$ due
 172 to Genesis does not allow to enter values in the 0 - 1 interval (see Supplemental Material 2).

173

174 3. RESULTS

175 **3.1 Gonadosomatic index (GSI).** In females, the GSI was not affected by BPA (Fig. 1a). However, in
 176 males, the GSI was increased in a dose-dependent manner (Fig. 1b).



177

178 **Figure 1.** Gonadosomatic Index, female (a) and male (b) treated with 10 and 20 $\mu\text{g/L}$ BPA. Data are reported as means \pm
 179 SEM. Superscript asterisks (*) denote statistically significant differences with respect to the control (CTRL): * ($p < 0.05$),
 180 ** ($p < 0.01$) (one-way ANOVA, $p < 0.05$, Dunnett's multiple comparison test).

181

182 3.2 Measurement of endocannabinoids (AEA, 2-AG) and AEA-like mediators (PEA, OEA) in testes 183 and ovaries.

184 In ovary, the levels of AEA were significantly increased following exposure to 10 μg BPA/L (Table 2).
 185 However, OEA was reduced in both BPA groups (10 and 20 $\mu\text{g/L}$). In testes, exposure to 20 μg BPA/L
 186 significantly reduced the levels of AEA and 2-AG, whereas the lowest concentration of BPA (10 $\mu\text{g/L}$)
 187 was without effect.

188 **Table 2.** Endocannabinoids and endocannabinoid-like levels in testes and ovaries.

	CTRL		10 $\mu\text{g/L}$		20 $\mu\text{g/L}$	
	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE
AEA	15.92 \pm 3.206	23.18 \pm 1.834	30.83 \pm 4.961 *	31.97 \pm 5.247	13.65 \pm 1.936	7.85 \pm 2.252 **
2-AG	3.08 \pm 0.563	6.86 \pm 1.411	2.96 \pm 0.545	7.50 \pm 1.266	2.75 \pm 0.751	2.56 \pm 0.549 *

OEA	0.21 ± 0.037	0.09 ± 0.032	0.03 ± 0.024**	0.06 ± 0.014	0.01 ± 0.004 **	0.08 ± 0.014
PEA	0.10 ± 0.014	0.10 ± 0.027	0.09 ± 0.013	0.09 ± 0.027	0.11 ± 0.009	0.10 ± 0.014

189

190

Table 2. Levels of endocannabinoids and endocannabinoid-like mediators in the testes and ovaries of zebrafish treated with 10 and 20 µg/L BPA. Data reported as means $\Delta Ct \pm SEM$. Superscript asterisks (*) denote statistically significant differences with respect to the control (CTRL): * (p<0.05), ** (p<0.01) (one-way ANOVA, Dunnett's multiple comparison test, p < 0.05). AEA expressed as pmol/g tissue, while 2-AG, PEA and OEA as pmol/mg tissue.

191

192

193

194

195

3.3 ECS gene transcript levels in gonads.

196

Ovaries. As shown in Table 3, exposure to BPA did not alter the transcript levels of genes coding for the cannabinoid receptors. However, 20 µg BPA/L significantly increased mRNA expression of the gene coding for the catabolic enzyme *faah* (Table 3).

197

198

199

Table 3. Relative gene expression of ECS components in zebrafish ovary.

GENE	CTRL	10 µg/L	20 µg/L
<i>cnr1</i>	2.70 ± 0.552	3.41 ± 0.578	1.96 ± 0.359
<i>trpv1</i>	4.23 ± 1.183	3.69 ± 1.106	2.04 ± 0.492
<i>gpr55</i>	1.69 ± 0.208	2.10 ± 0.539	1.91 ± 0.268
<i>napepld</i>	3.14 ± 1.042	5.27 ± 0.608	4.79 ± 0.378
<i>mgll</i>	2.10 ± 0.541	2.35 ± 0.146	3.87 ± 1.011
<i>faah</i>	2.89 ± 0.988	4.62 ± 0.881	9.67 ± 2.549*
<i>dagla</i>	2.68 ± 0.701	3.28 ± 0.585	3.27 ± 0.559

200

Table 3. RT-qPCR results for the zebrafish ovaries after 21-day exposure to 10 and 20 µg/L BPA. Data are expressed as means $\Delta Ct \pm SEM$. All data were normalized against the expression levels of *18s* and *rplp0*. Asterisks (*) denote statistically significant differences with respect to the control (CTRL): * p < 0.05; (one-way ANOVA, p < 0.05, Dunnett's multiple comparison test). Relative levels of mRNA are in arbitrary units (a.u.).

201

202

203

204

205

Testes. Exposure to BPA was without effect on the genes coding for the cannabinoid receptors. However, BPA had a biphasic effect on *mgll* mRNA levels; 10 µg BPA/L increased *mgll* transcript levels, whereas the highest concentration of BPA (20 µg/L) decreased them (Table 4).

206

207

208


Table 4. Relative gene expression of ECS components in zebrafish testes.

GENE	CTRL	10 µg/L	20 µg/L
<i>cnr1</i>	2.51 ± 0.491	2.32 ± 0.429	2.68 ± 0.661
<i>gpr55</i>	3.35 ± 0.801	5.57 ± 1.214	2.34 ± 0.591

<i>trpv1</i>	3.41 ± 0.633	3.44 ± 0.778	2.93 ± 0.798
<i>napepld</i>	1.84 ± 0.393	1.80 ± 0.405	1.64 ± 0.200
<i>mgll</i>	3.86 ± 0.319	6.24 ± 0.694*	1.32 ± 0.15*
<i>faah</i>	3.56 ± 0.587	3.70 ± 0.989	3.03 ± 1.009
<i>dagla</i>	3.18 ± 0.223	2.73 ± 0.329	2.45 ± 0.667

209 **Table 4.** RT-qPCR results of zebrafish testes treated with 10 and 20 µg/L BPA. Data are expressed as means ± SEM. All
 210 data were normalized against the expression levels of *rpl13* and *rplp0*. Asterisks (*) denote statistically significant
 211 differences with respect to the control (CTRL): * p < 0.05; (one-way ANOVA, p < 0.05, Dunnett's multiple comparison
 212 test). Relative levels of mRNA are reported in arbitrary units (a.u.).

213 Representation of the fold change of the genes analysed above (Figure 2).




214
 215 **Figure 2.** Heat maps for the ECS genes analysed in ovaries and testes. Red: up-regulation. Green: down-regulation. O:
 216 ovaries, T: testes.

217

218 3.4 Vitellogenin transcript levels in the liver.

219 The highest concentration of BPA increased the mRNA levels of vitellogenin (*vtg*) in the liver of both
 220 male (Figure 3a) and female (Figure 3b) zebrafish.




221

222 **Figure 3.** RT-qPCR results for *vtg* in zebrafish liver, in males (a) and females (b), following treatment with 10 and 20 µg/L
 223 BPA. Results are expressed as means $\Delta\text{Ct} \pm \text{SEM}$. All data are normalized against the expression levels of *rpl13* and *rplp0*
 224 (male) and *rplp0* and *18s* (female). Asterisks (*) denote statistically significantly differences with respect to the control

225 (CTRL): * ($p < 0.05$); (one-way ANOVA, $p < 0.05$, Dunnett's multiple comparison test). Levels of mRNA abundance are
226 reported in arbitrary units (a.u.).

227 228 **3.5 Relative area covered by spermatozoa and spermatogonia.**



229 The histological study of zebrafish testes revealed changes in zebrafish testicular development
230 following exposure to BPA (Figure 4a-c). Using image analysis, the area occupied by spermatogonia
231 and spermatozoa were quantified. The 20 μg BPA/L exposure induced a significant decrease of
232 spermatogonia area (Figure 4d) while no effect was observed on the area occupied by spermatozoa
233 (Figure 4e).



234
235 **Figure 4.** Histological sections of testis: CTRL group (a), 10 $\mu\text{g/L}$ BPA (b) and 20 $\mu\text{g/L}$ BPA (d). Eosin – Mayer's haematoxylin
236 staining. Sg: spermatogonia; Sc: spermatocyte; Sd: spermatid; Sz: spermatozoa. Scale bar: 20 μm . Percentage of zebrafish
237 testicular area occupied by spermatogonia (d) and spermatozoa (e). Data reported as means \pm SEM. Asterisks (*) denote
238 statistically significant differences with respect to the control (CTRL): ** $p < 0.01$, (one-way ANOVA, $p < 0.05$, Dunnett's
239 multiple comparison test).

240 241 **3.6 Changes in different classes of oocytes**

242 Using histological analysis, we quantified the proportion of previtellogenic, vitellogenic and mature
243 oocytes in the ovaries of zebrafish exposed to BPA (Figure 5a-c). The results demonstrate that
244 exposure to 10 μg BPA/L significantly augmented the proportion of oocytes in vitellogenic stage,
245 without affecting the percentage of previtellogenic and mature oocytes (Figure 5d-f).



246

247

248

249

250

251

252

253

4. DISCUSSION

254

255

256

257

258

259

260

261

262

Figure 5. Histological sections of zebrafish ovaries, CTRL (a), 10 µg/L BPA (b) and 20 µg/L BPA (c). Eosin – Mayer’s haematoxylin staining. Prev: previtellogenic oocytes; Vit: Vitellogenic oocytes; Mat: mature oocytes. Scale bar: 200 µm. Percentage of different classes of oocytes (d-f). Data are reported as mean ± SEM. Asterisks (*) denote statistically significant differences with respect to the control (CTRL): * p < 0.05, (one-way ANOVA, p < 0.05, Dunnett’s multiple comparison test).

263 appear that duration, window of the exposure, species differences and BPA concentrations are
264 important factors influencing the effects of BPA on the GSI [53]. In this context, in murine models,
265 different sensitivities to BPA were observed depending on the mouse strain [57].

266 Hepatic VTG, a yolk precursor protein and a key signal for female reproduction, is another marker for
267 testing the effects of BPA estrogenicity, commonly used for teleosts [58]. Usually, males have low
268 basal or undetectable VTG levels, but exposure to external compounds with estrogen-like activity can
269 stimulate VTG production. Thus, VTG in male fish has become a useful biomarker to assess the
270 presence of environmental compounds with estrogen-like activity. Indeed, several studies have
271 correlated abnormal values of VTG or *vtg* with BPA in different teleost species [19,27,53,55,59–62].
272 Specifically, Mandich and coworkers (2007) reported an increase of VTG in plasma in common carps
273 after 14 days of BPA treatment. With zebrafish, 21 day exposure was enough to find an increased
274 gene expression of hepatic *vtg* in both sexes, similar to what was observed in *Sparus aurata* after a
275 3-week BPA exposure [26]. Curiously, the increase of *vtg* transcript (20 µg BPA/L) does not match
276 with the group showing the highest percentage of vitellogenic oocytes (10 µg BPA/L). This may be
277 explained as 1) oocyte maturation is a complex process where VTG is not the only player and other
278 hormones (*i.e.* FSH, LH) are involved, 2) VTG is a complex protein which undergoes a strong post-
279 transcriptional process, and additionally, as cited in [63], the RT-qPCR results do not inform whether
280 the mRNA will be or not be translated into protein.

281

282 The regulation of endocannabinoid tone during the different phases of reproduction is crucial in
283 numerous species [31,33,35,64–66]. However, little information is available regarding the role of AEA
284 tone during oocyte maturation [67]. It is tempting to hypothesize that the AEA produced from the
285 granulosa cells of growing follicles play a role in the control of oocyte maturation in mammals [31].
286 In the aquatic model, *Xenopus laevis*, AEA is associated with K⁺ current supression in oocytes (stages
287 V or VI), and consequently, with the inhibition of oocyte responsiveness to gonadotropin and
288 progesterone required for final meiotic maturation [68]. Thus, we may assume that the augmented
289 AEA levels (10 µg BPA/L) may contribute to the signals inducing vitellogenesis but impairing the final
290 oocyte maturation. However, few information is available on the functional role of AEA in teleost
291 ovary, although our hypothesis would be consistent with a previous study [15] demonstrating that
292 10 µg BPA/L is associated with the down-regulation of oocyte maturation signals in female zebrafish.
293 Furthermore, in rodents and humans, the hormonal environment regulates the endocannabinoid
294 production in the gonads [38,69]. Additionally, we observed a decrease in ovarian OEA in the BPA

295 exposed groups. In this context, OEA, which is a AEA-related compound, may be involved in the
296 mammalian follicular maturation [70]. In addition, OEA also plays a role as an anti-inflammatory
297 signal molecule [71,72], suggesting an inflammatory process in the BPA ovaries. In this context, BPA-
298 induced inflammatory responses have been already reported in other species [73].

299

300 Concerning male ECS, reduction in AEA and 2-AG levels was observed following exposure to 20 µg
301 BPA/L in the testis. In male mouse germ cell cultures, the highest levels of 2-AG were observed in
302 spermatogonia and then, such levels were progressively decreased in spermatocytes and spermatids
303 [74]. The latter study is consistent with our findings where the lowest levels of 2-AG were found in
304 the BPA group associated with the reduced area of spermatogonia. In fact, CB2 signaling through 2-
305 AG contributes to the normal progression of the spermatogenesis [75]. Focusing on the other
306 endocannabinoid, AEA, Grimaldi and coworkers demonstrated a constant AEA concentration during
307 spermatogenesis (spermatogonia, spermatocyte and spermatids), whereas here we found a
308 decrease in testicular AEA following BPA exposure. Two factors may explain this finding: 1) the
309 observed reduction in AEA was the result of reduced spermatogonia induced by BPA, and/or 2) an
310 increased activity of FAAH (AEA catabolic enzyme) induced by the estrogen-like activity of BPA,
311 reduced the AEA levels. Based on the present results, we cannot distinguish between these two
312 possibilities. Though, similar results were obtained following E₂ treatment in mouse Sertoli cell [76].
313 Very few studies have quantified testicular cell types, and to date, there are no published
314 observations on the effects of environmental doses of BPA on testicular cell numbers. In zebrafish,
315 higher doses of BPA exclusively reduced the proportion of spermatocytes [18]. Interestingly, our
316 results are in agreement with a study in rats demonstrating that lower concentrations of BPA trigger
317 a reduction in the number of spermatogonia and spermatocytes [21,77]. Furthermore, in male
318 gonads, BPA can induce alterations in miRNAs [78–81] and epigenetic markers, as hyperacetylation
319 of histones and DNA hypermethylation [18], these changes can be also inherited by the offspring and
320 compromise early embryo development depending on the dose and the window of exposure [82,17].
321 Overall, the present results and those obtained by others in different species suggested that exposure
322 to BPA might be associated with low fertility in male zebrafish and reduced numbers of
323 spermatogonia and spermatogenesis.

324

325 5. CONCLUSIONS

326 21-day exposure to environmentally relevant concentrations of BPA induced changes on the ECS. As
327 far as we know, the correct functioning of the ECS is crucial for the normal progression of the
328 reproductive processes in both males and females. Despite of the ECS presence in teleost gonads and
329 its potential role in reproduction, further studies will be needed to fully understand its involvement
330 in fish oogenesis and spermatogenesis. Finally, our results suggested that the ECS may be used as a
331 biomarker for EDC activity related to reproduction.

332

333

6. REFERENCES

334

- 335 1. Rubin, B.S. Bisphenol A: an endocrine disruptor with widespread exposure and multiple effects. *J. Steroid Biochem. Mol. Biol.* **2011**, *127*, 27–34.
- 336 2. Vandenberg, L.N.; Maffini, M. V; Sonnenschein, C.; Rubin, B.S.; Soto, A.M. Bisphenol-A and the great divide: a review of
337 controversies in the field of endocrine disruption. *Endocr. Rev.* **2009**, *30*, 75–95.
- 338 3. Kang, J.-H.; Aasi, D.; Katayama, Y. Bisphenol A in the Aquatic Environment and Its Endocrine-Disruptive Effects on Aquatic
339 Organisms. *Crit. Rev. Toxicol.* **2007**, *37*, 607–625.
- 340 4. Flint, S.; Markle, T.; Thompson, S.; Wallace, E. Bisphenol A exposure, effects, and policy: A wildlife perspective. *J. Environ.*
341 *Manage.* **2012**, *104*, 19–34.
- 342 5. Crain, D.A.; Eriksen, M.; Iguchi, T.; Jobling, S.; Laufer, H.; LeBlanc, G.A.; Guillette, L.J. An ecological assessment of bisphenol-
343 A: Evidence from comparative biology. *Reprod. Toxicol.* **2007**, *24*, 225–239.
- 344 6. Powell, C.E.; Soto, A.M.; Sonnenschein, C. Identification and characterization of membrane estrogen receptor from MCF7
345 estrogen-target cells. *J. Steroid Biochem. Mol. Biol.* **2001**, *77*, 97–108.
- 346 7. Watson, C.S.; Bulayeva, N.N.; Wozniak, A.L.; Alyea, R.A. Xenoestrogens are potent activators of nongenomic estrogenic
347 responses. *Steroids* **2007**, *72*, 124–34.
- 348 8. Kuiper, G.G.; Lemmen, J.G.; Carlsson, B.; Corton, J.C.; Safe, S.H.; van der Saag, P.T.; van der Burg, B.; Gustafsson, J.A.
349 Interaction of estrogenic chemicals and phytoestrogens with estrogen receptor beta. *Endocrinology* **1998**, *139*, 4252–63.
- 350 9. Thomas, P.; Dong, J. Binding and activation of the seven-transmembrane estrogen receptor GPR30 by environmental
351 estrogens: a potential novel mechanism of endocrine disruption. *J. Steroid Biochem. Mol. Biol.* **2006**, *102*, 175–9.
- 352 10. Welshons, W. V.; Nagel, S.C.; vom Saal, F.S. Large Effects from Small Exposures. III. Endocrine Mechanisms Mediating Effects
353 of Bisphenol A at Levels of Human Exposure. *Endocrinology* **2006**, *147*, s56–s69.
- 354 11. Kinch, C.D.; Ibhazehiebo, K.; Jeong, J.-H.; Habibi, H.R.; Kurrasch, D.M. Low-dose exposure to bisphenol A and replacement
355 bisphenol S induces precocious hypothalamic neurogenesis in embryonic zebrafish. *Proc. Natl. Acad. Sci. U. S. A.* **2015**, *112*,
356 1475–80.
- 357 12. Sun, H.; Xu, L.-C.; Chen, J.-F.; Song, L.; Wang, X.-R. Effect of bisphenol A, tetrachlorobisphenol A and pentachlorophenol on
358 the transcriptional activities of androgen receptor-mediated reporter gene. *Food Chem. Toxicol.* **2006**, *44*, 1916–21.
- 359 13. Krüger, T.; Long, M.; Bonefeld-Jørgensen, E.C. Plastic components affect the activation of the aryl hydrocarbon and the
360 androgen receptor. *Toxicology* **2008**, *246*, 112–23.
- 361 14. Gaido, K.W.; Maness, S.C.; McDonnell, D.P.; Dehal, S.S.; Kupfer, D.; Safe, S. Interaction of methoxychlor and related
362 compounds with estrogen receptor alpha and beta, and androgen receptor: structure-activity studies. *Mol. Pharmacol.*
363 **2000**, *58*, 852–8.
- 364 15. Santangeli, S.; Maradonna, F.; Gioacchini, G.; Cobellis, G.; Piccinetti, C.C.; Dalla Valle, L.; Carnevali, O. BPA-Induced
365 Deregulation Of Epigenetic Patterns: Effects On Female Zebrafish Reproduction. *Sci. Rep.* **2016**, *6*, 21982.

- 366 16. Santangeli, S.; Consales, C.; Pacchierotti, F.; Habibi, H.R.; Carnevali, O. Transgenerational effects of BPA on female
367 reproduction. *Sci. Total Environ.* **2019**.
- 368 17. Lombó, M.; Fernández-Díez, C.; González-Rojo, S.; Herráez, M.P. Genetic and epigenetic alterations induced by bisphenol A
369 exposure during different periods of spermatogenesis: from spermatozoa to the progeny. *Sci. Rep.* **2019**, *9*, 1–13.
- 370 18. González-Rojo, S.; Lombó, M.; Fernández-Díez, C.; Herráez, M.P. Male exposure to bisphenol a impairs spermatogenesis and
371 triggers histone hyperacetylation in zebrafish testes. *Environ. Pollut.* **2019**, *248*, 368–379.
- 372 19. Molina, A.M.; Abril, N.; Morales-Prieto, N.; Monterde, J.G.; Lora, A.J.; Ayala, N.; Moyano, R. Evaluation of toxicological
373 endpoints in female zebrafish after bisphenol A exposure. *Food Chem. Toxicol.* **2018**, *112*, 19–25.
- 374 20. Migliaccio, M.; Chioccarelli, T.; Ambrosino, C.; Suglia, A.; Manfredola, F.; Carnevali, O.; Fasano, S.; Pierantoni, R.; Cobellis, G.
375 Characterization of Follicular Atresia Responsive to BPA in Zebrafish by Morphometric Analysis of Follicular Stage
376 Progression. *Int. J. Endocrinol.* **2018**, *2018*, 4298195.
- 377 21. Jin, P.; Wang, X.; Chang, F.; Bai, Y.; Li, Y.; Zhou, R.; Chen, L. Low dose bisphenol A impairs spermatogenesis by suppressing
378 reproductive hormone production and promoting germ cell apoptosis in adult rats. *J. Biomed. Res.* **2013**, *27*, 135–144.
- 379 22. Forner-Piquer, I.; Santangeli, S.; Maradonna, F.; Verde, R.; Piscitelli, F.; di Marzo, V.; Habibi, H.R.; Carnevali, O. Role of
380 Bisphenol A on the Endocannabinoid System at central and peripheral levels: Effects on adult female zebrafish.
381 *Chemosphere* **2018**, *205*, 118–125.
- 382 23. Hatef, A.; Zare, A.; Alavi, S.M.H.; Habibi, H.R.; Linhart, O. Modulations in androgen and estrogen mediating genes and
383 testicular response in male goldfish exposed to bisphenol A. *Environ. Toxicol. Chem.* **2012**, *31*, 2069–2077.
- 384 24. Toyama, Y.; Suzuki-Toyota, F.; Maekawa, M.; Ito, C.; Toshimori, K. Adverse effects of bisphenol A to spermiogenesis in mice
385 and rats. *Arch. Histol. Cytol.* **2004**, *67*, 373–81.
- 386 25. Chen, J.; Xiao, Y.; Gai, Z.; Li, R.; Zhu, Z.; Bai, C.; Tanguay, R.L.; Xu, X.; Huang, C.; Dong, Q. Reproductive toxicity of low level
387 bisphenol A exposures in a two-generation zebrafish assay: Evidence of male-specific effects. *Aquat. Toxicol.* **2015**, *169*,
388 204–14.
- 389 26. Forner-Piquer, I.; Fakriadis, I.; Mylonas, C.C.; Piscitelli, F.; Di Marzo, V.; Maradonna, F.; Calduch-Giner, J.; Pérez-Sánchez, J.;
390 Carnevali, O. Effects of Dietary Bisphenol A on the Reproductive Function of Gilthead Sea Bream (*Sparus aurata*) Testes. *Int.*
391 *J. Mol. Sci.* **2019**, *20*, 5003.
- 392 27. Mandich, A.; Bottero, S.; Benfenati, E.; Cevasco, A.; Erratico, C.; Maggioni, S.; Massari, A.; Pedemonte, F.; Viganò, L. In vivo
393 exposure of carp to graded concentrations of bisphenol A. *Gen. Comp. Endocrinol.* **2007**, *153*, 15–24.
- 394 28. Kundakovic, M.; Gudsruk, K.; Franks, B.; Madrid, J.; Miller, R.; Perera, F.; Champagne, F. Sex-specific epigenetic disruption
395 and behavioral changes following low-dose in utero bisphenol a exposure. *Proc. Natl. Acad. Sci. U. S. A.* **2013**, *110*, 9956–
396 9961.
- 397 29. Kinch, C.D.; Kurrasch, D.M.; Habibi, H.R. Adverse morphological development in embryonic zebrafish exposed to
398 environmental concentrations of contaminants individually and in mixture. *Aquat. Toxicol.* **2016**, *175*, 286–298.
- 399 30. Jordan, J.; Zare, A.; Jackson, L.J.; Habibi, H.R.; Weljie, A.M. Environmental Contaminant Mixtures at Ambient Concentrations
400 Invoke a Metabolic Stress Response in Goldfish Not Predicted from Exposure to Individual Compounds Alone. *J. Proteome*
401 *Res.* **2012**, *11*, 1133–1143.
- 402 31. El-Talatini, M.R.; Taylor, A.H.; Elson, J.C.; Brown, L.; Davidson, A.C.; Konje, J.C. Localisation and Function of the
403 Endocannabinoid System in the Human Ovary. *PLoS One* **2009**, *4*, e4579.
- 404 32. Battista, N.; Bari, M.; Rapino, C.; Trasatti, F.; D'Agostino, A.; Maccarrone, M. Regulation of female fertility by the
405 endocannabinoid system. *Hum. Fertil. (Camb).* **2007**, *10*, 207–16.
- 406 33. Wang, H.; Dey, S.K.; Maccarrone, M. Jekyll and Hyde: Two Faces of Cannabinoid Signaling in Male and Female Fertility.
407 *Endocr. Rev.* **2006**, *27*, 427–448.

- 408 34. Rossato, M.; Ion Popa, F.; Ferigo, M.; Clari, G.; Foresta, C. Human sperm express cannabinoid receptor Cb1, the activation of
409 which inhibits motility, acrosome reaction, and mitochondrial function. *J. Clin. Endocrinol. Metab.* **2005**, *90*, 984–91.
- 410 35. Cobellis, G.; Ricci, G.; Cacciola, G.; Orlando, P.; Petrosino, S.; Cascio, M.G.; Bisogno, T.; De Petrocellis, L.; Chioccarelli, T.;
411 Altucci, L.; et al. A gradient of 2-arachidonoylglycerol regulates mouse epididymal sperm cell start-up. *Biol. Reprod.* **2010**,
412 *82*, 451–8.
- 413 36. Chioccarelli, T.; Cacciola, G.; Altucci, L.; Lewis, S.E.M.; Simon, L.; Ricci, G.; Ledent, C.; Meccariello, R.; Fasano, S.; Pierantoni,
414 R.; et al. Cannabinoid receptor 1 influences chromatin remodeling in mouse spermatids by affecting content of transition
415 protein 2 mRNA and histone displacement. *Endocrinology* **2010**, *151*, 5017–29.
- 416 37. Lazzarin, N.; Valensise, H.; Bari, M.; Ubaldi, F.; Battista, N.; Finazzi-Agrò, A.; Maccarrone, M. Fluctuations of fatty acid amide
417 hydrolase and anandamide levels during the human ovulatory cycle. *Gynecol. Endocrinol.* **2004**, *18*, 212–8.
- 418 38. El-Talatini, M.R.; Taylor, A.H.; Konje, J.C. The relationship between plasma levels of the endocannabinoid, anandamide, sex
419 steroids, and gonadotrophins during the menstrual cycle. *Fertil. Steril.* **2010**, *93*, 1989–96.
- 420 39. Maccarrone, M.; Bari, M.; Di Rienzo, M.; Finazzi-Agrò, A.; Rossi, A. Progesterone activates fatty acid amide hydrolase (FAAH)
421 promoter in human T lymphocytes through the transcription factor Ikaros. Evidence for a synergistic effect of leptin. *J. Biol.*
422 *Chem.* **2003**, *278*, 32726–32.
- 423 40. Gorzalka, B.B.; Dang, S.S. Minireview: Endocannabinoids and Gonadal Hormones: Bidirectional Interactions in Physiology
424 and Behavior. *Endocrinology* **2012**, *153*, 1016–1024.
- 425 41. Zbucka-Kretowska, M.; Zbucki, R.; Parfieniuk, E.; Maslyk, M.; Lazarek, U.; Miltyk, W.; Czerniecki, J.; Wolczynski, S.; Kretowski,
426 A.; Ciborowski, M. Evaluation of Bisphenol A influence on endocannabinoid system in pregnant women. *Chemosphere* **2018**,
427 *203*, 387–392.
- 428 42. Martella, A.; Silvestri, C.; Maradonna, F.; Gioacchini, G.; Allara, M.; Radaelli, G.; Overby, D.R.; Di Marzo, V.; Carnevali, O.
429 Bisphenol a induces fatty liver by an endocannabinoid-mediated positive feedback loop. *Endocrinology* **2016**, *157*, 1751–
430 1763.
- 431 43. Forner-Piquer, I.; Mylonas, C.C.; Calduch-Giner, J.; Maradonna, F.; Gioacchini, G.; Allarà, M.; Piscitelli, F.; Di Marzo, V.; Pérez-
432 Sánchez, J.; Carnevali, O. Endocrine disruptors in the diet of male *Sparus aurata* : Modulation of the endocannabinoid
433 system at the hepatic and central level by Di-isononyl phthalate and Bisphenol A. *Environ. Int.* **2018**, *119*, 54–65.
- 434 44. Corrales, J.; Kristofco, L.A.; Steele, W.B.; Yates, B.S.; Breed, C.S.; Williams, E.S.; Brooks, B.W. Global Assessment of Bisphenol
435 A in the Environment. *Dose-Response* **2015**, *13*, 155932581559830.
- 436 45. Pfaffl, M.W. A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic Acids Res.* **2001**, *29*, e45.
- 437 46. Vandesompele, J.; De Preter, K.; Pattyn, F.; Poppe, B.; Van Roy, N.; De Paepe, A.; Speleman, F. Accurate normalization of
438 real-time quantitative RT-PCR data by geometric averaging of multiple internal control genes. *Genome Biol.* **2002**, *3*,
439 RESEARCH0034.
- 440 47. Forner-Piquer, I.; Santangeli, S.; Maradonna, F.; Rabbito, A.; Piscitelli, F.; Habibi, H.R.; Di, V. Disruption of the gonadal
441 endocannabinoid system in zebra fish exposed to diisononyl phthalate *. *Environ. Pollut.* **2018**, *241*, 1–8.
- 442 48. Di Marzo, V.; Goparaju, S.K.; Wang, L.; Liu, J.; Batkai, S.; Jarai, Z.; Fezza, F.; Miura, G.I.; Palmiter, R.D.; Sugiura, T.; et al.
443 Leptin-regulated endocannabinoids are involved in maintaining food intake. *Nature* **2001**, *410*, 822–825.
- 444 49. Qin, F.; Wang, L.; Wang, X.; Liu, S.; Xu, P.; Wang, H.; Wu, T.; Zhang, Y.; Zheng, Y.; Li, M.; et al. Bisphenol A affects gene
445 expression of gonadotropin-releasing hormones and type I GnRH receptors in brains of adult rare minnow *Gobiocypris*
446 *rarus*. *Comp. Biochem. Physiol. C. Toxicol. Pharmacol.* **2013**, *157*, 192–202.
- 447 50. Akingbemi, B.T.; Sottas, C.M.; Koulova, A.I.; Klinefelter, G.R.; Hardy, M.P. Inhibition of testicular steroidogenesis by the
448 xenoestrogen bisphenol A is associated with reduced pituitary luteinizing hormone secretion and decreased steroidogenic
449 enzyme gene expression in rat Leydig cells. *Endocrinology* **2004**, *145*, 592–603.

- 450 51. Hong, J.; Chen, F.; Wang, X.; Bai, Y.; Zhou, R.; Li, Y.; Chen, L. Exposure of preimplantation embryos to low-dose bisphenol A
451 impairs testes development and suppresses histone acetylation of StAR promoter to reduce production of testosterone in
452 mice. *Mol. Cell. Endocrinol.* **2016**, *427*, 101–11.
- 453 52. Zhang, Y.; Yuan, C.; Hu, G.; Li, M.; Zheng, Y.; Gao, J.; Yang, Y.; Zhou, Y.; Wang, Z. Characterization of four nr5a genes and
454 gene expression profiling for testicular steroidogenesis-related genes and their regulatory factors in response to bisphenol A
455 in rare minnow *Gobiocypris rarus*. *Gen. Comp. Endocrinol.* **2013**, *194*, 31–44.
- 456 53. Sohoni, P.; Tyler, C.R.; Hurd, K.; Caunter, J.; Hetheridge, M.; Williams, T.; Woods, C.; Evans, M.; Toy, R.; M. Gargas, A.; et al.
457 Reproductive Effects of Long-Term Exposure to Bisphenol A in the Fathead Minnow (*Pimephales promelas*). *Environ. Sci.*
458 *Technol.* **2001**, *35*, 2917–2925.
- 459 54. Sun, S.-X.; Wu, J.-L.; Lv, H.-B.; Zhang, H.-Y.; Zhang, J.; Limbu, S.M.; Qiao, F.; Chen, L.-Q.; Yang, Y.; Zhang, M.-L.; et al.
460 Environmental estrogen exposure converts lipid metabolism in male fish to a female pattern mediated by AMPK and mTOR
461 signaling pathways. *J. Hazard. Mater.* **2020**, *394*, 122537.
- 462 55. Hatef, A.; Alavi, S.M.H.; Abdulfatah, A.; Fontaine, P.; Rodina, M.; Linhart, O. Adverse effects of bisphenol A on reproductive
463 physiology in male goldfish at environmentally relevant concentrations. *Ecotoxicol. Environ. Saf.* **2012**, *76*, 56–62.
- 464 56. Wang, Q.; Yang, H.; Yang, M.; Yu, Y.; Yan, M.; Zhou, L.; Liu, X.; Xiao, S.; Yang, Y.; Wang, Y.; et al. Toxic effects of bisphenol A
465 on goldfish gonad development and the possible pathway of BPA disturbance in female and male fish reproduction.
466 *Chemosphere* **2019**, *221*, 235–245.
- 467 57. National Toxicology Program *National Toxicology Program's Report of the Endocrine Disruptors* ; 2001;
- 468 58. European Chemicals Agency *Substance Name: 4,4'-isopropylidenediphenol (Bisphenol A; BPA) EC Number: 201-245-8 CAS*
469 *Number: 80-05-7 SUPPORT DOCUMENT - 4,4'-ISOPROPYLIDENEDIPHENOL (BISPHENOL A, BPA)* ; 2017;
- 470 59. Kang, I.J.; Yokota, H.; Oshima, Y.; Tsuruda, Y.; Oe, T.; Imada, N.; Tadokoro, H.; Honjo, T. Effects of bisphenol a on the
471 reproduction of Japanese medaka (*Oryzias latipes*). *Environ. Toxicol. Chem.* **2002**, *21*, 2394–2400.
- 472 60. Zhao, F.; Wei, P.; Wang, J.; Yu, M.; Zhang, X.; Tian, H.; Wang, W.; Ru, S. Estrogenic effects associated with bisphenol a
473 exposure in male zebrafish (*Danio rerio*) is associated with changes of endogenous 17 β -estradiol and gene specific DNA
474 methylation levels. *Gen. Comp. Endocrinol.* **2017**, *252*, 27–35.
- 475 61. Zhang, Y.; Guan, Y.; Fan, X.; Wu, L.; Wang, Z. Bisphenol A regulates rare minnow testicular vitellogenin expression via
476 reducing its promoter Er recruitment. *Ecotoxicol. Environ. Saf.* **2018**, *147*, 423–429.
- 477 62. Larsen, B.K.; Bjørnstad, A.; Sundt, R.C.; Taban, I.C.; Pampanin, D.M.; Andersen, O.K. Comparison of protein expression in
478 plasma from nonylphenol and bisphenol A-exposed Atlantic cod (*Gadus morhua*) and turbot (*Scophthalmus maximus*) by
479 use of SELDI-TOF. *Aquat. Toxicol.* **2006**, *78*.
- 480 63. Bustin, S.A.; Benes, V.; Garson, J.A.; Hellemans, J.; Huggett, J.; Kubista, M.; Mueller, R.; Nolan, T.; Pfaffl, M.W.; Shipley, G.L.;
481 et al. The MIQE Guidelines: Minimum Information for Publication of Quantitative Real-Time PCR Experiments. *Clin. Chem.*
482 **2009**, *55*, 611–622.
- 483 64. Chioccarelli, T.; Cacciola, G.; Altucci, L.; Lewis, S.E.M.; Simon, L.; Ricci, G.; Ledent, C.; Meccariello, R.; Fasano, S.; Pierantoni,
484 R.; et al. Cannabinoid Receptor 1 Influences Chromatin Remodeling in Mouse Spermatids by Affecting Content of Transition
485 Protein 2 mRNA and Histone Displacement. *Endocrinology* **2010**, *151*, 5017–5029.
- 486 65. Battista, N.; Bari, M.; Rapino, C.; Trasatti, F.; D'Agostino, A.; Maccarrone, M. Regulation of female fertility by the
487 endocannabinoid system. *Hum. Fertil.* **2007**, *10*, 207–216.
- 488 66. Sun, X.; Wang, H.; Okabe, M.; Mackie, K.; Kingsley, P.J.; Marnett, L.J.; Cravatt, B.F.; Dey, S.K. Genetic loss of Faah
489 compromises male fertility in mice. *Biol. Reprod.* **2009**, *80*, 235–42.
- 490 67. Agirregoitia, E.; Ibarra-Lecue, I.; Totorikaguena, L.; Mendoza, R.; Expósito, A.; Matorras, R.; Urigüen, L.; Agirregoitia, N.
491 Dynamics of expression and localization of the cannabinoid system in granulosa cells during oocyte nuclear maturation.

- 492 *Fertil. Steril.* **2015**, *104*, 753–760.
- 493 68. Oz, M.; Yang, K.-H.; Dinc, M.; Shippenberg, T.S. The Endogenous Cannabinoid Anandamide Inhibits Cromakalim-Activated K⁺
- 494 Currents in Follicle-Enclosed *Xenopus* Oocytes. *J. Pharmacol. Exp. Ther.* **2007**, *323*, 547–554.
- 495 69. Bradshaw, H.B.; Allard, C. Endogenous cannabinoid production in the rat female reproductive tract is regulated by changes
- 496 in the hormonal milieu. *Pharmaceuticals* **2011**, *4*, 933–949.
- 497 70. Schuel, H.; Burkman, L.J.; Lippes, J.; Crickard, K.; Forester, E.; Piomelli, D.; Giuffrida, A. N-Acylethanolamines in human
- 498 reproductive fluids. *Chem. Phys. Lipids* **2002**, *121*, 211–227.
- 499 71. Xu, X.; Guo, H.; Jing, Z.; Yang, L.; Chen, C.; Peng, L.; Wang, X.; Yan, L.; Ye, R.; Jin, X.; et al. N-Oleylethanolamine Reduces
- 500 Inflammatory Cytokines and Adhesion Molecules in TNF- α -induced Human Umbilical Vein Endothelial Cells by Activating
- 501 CB2 and PPAR- α . *J. Cardiovasc. Pharmacol.* **2016**, *68*, 280–291.
- 502 72. Payahoo, L.; Khajebishak, Y.; Jafarabadi, M.A.; Ostadrahimi, A. Oleylethanolamide supplementation reduces inflammation
- 503 and oxidative stress in obese people: A clinical trial. *Adv. Pharm. Bull.* **2018**, *8*, 479–487.
- 504 73. Murata, M.; Kang, J.H. Bisphenol A (BPA) and cell signaling pathways. *Biotechnol. Adv.* **2018**, *36*, 311–327.
- 505 74. Grimaldi, P.; Orlando, P.; Di Siena, S.; Lolicato, F.; Petrosino, S.; Bisogno, T.; Geremia, R.; De Petrocellis, L.; Di Marzo, V. The
- 506 endocannabinoid system and pivotal role of the CB2 receptor in mouse spermatogenesis. *Proc. Natl. Acad. Sci. U. S. A.* **2009**,
- 507 *106*, 11131–6.
- 508 75. Di Giacomo, D.; De Domenico, E.; Sette, C.; Geremia, R.; Grimaldi, P. Type 2 cannabinoid receptor contributes to the
- 509 physiological regulation of spermatogenesis. *FASEB J.* **2016**, *30*, 1453–1463.
- 510 76. Rossi, G.; Gasperi, V.; Paro, R.; Barsacchi, D.; Cecconi, S.; Maccarrone, M. Follicle-Stimulating Hormone Activates Fatty Acid
- 511 Amide Hydrolase by Protein Kinase A and Aromatase-Dependent Pathways in Mouse Primary Sertoli Cells. *Endocrinology*
- 512 **2007**, *148*, 1431–1439.
- 513 77. Ullah, A.; Pirzada, M.; Jahan, S.; Ullah, H.; Turi, N.; Ullah, W.; Siddiqui, M.F.; Zakria, M.; Lodhi, K.Z.; Khan, M.M. Impact of
- 514 low-dose chronic exposure to bisphenol A and its analogue bisphenol B, bisphenol F and bisphenol S on hypothalamo-
- 515 pituitary-testicular activities in adult rats: A focus on the possible hormonal mode of action. *Food Chem. Toxicol.* **2018**, *121*,
- 516 24–36.
- 517 78. Avissar-Whiting, M.; Veiga, K.R.; Uhl, K.M.; Maccani, M.A.; Gagne, L.A.; Moen, E.L.; Marsit, C.J. Bisphenol A exposure leads
- 518 to specific microRNA alterations in placental cells. *Reprod. Toxicol.* **2010**, *29*, 401–406.
- 519 79. Veiga-Lopez, A.; Luense, L.J.; Christenson, L.K.; Padmanabhan, V. Developmental programming: Gestational bisphenol-A
- 520 treatment alters trajectory of fetal ovarian gene expression. *Endocrinology* **2013**, *154*, 1873–1884.
- 521 80. Ma, L.; Yu, H.; Wang, X.; Li, D.; Zhang, Y.; Pei, X.; Duan, Z.; Ma, M. The effects of maternal exposure to BPA during pregnancy
- 522 on the male reproductive system and the testicular microRNA expression profile. *Environ. Sci. Pollut. Res.* **2020**, 1–13.
- 523 81. Gao, G. zheng; Zhao, Y.; Li, H. xin; Li, W. Bisphenol A-elicited miR-146a-5p impairs murine testicular steroidogenesis through
- 524 negative regulation of Mta3 signaling. *Biochem. Biophys. Res. Commun.* **2018**, *501*, 478–485.
- 525 82. Lombó, M.; Getino-álvarez, L.; Depincé, A.; Labbé, C.; Herráez, M.P. Embryonic exposure to bisphenol a impairs primordial
- 526 germ cell migration without jeopardizing male breeding capacity. *Biomolecules* **2019**, *9*.

528 **Conflicts of interests:** none.

529

530 **Funding:** This work has received funding from *Progetti di Rilevante Interesse Nazionale* (PRIN) 2010

531 – 2011 (prot 2010W87LBJ) to OC.

Declaration of interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:

Author statement

Isabel Forner-Piquer: conceptualization, formal analysis, investigation, writing-original draft, visualization;

Silvia Beato: formal analysis, investigation, writing-original draft, visualization;

Fabiana Piscitelli: methodology, formal analysis;

Stefania Santangeli: formal analysis, investigation;

Vincenzo di Marzo: methodology, resources, writing - review, supervision;

Hamid R. Habibi: methodology, resources, writing - review funding acquisition;

Francesca Maradonna: conceptualization, investigation;

Oliana Carnevali: conceptualization, methodology, validation, resources, writing - review, supervision, project administration, funding acquisition.

Table S1. Additional information of primers.

GENE	Exon span	Amplicon size	Ta (°C)	TISSUE	ZFIN ID	Reference
Cannabinoid receptor 1	No	181	55	ovary, testis	ZDB-GENE-040312-3	[1]
Transient receptor potential cation channel, subfamily V, member 1	Yes	129	59	ovary, testis	ZDB-GENE-030912-8	[2]
G protein-coupled receptor 55	No	91	53	ovary, testis	ZDB-GENE-051113-260	[3]
N-acyl phosphatidylethanolamine phospholipase D	No	121	57	ovary, testis	ZDB-GENE-030131-3856	[1]
Monoglyceride lipase	No	118	53	ovary, testis	ZDB-GENE-031006-9	[1]
Fatty acid amide hydrolase	No	202	60	ovary, testis	ZDB-GENE-070619-3	[4]
Diacylglycerol lipase alpha	No	58	59	ovary, testis	ZDB-GENE-070619-1	[1]
Abhydrolase domain containing 4	Yes	132	60	ovary, testis	ZDB-GENE-050417-83	[4]
Vitellogenin 1	Yes	209	60	liver	ZDB-GENE-001201-1	[5]
Ribosomal subunit 18s	No	90	55	ovary, liver	ZDB-GENE-070410-9	[6]
Ribosomal protein large P0	No	160	60	ovary, testis, liver	ZDB-GENE-000629-1	[5]
Ribosomal protein L13a	Yes	147	59	testis, liver	ZDB-GENE-030131-168	[7]

REFERENCES

- Martella, A.; Silvestri, C.; Maradonna, F.; Gioacchini, G.; Allara, M.; Radaelli, G.; Overby, D.R.; Di Marzo, V.; Carnevali, O. Bisphenol a induces fatty liver by an endocannabinoid-mediated positive feedback loop. *Endocrinology* **2016**, *157*, 1751–1763.
- Forner-Piquer, I.; Santangeli, S.; Maradonna, F.; Rabbito, A.; Piscitelli, F.; Habibi, H.R.; Di, V. Disruption of the gonadal endocannabinoid system in zebra fish exposed to diisononyl phthalate *. *Environ. Pollut.* **2018**, *241*, 1–8.
- Martella, A.; Sepe, R.M.; Silvestri, C.; Zang, J.; Fasano, G.; Carnevali, O.; De Girolamo, P.; Neuhauss, S.C.F.; Sordino, P.; Di Marzo, V. Important role of endocannabinoid signaling in the development of functional vision and locomotion in zebrafish. *FASEB J.* **2016**, *30*, 4275–4288.
- Forner-Piquer, I.; Maradonna, F.; Gioacchini, G.; Santangeli, S.; Allarà, M.; Piscitelli, F.; Habibi, H.R.H.R.; Di Marzo, V.; Carnevali, O.; Allara, M.; et al. Dose-specific effects of di-isononyl phthalate on the endocannabinoid system and on liver of female zebrafish. *Endocrinology* **2017**, *158*, 3462–3476.
- Gioacchini, G.; Lombardo, F.; Merrifield, D.L.; Silvi, S.; Alberto, C.; Avella, M.A.; Carnevali, O. Effects of probiotic on zebrafish reproduction. *J. Aquac. Res. Dev.* **2011**, *s1*, 1–6.
- Santangeli, S.; Maradonna, F.; Gioacchini, G.; Cobellis, G.; Piccinetti, C.C.; Dalla Valle, L.; Carnevali, O. BPA-Induced Deregulation Of Epigenetic Patterns: Effects On Female Zebrafish Reproduction. *Sci. Rep.* **2016**, *6*, 21982.
- Tang, R.; Dodd, A.; Lai, D.; MCNABB, W.C.; LOVE, D.R. Validation of Zebrafish (*Danio rerio*) Reference Genes for Quantitative Real-time RT-PCR Normalization. *Acta Biochim. Biophys. Sin. (Shanghai)*. **2007**, *39*, 384–390.