

#### UNIVERSITÀ POLITECNICA DELLE MARCHE Repository ISTITUZIONALE

Long-term survival after surgical aortic valve replacement in patients aged 80 years and over

This is the peer reviewd version of the followng article:

Original

Long-term survival after surgical aortic valve replacement in patients aged 80 years and over / Malvindi, P. G.; Luthra, S.; Giritharan, S.; Kowalewski, M.; Ohri, S.. - In: EUROPEAN JOURNAL OF CARDIO-THORACIC SURGERY. - ISSN 1010-7940. - 60:3(2021), pp. 671-678. [10.1093/ejcts/ezab135]

Availability:

This version is available at: 11566/301706 since: 2024-12-03T15:31:20Z

Publisher:

Published DOI:10.1093/ejcts/ezab135

Terms of use:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. The use of copyrighted works requires the consent of the rights' holder (author or publisher). Works made available under a Creative Commons license or a Publisher's custom-made license can be used according to the terms and conditions contained therein. See editor's website for further information and terms and conditions. This item was downloaded from IRIS Università Politecnica delle Marche (https://iris.univpm.it). When citing, please refer to the published version.

#### Publisher copyright:

Oxford Academic- Postprint/Author's accepted Manuscript

This is a pre-copyedited, author-produced version of an article accepted for publication in EUROPEAN JOURNAL OF CARDIO-THORACIC SURGERY following peer review. The version of record of the above quoted article is available online at: https://doi.org/10.1093/ejcts/ezab135

# European Journal of Cardio-Thoracic Surgery

### Long-term survival after surgical aortic valve replacement in patients aged 80 and over --Manuscript Draft--

Manuscript Number:	EJCTS-2020-902681R2	
Full Title:	Long-term survival after surgical aortic valve replacement in patients aged 80 and over	
Article Type:	Original Article	
Order of Authors:	Pietro Giorgio Malvindi, MD, PhD	
	Suvitesh Luthra	
	Suresh Giritharan	
	Mariusz Kowalewski	
	Sunil Ohri	
Corresponding Author:	Pietro Giorgio Malvindi, MD, PhD University Hospital Southampton Southampton, UNITED KINGDOM	
Corresponding Author E-Mail:	pg.malvindi@hotmail.com	
Section/Category:	Valves	
Manuscript Classifications:	Clinical epidemiology; Valve disease	
Author Comments:	Southampton 08.02.2021	
	To the attention of	
	Matthias Siepe, Editor-in-Chief of European Journal of Cardio-Thoracic Surgery Thierry Carrel, Associate Editor of European Journal of Cardio-Thoracic Surgery Dear Editors, thank you for the time invested in our manuscript. The suggestions from the reviewers were of utmost importance in building up a meaningful discussion. Furthermore, the remarks of the Statistical Reviewer were always punctual and helped in providing a linear stat design and presentation of the results. We went through the last comments and hopefully we are able to provide adequate answers and changes. Whatever the outcome of the submissions, the review process of your Journal represents a valuable occasion of discussion and learning. Best regards Pietro G Malvindi Wessex Cardiothoracic Department University Hospital Southampton	
Abstract:	Objectives. Surgical aortic valve replacement can be safely performed in people aged 80 and older with early benefits on both symptomatic and prognostic ground. Whilst new approaches are advocated to treat this elderly and frail population, data on long-term outcomes are not available. Methods. We conducted a retrospective analysis of 1870 patients aged 80 and over who underwent first time surgical aortic valve replacement during the period 2000 – 2019. The Kaplan-Meier method was used to calculate survival and comparisons	

	among groups was performed by log-rank test. Cox analysis was used to determine independent risk factors for late mortality. Results. Patients' mean age was 84 years and 53% were male. Isolated aortic valve replacement was performed in 42% of the patients, CABG (n=956), mitral valve (n=94) or aortic surgery (n=69) were associated in the remaining cases. One hundred eighty-one patients (8%) sustained at least one postoperative complication (reopening for bleeding or tamponade 3%, renal replacement therapy 3%, new cerebral stroke 1.5%). In-hospital mortality was 3.2% in the overall population (60/1870) and 2.2% after isolated aortic valve replacement (18/790). Survival was 90%, 66%, 31%, 14% at 1-year, 5-years, 10-years and 15-years, respectively, and was similar to the expected survival of a sex- and age-matched population (log-rank p=0.96). A complicated postoperative course was an independent risk factor for mortality during the follow-up [hazard ratio 1.32 (1.03, 1.68), p=0.026]. Conclusions. Surgical aortic valve replacement can be performed with an acceptable early mortality rate and provides excellent long-term survival in people aged 80 and older.
Response to Reviewers:	Reviewer 1: The revised version of this paper is good. I do not have any more comments. Reviewer 2: With the additional information provided by the authors, complying with the reviewers' request, now the manuscript becomes a valuable piece of evidence for aortic valve replacement in patients aged 80 years and over.
	Reviewer 3: The authors responded well to the reviewers' comments. I agree that this manuscript could represent a benchmark for the forthcoming long-term results of non- surgical treatment of aortic valve. Now I agree that this manuscript could be accepted for publication in EJCTS. Answer. We want to thank you for your constructive reviews, they were all fundamental suggestions driving meaningful messages from our numbers and analyses.
	Reviewer 4: STATISTICAL REVIEW The authors took my comments into consideration. As a result of survival analyses with different starting points, readers might be confused. Hence, it is important to formulate the results precisely. Here are my comments: - As mentioned already in my review of the original version, the starting point of survival should be mentioned in the label of the time axis. According to the Statistical and data reporting guidelines https://emea01.safelinks.protection.outlook.com/?url=https%3A%2F%2Fdoi.org%2F10. 1093%2Fejcts%2Fezv168&data=04%7C01%7C%7C7b539a4510a940f0a3b908d 8cc1654aa%7C84df9e7fe9f640afb435aaaaaaaaaaaa%7C1%7C0%7C637483743562 922721%7CUnknovn%7CTWFpbGZsb3d8eyJWljoiMC4wLjAwMDAiLCJQIjoiV2luMzli LCJBTil6lk1haWwiLCJXVCI6Mn0%3D%7C1000&sdata=YtkwOCEBfiaXyrbTuyn0 9GfOyfl_gPubd2PoC3qgNklw%3D&reserved=0, the labels of Kaplan-Meier plots including the central figure should be "Time from surgery (years)" or "Time from discharge (years)" (supplemental figure 1A) instead of "years". - In the statistical analysis section, the starting points of survival should be mentioned. - The starting point should be mentioned when it is not obvious. In the section "Long- term outcomes", the starting point is surgery, then it switches to discharge, then goes back to surgery, before it switches to discharge (supplemental figure 1A) and surgery (supplemental figure 1B) again. The authors might change the order of analyses to improve readability. The starting point has to be mentioned after every switch. - To make figures and tables self contained, the starting point has to be mentioned in the titles or legends. Answer. We agree with this remark. Summarising, all the KM analyses have the starting point on the date of surgery, while the Cox analysis considers the time from discharge. We have punctually reported the starting point wherever necessary in the main text and legends. Regarding the figures we have opted for three different ways because of different data involv

	<ul> <li>-Figure 3/Central Figure: since there is no surgery date to be considered for the general population, we have opted for expanding the legend of the curve of the surgical population;</li> <li>-Supplemental Figure 1: similar considerations for Figure 3. In this case we have preferred to underline the starting point in the figure legends. Expanding the legends for 4 curves would have had a negative impact on the graph layout.</li> <li>Finally, we welcomed your idea of changing the order of the paragraph in "Long-term outcomes". It sounds more readable.</li> </ul>
	Changes. Page 8 lines 9 and 15 Page 10 lines 22-23 Page 11 lines 3-6 and 15 Page 17 lines 4 and 8 Page 21 line 1 Page 22 line 2 Suppl Page 1 lines 2 and 9
	- Supplemental figure 1B: The figure does not show "cumulative risk of mortality" but cumulative survival. Answer. Thank you, corrected. Changes. Suppl Page 1 line 9
	<ul> <li>Abstract, results: Text like "0.026; 1.317 (1.033, 1.680)" is incomprehensible. In the results, it might be replaced by something like "p=0.026; HR 1.32, 95% Cl 1.03 to 1.68". In the abstract, "(hazard ratio 1.32 (1.03,1.68), P=0.026)" would have only one word more than the incomprehensible text. Note that I would prefer HRs rounded to two decimal places.</li> <li>Answer. Thank you, corrected.</li> <li>Changes.</li> <li>Page 3 line 17</li> <li>Page 11 lines 17-20</li> <li>Page 21 Table</li> <li>Page 22 Table</li> </ul>
Order of Authors (with Contributor Roles):	Pietro Giorgio Malvindi, MD, PhD (Conceptualization; Data curation; Formal analysis; Investigation; Methodology; Validation; Writing – original draft)
	Suvitesh Luthra (Conceptualization; Data curation; Investigation; Methodology; Writing – original draft)
	Suresh Giritharan (Conceptualization; Data curation; Formal analysis; Investigation; Methodology; Validation; Writing – review & editing)
	Mariusz Kowalewski (Data curation; Formal analysis; Methodology; Software; Validation; Writing – review & editing)
	Sunil Ohri (Conceptualization; Investigation; Methodology; Project administration; Supervision; Validation; Writing – original draft)

#### Central image



1	Title: Long-term Survival After Surgical Aortic Valve Replacement in Patients Aged 80 and
2	Over
3	Running head: survival after aortic valve replacement
4	Pietro Giorgio Malvindi <sup>1</sup> , Suvitesh Luthra <sup>1</sup> , Suresh Giritharan <sup>1</sup> , Mariusz Kowalewski <sup>1,2</sup> , Sunil Ohri <sup>1</sup>
5	1 - Wessex Cardiothoracic Centre, University Hospital Southampton, Southampton – UK
6	2 - Clinical Department of Cardiac Surgery, Central Clinical Hospital of the Ministry of Interior and
7	Administration, Centre of Postgraduate Medical Education, Warsaw, Poland
8	
9	Word count: 4494
10	
11	
12	Corresponding author:
13	PG Malvindi
14	Wessex Cardiothoracic Centre
15	University Hospital Southampton
16	Tremona Road
17	SO16 6YD, Southampton – UK
18	Tel:0044(0)2380777222
19	e-mail:pg.malvindi@hotmail.com
20	

1	Visual abstract
2	
3	<b>Key question:</b> what is the long-term survival of people aged over 80 undergoing aortic valve surgery?
4	Key findings: survival is 30% at 10-year and similar to the expected survival of an age- and sex-
5	matched population
6	Take-home message: surgical aortic valve replacement is safe in octogenarian and older patients
7	and provide a good long-term survival

#### Abstract

Objectives. Surgical aortic valve replacement can be safely performed in people aged 80 and older
with early benefits on both symptomatic and prognostic ground. Whilst new approaches are
advocated to treat this elderly and frail population, data on long-term outcomes are not available.

Methods. We conducted a retrospective analysis of 1870 patients aged 80 and over who underwent
first time surgical aortic valve replacement during the period 2000 – 2019. The Kaplan-Meier method
was used to calculate survival and comparisons among groups was performed by log-rank test. Cox
analysis was used to determine independent risk factors for late mortality.

Results. Patients' mean age was 84 years and 53% were male. Isolated aortic valve replacement was 9 10 performed in 42% of the patients, CABG (n=956), mitral valve (n=94) or aortic surgery (n=69) were associated in the remaining cases. One hundred eighty-one patients (8%) sustained at least one 11 postoperative complication (reopening for bleeding or tamponade 3%, renal replacement therapy 3%, 12 13 new cerebral stroke 1.5%). In-hospital mortality was 3.2% in the overall population (60/1870) and 2.2% after isolated aortic valve replacement (18/790). Survival was 90%, 66%, 31%, 14% at 1-year, 14 15 5-years, 10-years and 15-years, respectively, and was similar to the expected survival of a sex- and 16 age-matched population (log-rank p=0.96). A complicated postoperative course was an independent risk factor for mortality during the follow-up [hazard ratio 1.32 (1.03, 1.68), p=0.026]. 17

Conclusions. Surgical aortic valve replacement can be performed with an acceptable early mortality
rate and provides excellent long-term survival in people aged 80 and older.

20 Abstract word count: 250

21 Keywords: aortic valve, aortic valve replacement, aortic valve prosthesis

#### 1 Introduction

Cardiac surgery has been increasingly performed in octogenarian and older patients during the last 2 decades (1)(2). The initial experiences reported a postoperative mortality up to 17% (3)(4)(5)(6), 3 4 however, more recent studies reported a progressive reduction of early mortality for these patients (7)(8). Particularly, surgical aortic valve replacement can be nowadays safely performed in patients 5 aged 80 and over with an acceptable in-hospital mortality rate (around 2%) (9)(10)(11) and the 6 7 provision of improved survival (12) and a satisfactory quality of life at mid-term follow-up (5)(7). Transcatheter techniques have shown similar procedural safety while providing a shorter hospital stay 8 9 and a quicker recovery (7)(11). These advantages have not translated so far into a better mid-term 10 survival (13), nevertheless, long-term results and the potential impact on the long run of postoperative complications such as acute kidney injury and bleeding, especially in surgical patients (8)(11), or a 11 high incidence of pacemaker implantation and paravalvular regurgitation in patients treated with 12 transcatheter procedures (14)(15), remain undetermined. 13

No long-term data is available for octogenarian and older patients who underwent surgical aortic valve replacement, for this reason we have reviewed our surgical experience of the last 20 years in order to provide a picture of the survival of this population and to study the risk factors associated with mortality during the follow-up.

18

#### **19** Materials and Methods

#### 20 Ethical statement

Approval was obtained for use of data (Safeguard System approval number SEV/0029, date
24.10.2018). Considering the type of the study involving anonymised and previously collected data,
patients' consent was waived.

1	Population
2	The internal database of Wessex Cardiothoracic Centre at UHS was searched to identify patients who
3	underwent aortic valve replacement during the period January 2000 - December 2019 using the
4	following criteria.
5	
6	Inclusion criteria
7	- Age≥80 years;
8	- First time sternotomy;
9	- Associated procedures including CABG, mitral valve surgery, tricuspid valve surgery,
10	aortic surgery;
11	Exclusion criteria
12	- Redo procedure (any previous cardiac operation);
13	
14	A total of 1870 patients fulfilled the criteria.
15	
16	Study design, data collection and outcomes
17	This study is a retrospective outcome evaluation from institutional records with prospective data entry
18	collected and used in compliance with institutional data protection and confidentiality policies. The
19	data were collected from the hospital database system, patients records and records of the general
20	practitioners.
21	The following data were collected:

22 Preoperative details:

1 -	Age;
-----	------

- 2 Gender;
- 3 History/diagnosis of systemic hypertension;
- 4 Diabetes Mellitus;
- 5 COPD;
- 6 Smoking history (ex-smoker or active smoker);
- 7 Creatinine>200 μmol/L;
- 8 Previous cerebral stroke;
- 9 Extracardiac arteriopathy;
- 10 Previous acute myocardial infarction;
- 11 NYHA class;
- 12 LVEF;
- 13 Haemodynamic of valve pathology (aortic stenosis, regurgitation);
- 14 Operation status (elective/urgent);
- 15 Logistic EuroSCORE.
- 16
- 17 Operative data:
- Associated procedures (CABG, MV surgery, TV surgery, aortic surgery);
- 19 Cross-clamp time;
- 20 Cardiopulmonary bypass time.

- 1 Postoperative outcomes:
- 2 In-hospital mortality (death before hospital discharge);
- 3 Reopening for bleeding/tamponade;
- 4 Renal replacement therapy/CVVHD;
- 5 New postoperative neurologic deficit (permanent stroke, TIA);
- 6 Deep sternal wound infection;
- 7 Permanent pacemaker implantation (pre-discharge and post-discharge);
- 8 Postoperative hospital length of stay;
- 9 Discharge destination.

- 11 Long-term outcomes:
- 12 Survival.
- 13

#### 14 Definitions

The preoperative data collected were as previously defined for EuroSCORE (16). All-cause mortality
was considered for survival during the follow-up. Postoperative outcomes (occurred before hospital
discharge) were recorded according to the VARC-2 criteria (17). Particularly, new neurologic deficit

18 was coded according to the criteria for "stroke and TIA" (17).

#### 19

20 Surgical techniques

All the operations were performed through midline sternotomy. Cardioplegic arrest was achieved with cold blood cardioplegia. Conventional aortic valve prostheses were used in all the cases; a tissue valve was implanted in 99.7% of the patients (1865/1870). Intraoperative TOE has been used routinely in the period 2010 - 2019. All the patients underwent a pre-discharge transthoracic
 echocardiogram and a clinic follow-up during the first three months since the operation.

3

#### 4 *Statistical analysis*

5 Continuous variables were presented as mean (SD) or median (1 IQR, 3 IQR). Categorical variables 6 were presented as number (%). Univariable comparisons of preoperative, operative and postoperative 7 variables were performed among the groups using the Student's t-test, Kruskall-Wallis test, or the  $\chi^2$ 8 or Fisher's exact test as appropriate.

9 Survival probabilities, with time starting on the date of surgery, were calculated using the Kaplan-

10 Meier method. Survival probabilities of a sex and age-matched English population were calculated

11 using Kaplan-Meier method and was based on data available on:

12 <u>https://www.ons.gov.uk/peoplepopulationandcommunity/birthsdeathsandmarriages/lifeexpectancies</u>

13 <u>/datasets/singleyearlifetablesuk1980to2018</u>.

14 Log-rank test was used to compare survival curves.

A Cox regression analysis, with time starting on the date of hospital discharge, was performed to 15 determine the hazard ratios (HRs) of long-term survival; a backward stepwise model with a 16 significance of p < 0.15 was used. The variables included were gender, age, NYHA III-IV, previous 17 myocardial infarction, diabetes mellitus, hypertension, COPD, history of cerebrovascular accident, 18 19 extracardiac arteriopathy, LVEF<30%, type of haemodynamic dysfunction (aortic valve stenosis or 20 regurgitation), associated procedures, CPB time, Cross-clamp time, period when operation was performed (2000-2009 or 2010-2019), occurrence or at least one postoperative complication OR 21 single complication, pacemaker implantation. 22

A p-value of  $\leq 0.05$  was considered statistically significant.

Statistical analyses were performed using the Stat-View Statistical Software Package 5.0 (SAS
 Institute, Inc., Cary, NC, USA), NCSS 2001 (Number Cruncher Statistical System, Kaysville, Utah)
 and Stata/MP version 13 (StataCorp, College Station, Texas 77845 USA).

4

#### 5

#### 6 **Results**

7 Preoperative characteristics and operative data

8 During the period 2000 – 2019, 1870 patients aged 80 and over underwent first time surgical aortic
9 valve replacement.

10 There was a progressive increase in the number of surgical aortic valve replacement procedures with 11 octogenarian and older patients accounting for almost the 30% of the overall population undergoing 12 aortic valve surgery. Table 1 reports the preoperative characteristics of our population; Figure 1 and 13 Supplemental Table 1 detail about patients' characteristics and number of procedures for every 5-14 year interval time.

Mean patients' age was 84 (SD: 3) years and among them 53% (987/1870) were male. Logistic 15 16 EuroSCORE was 13.3% (SD: 7.2%). Forty percent of the patients presented with a NYHA class III-IV: admission with advanced heart failure symptoms was more common at the beginning of our study 17 period (2000 - 2004) with 63% of the patients presenting a marked functional limitation compared 18 with the 31% of the cases in the period 2015 - 2019 (p<0.001). Similarly, preoperative creatinine> 19 200 µmol/L was more common in the early period (7%), while its prevalence declined to 3% in 717 20 patients operated during 2010-2014 and to 1% in 600 patients operated between 2015 and 2019 21 (Supplemental Table 1). 22

Isolated aortic valve replacement was performed in 42% of the patients (790/1870), details of the
 associated procedures and operative times are reported in Table 2.

3

#### 4 *Early outcomes*

The postoperative course before hospital discharge was complicated in 9.6% of the cases. Reopening
for bleeding or tamponade was necessary in 57 patients (3%); a new neurologic deficit was reported
in 47 patients (2.5%) and among them 28 suffered a new permanent cerebral stroke (1.5%); 57
patients (3.0%) required renal replacement therapy; a permanent pacemaker was implanted in 78
cases (4.1%) before hospital discharge.

Overall in-hospital mortality was 3.2% (60/1870) at a median time of 11 days (3, 30). In-hospital
mortality after isolated aortic valve replacement was 2.2% (18/790) at a median time of 14 days (4,
33). The mortality rate decreased significantly over time from 6% (9/150) in the period 2000 – 2004
to 1.5% (9/600) in the period 2015-2019 (p=0.006). Table 3 and Supplemental Table 1.

In 54% of the cases (977/1810), patients were discharged at home while in the remaining cases a
further period of rehabilitation, nursing or medical assistance was required. The overall median
postoperative length of stay was 10 (8, 15) days (Table 3 and Supplemental table 1).

During the first 6 months there were further 83 deaths among patients who were discharged fromhospital for an overall 6-month mortality of 7.6%.

19

20 Long-term outcomes

During a median follow-up of 4.6 (3.5, 11.0) years, there were 907 deaths including in-hospital
mortality. The survival probabilities at 1-year, 5-years, 10-years and 15-years from the date of surgery
were 90% (SE: 0.7%), 66% (SE: 1.2%), 31% (SE: 1.6%) and 14% (SE: 2.0%), respectively (Figure

2). There was no significant difference in survival between the study population and an age- and sex matched general English population (log-rank test p=0.96), Figure 3.

3 The survival for patients who experienced at least one postoperative complication (reopening, new 4 neurologic deficit, renal replacement therapy, deep sternal wound infection) was 71% (SE: 3.7%), 46% (4.2%) and 18% (SE: 4.2%) at 1-year, 5-years and 10-years from the date of surgery, 5 respectively. It was significantly lower (log rank test p=0.002) when compared with the survival of 6 7 patients who had an uncomplicated postoperative course, 92% (SE: 0.7%) at 1-year, 68% (SE: 1.3%) at 5-years, 33% (SE: 1.7%) at 10-years. Similarly, patients who were operated during the period 2000-8 9 2009 showed a significantly lower survival compared with patients who underwent surgery during 10 the period 2010-2019: 10-y survival 27% (SE: 1.9%) vs 41% (SE: 2.5%), respectively (log rank test, p=0.002). 11

Supplemental Figure 1 provides a graphical view of the impact of the occurrence of postoperative complication(s) and the time of surgery on 10-year survival when compared with the expected survival of an age- and sex-matched general English population.

A Cox regression, with follow-up starting at the discharge date, was performed to study independent predictors for late mortality in patients who were successfully discharged from hospital. Alongside co-existent preoperative comorbidities (previous myocardial infarction, extracardiac arteriopathy), a complicated postoperative course (p=0.026; HR 1.32, CI 1.03 to 1.68) and surgery in the first 10 years of the study period (p=0.003; HR 1.25, CI 1.08 to 1.46) were independent risk factors for mortality at the long-term (Table 4 and Table 5).

- 21
- 22
- 23
- 24

#### 1 Discussion

Nowadays cardiac surgery can be performed safely in octogenarians and older patients. Several
studies reported favourable early and mid-term outcomes across all the spectrum of cardiac diseases
and surgical procedures including CABG, aortic valve and aortic surgery, mitral valve surgery and
emergency operation (9)(10)(11)(18)(19).

Aortic valve surgery represents one of the most common operations undertaken in people aged over 6 7 80 years as, in our experience, they accounted for more than one fourth of the global cases of surgical 8 aortic valve replacement. In these patients, we have found an in-hospital mortality rate of 3.2% over a 20-year period with a progressive and significant reduction of the mortality rate across the years 9 10 (1.5% in the last period). Our data confirmed recent findings reporting an early mortality rate of 2% in octogenarian and older patients undergoing surgical aortic valve replacement (9)(10)(11). 11 12 Alongside this evidence of procedural safety, other studies have highlighted that the mid-term survival of elderly patients with aortic valve stenosis is dramatically improved by aortic valve 13 14 replacement (68% surgery vs. 22% medical therapy at 5-years) (12) and that this longer life-15 expectancy is characterised by an acceptable quality of life and independent functional status (6).

We have further explored the long-term outcome of these patients and reported a survival rate of 31% at 10-year and 14% at 15-year follow-up. No long-term data from similar surgical populations were available for a comparison, however, we found that the 10-year survival probability of our patients was not different from the survival rate of an English age- and sex-matched population.

Preoperative status and comorbidities play a major role in determining long-term survival. In more recent years, people aged 80 and over underwent surgery in a better functional status and less frequently presented with severe chronic kidney dysfunction and history of previous myocardial infarction, nevertheless the population operated in the last period of observation had a significantly lower mean logistic EuroSCORE value. Reasons for this shift in practice lie probably in an earlier referral driven by a new awareness of the benefits of surgery also in elderly people, the confidence in

surgical results and the availability of alternative interventional procedures associated with an established activity of the Heart Valve Team in evaluating frail, high risk and elderly patients. These changes in preoperative management and the possibility of undertaking a cardiac operation before the development of advanced heart failure symptoms, translated in a lower early mortality and a better long-term outcome.

Despite a low perioperative death rate, the first postoperative months were burdened by a significant 6 7 mortality. This represents a common finding (5)(6)(18) and underlines the vital importance of preserving the already reduced functional reserves of frail and comorbid patients. Prolonged operative 8 9 times has been extensively associated with perioperative mortality and morbidity in octogenarians 10 undergoing cardiac surgery (3)(8)(20). A complicated postoperative course may also expose these elderly patients to a difficult and prolonged recovery and has been associated with a worse outcome 11 during the first year since surgery with a mortality at 1-year of 67% in patients requiring postoperative 12 dialysis and 35% in patients suffering perioperative cerebral stroke (8). In our experience, any 13 deviation from an uncomplicated postoperative course was significantly associated with a worse 14 15 survival at long-term follow-up. Particularly, new permanent cerebral injury after surgery was an independent predictor of long-term mortality, as it can affect the postoperative recovery and patients' 16 quality of life and independence. 17

We recorded a lower in-hospital mortality and a better long-term survival in patients operated during the period 2010-2019. There were no significant technical changes in the intraoperative management of these patients throughout the study period; full sternotomy was the common surgical access and conventional stented prostheses were used. Noteworthy, the last 10 years have been characterised by the establishment and the progressive growing of a transcatheter aortic valve implantation programme.

The implementation of a TAVI programme has been associated with an increase in overall aortic valve intervention and surgical aortic valve replacement activities (21)(22). Similarly to our findings, a French administrative hospital-discharge database showed that the introduction of TAVI procedures
 translated in an earlier referral with an improved patients' preoperative clinical profile (22).

3 In French and German nationwide experiences the number of TAVI procedures is now exceeding 4 the surgical activity (22)(23) and this is particularly evident in patients aged over 80 years (22). A 5 less invasive procedure can undoubtedly be associated with a lower degree of postoperative morbidity 6 especially in high-risk elderly patients (11). Transcatheter procedures have been associated with a 7 lower risk of bleeding and postoperative renal failure compared with surgical aortic valve replacement 8 (11) and, according to our findings, this could translate in a potential benefit on survival. However, 9 several observational studies have failed to demonstrate, so far, a difference in early mortality after 10 TAVI vs surgical aortic valve replacement in elderly patients (11)(22)(23). No data are available of mid- and long-term outcomes after transcatheter procedures in low-risk octogenarians and, generally, 11 the impact of prosthesis degeneration and paravalvular regurgitation has not been investigated on the 12 long-term due to the inclusion of cohort of patients with reduced life-expectancy (survival<30% at 5-13 14 years) (24).

On these bases, in our practice, advanced age alone does not represent a sufficient factor to favour a transcatheter procedure over conventional surgery. In our experience, a better preoperative functional status and the careful selection of surgical candidates, while offering transcatheter procedures to higher-risk patients, were the two most important factors that ultimately led to a reduced incidence of postoperative complications, lower early mortality, better late outcomes and the opportunity to offer an interventional treatment (conventional surgery OR transcatheter procedure) to a wider population of elderly patients.

Within the limitations of a single-centre retrospective evaluation, we have reported in a large population an historical view of long-term outcomes of patients aged 80 and older who underwent surgical aortic valve replacement. This evidence represents a reliable picture of the contemporary surgical outcomes of aortic valve surgery in elderly patients and could represent a benchmark for the
 forthcoming long-term results of non-surgical interventional treatment of aortic valve disease.

3

#### 4 Conclusions

5 Conventional surgical aortic valve replacement during the last 20 years provided a safe and successful 6 treatment for aortic valve disease in elderly patients with an in-hospital mortality that in more recent 7 years was for both isolated and combined procedures below 2%. An early surgical referral before the 8 development of advanced signs of heart failure and the availability of alternative interventional 9 procedures and a careful patients' selection, contributed to a progressive and significant reduction of 10 postoperative complications and periprocedural mortality, and led to an improved long-term survival.

- 11 **Conflict of interest:** none declared
- 12 **Funding statement:** no fund

#### 13 Authors contribution:

Substantial contributions to the conception or design of the work; or the acquisition, analysis,
 or interpretation of data for the work;

16 2. Drafting the work or revising it critically for important intellectual content;

17 3. Final approval of the version to be published;

18 4. Agreement to be accountable for his/her contributions of the work in ensuring that questions

19 related to the accuracy or integrity of the work are appropriately investigated and resolved.

20 PGM: 1-2-3-4

- 21 SL: 1-2-3-4
- 22 SG: 1-2-3-4

- 1 MK: 1-2-3-4
- 2 SKO: 1-2-3-4
- 3
- J
- 4
- 5

#### 1 Figures

- Figure 1. Trend in aortic valve replacement in patients aged 80 years and over during the period 2000
   2019.
- Figure 2. Kaplan-Meier survival curve, with starting point on the date of surgery, for the overall
  population who underwent surgical aortic valve replacement.
- Figure 3. Survival curves of study population patients aged [80 90] years and of an age- and sexmatched general English population (log-rank test p=0.96) At 10-year follow-up, observed survival
  was 33% (SE: 1.6%) and expected survival 33% (SE: 1.1%). Time from surgery was considered for
  the observed survival.
- 10 [https://www.ons.gov.uk/peoplepopulationandcommunity/birthsdeathsandmarriages/lifeexpectancie
   s/datasets/singleyearlifetablesuk1980to2018].

### 1 Tables

# **Table 1.** Preoperative characteristics

Variables	Mean (SD) or Number (%) 3
Number of patients	1870
Gender M/F	987/883 (53%/47%)
Age (y)	84 (SD: 3)
Previous myocardial infarction	151 (8%)
NYHA class III-IV	739 (40%)
Diabetes Mellitus	178 (9%)
Hypertension	1223 (65%)
Smoking history	1015 (54%)
COPD	243 (13%)
Creatinine>200 µmol/L	51 (3%)
Cerebral stroke	187 (10%)
Extracardiac arteriopathy	125 (7%)
Aortic valve stenosis	1764 (94%)
LVEF<30%	94 (5%)
Acute infective endocarditis	12 (1%)
Emergency/salvage operation	46 (2%)
Logistic EuroSCORE %	13.3 (SD: 7.2)

# **Table 2.** Operative data

Variables	Mean (SD) or Number (%)
Isolated Aortic Valve replacement	790 (42%)
Associated CABG	956 (51%)
• Mean number of grafts	• 1.96 (SD: 0.93)
Associated Mitral Valve surgery	94 (5%)
• Mitral Valve repair	• 52 (3%)
Mitral Valve replacement	• 42 (2%)
Associated Tricuspid Valve repair	28 (2%)
Associated aortic surgery	69 (4%)
CPB times (minutes)	101 (SD: 40)
Crossclamp times (minutes)	73 (SD: 28)

# **Table 3.** Postoperative data

Variables	Median [IQR] or Number (%)
Complications	181/1870 (9.6%)
Re-opening for bleeding or tamponade	57 (3.0%)
Continuous veno-venous haemodialysis	57 (3.0%)
Neurologic deficit	47 (2.5%)
• Permanent cerebral stroke	• 28 (1.5%)
• Transient ischaemic attack	• 19 (1.0%)
Deep Sternal Wound Infection	9 (0.5%)
Permanent pacemaker implantation	123 (6.5%)
Pre discharge	• 78 (4.1%)
Post discharge	• 45 (2.4%)
Length of stay (days)	10 (8, 15)
Discharge destination	
Home/Rehabilitation	977 (54%) / 833 (46%)
In-Hospital Mortality	
Overall	3.2% (60/1870)
Isolated Aortic Valve replacement	2.2% (18/790)
Follow-up (years)	4.6 (3.5, 11.0)

- **Table 4.** Predictors of late mortality among survivors by Multivariable Cox analysis with time starting
- 2 on discharge date and backward stepwise variables selection [Schoenfeld test p=0.38].

Variables	p; HR (95% lower, 95% upper)
Gender male	0.001; 1.26 (1.09, 1.43)
Age (y)	<0.001; 1.07 (1.05, 1.10)
NYHA III-IV	0.14; 1.11 (0.97, 1.28)
Previous MI	0.001; 1.43 (1.15, 1.78)
Extracardiac arteriopathy	<0.001; 1.68 (1.31, 2.14)
CPB time (minutes)	0.008; 1.006 (1.001, 1.010)
Crossclamp time (minutes)	0.008; 0.992 (0.986, 0.998)
Operation date	
(2010 – 2019 reference)	
2000 - 2009	0.003; 1.26 (1.08, 1.46)
Postoperative complication(s)	0.026; 1.32 (1.03, 1.68)

- **Table 5.** Predictors of late mortality among survivors by Multivariable Cox analysis with time starting
- 2 on discharge date and backward stepwise variables selection including each complication
- 3 individually [Schoenfeld test p=0.57].

Variables	p; HR (95% lower, 95% upper)
Age (y)	<0.001; 1.08 (1.05, 1.10)
Gender male	0.001; 1.27 (1.10, 1.47)
NYHA III-IV	0.12; 1.19 (0.97, 1.29)
Previous MI	<0.001; 1.45 (1.16, 1.80)
Extracardiac arteriopathy	<0.001; 1.69 (1.33, 2.16)
LVEF<30%	0.12; 1.25 (0.94, 1.66)
CPB time (minutes)	0.016; 1.005 (1.001 – 1.010)
Crossclamp time (minutes)	0.011; 0.992 (0.986, 0.998)
Operation date	
(2010 – 2019 reference)	
2000 - 2009	0.037; 1.25 (1.08, 1.45)
Postoperative cerebral stroke	<0.001; 2.75 (1.69, 4.46)
Renal replacement therapy	0.075; 1.65 (0.95, 2.87)

1 References	
--------------	--

2	1.	Bridges CR, Edwards FH, Peterson ED, Coombs LP, Ferguson TB. Cardiac surgery in
3		nonagenarians and centenarians. J Am Coll Surg. 2003;197:347-56
4	2.	Akins CW, Daggett WM, Vlahakes GJ, Hilgenberg AD, Torchiana DF, Madsen JC, et al.
5		Cardiac operations in patients 80 years old and older. Ann Thorac Surg. 1997;64:606-14
6	3.	Kolh P, Kerzmann A, Lahaye L, Gerard P, Limet R. Cardiac surgery in octogenarians. Peri-
7		operative outcome and long-term results. Eur Heart J. 2001;22:1235-1243
8	4.	Elayda MA, Hall RJ, Reul RM, Alonzo DM, Gillette N, Reul GJ Jr, et al. Aortic valve
9		replacement in patients 80 years and older. Operative risks and long-term results. Circulation.
10		1993;88:II11-6
11	5.	Krane M, Voss B, Hiebinger A, Deutsch MA, Wottke M, Hapfelmeier A, et al. Twenty years
12		of cardiac surgery in patients aged 80 years and older: risks and benefits. Ann Thorac Surg.
13		2011;91:506-13
14	6.	Chaturvedi RK, Blaise M, Verdon J, Iqbal S, Ergina P, Cecere R, et al. Cardiac surgery in
15		octogenarians: long-term survival, functional status, living arrangements, and leisure activities.
16		Ann Thorac Surg. 2010;89:805-10
17	7.	Hirji SA, Ramirez-Del Val F, Kolkailah AA, Ejiofor JI, McGurk S, Chowdhury R, et al.
18		Outcomes of surgical and transcatheter aortic valve replacement in the octogenarians-surgery
19		still the gold standard? Ann Cardiothorac Surg. 2017;6:453-462
20	8.	Atladottir HO, Modrau IS, Jakobsen CJ, Torp-Pedersen CT, Gissel MS, Nielsen DV. Impact of
21		perioperative course during cardiac surgery on outcomes in patients 80 years and older. J
22		Thorac Cardiovasc Surg. 2020 Mar 19:S0022-5223(20)30570-5. doi:
23		10.1016/j.jtcvs.2020.03.005
24	9.	El-Essawi A, Follis M, Brouwer R, Breitenbach I, Groeger S, Anssar M, et al. Is aortic valve
25		replacement with a minimally invasive extracorporeal circuit a contemporary option for
26		octogenarians? Interactive Cardiovasc Thorac Surg. 2020;31:56-62

1	10.	Timek TA, Turfe Z, Hooker RL, Davis AT, Willekes CL, Murphy ET, et al. Aortic valve
2		replacement in octogenarians with prior cardiac surgery. Ann Thorac Surg. 2015;99:518-23
3	11.	Sheng SP, Strassle PD, Arora S, Kolte D, Ramm CJ, Sitammagari K, et al. In-Hospital
4		Outcomes After Transcatheter Versus Surgical Aortic Valve Replacement in Octogenarians. J
5		Am Heart Assoc. 2019 22;8:e011206. doi: 10.1161/JAHA.118.011206
6	12.	Varadarajan P, Kapoor N, Bansal RC, Pai RG. Survival in elderly patients with severe aortic
7		stenosis is dramatically improved by aortic valve replacement: results from a cohort of 277
8		patients aged ≥80 years. Eur J Cardiothorac Surg. 2006;30:722-727
9	13.	Makkar RR, Thourani VH, Mack MJ, Kodali SK, Kapadia S, Webb JG, et al; PARTNER 2
10		Investigators. Five-Year Outcomes of Transcatheter or Surgical Aortic-Valve Replacement. N
11		Engl J Med. 2020;382:799-809
12	14.	Barbanti M, Costa G, Zappulla P, Todaro D, Picci A, Rapisarda G, et al. Incidence of Long-
13		Term Structural Valve Dysfunction and Bioprosthetic Valve Failure After Transcatheter Aortic
14		Valve Replacement. J Am Heart Assoc. 2018;7(15):e008440. doi: 10.1161/JAHA.117.008440
15	15.	Søndergaard L, Ihlemann N, Capodanno D, Jørgensen TH, Nissen H, Kjeldsen BJ, et al.
16		Durability of Transcatheter and Surgical Bioprosthetic Aortic Valves in Patients at Lower
17		Surgical Risk. J Am Coll Cardiol. 2019 Feb 12;73(5):546-553. doi: 10.1016/j.jacc.2018.10.083
18	16.	Nashef SA, Roques F, Sharples LD, Nilsson J, Smith C, Goldstone AR, et al. EuroSCORE II.
19		Eur J Cardiothorac Surg. 2012;41:734-44
20	17.	Kappetein AP, Head SJ, Généreux P, Piazza N, van Mieghem NM, Blackstone EH, et al.
21		Updated standardized endpoint definitions for transcatheter aortic valve implantation: the Valve
22		Academic Research Consortium-2 consensus document. J Am Coll Cardiol. 2012;60:1438-54
23	18.	Chivasso P, Bruno VD, Farid S, Malvindi PG, Modi A, Benedetto U, et al. Predictors of survival
24		in octogenarians after mitral valve surgery for degenerative disease: The Mitral Surgery in
25		Octogenarians study. J Thorac Cardiovasc Surg. 2018;155:1474-1482

1	19.	Igarashi T, Sato Y, Satokawa H, Takase S, Wakamatsu H, Seto Y, et al. The results of an entry-
2		oriented strategy for acute type A aortic dissection in octogenarians: an 18-year experience. Eur
3		J Cardiothorac Surg. 2020 Jul 23:ezaa195. doi: 10.1093/ejcts/ezaa195
4	20.	Rohde SL, Baker RA, Tully PJ, Graham S, Cullen H, Knight JL. Preoperative and intraoperative
5		factors associated with long-term survival in octogenarian cardiac surgery patients. Ann Thorac
6		Surg. 2010;89:105-11
7	21.	Grant SW, Devbhandari MP, Grayson AD, Dimarakis I, Kadir I, Saravanan DM, et al. What is
8		the impact of providing a transcatheter aortic valve implantation service on conventional aortic
9		valve surgical activity: patient risk factors and outcomes in the first 2 years. Heart.
10		2010;96:1633-7
11	22.	Nguyen V, Michel M, Eltchaninoff H, Gilard M, Dindorf C, Iung B, et al. Implementation of
12		Transcatheter Aortic Valve Replacement in France. J Am Coll Cardiol. 2018;71:1614-27
13	23.	Eggebrecht H, Mehta R. Transcatheter aortic valve implantation (TAVI) in Germany: more
14		than 100,000 procedures and now the standard of care for elderly. EuroIntervention.
15		2019;14:1549-52
16	24.	Barbanti M, Costa G, Zappulla P, Todaro D, Picci A, Rapisarda G, et al. Incidence of Long-
17		Term Structural Valve Dysfunction and Bioprosthetic Valve Failure After Transcatheter Aortic
18		Valve Replacement. J Am Heart Assoc. 2018 Aug 7;7(15):e008440.doi:
19		10.1161/JAHA.117.008440







Supplementary Figure 1

Click here to access/download Supplementary material Figure 4.2.pdf Supplementary material

Click here to access/download Supplementary material Supplementary materials revised 2.docx