

The role of dietary polyphenols in the control of chronic noncommunicable diseases

Yasmany Armas Díaz¹  | Maria Soledad Ferreiro Cotorruelo² | Maurizio Battino^{1,3,4} 

¹Faculty of Medicine, Department of Clinical Sciences, Polytechnic University of Marche, Ancona, Italy

²Direzione Medica Ospedaliera, Azienda Ospedaliera-Universitaria delle Marche, Ancona, Italy

³International Joint Research Laboratory of Intelligent Agriculture and Agri-products Processing, Jiangsu University, Zhenjiang, China

⁴Research Group on Food, Nutritional Biochemistry and Health, Universidad Europea del Atlantico, Santander, Spain

Correspondence

Maurizio Battino, Facoltà di Medicina,
Dipartimento di Scienze Cliniche
Specialistiche, Università Politecnica delle
Marche, Ancona 60131, Italy.
Email: m.a.battino@univpm.it

Abstract

Noncommunicable chronic diseases (NCDs) are among the leading causes of death and disability worldwide. The most common NCDs are cancer, obesity, cardiovascular diseases, and diabetes. Nowadays, they represent one of the greatest challenges health systems face worldwide. The increase in the consumption of polyphenol-rich foods could contribute to the reduction of these pathologies, due to their antioxidant, anti-inflammatory, anticancer, immunomodulatory, and cardiovascular protective properties, among others. This review aims to highlight some studies carried out in recent years to enhance the possible benefits of a diet rich in polyphenols in the prevention or treatment of NCDs.

KEYWORDS

anti-cancer, anti-diabetic, anti-obesity, cardioprotective, chronic noncommunicable diseases, polyphenols

1 | INTRODUCTION

Polyphenols are natural compounds that can be found in fruits, vegetables, cereals, and beverages. The production of these in plants is mainly due to exposure to different types of stress or infections, so they can also be considered secondary metabolites (Abbas et al., 2017).

Polyphenolic compounds are grouped into four main groups, based on their chemical structure, including flavonoids, stilbenes, lignans, and phenolic acids (Figure 1). They are found in abundance in foods such as red grapes, berries, coffee, tea, cereal grains, citrus

fruits, soybeans, legumes, and apples, among others. The most studied polyphenols are anthocyanins, ellagitannins, quercetin, resveratrol, and curcumin (Tsao, 2010).

A number of studies at cellular, preclinical, and clinical level have demonstrated the possible biological activity attributed to polyphenols. Anti-inflammatory, immunomodulatory, antioxidant, cardiovascular protective, anticancer, and antimicrobial are some of the properties exerted by these compounds (Figure 1). A short revision on PubMed was conducted to highlight, once again, the positive role of polyphenols in the prevention or treatment of Noncommunicable chronic diseases (NCDs).

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2023 The Authors. *Food Safety and Health* published by John Wiley & Sons Australia, Ltd on behalf of International Association of Dietetic Nutrition and Safety.




FIGURE 1 Polyphenols classification and health beneficial effects.

2 | CARDIOPROTECTIVE EFFECT

Dietary choice plays a central role in the development of cardiovascular diseases (CVD), and their potential mechanisms of action are diverse. The positive effect of polyphenols on CVD is mainly related with their possibility of improving lipid metabolism, decreasing inflammatory signaling, and helping in the regeneration of vascular endothelium. In addition, they reduce the oxidative stress and have a positive effect on survival signaling, calcium homeostasis, and sirtuin (SIRT) activity (Kiyimba et al., 2023; Torres-Fuentes et al., 2022).

Plenty laboratories have focused their effort to elucidate the role of polyphenol extracts or isolated compounds on CVD. Regarding this, the mean systolic and diastolic blood pressure was significantly lower in patients with a diet rich in polyphenols in a study carried out in Italy (Vitale et al., 2018). Ferguson et al. (2022) assayed the effects of a polyphenol-rich dietary supplement containing *Pinus massoniana* bark extract (PMBE) on systolic (SBP) and diastolic blood pressure in healthy Australian adults. SBP reduced significantly in individuals with moderately high SBP at the baseline following 12 weeks PMBE supplementation. The lowering of blood pressure was associated mainly to the fact that polyphenols can decrease the expression of nuclear factor- κ B (NF- κ B), improving enzymatic activities of catalase, glutathione peroxidase, and superoxide dismutase and activating the redox-sensitive phosphoinositide 3-kinase/Akt pathway (Davinelli & Scapagnini, 2016).

On the other hand, to get deeper on how polyphenols can reduce the incidence of CVD, some studies have been conducted using isolated compounds. For instance, after several studies, it is proposed that flavonoids can specifically act as signaling molecules. This effect seems to be related with the fact that they can interact with some specific cellular receptors or proteins that are involved in regulatory pathways, resulting in physiological responses or regulation of gene expression. These compounds have also been related to the regulation of nitric oxide (NO) levels in the vascular endothelium through the inhibition of superoxide production. These aspects demonstrate

their positive action on endothelial function and the regulation of vasodilator processes associated with blood flow (Galiniak et al., 2019; Godos et al., 2019; Zhao et al., 2023).

Furthermore, other clinical and laboratory investigations have confirmed the possible positive benefits of some flavonoids in the regulation of blood pressure and endothelial function (Qian et al., 2017). Administration of cocoa flavanol significantly reduced CVD death by 27% in US healthy adults (Sesso et al., 2022).

Evidence from meta-analyses suggests that anthocyanin-rich foods like berries can improve endothelial function, blood pressure, and arterial stiffness, decrease fasting and postprandial glucose levels and reduce LDL cholesterol (Fairlie-Jones et al., 2017; Giampieri et al., 2023; LiPing et al., 2017; Wood et al., 2019).

Atherosclerotic plaque formation is the key step that underlies cardiometabolic diseases. Several studies have showed that polyphenols treatment reduces atherosclerotic plaque formation (Clark et al., 2022). For instance, gallic acid (GA), a phenolic acid abundant in blackberries, reduced atherosclerosis in male but not in female ApoE^{-/-} mice. Specifically, GA treatment in male mice restored microbiome dysbiosis and reduced serum IL3 and IL12 levels and spleen weight (Clark et al., 2022).

Another effect showed by polyphenols involved the angiotensin II (Ang II) signaling. Activation of Ang II is linked to the onset and progression of CVDs. Polyphenols extract from berries attenuated Ang II-induced senescence through a Nox1-dependent mechanism in vascular smooth muscle cells (Feresin et al., 2016).

Other class of polyphenols that have been extensively studied are stilbenes, in particular, resveratrol. The last one, present in red wine, grapes, and nuts, has also been reported to possess cardioprotective activity, mainly attributed to its anti-inflammatory properties. In vitro and in vivo studies have shown that it can inhibit the enzyme cyclooxygenase, inactivate the gamma receptor, and induce endothelial NO synthase production (Galiniak et al., 2019). In addition, it can also inhibit the NF- κ B pathway counteracting the overexpression of the vascular and intracellular

adhesion molecules (VCAM1 and ICAM1) in endothelial cells (Deng et al., 2011).

Taking in account the information above, it is notable that the development of CVD is influenced by dietary behavior. Lifestyle is an important risk factor for the development of CVD. Due to the fact that treatment available for CVD can cause side effects, the development of safer therapeutics is necessary, and the intake food rich in polyphenols can be one of them. In summary, foods rich in polyphenols and their isolated bioactive compounds may be effective in reducing cardiometabolic risks and improving lifespan. Important to notice is that effects on clinical cardiovascular events are untested, so more research is needed to understand the mechanism and specific plants' phytochemicals responsible for treating CVD.

3 | ANTICANCER ACTIVITY

The anticancer effect of polyphenols is largely attributed to the possible dynamic regulation of cell death programming pathways, such as apoptosis and autophagy. They also inhibit cell proliferation, interrupting the cell cycle. The anti-inflammatory properties associated with those also favor the inhibition of cancer progression or delay its appearance (Patra et al., 2021; Sharma et al., 2018). Studies have shown the direct participation of polyphenols in the control of the expression of oncogenes and the promotion of tumor suppressor genes for the subsequent inhibition of cancer progression (Patra et al., 2021).

It is well-known that polyphenols may be able to modulate cytokine and chemokine production and immune cell activation. These properties justify their positive action in the chemoprevention, the cytotoxic action against cancer cells, and the induction of greater sensitization to chemoradiotherapies (Mileo et al., 2019). The positive effect of polyphenols has been shown in a plenty number of cancer cells such as colorectal, lung, breast, and ovarian, among others.

The cultivation of polyphenols extracted from Manuka honey on human colon adenocarcinoma (HCT-116) and colon metastasis (LoVo) cell lines demonstrated strong inhibition on cell growth by inducing apoptosis through the upregulation of p53, cleaved-PARP and caspase-3, and the activation of caspase-9 and caspase-8. Also, the treatment caused the arrest in the cell cycle and the reduction of proliferation in a dose-dependent manner (Afrin et al., 2018).

Targeting the metabolic pathways of cancer stem cells (CSCs) could be a possible treatment to increase the survival rate of patients affected by cancer. The application of an extract of Manuka honey polyphenols reduced the entire culture of spheroids deriving from the HCT-116 cell line. The treatment also affected the morphological parameters and induced apoptosis and intracellular accumulation of ROS. In addition, the self-renewal capacity of these cells was affected (Cianciosi et al., 2020).

The application of a strawberry extract rich in polyphenols reduced the survival of the aggressive breast cancer cell (A17), affecting their migration, adhesion, invasion, and the cell cycle. This positive activity was also shown in the female FVB/N mice assay (Amatori et al., 2016). On the other hand, adoption of a

Mediterranean diet by a population of Italian women tended to decrease by 30%–50% the incidence of breast cancer (Villarini et al., 2016). Giampieri et al. (2019) and Greco et al. (2023) suggested that strawberry polyphenols could be used in the development of therapeutic and/or preventive supplements for uterine leiomyomas.

Gynecological cancers development has been shown affected by polyphenols treatments. Regarding this topic, El-Kott et al. (2020) demonstrated that kaempferol promote cell apoptosis in A2780 ovarian cancer cells by triggering ER stress-mediated cytotoxic autophagy. The treatment also caused inhibition of the PI3K/Akt signaling pathway, which seems to be the way kaempferol enhances the sensitivity of ovarian cancer cells to chemotherapy.

Kaempferol also showed chemotherapeutic efficacy against nonsmall cell lung cancer (A549 and NCIH460 cell lines). This flavonoid hampered the Nrf2 signaling pathway which induced intracellular accumulation of ROS. Also, it inhibited the proliferation and affected the cell cycle. Taken together, kaempferol induced apoptosis and inhibited the proliferation of these cancer cell lines (Fouzder et al., 2021).

Curcumin also has well-known exerting healing properties and antitumor activity. Its action in controlling cancer development is mainly due to its activities by modulating various cellular processes, including apoptosis, angiogenesis, autophagy, metastasis, and epithelial-mesenchymal transition. Recently, its therapeutic effects have been attributed to alterations of noncoding RNAs, such as circ-HN1, circ-PRKCA, and circPLEKHM3 (Si et al., 2023; Sun & Fang, 2021; Wu et al., 2022; Xu et al., 2021).

As happened with others NCDs, polyphenols extracted from tea have been studied extensively in cancer treatments. For instance, in Chandrani Fouzder squamous cell carcinoma (KYSE 150), epigallocatechin gallate (EGCG) treatment activated methylation silencing genes and inhibited DNA methyltransferase activity (Ding et al., 2020).

The development of cancer is a microevolutionary and multistage procedure. One of the ways to decrease its incidence or improve life quality of people affected could be the intake of food rich in bioactive compounds, specifically polyphenols. Polyphenols provide a potentially beneficial health-promoting effect by decreasing the growth and development of cancer cells and chemoresistance. Some of these have been well studied, but more research studies have to be conducted to understand the precise molecular basis of the disease and to develop more efficient therapeutic options. Even though this paper shows few results about in vivo studies, several of these have been conducted in animal and human models, but still the information is not enough to be sure about the incidence of polyphenols in the human body.

4 | ANTI-OBESITY EFFECT

The proper lipid metabolism pathway is essential for the prevention of several chronic metabolic diseases such as obesity. Epidemiological evidence suggests the positive effect of polyphenols found in the diet, on overweight and obesity development (Chen et al., 2022).

The AMPK signaling pathway has been widely studied to target lipid metabolism. Regarding this, Wang et al. (2014) reported that polyphenols extracted from tea, curcumin, and resveratrol activated AMPK; therefore, accumulation of fat in adipocytes was diminished. In addition, they noticed the downregulation of some adipogenic genes, such as stearoyl Co-A desaturase-1 and fatty acid synthase. Other studies showed that the application of a strawberry extract in liver cancer cells decreased levels of total cholesterol, LDL, and triglycerides and stimulated the p-AMPK/AMPK expression (Forbes-Hernández et al., 2017). Recently, treatment of a murine 3T3-L1 fibroblast cell line with a mixture of polyphenols and micro-nutrients reduced lipid accumulation, and this effect was mediated via activation of the AMPK signaling pathway (Pacifci et al., 2023).

Epidemiological studies have shown that tea polyphenols could induce the reduction of waist circumference and body mass index (BMI) (Hiroki T et al., 2017). Similar results were reported by Grosso et al. (2016) where BMI, waist circumference, blood pressure, and triglycerides were significantly lower among people with a high intake of polyphenol-rich foods (Grosso et al., 2016). Polyphenols extracted from berry fruits showed a positive effect on the process related with the dissipation of energy through the production of heat by improving mitochondrial function and the biogenesis of brown adipose tissue (Jiang et al., 2021). Batista-Jorge et al. (2020) conducted a study where resveratrol was orally administrated to obese patients. After 12 weeks of treatment, HDL levels increased, triglycerides decreased 36%, and total cholesterol and VLDL improved.

A double-blind, placebo-controlled, randomized trial with 35 adults with obesity was carried out by Jamar et al. (2020). After 6 weeks of júcara pulp administration, body fat was reduced and HDL increased. Romain et al. (2021) got similar results when they administrated a pull of polyphenols extracted from a Mediterranean diet to overweight and obese but otherwise healthy individuals. Body fat mass decreased and health-related quality of life improved significantly. People with intellectual disabilities seem to be more susceptible to obesity issues. Regarding this matter, Xicota et al. (2020) studied the effect of encapsulated green tea extract, mainly EGCG, on body weight in young Down syndrome. The main results were lower body weight, less increase in BMI, and decrease in plasma lipid concentrations in the treated group compared with placebo.

Finally, in postmenopausal women with normal weight, it was found that the risk of being overweight and obese decreased 35% and 88%, respectively, for women with higher intakes of wine, relative to nondrinkers (Thomson et al., 2012).

To resume, the incidence of overweight and obesity is climbing worldwide at exponential levels which has caused negative effects on human health systems and economic development. Due to these issues, many efforts have been put to elucidate the main metabolic pathways that can regulate the progress of this pathology. In this sense, researchers have looked into the potentials of natural products such as fresh vegetables and fruits for resolving obesity due to their high efficiencies and few undesirable outcomes. Important to notice is that nutrition by itself is not enough to burn out this problem, but it is one of the most important solutions.

5 | ANTIDIABETIC EFFECT

Lifestyle and dietary habit changes have cost a high prevalence of diabetes in recent decades. Fruits and vegetables are a good source of bioactive compounds highly recommended in the medical nutrition therapy to prevent or reduce the incidence of type 2 diabetes mellitus (T2DM). During last years, many groups have focused their work on the role of polyphenols in the development of this disease.

As it is well-known, mTORC1 hyperactivation is a key event in the progression of T2DM (García-Aguilar & Guillén, 2022). Several polyphenols, such as resveratrol, curcumin, and EGCG, have been related with the downregulation of mTORC1 through AMPK-dependent and AMPK-independent mechanisms being essential in the survival of pancreatic β cells (García-Aguilar et al., 2021). Furthermore, the administration of curcumin to male rats fed with high-fat diet and streptozotocin for 12 weeks prevented the liver damage induced by hyperglycemia and hyperlipidemia, mainly due to their antiapoptotic and antioxidant activity (Xia et al., 2020).

Also recently, García-Díez et al. (2022) found that epicatechin (EC) was able to decrease lipid accumulation and glucose absorption through AKT and AMPK pathway activation in cardiac H9c2 cells. In addition, the coinubation increased insulin receptor levels and expression of the glucose transporter GLUT-4.

Human gut microbiota are also suggested to be involved in T2DM development, and some authors have found positive effect on the interaction between polyphenols and microbiota. In this sense, it is proposed that phenolic compounds could affect the release of intestinal hormones and the composition of the intestinal microbiota, which, in turn, are directly or indirectly involved in the regulation of metabolic homeostasis (Wang et al., 2021). On the other hand, the digestion of carbohydrates in the small intestine can be reduced by some polyphenols through the lowering of glucose absorption by inhibition of GLUTs, the activity of related enzymes, and enhancing the uptake of glucose by promoting the translocation of GLUTs to the plasma membrane in tissue cells (Chen et al., 2022).

A wide range of human intervention trials and preclinical studies have also shown the possible antidiabetic properties of polyphenols, but evidence is still limited and inconsistent (Menezes et al., 2022). For instance, prospective analysis in the context of the PREDIMED-Plus trial conducted by Menezes et al. (2022) showed that a diet with higher polyphenols lowered the levels of glucose and HbA1c after 1 year in a senior population with T2DM or are at high risk of developing T2DM. In another study carried out by Grabež et al. (2022), they observed that polyphenols extracted from pomegranate peel decreased oxidative stress biomarkers such as O_2^{-2} , NO^{-2} , TBARS, inflammatory factors (IL-6, TNF- α , hsCRP), and homocysteine in overweight patients with T2DM. Besides, this treatment also increased total antioxidant capacity.

Few years ago, Shoji et al. (2017) also observed that the intake of polyphenols extracted from apples, by the Japanese population between 30 and 60 years, significantly contributed to glucose tolerance by the body. Furthermore, a study conducted in a Polish population between 45 and 69 years old found that a higher intake of a variety

of polyphenols could lower the risk of developing T2DM (Grosso et al., 2017). The main reason attributed to this behavior was the action of flavonoids, which are thought to regulate the biological pathways related to the development of T2DM, by improving endothelial function (Munir et al., 2013).

Even though a lot of effort has to be put to clarify the positive effect of polyphenols in the prevention and treatment of DMT2, it

can be concluded that dietary polyphenols might be a good choice to maintain and protect pancreatic beta cells, prolonging its lifespan and avoiding the occurrence of diabetes.

Several years ago, the positive effect of polyphenols in the development and treatment of NCDs was associated mainly to their antioxidant and radical scavenger activity. However, nowadays, it is well established that they also eject their action through the up/

TABLE 1 Resume of different ways by which polyphenol-rich extract or isolated compounds exert their positive effect on prevention or treatment of NCDs.

Cardioprotective	Polyphenol-rich extract	Lowering blood pressure Improvement of endothelial function Improvement of the lipid profile Reduction of oxidative stress and inflammation Antithrombotic effect	Szczepańska et al. (2022), Haş et al. (2023)
Lipid metabolism regulation	Polyphenols	Improvements in vascular function through antioxidative and anti-inflammatory properties	Yaskolka Meir and Tsaban (2022), Romain et al. (2021)
		Reduction of lipid peroxidation and adipose tissue lipolysis	Chen et al. (2022)
		Increasing muscle mitochondrial respiration, energy consumption, and fat oxidation	Jiang et al. (2021)
	Stillbenes and flavonoids	Increase of α -diversity of gut microbiome which may mediate the reduction of BMI	Mompeo et al. (2020), Le Roy et al. (2020)
Antidiabetic	Anthocyanin	Improve postprandial insulin and possible glucose response	Solverson et al. (2019)
	Flavonoids	Slows down the absorption of glucose in the intestine	Kerimi et al. (2019)
	Polyphenol-rich extract	Reduction of intracellular ROS and increased survival of pancreatic β -cells	Xu et al. (2018)
		Reduction of β -glucosidase, β -glucuronidase, and β -galactosidase activity	Berendika et al. (2022)
		Decreased α -glucosidase activity in the small intestine	Ramos et al. (2022)
	Cyanidin	Reduction of the formation of AGEs by increasing Glo-1 activity	Suantawee et al. (2020)
Anticancer activity	Resveratrol and curcumin	Regulating of apoptosis genes, such as CASP3, CASP7, and FAS	Chimento et al. (2023)
	Naringenin	Inhibition of cancer progression via apoptosis induction, cell cycle arrest, angiogenesis hindrance, and modification of various signaling pathways including Wnt/ β -catenin, NF- κ B, and TGF- β pathways	Motallebi et al. (2022)
	Polyphenol-rich extract	Increase in ROS production and apoptosis induction	Trisha et al. (2022)
		Suppressed cell proliferation to increase the sensitivity of chemotherapy drugs	Maleki Dana et al. (2022)
		Activation of tumor suppressor genes, inhibition of angiogenesis, PI3K/AKT pathways, STAT3, transcription factor AP-1, and Nrf2 and also suppressed Wnt/ β -catenin signaling	Maugeri et al. (2023)
		Prevention/inhibition of the angiogenesis process	Kim et al. (2023)
	Curcumin	Alterations of noncoding RNAs	Si et al. (2023)
	Kaempferol	Restriction of the inflammation via suppressing the pro-inflammatory cytokines as well as chemokines production	Almatroudi et al. (2023)

downregulation of other cell-signaling pathways that have no connection to ROS production. Overall, the published data available about the effect of dietary polyphenols are not enough and in some cases contradictory. Recent metadata analyses showed that there are still experimental strategies that should be improved. For instance, one of the most important issues to overcome is the metabolic transformation and bioavailability of the polyphenols. During the digestion and absorption, they can suffer some modifications; therefore, the specific metabolites formed in vivo may be different when they reach the specific target tissues.

Despite the need of the improvement of experiments to target the incidence of polyphenols in the development of NCDs, the intake of polyphenols seems to be helpful in their prevention or treatment. Table 1 shows a resume of the positive effect that can be attributed to the intake of polyphenols.

6 | CONCLUSION

In summary, a diet rich in polyphenols is associated with the maintenance of good well-being and with the prevention of many NCDs. Polyphenols stand out due to their possible antioxidant, anticancer, antidiabetic, anti-inflammatory, etc. properties. Although there are numerous studies that demonstrate their benefits, it is necessary to continue with research studies because the nutritional amount for human health is not clear since exposure in the normal diet occurs at lower concentrations than that in the laboratory; their interaction with the intestinal microbiota can affect functionality, and the clinical studies carried out cannot always be generalized to the entire population.

AUTHOR CONTRIBUTIONS

Yasmany Armas Díaz: Conceptualization; Writing – original draft.
Maria Soledad Ferreiro Cotorruelo: Validation; Visualization.
Maurizio Battino: Funding acquisition; Project administration; Visualization

ACKNOWLEDGMENTS

The work received no funding.

CONFLICT OF INTEREST STATEMENT

The authors have no conflict of interest.

DATA AVAILABILITY STATEMENT

No datasets were generated or analyzed during the current study.

ORCID

Yasmany Armas Díaz  <https://orcid.org/0000-0003-2772-2225>

Maurizio Battino  <https://orcid.org/0000-0002-7250-1782>

REFERENCES

Abbas, M., Saeed, F., Anjum, F. M., Afzaal, M., Tufail, T., Bashir, M. S., Ishtiaq, A., Hussain, S., & Suleria, H. A. R. (2017). Natural polyphenols: An

overview. *International Journal of Food Properties*, 20(8), 1689–1699. <https://doi.org/10.1080/10942912.2016.1220393>

Afrin, S., Giampieri, F., Gasparrini, M., Forbes-Hernández, T. Y., Cinciosi, D., Reboredo-Rodríguez, P., Amici, A., Quiles, J. L., & Battino, M. (2018). The inhibitory effect of Manuka honey on human colon cancer HCT-116 and LoVo cell growth. Part 1: The suppression of cell proliferation, promotion of apoptosis and arrest of the cell cycle. *Food and Function*, 9(4), 2145–2157. <https://doi.org/10.1039/c8fo00164b>

Almatroudi, A., Allemailem, K. S., Alwanian, W. M., Alharbi, B. F., Alrumaihi, F., Khan, A. A., Almatroodi, S. A., & Rahmani, A. H. (2023). Effects and mechanisms of kaempferol in the management of cancers through modulation of inflammation and signal transduction pathways. *International Journal of Molecular Sciences*, 24(10), 8630. <https://doi.org/10.3390/ijms24108630>

Amatori, S., Mazzoni, L., Alvarez-Suarez, J. M., Giampieri, F., Gasparrini, M., Forbes-Hernandez, T. Y., Afrin, S., Errico Provenzano, A., Persico, G., Mezzetti, B., Amici, A., Fanelli, M., & Battino, M. (2016). Polyphenol-rich strawberry extract (PRSE) shows in vitro and in vivo biological activity against invasive breast cancer cells. *Scientific Reports*, 6(1), 30917. <https://doi.org/10.1038/srep30917>

Batista-Jorge, G. C., Barcala-Jorge, A. S., Silveira, M. F., Lelis, D. F., Andrade, J. M. O., de Paula, A. M. B., Guimarães, A. L. S., & Santos, S. H. S. (2020). Oral resveratrol supplementation improves Metabolic Syndrome features in obese patients submitted to a lifestyle-changing program. *Life Sciences*, 256, 117962. <https://doi.org/10.1016/j.lfs.2020.117962>

Berendika, M., Domjanić Drozdek, S., Odeh, D., Oršolić, N., Dragičević, P., Sokolović, M., Garofulić, I. E., Đikić, D., & Jurčević, I. L. (2022). Beneficial effects of Laurel (*Laurus nobilis* L.) and Myrtle (*Myrtus communis* L.) extract on rat health. *Molecules*, 27(2), 581. <https://doi.org/10.3390/molecules27020581>

Chen, L., Pu, Y., Xu, Y., He, X., Cao, J., Ma, Y., & Jiang, W. (2022). Anti-diabetic and anti-obesity: Efficacy evaluation and exploitation of polyphenols in fruits and vegetables. *Food Research International*, 157, 111202. <https://doi.org/10.1016/j.foodres.2022.111202>

Chimento, A., D'Amico, M., De Luca, A., Conforti, F. L., Pezzi, V., & De Amicis, F. (2023). Resveratrol, epigallocatechin gallate and curcumin for cancer therapy: Challenges from their pro-apoptotic properties. *The Life*, 13(2), 261. <https://doi.org/10.3390/life13020261>

Cienciosi, D., Forbes-Hernández, T. Y., Ansary, J., Gil, E., Amici, A., Bompadre, S., Simal-Gandara, J., Giampieri, F., & Battino, M. (2020). Phenolic compounds from Mediterranean foods as nutraceutical tools for the prevention of cancer: The effect of honey polyphenols on colorectal cancer stem-like cells from spheroids. *Food Chemistry*, 325, 126881. <https://doi.org/10.1016/j.foodchem.2020.126881>

Clark, M., Centner, A. M., Ukhonov, V., Nagpal, R., & Salazar, G. (2022). Gallic acid ameliorates atherosclerosis and vascular senescence and remodels the microbiome in a sex-dependent manner in ApoE(-/-) mice. *The Journal of Nutritional Biochemistry*, 110, 109132. <https://doi.org/10.1016/j.jnutbio.2022.109132>

Davinelli, S., & Scapagnini, G. (2016). Polyphenols: A promising nutritional approach to prevent or reduce the progression of prehypertension. *High Blood Pressure and Cardiovascular Prevention*, 23(3), 197–202. <https://doi.org/10.1007/s40292-016-0149-0>

Deng, Y. H., Alex, D., Huang, H. Q., Wang, N., Yu, N., Wang, Y. T., Leung, G. P., & Lee, S. M. (2011). Inhibition of TNF-α-mediated endothelial cell-monocyte cell adhesion and adhesion molecules expression by the resveratrol derivative, trans-3,5,4'-trimethoxy-stilbene. *Phytotherapy Research*, 25(3), 451–457. <https://doi.org/10.1002/ptr.3279>

Ding, S., Xu, S., Fang, J., & Jiang, H. (2020). The protective effect of polyphenols for colorectal cancer. *Frontiers in Immunology*, 11, 1407. <https://doi.org/10.3389/fimmu.2020.01407>

- El-Kott, A. F., Shati, A. A., Al-Kahtani, M. A., & Alharbi, S. A. (2020). Kaempferol induces cell death in A2780 ovarian cancer cells and increases their sensitivity to cisplatin by activation of cytotoxic endoplasmic reticulum-mediated autophagy and inhibition of protein kinase B. *Folia Biologica*, 66(1), 36–46.
- Fairlie-Jones, L., Davison, K., Fromentin, E., & Hill, A. (2017). The effect of anthocyanin-rich foods or extracts on vascular function in adults: A systematic review and meta-analysis of randomised controlled trials. *Nutrients*, 9(8), 908. <https://doi.org/10.3390/nu9080908>
- Feresin, R. G., Huang, J., Klarich, D. S., Zhao, Y., Pourafshar, S., Arjmandi, B. H., & Salazar, G. (2016). Blackberry, raspberry and black raspberry polyphenol extracts attenuate angiotensin II-induced senescence in vascular smooth muscle cells. *Food and Function*, 7(10), 4175–4187. <https://doi.org/10.1039/c6fo00743k>
- Ferguson, J. J. A., Oldmeadow, C., Bentley, D., Eslick, S., & Garg, M. L. (2022). Effect of a polyphenol-rich dietary supplement containing *Pinus massoniana* bark extract on blood pressure in healthy adults: A parallel, randomized placebo-controlled trial. *Complementary Therapies in Medicine*, 71, 102896. <https://doi.org/10.1016/j.ctim.2022.102896>
- Forbes-Hernández, T. Y., Giampieri, F., Gasparrini, M., Afrin, S., Mazzoni, L., Cordero, M. D., Mezzetti, B., Quiles, J. L., & Battino, M. (2017). Lipid accumulation in HepG2 cells is attenuated by strawberry extract through AMPK activation. *Nutrients*, 9(6), 621. <https://doi.org/10.3390/nu9060621>
- Fouzder, C., Mukhuty, A., & Kundu, R. (2021). Kaempferol inhibits Nrf2 signalling pathway via downregulation of Nrf2 mRNA and induces apoptosis in NSCLC cells. *Archives of Biochemistry and Biophysics*, 697, 108700. <https://doi.org/10.1016/j.abb.2020.108700>
- Galiniak, S., Aebischer, D., & Bartusik-Aebischer, D. (2019). Health benefits of resveratrol administration. *Acta Biochimica Polonica*. <https://doi.org/10.18388/abp.2018.2749>
- García-Aguilar, A., & Guillén, C. (2022). Targeting pancreatic beta cell death in type 2 diabetes by polyphenols. *Frontiers in Endocrinology*, 13, 1052317. <https://doi.org/10.3389/fendo.2022.1052317>
- García-Aguilar, A., Palomino, O., Benito, M., & Guillén, C. (2021). Dietary polyphenols in metabolic and neurodegenerative diseases: Molecular targets in autophagy and biological effects. *Antioxidants*, 10(2), 142. <https://doi.org/10.3390/antiox10020142>
- García-Díez, E., López-Oliva, M. E., Pérez-Jiménez, J., Martín, M. A., & Ramos, S. (2022). Metabolic regulation of (-)-epicatechin and the colonic metabolite 2,3-dihydroxybenzoic acid on the glucose uptake, lipid accumulation and insulin signalling in cardiac H9c2 cells. *Food and Function*, 13(10), 5602–5615. <https://doi.org/10.1039/d2fo00182a>
- Giampieri, F., Cinciosi, D., Alvarez-Suarez, J. M., Quiles, J. L., Forbes-Hernández, T. Y., Navarro-Hortal, M. D., Machi, M., Casanova, R. d. J. P., Espinosa, J. C. M., Chen, X., Zhang, D., Bai, W., Lingmin, T., Mezzetti, B., Battino, M., & Diaz, Y. A. (2023). Anthocyanins: What do we know until now? *Journal of Berry Research*, 13, 1–6. <https://doi.org/10.3233/JBR-220087>
- Giampieri, F., Islam, M. S., Greco, S., Gasparrini, M., Forbes Hernandez, T. Y., Delli Carpini, G., Giannubilo, S. R., Ciavattini, A., Mezzetti, B., Mazzoni, L., Capocasa, F., Castellucci, M., Battino, M., & Ciarmela, P. (2019). Romina: A powerful strawberry with in vitro efficacy against uterine leiomyoma cells. *Journal of Cellular Physiology*, 234(5), 7622–7633. <https://doi.org/10.1002/jcp.27524>
- Godos, J., Vitale, M., Micek, A., Ray, S., Martini, D., Del Rio, D., Riccardi, G., Galvano, F., & Grosso, G. (2019). Dietary polyphenol intake, blood pressure, and hypertension: A systematic review and meta-analysis of observational studies. *Antioxidants*, 8(6), 152. <https://doi.org/10.3390/antiox8060152>
- Grabež, M., Škrbić, R., Stojiljković, M. P., Vučić, V., Grujić, V. R., Jakovljević, V., Djurić, D. M., Suručić, R., Šavikin, K., Bigović, D., & Vasiljević, N. (2022). A prospective, randomized, double-blind, placebo-controlled trial of polyphenols on the outcomes of inflammatory factors and oxidative stress in patients with type 2 diabetes mellitus. *Rapid Communications in Mass Spectrometry*, 23(2), 057. <https://doi.org/10.31083/j.rcm.2302057>
- Greco, S., Pellegrino, P., Giampieri, F., Capocasa, F., Delli Carpini, G., Battino, M., Mezzetti, B., Giannubilo, S. R., Ciavattini, A., & Ciarmela, P. (2023). The in vitro effects of Romina strawberry extract on 3D uterine leiomyosarcoma cells. *Nutrients*, 15(11), 2557. <https://doi.org/10.3390/nu15112557>
- Grosso, G., Micek, A., Castellano, S., Pajak, A., & Galvano, F. (2016). Coffee, tea, caffeine and risk of depression: A systematic review and dose-response meta-analysis of observational studies. *Molecular Nutrition and Food Research*, 60(1), 223–234. <https://doi.org/10.1002/mnfr.201500620>
- Grosso, G., Stepaniak, U., Micek, A., Kozela, M., Stefler, D., Bobak, M., & Pajak, A. (2017). Dietary polyphenol intake and risk of type 2 diabetes in the Polish arm of the Health, Alcohol and Psychosocial factors in Eastern Europe (HAPIEE) study. *The British Journal of Nutrition*, 118(1), 60–68. <https://doi.org/10.1017/s0007114517001805>
- Haş, I. M., Teleky, B. E., Vodnar, D. C., Ştefănescu, B. E., Tit, D. M., & Niţescu, M. (2023). Polyphenols and cardiometabolic health: Knowledge and concern among Romanian people. *Nutrients*, 15(10), 2281. <https://doi.org/10.3390/nu15102281>
- Hiroki, T., Monira, P., Shingo, G., Mamoru, I., & Yoriyuki, N. (2017). Beneficial effects of plant polyphenols on obesity. *Obesity and Control Therapies*, 4(3), 1–16.
- Jamar, G., Santamarina, A. B., Flygare, A. C., Gagliardi, A., de Rosso, V. V., Dourado, V. Z., & Pisani, L. P. (2020). Effects of the juçara fruit supplementation on metabolic parameters in individuals with obesity: A double-blind randomized controlled trial. *The Journal of Nutritional Biochemistry*, 83, 108430. <https://doi.org/10.1016/j.jnutbio.2020.108430>
- Jiang, H., Zhang, W., Li, X., Xu, Y., Cao, J., & Jiang, W. (2021). The anti-obesogenic effects of dietary berry fruits: A review. *Food Research International*, 147, 110539. <https://doi.org/10.1016/j.foodres.2021.110539>
- Kerimi, A., Gauer, J. S., Crabbe, S., Cheah, J. W., Lau, J., Walsh, R., Can-calón, P. F., & Williamson, G. (2019). Effect of the flavonoid hesperidin on glucose and fructose transport, sucrase activity and glycaemic response to orange juice in a crossover trial on healthy volunteers. *The British Journal of Nutrition*, 121(7), 782–792. <https://doi.org/10.1017/s0007114519000084>
- Kim, K. H., Ki, M. R., Min, K. H., & Pack, S. P. (2023). Advanced delivery system of polyphenols for effective cancer prevention and therapy. *Antioxidants*, 12(5), 1048. <https://doi.org/10.3390/antiox12051048>
- Kiyimba, T., Yiga, P., Bamuwamy, M., Ogowok, P., Van der Schueren, B., & Matthys, C. (2023). Efficacy of dietary polyphenols from whole foods and purified food polyphenol extracts in optimizing cardiometabolic health: A meta-analysis of randomized controlled trials. *Advances in Nutrition*, 14(2), 270–282. <https://doi.org/10.1016/j.advnut.2023.01.002>
- Le Roy, C. I., Wells, P. M., Si, J., Raes, J., Bell, J. T., & Spector, T. D. (2020). Red wine consumption associated with increased gut microbiota α -diversity in 3 independent cohorts. *Gastroenterology*, 158(1), 270–272. <https://doi.org/10.1053/j.gastro.2019.08.024>
- LiPing, Y., WenHua, L., ZhiCheng, D., YuMing, C., Dan, L., ShiZhou, D., ZhaoMin, L., & LiLi, Y. (2017). Effects of anthocyanins on cardiometabolic health: A systematic review and meta-analysis of randomized controlled trials. *Advances in Nutrition*, 8(5), 684–693. <https://doi.org/10.3945/an.116.014852>
- Maleki Dana, P., Sadoughi, F., Asemi, Z., & Yousefi, B. (2022). The role of polyphenols in overcoming cancer drug resistance: A comprehensive review. *Cellular and Molecular Biology Letters*, 27(1), 1. <https://doi.org/10.1186/s11658-021-00301-9>
- Maugeri, A., Calderaro, A., Patanè, G. T., Navarra, M., Barreca, D., Cirmi, S., & Felice, M. R. (2023). Targets involved in the anti-cancer activity of

- quercetin in breast, colorectal and liver neoplasms. *International Journal of Molecular Sciences*, 24(3), 2952. <https://doi.org/10.3390/ijms24032952>
- Menezes, R., Matafome, P., Freitas, M., & García-Conesa, M. T. (2022). Updated information of the effects of (poly)phenols against type-2 diabetes mellitus in humans: Reinforcing the recommendations for future research. *Nutrients*, 14(17), 3563. <https://doi.org/10.3390/nu14173563>
- Mileo, A., Nisticò, P., & Miccadei, S. (2019). Polyphenols: Immunomodulatory and therapeutic implication in colorectal cancer. *Frontiers in Immunology*, 10, 729. <https://doi.org/10.3389/fimmu.2019.00729>
- Mompeo, O., Spector, T. D., Matey Hernandez, M., Le Roy, C., Ista, G., Le Sayec, M., Mangino, M., Jennings, A., Rodriguez-Mateos, A., Valdes, A. M., & Menni, C. (2020). Consumption of stilbenes and flavonoids is linked to reduced risk of obesity independently of fiber intake. *Nutrients*, 12(6), 1871. <https://doi.org/10.3390/nu12061871>
- Motallebi, M., Bhia, M., Rajani, H. F., Bhia, I., Tabarraei, H., Mohammadkhani, N., Pereira-Silva, M., Kasaii, M. S., Nouri-Majid, S., Mueller, A.-L., Veiga, F. J. B., Paiva-Santos, A. C., & Shakibaei, M. (2022). Naringenin: A potential flavonoid phytochemical for cancer therapy. *Life Sciences*, 305, 120752. <https://doi.org/10.1016/j.lfs.2022.120752>
- Munir, K. M., Chandrasekaran, S., Gao, F., & Quon, M. J. (2013). Mechanisms for food polyphenols to ameliorate insulin resistance and endothelial dysfunction: Therapeutic implications for diabetes and its cardiovascular complications. *American Journal of Physiology. Endocrinology and Metabolism*, 305(6), 679–686. <https://doi.org/10.1152/ajpendo.00377.2013>
- Pacifici, F., Malatesta, G., Mammi, C., Pastore, D., Marzolla, V., Ricordi, C., Chierighin, F., Infante, M., Donadel, G., Curcio, F., Noce, A., Rovella, V., Lauro, D., Tesaro, M., Di Daniele, N., Garaci, E., Caprio, M., & Della-Morte, D. (2023). A novel mix of polyphenols and micronutrients reduces adipogenesis and promotes white adipose tissue browning via UCP1 expression and AMPK activation. *Cells*, 12(5), 714. <https://doi.org/10.3390/cells12050714>
- Patra, S., Pradhan, B., Nayak, R., Behera, C., Das, S., Patra, S. K., Efferth, T., Jena, M., & Bhutia, S. K. (2021). Dietary polyphenols in chemoprevention and synergistic effect in cancer: Clinical evidences and molecular mechanisms of action. *Phytomedicine*, 90, 153554. <https://doi.org/10.1016/j.phymed.2021.153554>
- Qian, Y., Babu, P. V. A., Symons, J. D., & Jalili, T. (2017). Metabolites of flavonoid compounds preserve indices of endothelial cell nitric oxide bioavailability under glucotoxic conditions. *Nutrition and Diabetes*, 7(9), e286. <https://doi.org/10.1038/nutd.2017.34>
- Ramos, L. V., da Costa, T. H. M., & Arruda, S. F. (2022). The effect of coffee consumption on glucose homeostasis and redox-inflammatory responses in high-fat diet-induced obese rats. *The Journal of Nutritional Biochemistry*, 100, 108881. <https://doi.org/10.1016/j.jnutbio.2021.108881>
- Romain, C., Chung, L. H., Marín-Cascales, E., Rubio-Arias, J. A., Gaillet, S., Laurent, C., Morillas-Ruiz, J. M., Martínez-Rodríguez, A., Alcaraz, P. E., & Cases, J. (2021). Sixteen weeks of supplementation with a nutritional quantity of a diversity of polyphenols from foodstuff extracts improves the health-related quality of life of overweight and obese volunteers: A randomized, double-blind, parallel clinical trial. *Nutrients*, 13(2), 492. <https://doi.org/10.3390/nu13020492>
- Sesso, H. D., Manson, J. E., Aragaki, A. K., Rist, P. M., Johnson, L. G., Friedenberg, G., Copeland, T., Clar, A., Mora, S., Moorthy, M. V., Sarkissian, A., Carrick, W. R., & Anderson, G. L. (2022). Effect of cocoa flavanol supplementation for the prevention of cardiovascular disease events: The COcoa supplement and multivitamin outcomes study (COSMOS) randomized clinical trial. *The American Journal of Clinical Nutrition*, 115(6), 1490–1500. <https://doi.org/10.1093/ajcn/nqac055>
- Sharma, A., Kaur, M., Katnoria, J., & Nagpal, A. (2018). Polyphenols in food: Cancer prevention and apoptosis induction. *Current Medicinal Chemistry*, 25(36), 4740–4757. <https://doi.org/10.2174/0929867324666171006144208>
- Shoji, T., Yamada, M., Miura, T., Nagashima, K., Ogura, K., Inagaki, N., & Maeda-Yamamoto, M. (2017). Chronic administration of apple polyphenols ameliorates hyperglycaemia in high-normal and borderline subjects: A randomised, placebo-controlled trial. *Diabetes Research and Clinical Practice*, 129, 43–51. <https://doi.org/10.1016/j.diabres.2017.03.028>
- Si, L., Zhang, L., Xing, S., Fang, P., Tian, X., Liu, X., & Xv, X. (2023). Curcumin as a therapeutic agent in cancer therapy: Focusing on its modulatory effects on circular RNAs. *Phytotherapy Research*, 37(7), 3121–3134. <https://doi.org/10.1002/ptr.7863>
- Solverson, P. M., Henderson, T. R., Debelo, H., Ferruzzi, M. G., Baer, D. J., & Novotny, J. A. (2019). An anthocyanin-rich mixed-berry intervention may improve insulin sensitivity in a randomized trial of overweight and obese adults. *Nutrients*, 11(12), 2876. <https://doi.org/10.3390/nu11122876>
- Suantawee, T., Thilavech, T., Cheng, H., & Adisakwattana, S. (2020). Cyanidin attenuates methylglyoxal-induced oxidative stress and apoptosis in INS-1 pancreatic β -cells by increasing glyoxalase-1 activity. *Nutrients*, 12(5), 1319. <https://doi.org/10.3390/nu12051319>
- Sun, S., & Fang, H. (2021). Curcumin inhibits ovarian cancer progression by regulating circ-PLEKHM3/miR-320a/SMG1 axis. *Journal of Ovarian Research*, 14(1), 158. <https://doi.org/10.1186/s13048-021-00916-8>
- Szczeńska, E., Białek-Dratwa, A., Janota, B., & Kowalski, O. (2022). Dietary therapy in prevention of cardiovascular disease (CVD)-tradition or modernity? A review of the latest approaches to nutrition in CVD. *Nutrients*, 14(13), 2649. <https://doi.org/10.3390/nu14132649>
- Thomson, C. A., Wertheim, B. C., Hingle, M., Wang, L., Neuhauser, M. L., Gong, Z., Garcia, L., Stefanick, M. L., & Manson, J. E. (2012). Alcohol consumption and body weight change in postmenopausal women: Results from the women's health initiative. *International Journal of Obesity*, 36(9), 1158–1164. <https://doi.org/10.1038/ijo.2012.84>
- Torres-Fuentes, C., Suárez, M., Aragonès, G., Mulero, M., Ávila-Román, J., Arola-Arnal, A., Salvadó, M. J., Arola, L., Bravo, F. I., & Muguerza, B. (2022). Cardioprotective properties of phenolic compounds: A role for biological rhythms. *Molecular Nutrition & Food Research*, 66(21), e2100990. <https://doi.org/10.1002/mnfr.202100990>
- Trisha, A. T., Shakil, M. H., Talukdar, S., Rovina, K., Huda, N., & Zzaman, W. (2022). Tea polyphenols and their preventive measures against cancer: Current trends and directions. *Foods*, 11(21), 3349. <https://doi.org/10.3390/foods11213349>
- Tsao, R. (2010). Chemistry and biochemistry of dietary polyphenols. *Nutrients*, 2(12), 1231–1246. <https://doi.org/10.3390/nu2121231>
- Villarini, M., Lanari, C., Nucci, D., Gianfredi, V., Marzulli, T., Berrino, F., Borgo, A., Bruno, E., Gargano, G., Moretti, M., & Villarini, A. (2016). Community-based participatory research to improve life quality and clinical outcomes of patients with breast cancer (DianaWeb in Umbria pilot study). *BMJ Open*, 6(6), e009707. <https://doi.org/10.1136/bmjopen-2015-009707>
- Vitale, M., Masulli, M., Rivellese, A. A., Bonora, E., Cappellini, F., Nicolucci, A., Squatrito, S., Antenucci, D., Barrea, A., Bianchi, C., Bianchini, F., Fontana, L., Fornengo, P., Giorgino, F., Gnasso, A., Mannucci, E., Mazzotti, A., Nappo, R., Palena, A. P., ... Vaccaro, O. (2018). Dietary intake and major food sources of polyphenols in people with type 2 diabetes: The TOSCA.IT Study. *European Journal of Nutrition*, 57(2), 679–688. <https://doi.org/10.1007/s00394-016-1355-1>
- Wang, S., Moustaid-Moussa, N., Chen, L., Mo, H., Shastri, A., Su, R., Bapat, P., Kwun, I., & Shen, C.-L. (2014). Novel insights of dietary polyphenols and obesity. *The Journal of Nutritional Biochemistry*, 25(1), 1–18. <https://doi.org/10.1016/j.jnutbio.2013.09.001>
- Wang, Y., Alkhalid, H., & Liu, D. (2021). The emerging role of polyphenols in the management of type 2 diabetes. *Molecules*, 26(3), 703. <https://doi.org/10.3390/molecules26030703>

- Wood, E., Hein, S., Heiss, C., Williams, C., & Rodriguez-Mateos, A. (2019). Blueberries and cardiovascular disease prevention. *Food & Function*, 10(12), 7621–7633. <https://doi.org/10.1039/c9fo02291k>
- Wu, Z., Zhao, X., Sun, Y., & Yu, H. (2022). Curcumin suppresses colorectal cancer development with epithelial-mesenchymal transition via modulating circular RNA HN1/miR-302a-3p/PIK3R3 axis. *Journal of Physiology and Pharmacology*, 73(2). <https://doi.org/10.26402/jpp.2022.2.05>
- Xia, Z. H., Chen, W. B., Shi, L., Jiang, X., Li, K., Wang, Y. X., & Liu, Y. Q. (2020). The underlying mechanisms of curcumin inhibition of hyperglycemia and hyperlipidemia in rats fed a high-fat diet combined with STZ treatment. *Molecules*, 25(2), 271. <https://doi.org/10.3390/molecules25020271>
- Xicota, L., Rodríguez, J., Langohr, K., Fitó, M., Dierssen, M., & De La Torre, R. (2020). Effect of epigallocatechin gallate on the body composition and lipid profile of down syndrome individuals: Implications for clinical management. *Clinical Nutrition*, 39(4), 1292–1300. <https://doi.org/10.1016/j.clnu.2019.05.028>
- Xu, J., Wang, X., Yue, J., Sun, Y., Zhang, X., & Zhao, Y. (2018). Polyphenols from acorn leaves (*Quercus liaotungensis*) protect pancreatic beta cells and their inhibitory activity against α -glucosidase and protein tyrosine phosphatase 1B. *Molecules*, 23(9), 2167. <https://doi.org/10.3390/molecules23092167>
- Xu, X., Zhang, X., Zhang, Y., & Wang, Z. (2021). Curcumin suppresses the malignancy of non-small cell lung cancer by modulating the circ-PRKCA/miR-384/ITGB1 pathway. *Biomedicine and Pharmacotherapy*, 138, 111439. <https://doi.org/10.1016/j.biopha.2021.111439>
- Yaskolka Meir, A., & Tsaban, G. (2022). Dietary strategies to reduce obesity burden-polyphenols as a game-changer? *Healthcare*, 10(12), 2430. <https://doi.org/10.3390/healthcare10122430>
- Zhao, Y., Li, D., & Huang, T. (2023). Associations of dietary flavonoids and subclasses with total and cardiovascular mortality among 369,827 older people: The NIH-AARP Diet and Health Study. *Atherosclerosis*, 365, 1–8. <https://doi.org/10.1016/j.atherosclerosis.2022.12.006>

How to cite this article: Armas Díaz, Y., Ferreiro Cotorruelo, M. S., & Battino, M. (2023). The role of dietary polyphenols in the control of chronic noncommunicable diseases. *Food Safety and Health*, 1(1), 13–21. <https://doi.org/10.1002/fsh3.12013>