



UNIVERSITÀ POLITECNICA DELLE MARCHE
Repository ISTITUZIONALE

Nicotinamide N-methyltransferase gene silencing enhances chemosensitivity of melanoma cell lines

This is the peer reviewed version of the following article:

Original

Nicotinamide N-methyltransferase gene silencing enhances chemosensitivity of melanoma cell lines / Campagna, R; Salvolini, E; Pompei, V; Pozzi, V; Salvucci, A; Molinelli, E; Brisigotti, V; Sartini, D; Campanati, A; Offidani, A; Emanuelli, M. - In: PIGMENT CELL & MELANOMA RESEARCH. - ISSN 1755-148X. - STAMPA. - 34:6(2021), pp. 1039-1048. [10.1111/pcmr.12993]

Availability:

This version is available at: 11566/290822 since: 2024-04-11T11:01:13Z

Publisher:

Published

DOI:10.1111/pcmr.12993

Terms of use:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. The use of copyrighted works requires the consent of the rights' holder (author or publisher). Works made available under a Creative Commons license or a Publisher's custom-made license can be used according to the terms and conditions contained therein. See editor's website for further information and terms and conditions.

This item was downloaded from IRIS Università Politecnica delle Marche (<https://iris.univpm.it>). When citing, please refer to the published version.

note finali coverage

(Article begins on next page)

This is the peer reviewed version of the following article: Nicotinamide N-methyltransferase gene silencing enhances chemosensitivity of melanoma cell lines / Campagna, R; Salvolini, E; Pompei, V; Pozzi, V; Salvucci, A; Molinelli, E; Brisigotti, V; Sartini, D; Campanati, A; Offidani, A; Emanuelli, M. - In: PIGMENT CELL & MELANOMA RESEARCH. - ISSN 1755-148X. - STAMPA. - 34:6(2021), pp. 1039-1048. © 2021 John Wiley & Sons A/S. Published by John Wiley & Sons Ltd . Published by John Wiley & Sons Ltd, which has been published in final form at <https://dx.doi.org/10.1111/pcmr.12993>. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions <https://authorservices.wiley.com/author-resources/Journal-Authors/licensing/self-archiving.html>. This article may not be enhanced, enriched or otherwise transformed into a derivative work, without express permission from Wiley or by statutory rights under applicable legislation. Copyright notices must not be removed, obscured or modified. The article must be linked to Wiley's version of record on Wiley Online Library and any embedding, framing or otherwise making available the article or pages thereof by third parties from platforms, services and websites other than Wiley Online Library must be prohibited.



Nicotinamide N-Methyltransferase Gene Silencing Enhances Chemosensitivity of Melanoma Cell Lines.

Journal:	<i>Pigment Cell & Melanoma Research</i>
Manuscript ID	21-O-035.R2
Manuscript Type:	Original Article
Date Submitted by the Author:	11-May-2021
Complete List of Authors:	<p>Campagna, Roberto; Università Politecnica delle Marche, Department of Clinical Sciences Salvolini, Eleonora; Università Politecnica delle Marche, Department of Clinical Sciences Pompei, Veronica; Università Politecnica delle Marche, Department of Clinical Sciences Pozzi, Valentina; Università Politecnica delle Marche, Department of Clinical Sciences Salvucci, Alessia; Università Politecnica delle Marche, Department of Clinical Sciences Molinelli, Elisa; Università Politecnica delle Marche, Department of Clinical and Molecular Sciences Brisigotti, Valerio; Università Politecnica delle Marche, Department of Clinical and Molecular Sciences Sartini, Davide; Università Politecnica delle Marche, Department of Clinical Sciences Campanati, Anna; Università Politecnica delle Marche, Department of Clinical and Molecular Sciences Offidani, Annamaria; Università Politecnica delle Marche, Department of Clinical and Molecular Sciences Emanuelli, Monica; Università Politecnica delle Marche, Department of Clinical Sciences; Università Politecnica delle Marche, New York-Marche Structural Biology Center (NY-MaSBiC)</p>
Keywords:	Nicotinamide N-methyltransferase, Cutaneous melanoma, Cell proliferation, Cell migration, Chemosensitivity

TITLE

Nicotinamide N-Methyltransferase Gene Silencing Enhances Chemosensitivity of Melanoma Cell Lines

ABSTRACT

Melanoma accounts for less than 5% of all cutaneous neoplasms but is responsible for the greater part of skin cancer-related deaths. Therefore, the identification of molecules that could serve as therapeutic target is urgent. This study focused on the enzyme nicotinamide N-methyltransferase (NNMT). The effect of NNMT knockdown on cell proliferation and migration of A375 melanoma cells was evaluated by MTT and wound healing assays, respectively. Viability of A375 cells downregulating NNMT was also explored under treatment with dacarbazine, a chemotherapeutic drug approved for advanced melanoma treatment. The impact of enzyme knockdown on cell proliferation and chemosensitivity was also investigated in WM-115 melanoma cells. Results obtained demonstrated that NNMT silencing led to a significant reduction of cell proliferation and migration of A375 cells. Moreover, enzyme downregulation was associated with an increase of melanoma cells sensitivity to treatment with dacarbazine. Analogous effects induced by enzyme knockdown on cell proliferation and chemosensitivity were also found in WM-115 cell line. Our data seem to demonstrate that NNMT could represent a promising molecular target for the effective treatment of this form of skin cancer.

KEYWORDS

Nicotinamide N-methyltransferase; cutaneous melanoma; cell proliferation; cell migration; chemosensitivity.

1. INTRODUCTION

Melanoma arises from the transformation of cutaneous or mucous melanocytes. Cutaneous malignant melanoma represents less than 5% of all malignant skin neoplasms but, despite the low incidence, it is responsible for the majority of skin cancer-related deaths, due to its tendency to metastasize [1,2]. Its incidence varies greatly between countries and shows a steady increase [3]. Ultraviolet radiation exposure represents the main environmental risk factor, although host risks factors, such as genetic predisposition and the number of nevi, play an important role in tumor development [3].

If melanoma is diagnosed at an early stage, surgical excision is curative in the majority of cases [4]. Unfortunately, approximately 10% of neoplasms are detected at an advanced stage. These lesions are more difficult to treat although several therapeutic agents have been recently approved [3-5]. Despite progress in advanced malignant melanoma therapy, the patient prognosis remains poor [6]. The identification of reliable biomarkers represents therefore a primary goal.

In the present study, we focused on nicotinamide N-methyltransferase (NNMT), a cytosolic enzyme catalyzing the N-methylation of nicotinamide, pyridines, and other structural analogues [7], which was previously found to be upregulated in many solid tumors, among which cutaneous [8] and oral melanoma [9]. Moreover, a reduced proliferation and tumorigenicity of several cell lines, associated with NNMT downregulation, was also reported [10-12], while enzyme overexpression led to an increase of cell viability

[13]. Taken together, these results indicate that NNMT may be involved in tumor initiation and growth, thus supporting its potential as a novel diagnostic and prognostic biomarker, as well as an interesting therapeutic target.

Our aim was to analyze enzyme expression levels in human skin melanoma A375 cell line by Real-Time PCR and Western blot, and to evaluate the effect of NNMT gene silencing on cell proliferation and migration by means of MTT and wound healing assays, respectively. Subsequently, proliferation of NNMT downregulating A375 and control cells was evaluated after treatment with dacarbazine, in order to explore the potential involvement of the enzyme in sensitivity of skin melanoma cells to chemotherapy. The impact of NNMT knockdown on cell proliferation and chemosensitivity was further confirmed in WM-115 melanoma cell line.

2. MATERIALS AND METHODS

2.1 Cell lines and reagents

The human malignant melanoma cell line A375 was purchased from the American Type Culture Collection, while WM-115 melanoma cells were kindly provided by Dr. Claudia Matteucci, Department of Experimental Medicine and Surgery, University of Rome "Tor Vergata", Rome, Italy. Cells were cultured in Dulbecco's Modified Eagle's Medium (D-MEM) (Lonza, Basel, Switzerland) supplemented with L-glutamine 2mM, 10% fetal bovine serum (FBS) and 50µg/ml gentamicin, at 37°C in a humidified 5% CO₂ incubator.

2.2 shRNA-mediated gene silencing of NNMT

To achieve NNMT knockdown in A375 and WM-115 cells, a set of pLKO.1 vectors containing stem-loop cassettes encoding for different short hairpin RNAs (shRNAs) targeting different region of NNMT mRNA was used (Sigma-Aldrich, Saint Louis, MO), according to what previously reported [12].

The day before transfection, cells were seeded in 24-well plates (4×10^4 cells/well) with different plasmids (0.5µg for each well) coding each a specific shRNA against NNMT (pLKO.1-164, pLKO.1-330, pLKO.1-711), or with a combination of all three plasmids (0.5µg final DNA amount, named as pLKO.1-mix). Control cells were transfected with an empty vector (pLKO.1-puro) or treated with transfection reagent only (Mock).

Transfection was performed by using FuGENE HD Transfection reagent (Promega, Madison, WI, USA) according to manufacturer's instruction. Forty-eight hours from the beginning of transfection, cellular clones stably downregulating NNMT started to be selected by cultivating cells with complete medium containing 1µg/ml puromycin. For all subsequent experiments, puromycin resistant cells were maintained in complete selection medium

The efficiency of NNMT silencing in A375 and WM-115 cells was evaluated by Real-Time PCR and Western blot analysis.

2.3 Real-Time PCR

1
2
3 Total RNA isolation, reverse transcription and Real-Time PCR assay were performed according to the schemes
4 described elsewhere [11].
5

6 Cell pellets (1×10^6 cells) were homogenized in a lysis buffer and total RNA was isolated with SV Total
7 RNA Isolation System (Promega, Madison, WI, USA), according to manufacturer's protocol. Quantity and
8 quality of RNA were spectrophotometrically evaluated at 260nm and 280nm. Total RNA (2ug) was reverse
9 transcribed with M-MLV Reverse Transcriptase (Promega), using random primers in a total volume of 25 μ l for
10 60 min at 37°C.
11
12
13

14 The quantitative evaluation of NNMT mRNA expression was carried out using Real-Time PCR assay
15 performed with CFX96 Real-Time PCR Detection System (Bio-Rad Laboratories, Hercules, CA, USA).
16 cDNA, generated as described above, was used as template. The primer oligonucleotides sequences were 5'-
17 GAATCAGGCTTCACCTCCAA-3' (forward) and 5'-TCACACCGTCTAGGCAGAAT-3' (reverse) for
18 NNMT, and 5'-TCACACCGTCTAGGCAGAAT-3' (forward) and 5'-AGCACTGTGTTGGCGTACAG-3'
19 (reverse) for β -actin. Genes were run in duplicate using SsoFaste EvaGreen Supermix (Bio-Rad Laboratories)
20 for 40 cycles at 95°C for 30 s and 58°C for 30s. All samples were tested in triplicate with the reference gene
21 β -actin for data normalization to correct for variations in RNA quality and quantity. Direct detection of
22 amplification products was monitored by measuring the fluorescence produced by EvaGreen. The
23 measurements were then plotted against cycle numbers. Threshold cycle (Ct) was defined as the cycle number
24 at which the first fluorescence increase above the threshold was detected. Changes in relative NNMT
25 expression were calculated by $2^{-\Delta(\Delta Ct)}$, where $\Delta Ct = Ct(\text{NNMT}) - Ct(\beta\text{-actin})$ and $\Delta(\Delta Ct) = \Delta Ct(\text{pLKO.1-164,}$
26 $\text{pLKO.1-330, pLKO.1-711, pLKO.1-mix or pLKO.1-puro}) - \Delta Ct(\text{Mock})$.
27
28
29
30
31
32
33
34
35

36 2.4 Western blot

37 A Western blot assay was used to evaluate NNMT protein levels. Cells homogenization and Western blot
38 procedure were performed as previously described [12].
39
40

41 Cell pellets (2×10^6 cells) were suspended in 200 μ l of lysis buffer (phosphate buffered saline containing
42 1% Nonidet P-40, 0.1% sodium dodecyl sulphate, 1mM phenylmethylsulphonyl fluoride and 2 μ g/ml aprotinin)
43 and homogenized by passing (3–10 times) through a 30G needle attached to a 1ml syringe. Lysates were
44 centrifuged at 16000xg at 4°C for 10 min and supernatants represented protein extract. 50 μ g of each sample
45 were subjected to 15% sodium dodecyl sulphate-polyacrylamide gel electrophoresis and transferred to
46 polyvinylidene fluoride membranes. After blocking and washing, membranes were incubated with rabbit
47 polyclonal antibody against NNMT (Sigma-Aldrich) (1:1000 dilution) or with rabbit polyclonal antibody
48 against β -actin (Sigma-Aldrich) (1:8000 dilution) for 1h, followed by an incubation with horseradish
49 peroxidase-conjugated goat anti-rabbit IgG (Sigma-Aldrich) (1:150000 dilution) for 1h. An incubation (5 min)
50 with Super-Signal West Femto Maximum Sensitivity Substrate (Thermo Fisher Scientific, Waltham, MA,
51 USA) allowed the detection of the NNMT-related chemiluminescent signals.
52
53
54
55
56
57
58
59
60

1
2
3 Chemiluminescent bands were then acquired using a ChemiDoc XRS+ System (Bio-Rad Laboratories,
4 Hercules, CA, USA), and signal intensity was quantified using Image Lab Software (Bio-Rad Laboratories,
5 Hercules, CA, USA). Each experiment was performed in triplicate and independently repeated three times.
6
7
8

9 **2.5 Protein concentration assay**

10 Protein concentration was evaluated through the Bradford method using bovine serum albumin as standard.
11
12
13

14 **2.6 MTT assay**

15 Cell proliferation was evaluated using a colorimetric assay that quantifies the conversion of 3-(4,5-
16 dimethylthiazol-2-yl)-2,5-diphenyl tetrazolium bromide (MTT) to insoluble formazan by dehydrogenase
17 enzymes of undamaged mitochondria of living cells. The MTT assay was performed in accordance with what
18 previously described [12].
19
20
21

22 Briefly, cells downregulating NNMT (pLKO.1-164, pLKO.1-330, pLKO.1-711 and pLKO.1-mix for
23 A375 cell line; pLKO.1-711 for WM-115 cells), as well as controls (pLKO.1-puro and Mock) cells, were
24 seeded in 96-well plates (1×10^3 cells/well for A375 cells; 3.5×10^3 cells/well for WM-115 cell line). The day
25 after seeding, cell proliferation was evaluated (0h, 24h, 48h and 72h) by measuring the conversion of the
26 tetrazolium salt MTT to formazan crystals. 100 μ l D-MEM containing 8.4 μ l MTT reagent (5mg/ml in
27 phosphate buffered saline) were added to cells. After 4h of incubation at 37°C, medium was discarded,
28 formazan crystals were dissolved by addition of 200 μ l of dimethyl sulfoxide and the reaction product was
29 quantified by assessing the absorbance at 540nm.
30
31
32
33
34

35 Each experiment was repeated three times. Results were expressed as percentage of the control (control
36 equals 100% and corresponds to the absorbance value of each sample at 0h) and presented as mean values \pm
37 standard deviation of three independent experiments performed in triplicate.
38
39
40

41 **2.7 Monolayer wound healing assay**

42 In order to evaluate migration capability, A375 cells were seeded into 6-well plates (3×10^5 cells/well) and
43 allowed to grow up to 100% confluency. The wound on cells monolayers was made using sterile 200 μ l pipette
44 tips. After scratching, cell debris were removed by washing cell monolayers three times and medium was
45 replaced with D-MEM supplemented with L-glutamine 2mM and 0.5% FBS. Wounded monolayers were
46 monitored under a microscope equipped with a camera and photographed at 0h, 4h, 8h, 24h and 28h after
47 scratching. Images were analyzed by the software ImageJ (Rasband, W.S., ImageJ, U. S. National Institutes of
48 Health, Bethesda, Maryland, USA, <https://imagej.nih.gov/ij/>, 1997-2018). Each experiment was run in
49 triplicate and independently replicated three times.
50
51
52
53
54
55

56 **2.8 Chemotherapeutic treatment**

57 Cells were seeded in 96-well plates (1×10^3 cells/well for A375 cell line; 3.5×10^3 cells/well for WM-115 cells).
58 The day after seeding, A375 and WM-115 cells started to be treated with complete medium, containing
59
60

(Sigma-Aldrich) 100 and 10 μ g/ml dacarbazine, respectively, previously dissolved in HCl 1M at 50mg/ml stock solution. Untreated cells were grown in complete medium containing HCl only, diluted at 2mM final concentration. This HCl concentration value (which did not affect the intracellular pH) corresponds to that obtained in samples subjected to treatment with chemotherapeutic drug. MTT assay was used to estimate the cell proliferation in untreated samples (0h) and at different time points (24h, 48h and 72h) after starting treatment with drug.

2.9 Statistical analysis

Data analysis was performed using GraphPad Prism software version 8.00 for Windows (GraphPad Software, San Diego, USA). Differences between groups were established using two-way analysis of variance (ANOVA). A p-value <0.05 was considered as statistically significant.

3 RESULTS

3.1 Efficiency of shRNA-mediated gene silencing of NNMT

In order to achieve enzyme knockdown, A375 and WM-115 cell lines were transfected with plasmids encoding shRNA targeting a different region of NNMT. Control cells were transfected with an empty vector (pLKO.1-puro) or treated with transfection reagent only (Mock). The effect of shRNA treatment on enzyme expression was determined by measuring the NNMT mRNA and protein levels, through Real-Time PCR and Western blot, respectively.

Results obtained showed that A375 cells transfected with plasmids encoding shRNA targeting NNMT mRNA displayed a significant ($p < 0.05$) enzyme downregulation (0.48 \pm 0.02 for pLKO.1-164, 0.49 \pm 0.05 for pLKO.1-330, 0.37 \pm 0.03 for pLKO.1-711 and 0.56 \pm 0.03 for pLKO.1-mix) compared to Mock (1.00 \pm 0.07) or to cells treated with pLKO.1-puro empty vector (1.04 \pm 0.03) (Figure 1a). Western blot analysis confirmed a decreased NNMT expression in cells transfected with pLKO.1-164, pLKO.1-330, pLKO.1-711 or pLKO.1-mix compared with those treated with empty vector (pLKO.1-puro) or with transfection reagent only (Mock) (Figure 1b and c).

Analogously, WM-115 transfected with pLKO.1-711 exhibited significantly ($p < 0.05$) lower NNMT mRNA levels (0.24 \pm 0.01) compared with controls (1.00 \pm 0.07 for mock; 1.04 \pm 0.03 for pLKO.1-puro) (Figure 1d).

3.2 Effect of shRNA-mediate gene silencing of NNMT on cell proliferation and migration

To explore the role of NNMT in cancer cell metabolism and evaluate the biological effect of enzyme downregulation, cell proliferation and migration were analyzed at different time points.

The influence of NNMT knockdown on cell proliferation was assessed by MTT assay, and results of colorimetric assay were expressed as relative cell proliferation referred to the control (absorbance at 0h and equal to 100%). As shown in Figure 2a, A375 cells treated with empty vector (pLKO.1-puro) showed no significant difference in cell proliferation with respect to Mock. Interestingly, enzyme downregulation

determined a significant ($p < 0.05$) decrease of cell proliferation at 24h, 48h and 72h, for all cellular clones stably encoding shRNAs against NNMT mRNA (pLKO.1-164, pLKO.1-330, pLKO.1-711 and pLKO.1-mix) compared with controls (pLKO.1-puro and Mock). Moreover, cell proliferation was significantly ($p < 0.05$) decreased in WM-115 cell line downregulating NNMT compared with controls (Figure 2b).

Monolayer wound healing assay was performed to monitor the migration ability of A375 cells. For each sample, results were reported as percentage of wound recovery with respect to 0h time point. Compared with Mock or cells treated with empty vector (pLKO.1-puro), the migration rate of A375 cells was significantly ($p < 0.05$) decreased after NNMT silencing at all examined time points (4h, 8h, 12h, 24h and 28h) (Figure 3, panels a and b).

3.3 NNMT influence on A375 and WM-115 cells sensitivity to chemotherapeutic treatment

MTT assay was used to evaluate the effect of treatment with dacarbazine on cell proliferation. Cell treatment with medium containing 2mM HCl (Untreated) determined no significant difference in cell proliferation compared to cells treated with complete medium only (data not shown). At 48h and 72h time points, treatment with dacarbazine significantly ($p < 0.05$) reduced proliferation of both A375 cells transfected with plasmids encoding shRNAs targeting NNMT mRNA (pLKO.1-164, pLKO.1-330, pLKO.1-711 and pLKO.1-mix) and Mock.

Interestingly, the decrease of cell proliferation upon treatment with dacarbazine was markedly enhanced in NNMT downregulating A375 cells compared with that detected in Mock (Figure 4a-b). Similarly, NNMT silenced WM-115 cells (pLKO.1-711) showed an increased sensitivity to treatment with dacarbazine with respect to control (Mock) (Figure 4e).

4 DISCUSSION

Cutaneous malignant melanoma (CMM) represents a widespread form of cancer associated with a high mortality rate. Excisional surgery is the first therapeutic option for early-stages CMM, followed by lymph node dissection, if appropriate [14]. However, about 9% of patients displays regional metastatic disease, while 4% of patients are diagnosed showing distant metastasis [15]. These patients display a poor prognosis, since their tumors are disseminated and unresectable. Moreover, unlike other solid tumors, advanced CMM typically responds weakly to traditional chemotherapy [14]. However, due to the lack of other effective therapeutic strategies, chemotherapy is still the best options for these patients, if affected with melanomas that do not harbor somatic mutations, such as those related with BRAF, NRAS, or cKIT. These subjects, in fact, cannot receive benefit from targeted immunotherapy based on specific inhibitors and without chemotherapy undergo disease progression [16].

The alkylating agent dacarbazine is used for treatment of advanced-stage CMM. This molecule gained FDA approval for the melanoma treatment in 1975 and therefore represents a longstanding chemotherapeutic drug. Several phase I and phase II clinical trials demonstrated partial and complete response rates in 15–28% and 3–5% treated patients, respectively, and durable responses were found in <2% subjects, only [17]. In order

1
2
3 to try to increase the efficacy of dacarbazine, a number of clinical studies have been carried out by using this
4 compound combined with immunotherapies or other chemotherapeutic agents [18]. However, no improvement
5 in overall survival was observed, but only an increase of side effects and the reduction of life quality [18].
6 Therefore, despite poor response rates, the overall outcome from these studies demonstrated that dacarbazine
7 is still the standard chemotherapeutic option for patients with metastatic CMM [16,19-20]. In the light of these
8 considerations, it is urgent to identify molecules, involved in melanoma initiation and progression, which are
9 associated with resistance to chemotherapy, aiming to set up the basis for a targeted treatment of this neoplasm.
10 The purpose of this research work was to explore the role played by NNMT in melanoma cell metabolism,
11 focusing on aspects related to the possibility of the enzyme to affect cell sensitivity to chemotherapeutic
12 treatment.
13

14
15 An overexpression of NNMT was found in numerous neoplasms, including clear cell renal cell
16 carcinoma (ccRCC) [21,22], oral squamous cell carcinoma (OSCC) [23-25], non-small cell lung cancer
17 (NSCLC) [26], bladder urothelial carcinoma (BUC) [27,28], and recently CMM [8] and basal cell carcinoma
18 (BCC) [29].

19
20 NNMT upregulation in ccRCC displayed to be inversely related with the size of tumor, leading to
21 hypothesize that the enzyme may be implicated in cancer progression [21]. In OSCC, NNMT overexpression
22 was negatively correlated with pT, lymph node metastasis, pathological staging and histological grading, thus
23 supporting its potential contribution in tumor growth and differentiation [23,24]. In pathological samples
24 obtained from CMM patients, an inverse correlation was found between enzyme levels and clinicopathological
25 parameters, such as Breslow thickness, Clark level, the presence/number of mitoses and ulceration [8]. The
26 recent study, showing NNMT upregulation in BCC compared with healthy tissue margin, revealed that enzyme
27 expression was significantly enhanced in nodular forms with respect to the infiltrative subtype, thus
28 highlighting an interesting inverse correlation between NNMT levels and tumor aggressiveness [29]. The
29 above results, taken together, seem to suggest that the enzyme could represent a candidate as molecular
30 biomarker for both diagnosis and prognosis in the above-mentioned neoplasms.
31

32
33 NNMT expression was also found to be significantly higher in exfoliated cells from urines of patients
34 with BUC compared with those of controls. Furthermore, statistical analyzes revealed an inverse correlation
35 between tumor enzyme levels and histological grade, demonstrating the excellent diagnostic accuracy of a
36 potential NNMT-based urine test [28]. Similarly, preliminary results indicated an upregulation of NNMT in
37 saliva samples collected from OSCC patients compared with those of healthy subjects, strongly suggesting its
38 potential application as a new salivary diagnostic biomarker for early and non-invasive detection of oral cancer
39 [25].
40

41
42 The mRNA and protein levels of the NNMT, as well as its catalytic activity were also determined in
43 several human tumor cell lines and corresponding cell populations enriched with cancer stem cells (CSCs).
44 Data obtained revealed that enzyme expression was markedly increased in CSC-enriched populations
45 compared with parental counterparts [30,31]. Given the role played by CSCs in tumor initiation, progression
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 and relapse, as well as in resistance to radio- and chemotherapy, these results strongly highlight the leading
4 function exerted by NNMT in the metabolism of the cancer cells.
5

6 To date, it is not clear the exact mechanism by which NNMT may promote tumorigenesis and tumor
7 progression. It has been proposed that NNMT can affect the methylation state of tumor cells not only by
8 altering the ratio SAM:SAH ratio, but more likely by storing methylation units in tumor cells in form of MNA,
9 thus fine-tuning the methylation state of cancer cells [32]. Furthermore, NNMT acts as a master regulator of
10 the homeostasis of nicotinamide, a key precursor of the NAD⁺. Therefore, the catalytic activity of the NNMT
11 can control the amount of nicotinamide inside the cell available for the energy metabolism, thus influencing
12 and modulating multiple pathways linked to both cellular survival and death [33]. Some of them, involving
13 SIRT1 and the PARPs enzymes, are crucial for the correct DNA repair which prevents the neoplastic
14 transformation [33, 34].
15

16 Results obtained in the present study clearly demonstrated that NNMT knockdown led to a significant
17 reduction of cell proliferation and migration of melanoma cell lines. Moreover, the enzyme downregulation in
18 melanoma cells was associated with an increased sensitivity to treatment with dacarbazine. In the context of
19 studies carried out to explain the significance of NNMT upregulation in solid tumors and to clarify the impact
20 of such dysregulation within cellular events promoting malignant transformation, a great contribution was
21 made from our research group. Enzyme downregulation was significantly associated with a decrease of cell
22 viability and anchorage-independent cell growth of HeLa-derived KB cancer cells [10], PE/CA PJ-15 OSCC
23 cells [11] and NSCLC tumor cell line A549 [35]. Further-more, NNMT knockdown led to a considerable
24 reduction of tumor formation capacity of PE/CA-PJ15 cells upon subcutaneous transplantation into athymic
25 mice [11]. Conversely, the induction of enzyme overexpression in HSC-2 OSCC cell line was significantly
26 associated with enhanced cell proliferation [13].
27

28 Treatment with short interfering RNAs against NNMT mRNA led to a marked reduction of chemotaxis
29 and chemokinesis of the human BUC cell line 253J, thus highlighting the crucial role exerted by the enzyme
30 in cellular events promoting cancer cell invasion and metastasis [36]. shRNA-mediated gene silencing of
31 NNMT significantly reduced tumor formation and metastatic spread of 786O ccRCC cells subcutaneously
32 transplanted into athymic mice, through the activation of the PI3K/Akt/SP1/MMP-2 pathway [37]. NNMT
33 knockdown in Bcap-37 and MDA-MB-231 human breast cancer cells, originally exhibiting high enzyme
34 levels, was associated with the induction of apoptosis via the mitochondria-mediated pathway and led to a
35 significant reduction of cell growth in vitro, as well as tumor formation in vivo. Converse effects on cellular
36 phenotype emerged after the induction of enzyme upregulation in MCF-7 and SK-BR-3 breast cancer cell lines
37 displaying no endogenous NNMT expression [38]. By mimicking the above reported experimental procedure,
38 NNMT role was explored in SW480 and HT-29 colorectal cancer (CRC) cell lines, lacking and showing high
39 constitutive enzyme levels, respectively. Results obtained demonstrated that NNMT was able to enhance cell
40 proliferation and colony formation ability, inhibit apoptotic pathway, promote the progression of cell cycle,
41 and lead to an increase of ATP levels [39]. Furthermore, NNMT downregulation was significantly associated
42 with reduction of cell growth and migration, as well as invasive capacity of the human pancreatic cancer cell
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 line PANC-1, thus highlighting enzyme involvement in cell proliferation and meta-static potential of cancer
4 cell [40].

5
6 In the human NSCLC H1993 cells, which displayed high endogenous NNMT expression levels and
7 were resistant to the treatment with the inhibitor of the tyrosine kinase domain of the EGFR gefitinib, enzyme
8 knockdown significantly enhanced sensitivity to this molecule and suppressed colony formation ability [41].
9 The induction of NNMT overexpression in PC3 prostate cancer cell line enhanced cell proliferation, invasive
10 capacity and migration ability, by increasing mRNA levels of histone deacetylase SIRT1. On the contrary,
11 treatment with nicotinamide reversed the above reported effects, since suppressed the expression of SIRT1 and
12 markedly decreased the invasive capacity of PC3 cells [42]. NNMT upregulation in BGC-823 gastric cancer
13 cells led to the induction of transforming growth factor- β 1 (TGF- β 1) expression, with consequent activation
14 of TGF- β 1/Smad signaling, which in turn promotes epithelial-mesenchymal transition (EMT) as a key event
15 in tumor invasion and metastasis [43]. Similarly, treatment of EC9706 and TE1 esophageal squamous cell
16 carcinoma cells with short interfering RNAs targeted to NNMT mRNA significantly suppressed cell viability
17 and migration, induced cell cycle arrest, promoted apoptosis and inhibited EMT via Wnt/ β -catenin pathway
18 [44]. Taken together, all these results seem to indicate that NNMT is a fundamental mediator of crucial events
19 inducing malignant transformation, thus representing a potential molecular target for cancer therapy.

20
21 A few studies evaluated the ability of NNMT to influence the efficacy of drugs used for
22 chemotherapeutic treatment. Enzyme downregulation diminished resistance to 5-fluorouracile (5-FU) in CRC
23 HT-29 cells, while the opposite effect was induced in CRC cell line SW480 upon NNMT overexpression. The
24 ability of NNMT to increase chemosensitivity was found to be mediated by the enzyme reaction product N1-
25 methylnicotinamide, which reduces the production of reactive oxygen species induced by 5-FU treatment [45].
26 Furthermore, a recent study carried out in esophageal squamous cell carcinoma cell lines showed that NNMT
27 knockdown increases 5-FU sensitivity of TE1 cells via suppressing Warburg effect, while overexpression of
28 NNMT in EC1 and Eca109 cells increased the 5-FU sensitivity, thus suggesting that NNMT might be a
29 potential therapeutic target to enhance the therapeutic efficacy of 5-FU [46]. Moreover, ectopic expression of
30 NNMT in MDA-MB-231 breast cancer cells significantly inhibited apoptosis and suppression of colony
31 formation induced by adriamycin and paclitaxel, through the induction SIRT1 expression [47].

32
33 Results obtained in this work, seems to suggest that NNMT is somehow involved in mechanisms
34 promoting melanoma cell resistance to chemotherapy. In this light, further studies will clarify whether the
35 combination of enzyme inhibitors with dacarbazine could enhance sensitivity of A375 cells to
36 chemotherapeutic treatment. Indeed, cloning, expression and purification of human recombinant NNMT
37 allowed to determine the crystal structure of the enzyme, as well as to identify the main amino acid residues
38 involved in catalysis exerted by NNMT [7], thus leading to design and assay new and effective NNMT
39 inhibitors [48].

40
41 To the best of our knowledge, this is the first study to evaluate the role played by NNMT in cellular
42 events featuring melanoma progression and spread, as well as to explore its potential involvement in molecular
43 processes affecting chemoresistance. Although further analyses are necessary to give a deeper insight into the
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

mechanisms by which the enzyme could participate to melanoma tumorigenesis, our results seem to demonstrate that NNMT could represent an interesting molecular target for effective treatment of this form of skin cancer.

ACKNOWLEDGMENTS

We would like to offer our sincere thanks to Dr. Claudia Matteucci (Department of Experimental Medicine and Surgery, University of Rome “Tor Vergata”, Rome, Italy), for providing with WM-115 melanoma cell line.

CONFLICTS OF INTEREST

The authors declare no conflict of interest.

REFERENCES

1. Ossio, R., Roldán-Marín, R., Martínez-Said, H., Adams, D.J., Robles-Espinoza, C.D. (2017). Melanoma: a global perspective. *Nat Rev Cancer*, 17, 393-394.
2. Erdei, E., Torres, S.M. A new understanding in the epidemiology of melanoma. (2010). *Exp Rev Anticancer Ther*, 10, 1811–1823.
3. Leonardi, G., Falzone, L., Salemi, R., Zanghi, A., Spandidos, D.A., Mccubrey, J.A., Candido, S., Libra, M. (2018). Cutaneous melanoma: From pathogenesis to therapy. *Int J Oncol*, 52, 1071–1080.
4. Ross, M.I., Gershenwald, J.E. (2011). Evidence-based treatment of early-stage melanoma. *J Surg Oncol*, 104, 341–353.
5. Malas, S., Harrasser, M., Lacy, K.E., Karagiannis, S.N. (2014). Antibody therapies for melanoma: New and emerging opportunities to activate immunity. *Oncol Rep*, 32, 875–886.
6. Wiesner, T., Zbyszewski, A. (2018). Progress in the diagnosis and therapy of melanoma. *Br J Dermatol*, 178, 12-14.
7. Peng, Y., Sartini, D., Pozzi, V., Wilk, D., Emanuelli, M., Yee VC. (2011). Structural basis of substrate recognition in human nicotinamide N-methyltransferase. *Biochemistry*, 50, 7800-7808.
8. Ganzetti, G., Sartini, D., Campanati, A., Rubini, C., Molinelli, E., Brisigotti, V., Cecati, M., Pozzi, V., Campagna, R., Offidani, A., Emanuelli, M. (2018). Nicotinamide N-methyltransferase: potential involvement in cutaneous malignant melanoma. *Melanoma Res*, 28, 82-88.
9. Mascitti, M., Santarelli, A., Sartini, D., Rubini, C., Colella, G., Salvolini, E., Ganzetti, G., Offidani, A., Emanuelli, M. (2019). Analysis of nicotinamide N-methyltransferase in oral malignant melanoma and potential prognostic significance. *Melanoma Res*, 29, 151-156.
10. Pozzi, V., Mazzotta, M., Lo Muzio, L., Sartini, D., Santarelli, A., Renzi, E., Rocchetti, R., Tomasetti, M., Ciavarella, D., Emanuelli, M. (2011). Inhibiting proliferation in KB cancer cells by RNA interference-mediated knockdown of nicotinamide N-methyltransferase expression. *Int J Immunopathol Pharmacol*, 24, 69-77.

11. Pozzi, V., Sartini, D., Morganti, S., Giuliante, R., Di Ruscio, G., Santarelli, A., Rocchetti, R., Rubini, C., Tomasetti, M., Giannatempo, G., Orlando, F., Provinciali, M., Lo Muzio, L., Emanuelli, M. (2013). RNA-mediated gene silencing of nicotinamide N-methyltransferase is associated with decreased tumorigenicity in human oral carcinoma cells. *PLoS One*, 8, e71272.
12. Sartini, D., Seta, R., Pozzi, V., Morganti, S., Rubini, C., Zizzi, A., Tomasetti, M., Santarelli, L., Emanuelli, M. (2015). Role of nicotinamide N-methyltransferase in non-small cell lung cancer: in vitro effect of shRNA-mediated gene silencing on tumourigenicity. *Biol Chem*, 396, 225-234.
13. Seta, R., Mascitti, M., Campagna, R., Sartini, D., Fumarola, S., Santarelli, A., Giuliani, M., Cecati, M., Muzio, L.L., Emanuelli, M. (2019). Overexpression of nicotinamide N-methyltransferase in HSC-2 OSCC cell line: effect on apoptosis and cell proliferation. *Clin Oral Investig*, 23, 829-838.
14. Coit, D.G., Thompson, J.A., Algazi, A., Andtbacka, R., Bichakjian, C.K., Carson, W.E., Daniels, G.A., DiMaio, D., Ernstoff, M., Fields, R.C., et al. (2016). Melanoma, Version 2.2016, NCCN Clinical Practice Guidelines in Oncology. *J. Natl. Compr. Cancer Netw*. JNCCN, 14, 450–473.
15. Siegel, R.L., Miller, K.D., Jemal, A. (2015). Cancer statistics, 2015. *CA. Cancer J. Clin*, 65, 5–29.
16. Wilson, M.A., Schuchter, L.M. (2016). Chemotherapy for Melanoma. *Cancer Treat Res*, 167, 209-229.
17. Crosby, T., Fish R., Coles, B., Mason, M. D. (2000). Systemic treatments for metastatic cutaneous melanoma. *Cochrane Database Syst Rev* 2:CD001215.
18. Lui, P., Cashin, R., Machado, M., Hemels, M., Corey-Lisle, P.K., Einarson, T.R. (2007). Treatments for metastatic melanoma: Synthesis of evidence from randomized trials. *Cancer Treat Rev*, 33, 665–680.
19. Chapman, P.B., Hauschild, A., Robert, C., Haanen, J.B., Ascierto, P., Larkin, J., Dummer, R., Garbe, C., Testori, A., Maio, M., Hogg, D., Lorigan, P., Lebbe, C., Jouary, T., Schadendorf, D., Ribas, A., O'Day, S.J., Sosman, J.A., Kirkwood, J.M., Eggermont, A.M., Dreno, B., Nolop, K., Li, J., Nelson, B., Hou, J., Lee, R.J., Flaherty, K.T., McArthur, G.A., BRIM-3 Study Group. (2011). Improved survival with vemurafenib in melanoma with BRAF V600E mutation. *N Engl J Med*, 364, 2507–2516.
20. Eggermont, A.M., Kirkwood, J.M. (2004) Re-evaluating the role of dacarbazine in metastatic melanoma: what have we learned in 30 years? *Eur J Cancer*, 40, 1825–1836.
21. Sartini, D., Muzzonigro, G., Milanese, G., Pierella, F., Rossi, V., Emanuelli, M. (2006). Identification of nicotinamide N-methyltransferase as a novel tumor marker for renal clear cell carcinoma. *J Urol*, 176, 2248-2254.
22. Campagna, R., Cecati, M., Pozzi, V., Fumarola, S., Pompei, V., Milanese, G., Galosi, A.B., Sartini, D., Emanuelli, M. (2018). Involvement of transforming growth factor beta 1 in the transcriptional regulation of nicotinamide N-methyltransferase in clear cell renal cell carcinoma. *Cell Mol Biol (Noisy-le-grand)*, 64, 51-55.

- 1
2
3 23. Sartini, D., Santarelli, A., Rossi, V., Goteri, G., Rubini, C., Ciavarella, D., Lo Muzio, L., Emanuelli,
4 M. (2007). Nicotinamide N-methyltransferase upregulation inversely correlates with lymph node
5 metastasis in oral squamous cell carcinoma. *Mol Med*, 13, 415-421.
6
7
- 8 24. Emanuelli, M., Santarelli, A., Sartini, D., Ciavarella, D., Rossi, V., Rubini, C., Lo Muzio, L. (2010).
9 Nicotinamide N-Methyltransferase upregulation correlates with tumour differentiation in oral
10 squamous cell carcinoma. *Histol Histopathol*, 25, 15-20.
11
- 12 25. Sartini, D., Pozzi, V., Renzi, E., Morganti, S., Rocchetti, R., Rubini, C., Santarelli, A., Lo Muzio, L.,
13 Emanuelli, M. (2012). Analysis of tissue and salivary nicotinamide N-methyltransferase in oral
14 squamous cell carcinoma: basis for the development of a noninvasive diagnostic test for early-stage
15 disease. *Biol Chem*, 393, 505-511.
16
17
- 18 26. Sartini, D., Morganti, S., Guidi, E., Rubini, C., Zizzi, A., Giuliante, R., Pozzi, V., Emanuelli, M.
19 (2013). Nicotinamide N-methyltransferase in non-small cell lung cancer: promising results for targeted
20 anti-cancer therapy. *Cell Biochem Biophys*, 67, 865-873.
21
22
- 23 27. Sartini, D., Muzzonigro, G., Milanese, G., Pozzi, V., Vici, A., Morganti, S., Rossi, V., Mazzucchelli,
24 R., Montironi, R., Emanuelli, M. (2013). Upregulation of tissue and urinary nicotinamide N-
25 methyltransferase in bladder cancer: potential for the development of a urine-based diagnostic test.
26 *Cell Biochem Biophys*, 65, 473-483.
27
28
- 29 28. Pozzi, V., Di Ruscio, G., Sartini, D., Campagna, R., Seta, R., Fulvi, P., Vici, A., Milanese, G.,
30 Brandoni, G., Galosi, A.B., Montironi, R., Cecati, M., Emanuelli, M. (2018). Clinical performance
31 and utility of a NNMT-based urine test for bladder cancer. *Int J Biol Markers*, 33, 94-101.
32
33
- 34 29. Pompei, V., Salvolini, E., Rubini, C., Lucarini, G., Molinelli, E., Brisigotti, V., Pozzi, V., Sartini, D.,
35 Campanati, A., Offidani, A., Emanuelli, M. (2019). Nicotinamide N-methyltransferase in
36 nonmelanoma skin cancers. *Eur J Clin Invest*, 49, e13175.
37
38
- 39 30. Pozzi, V., Sartini, D., Rocchetti, R., Santarelli, A., Rubini, C., Morganti, S., Giuliante, R., Calabrese,
40 S., Di Ruscio, G., Orlando, F., Provinciali, M., Saccucci, F., Lo Muzio, L., Emanuelli, M. (2015).
41 Identification and characterization of cancer stem cells from head and neck squamous cell carcinoma
42 cell lines. *Cell Physiol Biochem*, 36, 784-798.
43
44
- 45 31. Pozzi, V., Salvolini, E., Lucarini, G., Salvucci, A., Campagna, R., Rubini, C., Sartini, D., Emanuelli,
46 M. (2020). Cancer stem cell enrichment is associated with enhancement of nicotinamide N-
47 methyltransferase expression. *IUBMB Life*. 72, 1415-1425.
48
49
- 50 32. Ulanovskaya OA, Zuhl AM, Cravatt BF. (2013) NNMT promotes epigenetic remodeling in cancer by
51 creating a metabolic methylation sink. *Nat Chem Biol*, 9:300-306.
52
53
- 54 33. Pissios P. (2017) Nicotinamide N-Methyltransferase: More Than a Vitamin B3 Clearance Enzyme.
55 *Trends Endocrinol Metab*, 28, 340-353.
56
- 57 34. Mateuszuk Ł, Campagna R, Kutryb-Zajac B, Kuś K, Słominska EM, Smolenski RT, Chlopicki S.
58 (2020) Reversal of endothelial dysfunction by nicotinamide mononucleotide via extracellular
59 conversion to nicotinamide riboside. *Biochem Pharmacol*, 178:114019.
60

- 1
2
3 35. Sartini, D., Seta, R., Pozzi, V., Morganti, S., Rubini, C., Zizzi, A., Tomasetti, M., Santarelli, L.,
4 Emanuelli, M. (2015). Role of nico-tinamide N-methyltransferase in non-small cell lung cancer: in
5 vitro effect of shRNA-mediated gene silencing on tumour-igenicity. *Biol Chem*, 396, 225-234.
6
7
8 36. Wu, Y., Siadaty, M.S., Berens, M.E., Hampton, G.M., Theodorescu, D. (2008). Overlapping gene
9 expression profiles of cell migration and tumor invasion in human bladder cancer identify
10 metallothionein 1E and nicotinamide N-methyltransferase as novel regulators of cell migration.
11 *Oncogene*, 27, 6679-6689.
12
13
14 37. Tang, S.W., Yang, T.C., Lin, W.C., Chang, W.H., Wang, C.C., Lai M.K., Lin, J.Y. (2011).
15 Nicotinamide N-methyltransferase induces cellular invasion through activating matrix
16 metalloproteinase-2 expression in clear cell renal cell carcinoma cells. *Carcinogenesis*, 32, 138-145.
17
18
19 38. Zhang, J., Wang, Y., Li, G., Yu, H., Xie, X. (2014). Down-regulation of nicotinamide N-
20 methyltransferase induces apoptosis in hu-man breast cancer cells via the mitochondria-mediated
21 pathway. *PLoS One*, 9, e89202.
22
23
24 39. Xie, X., Yu, H., Wang, Y., Zhou, Y., Li, G., Ruan, Z., Li, F., Wang, X., Liu, H., Zhang, J. (2014).
25 Nicotinamide N-methyltransferase enhances the capacity of tumorigenesis associated with the
26 promotion of cell cycle progression in human colorectal cancer cells. *Arch Biochem Biophys*, 564, 52-
27 66.
28
29
30 40. Yu, T., Wang, Y.T., Chen, P., Li, Y.H., Chen, Y.X., Zeng, H., Yu, A.M., Huang, M., Bi, H.C. (2015).
31 Effects of nicotinamide N-methyltransferase on PANC-1 cells proliferation, metastatic potential and
32 survival under metabolic stress. *Cell Physiol Biochem*, 35, 710-721.
33
34
35 41. Bach, D.H., Kim, D., Bae, S.Y., Kim, W.K., Hong, J.Y., Lee, H.J., Rajasekaran, N., Kwon, S., Fan,
36 Y., Luu, T.T., Shin, Y.K., Lee, J., Lee, S.K. (2018). Targeting Nicotinamide N-Methyltransferase and
37 miR-449a in EGFR-TKI-Resistant Non-Small-Cell Lung Cancer Cells. *Mol Ther Nucleic Acids*, 11,
38 455-467.
39
40
41 42. You, Z., Liu, Y., Liu, X. (2018). Nicotinamide N-methyltransferase enhances the progression of
42 prostate cancer by stabilizing sirtuin 1. *Oncol Lett*, 15, 9195-9201.
43
44
45 43. Liang, L., Zeng, M., Pan, H., Liu, H., He, Y. (2018). Nicotinamide N-methyltransferase promotes
46 epithelial-mesenchymal transition in gastric cancer cells by activating transforming growth factor- β 1
47 expression. *Oncol Lett*, 15, 4592-4598.
48
49
50 44. Cui, Y., Zhang, L., Wang, W., Ma, S., Liu, H., Zang, X., Zhang, Y., Guan, F. (2019). Downregulation
51 of nicotinamide N-methyltransferase inhibits migration and epithelial-mesenchymal transition of
52 esophageal squamous cell carcinoma via Wnt/ β -catenin pathway. *Mol Cell Biochem*, 460, 93-103.
53
54
55 45. Xie, X., Liu, H., Wang, Y., Zhou, Y., Yu, H., Li, G., Ruan, Z., Li, F., Wang, X., Zhang, J. (2016).
56 Nicotinamide N-methyltransferase enhances resistance to 5-fluorouracil in colorectal cancer cells
57 through inhibition of the ASK1-p38 MAPK pathway. *Oncotarget*, 7, 45837-45848.
58
59
60 46. Cui, Y., Yang, D., Wang, W., Zhang, L., Liu, H., Ma, S., Guo, W., Yao, M., Zhang, K., Li, W., Zhang,
Y., Guan, F. (2020). Nicotina-mide N-methyltransferase Decreases 5-fluorouracil Sensitivity in

Human Esophageal Squamous Cell Carcinoma Through Metabolic Reprogramming and Promoting the Warburg Effect. *Mol Carcinog*, 59, 940-954.

47. Wang, Y., Zeng, J., Wu, W., Xie, S., Yu, H., Li, G., Zhu, T., Li, F., Lu, J., Wang, G.Y., Xie, X., Zhang, J. (2019). Nicotinamide N-methyltransferase enhances chemoresistance in breast cancer through SIRT1 protein stabilization. *Breast Cancer Res*, 21, 64.
48. van Haren, M.J., Taig, R., Kuppens, J., Sastre Toraño, J., Moret, E.E., Parsons, R.B., Sartini, D., Emanuelli, M., Martin, N.I. (2017). Inhibitors of nicotinamide N-methyltransferase designed to mimic the methylation reaction transition state. *Org Biomol Chem*. 15, 6656-6667.

FIGURE LEGEND

Figure 1. Evaluation of NNMT silencing. A375 and WM-115 cells were treated with different shRNA plasmids against NNMT (pLKO.1-164, pLKO.1-330, pLKO.1-711 and pLKO.1-mix), with empty vector (pLKO.1-puro) or with transfection reagent only (Mock). NNMT expression was evaluated at mRNA (panel a for A375 cells; panel d for WM-115 cells) and protein level (for A375 cells, blot is reported in panel b and densitometry is reported in panel c) by Real Time PCR and Western blot, respectively. Values are expressed as mean \pm standard deviation (* $p < 0.05$).

Figure 2. In vitro effect of NNMT silencing on cell proliferation. A375 (panel a) and WM-115 (panel b) cell proliferation was analyzed by MTT assay in Mock and NNMT downregulating cells (pLKO.1-164, pLKO.1-330, pLKO.1-711 and pLKO.1-mix) at 0h, 24h, 48h and 72h. Values are expressed as mean \pm standard deviation (* $p < 0.05$; ns = not significant).

Figure 3. In vitro effect of NNMT silencing on cell migration. A375 cells were subjected to wound healing assay to evaluate their migration potential. A375 cells transfected with plasmids (pLKO.1-164, pLKO.1-330, pLKO.1-711, pLKO.1-mix or pLKO.1-puro) and Mock were photographed immediately after scratch (0h) and at different time points, ranging between 4h and 28h. For each sample, cell migration ability was evaluated at different time points by measuring percentage of wound recovery compared with 0h. Panel a displays images of wound recovery of Mock, pLKO.1-puro, pLKO.1-164, pLKO.1-330, pLKO.1-711 and pLKO.1-mix. Panel b reports bar diagram showing time course of wound recovery of all samples. Values are expressed as mean \pm standard deviation (* $p < 0.05$).

Figure 4. Effect of chemotherapy treatment on melanoma cell proliferation. MTT assay was used to evaluate the effect of dacarbazine on proliferation of mock and NNMT downregulating cells (panels a-d for A375; panel e for WM-115). Measurements were performed at different time points (0h, 24h, 48h and 72h). All values are expressed as mean \pm standard deviation (* $p < 0.05$).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review




Figure 1. Evaluation of NNMT silencing. A375 and WM-115 cells were treated with different shRNA plasmids against NNMT (pLKO.1-164, pLKO.1-330, pLKO.1-711 and pLKO.1-mix), with empty vector (pLKO.1-puro) or with transfection reagent only (Mock). NNMT expression was evaluated at mRNA (panel a for A375 cells; panel d for WM-115 cells) and protein level (for A375 cells, blot is reported in panel b and densitometry is reported in panel c) by Real Time PCR and Western blot, respectively. Values are expressed as mean \pm standard deviation (* $p < 0.05$).

30x54mm (300 x 300 DPI)




Figure 2. In vitro effect of NNMT silencing on cell proliferation. A375 (panel a) and WM-115 (panel b) cell proliferation was analyzed by MTT assay in Mock and NNMT downregulating cells (pLKO.1-164, pLKO.1-330, pLKO.1-711 and pLKO.1-mix) at 0h, 24h, 48h and 72h. Values are expressed as mean \pm standard deviation (* $p < 0.05$; ns = not significant).




Figure 3. In vitro effect of NNMT silencing on cell migration. A375 cells were subjected to wound healing assay to evaluate their migration potential. A375 cells transfected with plasmids (pLKO.1-164, pLKO.1-330, pLKO.1-711, pLKO.1-mix or pLKO.1-puro) and Mock were photographed immediately after scratch (0h) and at different time points, ranging between 4h and 28h. For each sample, cell migration ability was evaluated at different time points by measuring percentage of wound recovery compared with 0h. Panel a displays images of wound recovery of Mock, pLKO.1-puro, pLKO.1-164, pLKO.1-330, pLKO.1-711 and pLKO.1-mix. Panel b reports bar diagram showing time course of wound recovery of all samples. Values are expressed as mean \pm standard deviation (* $p < 0.05$).




Figure 4. Effect of chemotherapy treatment on melanoma cell proliferation. MTT assay was used to evaluate the effect of dacarbazine on proliferation of mock and NNMT downregulating cells (panels a-d for A375; panel e for WM-115). Measurements were performed at different time points (0h, 24h, 48h and 72h). All values are expressed as mean \pm standard deviation ($*p < 0.05$).