



UNIVERSITÀ POLITECNICA DELLE MARCHE  
Repository ISTITUZIONALE

Post-digestate composting benefits and the role of enzyme activity to predict trace element immobilization and compost maturity

This is the peer reviewed version of the following article:

*Original*

Post-digestate composting benefits and the role of enzyme activity to predict trace element immobilization and compost maturity / Biyensa, Gurmessa; Cocco, S.; Ashworth, J. A.; Foppa Pedretti, E.; Ilari, A.; Cardelli, V.; Fornasier, F.; Ruello, Maria Letizia; Corti, G.. - In: BIORESOURCE TECHNOLOGY. - ISSN 0960-8524. - ELETTRONICO. - 338:(2021). [10.1016/j.biortech.2021.125550]

*Availability:*

This version is available at: 11566/295363 since: 2024-10-24T15:42:46Z

*Publisher:*

*Published*

DOI:10.1016/j.biortech.2021.125550

*Terms of use:*

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. The use of copyrighted works requires the consent of the rights' holder (author or publisher). Works made available under a Creative Commons license or a Publisher's custom-made license can be used according to the terms and conditions contained therein. See editor's website for further information and terms and conditions.

This item was downloaded from IRIS Università Politecnica delle Marche (<https://iris.univpm.it>). When citing, please refer to the published version.

(Article begins on next page)

# Bioresource Technology

## Post-digestate composting benefits and the role of enzyme activity to predict trace element immobilization and compost maturity

--Manuscript Draft--

|                              |  |
|------------------------------|--|
| <b>Manuscript Number:</b>    | BITE-D-21-03871R1  |
| <b>Article Type:</b>         | Original research paper  |
| <b>Keywords:</b>             | C/N ratio; compost maturity, food processing waste, maize silage, poultry litter   |
| <b>Corresponding Author:</b> | Biyensa Gurmessa<br>Marche Polytechnic University<br>ITALY   |
| <b>First Author:</b>         | Biyensa Gurmessa   |
| <b>Order of Authors:</b>     | Biyensa Gurmessa<br>Stefania Cocco<br>Amanda J. Ashworth<br>Ester Foppa Pedretti<br>Alessio Ilari<br>Valeria Cardelli<br>Flavio Fornasier<br>Maria Letizia Ruello<br>Giuseppe Corti  |
| <b>Abstract:</b>             | <p>The current study evaluated the quality of agricultural waste digestate by composting or co-composting with biogas feedstock (maize silage, food processing waste, or poultry litter). Temperature, phytotoxicity, C/N ratio, water extractable trace elements and 14 enzyme activities were monitored for 90 days. Temperature dropped earlier in digestate and maize silage co-composting pile, reducing time to maturity by 20 days. Composting and co-composting reduced phytotoxicity and C/N ratio, but increased immobilization of Al, Ba, Fe, Zn, and Mn at least by 40% in all piles. All the enzyme activities, except arylsulfatase and <math>\alpha</math>-glucosidase, were increased at the maturity phase and negatively correlated with organic matter content and most of trace elements. Post-digestate composting or co-composting with biogas feedstock is a promising strategy to improve digestate quality for fertilizer use, and selected enzyme activities can be indicators of compost maturity and immobilization of trace elements.</p> |

07/07/2021

RE: Submission of revised paper

Dear Prof. HUU HAO NGO,

Thanks for inviting us to resubmit our manuscript after revision. We have carefully considered reviewers' comments and made changes to the manuscript, and we believe the current version of the manuscript is significantly improved. We have also modified the title a little bit. Revisions are made following points raised by reviewers, and response is provided under each inquiry. We used red font to indicate the changes we made in the manuscript.

Best regards,

Biyensa

Dear Prof. HUU HAO NGO,

Thanks for inviting us to resubmit our manuscript after revision. We have carefully considered reviewers' comments and made changes to the manuscript, and we believe the current version of the manuscript is significantly improved. We have also modified the title a little bit. Revisions are made following points raised by reviewers, and response is provided under each inquiry. We used red font to indicate the changes we made in the manuscript. Our responses to reviewers are as follows:

Reviewer #1: This paper studied the quality of post-digestate compost quality and dynamics of enzyme activities on predicting compost maturity and trace elements immobilization. The experiments were carefully carried out, and good results were obtained. This research is of great significance to the utilization of solid digestate. Therefore, the paper is acceptable for publication in Bioresource Technology. However, before acceptance, please consider the following comments and suggestions.

Response: We are grateful to the reviewer for the revisions and suggestions for the manuscript. We have made significant changes to the manuscript following the points of improvements requested by this Reviewer.

1. The contents of manuscript should be checked again, especially the grammar and punctuation.

Response: Thanks. This paper has been thoroughly checked throughout the manuscript and corrections were made where necessary.

2. Awkward sentence make reader confused.

Response: Since we have made major changes to the manuscript, sentences were seemingly confusing have been changed. However, if reviewer specifies which statement needs to be corrected, we will address it in the next round of review.

3. A clear and concise graphical abstract is needed to show readers the main information of the research work.

Response: Thanks. A graphical abstract is now included to this revised manuscript.

4. The format of reference should be carefully checked again.

Response: References have been thoroughly checked and corrections are made where necessary, according to the guidelines.

Reviewer #2: The manuscript number BITE-D-21-03871 entitled "Enzyme activities predict trace elements immobilization and compost maturity during agricultural waste digestate composting" by Gurmessa et al looks at impact of co-composting on post-digestate quality. The study also probes into some fourteen enzyme activities. The presented work addresses one of the important problems of repurposing anaerobic digestion waste (digestate) which is a need of an hour. However, in my opinion, the presented manuscript does NOT seem fit for publication because of the following reasons:

Response: We are thankful to the reviewer for the complements and review suggestions. We admit there were mess up in sections order and apologize for the inconvenience. We have carefully addressed the suggestions by the reviewer and the manuscript now complies with BITE author guidelines . Authors hope that the amended manuscript it deemed more accessible.

1. The manuscript does NOT seem to have followed BITE author guidelines:

Response: Thanks! We have reviewed and made changes to manuscript to comply with the BITE author guidelines.

2. The structure of the manuscript seems a little unusual with first section being "Material and Methods".

Response: Right! We have re-organized the paper accordingly.

3. The sub-section numbering seems totally messed up

Response: This is also corrected in the current version of the MS; thank you for your attention to this.

4. Abstract is poorly written:

Response: Thanks. Abstract is re-written in the current version with significant improvements.

5. Phrases like "The current study set out to.." seem a little unusual from academic writing pointing of view

Response: This is changed to "This study evaluated..." in the revised MS.

6. Abstract demands at least a cursorily glimpse at the methodology adapted - which seems lacking here.

Response: Thanks! We have included the following into the abstract to briefly show the method:

"The current study evaluated the quality of agricultural waste digestate by composting or co-composting with biogas feedstock (maize silage, food processing waste, or poultry litter). Temperature, phytotoxicity, C/N ratio, water extractable trace elements and 14 enzyme activities were monitored for 90 days."

7. There is no closing sentence(s) on future directions, likely impact etc

Response: Thanks! We have now included concluding remark at the end of the abstract:

“Post-digestate composting or co-composting with biogas feedstock is a promising strategy to improve digestate quality for fertilizer use, and selected enzyme activities can be indicators of compost maturity and immobilization of trace elements.”

8. Conclusion is also poorly written with NO concrete/quantifiable information.

Response: Conclusion is now rewritten, and significant changes are made. Some quantifiable information is included. In addition, number of words is reduced to less than 100.

9. Too much information compressed in figures 1-5 making data comprehension challenging. Authors might consider splitting them in multiple sub-figures and present only relevant information upfront. For instance, in my opinion, Fig. 5-a could better suit in the supplementary rather than mainframe. This might also help readers grasp information in Fig. 5-b and c effectively.

Response: Thanks. As suggested, authors moved Fig.5 A to supplementary files. Accordingly, Fig 5 now has only two sub-figures. We understand the concern of the reviewer, however, keeping all Fig.4 together is important for reader to understand relationships or differences between activities of the different enzymes. However, if it still is of interest to separate, we can make this change in the next round of revision.

Reviewer # 6: Please carefully check if the paper meets the following points. **NON-COMPLIANCE OF THESE POINTS WILL CAUSE A SIGNIFICANT DELAY OF THE PUBLICATION AND MIGHT LEAD REJECTION OF THE MANUSCRIPT.**

Response: Thanks. We have checked all the points and made corrections (e.g., conclusions part) where and the paper meets the requirements. We have made changes to conclusions part which previously did not meet the requirement.

- No E-supplement figures and tables (e.g. Fig S1..Table S1,.etc.) quoted in text directly;
  - Checked. There are no E-supplement figures quoted as Fig S1, Table S1, etc.
- Maximum five (including tables/figures/sub-figures) in Supplementary Material;
  - Checked. Only one supplementary material is provided.
- Only ONE PERSON as corresponding author;
  - Only one person is provided as corresponding author for the paper.
- No abbreviations in the title;
  - There is no abbreviation in the title.
- Maximum 150 words in Abstract;
  - Abstract is limited to 143 words.
- Maximum 100 words in Conclusion;
  - This is limited to 100 words.

- Combined results and discussion;
  - This is in accordance.
- Maximum 85 characters, including space for each item of highlights;
  - Checked. Every highlight does not exceed 85 characters including space
- No list of abbreviations or Nomenclatures, except huge number of abbreviations, i.e. in kinetic and modelling works;
  - Checked. No list of abbreviations provided.
- No conventional spectra (X-ray, FTIR, UV, NMR, etc.), SEM photographs, one column simple data table, biochemical (having chemical structures) pathways, simple one-line drawing;
  - Checked. No item of these types is provided in the manuscript.
- No usage of first person (we, our, us);
  - Checked. No usage of such pronouns is found in the manuscript.
- No full justification (e.g. no usage of a constant right-hand margin);
  - Checked. Full justification avoided.
- Maximum 6 Tables and 6 Figures in Research paper,
  - The paper conforms this limitation: has only 5 figures and 3 tables.
- Numbered references in the list of References (to check if followed the limit set by BITE, i.e., Short Communication – Max. 25; Research Article – Max. 50; Review Article- Max. 150);
  - Checked. Number of references is 43.
- No usage of non-English references;
  - Checked. Non-English reference is not used.
- No usage of the Track Changes feature in Microsoft Word in the revised version
  - Checked. Track changes are avoided.

### Highlights

- Pilot level aerobic co-composting of digestate and biogas feedstocks was conducted.
- Composting reduced C/N ratio, release of trace elements, and phytotoxicity.
- Enzyme activities were influenced by OM content and release of trace elements.
- Enzyme activities could be compost maturity indicators.



1 **Post-digestate composting benefits and the role of enzyme activity to**  
2 **predict trace element immobilization and compost maturity**

3  
4 Biyensa Gurmessa<sup>1\*</sup>, Stefania Cocco<sup>1</sup>, Amanda J. Ashworth<sup>2</sup>, Ester Foppa Pedretti<sup>1</sup>, Alessio Ilari<sup>1</sup>,  
5 Valeria Cardelli<sup>1</sup>, Flavio Fornasier<sup>3</sup>, Maria Letizia Ruello<sup>4</sup>, Giuseppe Corti<sup>1</sup>

6  
7 <sup>1</sup>Department of Agricultural, Food and Environmental Sciences, Università Politecnica delle  
8 Marche, 60131 Ancona, Italy

9 <sup>2</sup>USDA-ARS, Poultry Production and Product Safety Research Unit, 1260 W. Maple St.  
10 Fayetteville, AR 72701, USA

11 <sup>3</sup>Consiglio per la Ricerca e la Sperimentazione in Agricoltura, Centro di Ricerca per lo Studio delle  
12 Relazioni tra Pianta e Suolo (C.R.E.A.-R.P.S.), 34170 Gorizia, Italy

13 <sup>4</sup>Department of Materials, Environmental Sciences and Urban Planning, Università Politecnica  
14 delle Marche, 60131 Ancona, Italy

15

16

17 \*Corresponding author: [biyensa@gmail.com](mailto:biyensa@gmail.com)

18

19

20

1 Abstract

2 The current study evaluated the quality of agricultural waste digestate by composting or co-  
3 composting with biogas feedstock (maize silage, food processing waste, or poultry litter).  
4 Temperature, phytotoxicity, C/N ratio, water extractable trace elements and 14 enzyme activities  
5 were monitored for 90 days. Temperature dropped earlier in digestate and maize silage co-  
6 composting pile, reducing time to maturity by 20 days. Composting and co-composting reduced  
7 phytotoxicity and C/N ratio, but increased immobilization of Al, Ba, Fe, Zn, and Mn at least by  
8 40% in all piles. All the enzyme activities, except arylsulfatase and  $\alpha$ -glucosidase, were  
9 increased at the maturity phase and negatively correlated with organic matter content and most of  
10 trace elements. Post-digestate composting or co-composting with biogas feedstock is a promising  
11 strategy to improve digestate quality for fertilizer use, and selected enzyme activities can be  
12 indicators of compost maturity and immobilization of trace elements.

13  
14 Keywords: C/N ratio; compost maturity, food processing waste, maize silage, poultry litter

15  
16  
17  
18  
19  
20  
21  
22  
23

# 1. Introduction

In Europe, solid digestate, a byproduct of biogas production, is often directly used as fertilizer although legal status differs among member States. Some have policies that encourage direct use as fertilizer (Beggio et al., 2019; Tambone et al., 2015), while others do not consider it a fertilizer because of possible associated environmental risks such as ammonia emissions, odor, and high content of volatile fatty acids (Nkoa, 2014; PiotrZeng et al., 2016). Low nutrient supply and load of trace elements are other disadvantages associated with direct usage of digestate as fertilizer (Kupper et al., 2014; Torres-Climent et al., 2015). Digestates, especially those originating from manure or a mixture of manure and food processing wastes, can also be source of weeds, pathogens, pharmaceutical residues, and antibiotic resistant genes (Gurmessa et al., 2020).

Trace element loading with digestate at the farm-level could be high owing to continuous land applications as fertilizer (Kupper et al., 2014), thereby posing environmental risks. To avoid these and improve the quality of digestate for its use as an amendment, post-digestate treatment has been proposed (Bustamante et al., 2012; Rehl and Müller, 2011). It can be one possible treatment that has advantages in terms of increasing nutrient content, attenuating trace elements release, reducing the volume of biomass, and mitigating overall environmental risks. For instance, Karwal and Kaushik (2020) found significant reduction of Cd, Cu, Pb, and Zn contents with composting.

Thus, effective post-digestate composting may be sought for industrial level composting, which could involve the use of an appropriate co-composting material that is inexpensive and locally available. Moreover, it is essential to understand the benefits of composting digestate in terms of nutrient release, immobilization of trace elements, and phytotoxicity reduction. In

1 addition to the commonly known compost monitoring indicators such as temperature, C/N ratio,  
2 pH, total solids content, total heavy metals content, and germination index (Tang et al., 2020),  
3 there is lack of knowledge on the dynamics of enzyme activities during post-digestate  
4 composting, which could be indicators of compost functioning, fate of trace elements, and  
5 maturity (a ready to use stable compost). Enzyme activities during composting were reported to  
6 be one of the best indicators of compost quality and maturity when the activities tend to increase  
7 (Wan et al., 2020) or become stable (Mondini et al., 2004) during the maturity phase. However,  
8 findings were not consistent, and variations can be due to the differences in the sources of  
9 composting materials, the composting period, and enzyme activities.

10 Thus, the current study is aimed at understanding the role of post-digestate composting and  
11 **co-composting with biogas feedstock** on the final compost quality in terms of nutrient release,  
12 immobilization of trace elements, and phytotoxicity. The study is also aimed at investigating: *i*)  
13 the role of co-composting materials on post-digestate compost quality and maturity, *ii*) the trends  
14 of enzyme activities during composting and their use as proxy to define maturity of post-  
15 digestate compost, and *iii*) the relationship between enzyme activities and soluble nutrients and  
16 trace elements content during composting.

## 17 2. Materials and Methods

### 18 2.1. Composting, monitoring, and sampling

19 The digestate and co-composting materials were obtained from an industrial biogas plant in  
20 the Marche region, Italy. Digestate used is a byproduct of anaerobic digestion from a biogas  
21 plant that uses 10% poultry litter (85:15 chicken manure: wheat straw ratio; moisture  $\approx$ 58%) and  
22 90% of other biomass sources such as maize silage and food processing wastes (**byproducts of**  
23 **cereal mill and fruit**).

1 Five composting piles, 300 kg each (on wet basis) were prepared as follows: 1) solid digestate  
2 only (D00); 2) solid digestate + food processing waste (DCB); 3) solid digestate + maize silage  
3 (DMS); 4. solid digestate + poultry litter (DPL); 5) solid digestate + maize silage + poultry litter  
4 (DMP). Solid digestate was the target composting material and constituted 80% of the total  
5 biomass (w/w) of the piles; the other 20% was made by the co-composting material. The mix-  
6 ratio is described in Table 1.

7 The composting materials were mixed thoroughly and stacked inside high-density  
8 polyethylene boxes, each having a volume of 1 m<sup>3</sup>. To allow air movement, four holes of 3 cm  
9 diameter were drilled at 5 cm and 50 cm from the base on the corners of the box: two from one  
10 side and two on the opposite side. Two hard plastic tubes of 2 m length, each with holes every 20  
11 cm, were inserted diagonally from the bottom hole of one corner to the upper hole of the other  
12 corner of the box. Piles were turned weekly during the thermophilic phase, every two weeks  
13 during the mesophilic period, and every three weeks during the maturity period, following  
14 standard industry procedures. Composting lasted for 90 days, and temperature was monitored  
15 daily from 4 to 5 o'clock p.m. using a temperature probe (A.M. Leonard Backyard Compost  
16 Thermometer) 30 cm depths in each pile; temperature monitoring stopped after 50 days of  
17 composting, when similar readings were observed across piles.

18 Three 1 kg samples were collected from every pile at ≈20 cm depths at 0, 7, 14, 21, 35, 49,  
19 70, and 90 days of composting. The samples were dried at 40 °C and ground until it passed all  
20 through a 2 mm sieve. For each fragmented sample, an aliquot of about 100 g was stored at 4 °C  
21 for < 10 days to be analyzed for enzyme activities, while the rest of the sample was stored at  
22 room temperature for the physicochemical analyses.

## 2.2. Main physicochemical characteristics

The pH was determined potentiometrically in H<sub>2</sub>O (1:8 w/v) after one night of contact time. Total solid (TS) content was estimated as the fraction of dry mass remaining after samples were dried at 105 °C for 24 h. The organic matter (OM) content was determined as the loss on ignition at 550 °C until a constant weight was reached (Heiri et al., 2001). Total C and N were determined by the dry combustion method using a CHNS analyzer (EA-1110, Carlo Erba Instruments, Milan, Italy).

## 2.3. Enzyme activities

The activities of 14 enzymes were determined according to the method described in Cardelli et al. (2019). Briefly, 150 mg of specimen was placed in a 2-ml Eppendorf tube with glass beads containing 1.2 ml of 50 mM tris-HCl solution at pH 7 containing 2% lysozyme as a desorbing protein. The tube was then subjected to bead-beating (3 min, 30 strokes s<sup>-1</sup>) using a Retsch MM400 mill (Haan, Germany) and centrifuged for 5 min at 20,000 g. Enzyme activities were analyzed fluorometrically in microplates using 4-methyl-umbelliferyl and 7-amino-4-methyl coumarine conjugated surrogate substrates (Sigma, St. Louis, MO, USA) at three pH ranges in three different solutions. Activities of acid phosphomonoesterase,  $\alpha$ -glucosidase, arylsulfatase,  $\beta$ -glucosidase,  $\beta$ -galactosidase, cellulase, chitinase, glucuronidase, and xylosidase were determined in 200 mM MES (morpholineptansulfonic acid) solution at pH 5.8; whereas, the activities of leucine aminopeptidase, lipase nonanoate-esterase, pyrophosphatase-phosphodiesterase, and phosphodiesterase were determined in 100 mM tris-HCl solution at pH 7.5. The alkaline phosphomonoesterase activity was determined in 100 mM tris-HCl solution at pH 9.0.

1        2.4. Water-extractable elements

2        Ten g of each sample was added to 100 ml of distilled water (1:10 w/v) and shaken for about  
3        1 h. Then, the suspension was centrifuged for five minutes at 300 g and the solution was filtered  
4        using a Whatman 42 filter. The amounts of macronutrients (Ca, K, Mg, P, and S) and trace  
5        elements (Al, Ba, Cd, Cu, Fe, Mn, Ni, Pb, and Zn) in the extracted solution were determined  
6        using Inductively Coupled Plasma Spectroscopy.

7

8        2.5. Phytotoxicity test

9        Germination index (GI) was used to evaluate the phytotoxicity of the original solid digestate  
10        (experimental control) and of the composting digestate at 7, 21, 51, and 90 days of composting.  
11        The experiment was conducted in a completely randomized design with three replications. One g  
12        of digestate or composting material was added to 5 ml deionized water in a 50 ml tube and  
13        shaken for 2 hours; the suspension was then centrifuged for 5 minutes at 300 g, and the solution  
14        filtered through a Whatman 42 filter. The germination test was conducted on petri dishes of 8.5  
15        cm diameter using cress (*Lepidium sativum* L.) seeds, which are highly recommended because of  
16        their high sensitive to even low toxicity (Luo et al., 2018). Two Whatman™ 1442-125 grade 42  
17        filter papers were moistened by applying 1 ml of the extract solution, whereas deionized water  
18        was used as a control. One filter paper was placed on the petri dish and 10 seeds were placed per  
19        dish. Then, the second filter paper was used to cover seeds to protect them from moisture loss.  
20        All Petri dishes were closed, sealed to avoid moisture loss, and incubated in the dark at  
21        temperature of 25 °C. Germinated seeds were counted after 48 hours of incubation, and root  
22        length was measured using a digital caliper. GI was estimated as the percentage of seeds

1 germinated multiplied by the average root length in the treated Petri dishes in relation to the  
2 number of seeds germinated multiplied by the average root length in the control Petri dishes.

3

## 4 2.6. Statistical analysis

5 Statistical analysis and visualizations were conducted using R packages. **Pearson's**  
6 **correlation analysis was conducted and plotted between enzyme activities and other chemical**  
7 **compositions using the CorLevel package in R.** Redundancy analysis (RDA) was conducted  
8 using Vegan package (Oksanen et al., 2019) in R for Windows, v. 4.0.1. and **plotted using**  
9 **ggpot2** to understand the relationship between enzyme activities (response variables) and **pH,**  
10 **OM, total C, and water extractable macronutrients and trace elements (environmental variables).**  
11 The environmental variables were log transformed, whereas response variables were Hellinger  
12 transformed. PERMANOVA test was run to evaluate the statistical significance of the interaction  
13 effects of the environmental variables on response variables. The significance of composting on  
14 release and immobilization of elements were computed as the additional amount released or the  
15 amount reduced in the final composts as percentage of the initial concentration.

16

## 17 3. Results and discussion

### 18 3.1. Physicochemical characteristics of the digestate and co-composting materials

19 In Table 2, characteristics of digestate and co-composting materials are reported. The solid  
20 digestate had sub-alkaline pH (8.1), which was similar to previously studied digestate originating  
21 from feedstock mixtures containing pig-slurry, but higher than that of cattle slurry (Albuquerque  
22 et al., 2012a). In contrast, **the** pH of poultry litter was neutral, while it was slightly acidic for  
23 maize silage and food processing waste. Digestate and maize silage had the lowest TS content



1 (38-41%), while food processing waste **was** the greatest (94%). OM content was the highest in  
2 food processing waste and maize silage, but the organic C content was similar in the four  
3 materials. Digestate mean C/N ratio was  $\approx 26$  (similar to that of maize silage), and exceeded the  
4 limit of  $<25$  in use in several countries including Italy (Tambone et al., 2015) for direct use as  
5 fertilizer. This result was greater than **that of** by Albuquerque et al. (2012b), indicating possible  
6 variations among the different sources of digestate. Digestates with C/N values greater than 20,  
7 meaning with excess of degradable organic C, could lead to immobilization of N (Teglia et al.,  
8 2011a), and may undermine the agronomic benefits of direct use of digestate. The lowest C/N  
9 ratio of the poultry litter was due to **greater** total N content. Water extractable macro-nutrients  
10 and trace elements mainly abounded in the poultry litter. Instead, digestate showed the highest  
11 content of Fe and the lowest of Mg, P (together with maize silage), and Ba.

### 12 3.2. Changes of Temperature, pH, OM, total N, C/N, and GI during composting

13 In Fig. 1, temperature, pH, OM, total C, total N, C/N ratio, and GI changes are presented.  
14 These variables govern compost dynamics over the composting period and are also indicators of  
15 maturity. Except for D00, temperatures of all other piles increased to above 60 °C in the first two  
16 weeks of composting, but dropped quickly for DMS, thus reducing time to maturity by about 20  
17 days compared to the control. The thermophilic phase lasted for about 49 days for D00, DCB,  
18 DPL, and DMP, and maturity was delayed. The pH decreased from alkaline to neutral in the  
19 control pile, while it increased from slightly acidic to neutral in DCB. The pH changes in other  
20 piles were minimal. Both total C and OM showed a decreasing trend in all the piles, but the rate  
21 was **lowest** in D00 compared to the rest of the piles, suggesting the significance of the co-  
22 composting materials. Total N increased throughout the composting period with higher rates in  
23 all the piles with co-composting materials compared to the D00 pile. The C/N ratio decreased

1 over the experimental period in all piles, reaching the value of 17 or lower in the final composts,  
2 **indicating compost** maturity.

3 GI increased in all piles over the composting period, but reached more than 60% only in  
4 DCB, DMS, and DMP, suggesting the significance of co-composting for reducing phytotoxicity  
5 of digestate. However, even after 90 days of composting, compared to the other piles, GI of D00  
6 did not reach a satisfactory level, indicating the possible occurrence of phytotoxicity, especially  
7 when sensitive crops or vegetables are grown using digestate as fertilizer. A previous study also  
8 reported possible ecotoxicity of digestate (Tigini et al., 2016), and our results suggest post-  
9 composting digestate, **and** co-composting materials **may help** mitigate phytotoxicity. A mature  
10 compost should have GI of 50% or more (Bernal et al., 2009), but it is commonly recommended  
11 to be 60% or more, which is generally believed to be indicator for a low toxicity level (Tambone  
12 et al., 2015), although there are differences among the seed types used for the test. The relatively  
13 low GI found in this study **was similar** to previous findings can be linked to the sensitivity to  
14 toxicity of the cress seeds used for the test, which is high even at low toxicity levels (Luo et al.,  
15 2018).

16

### 17 3.3. Composting effect on macronutrients release

18 The trends of water extractable values showed disparities among macronutrients during the  
19 composting period (Fig. 2). K and P showed an increasing trend in all the piles, while Mg  
20 showed an increasing trend only in D00. Instead, Ca displayed an inverse relationship in DCB  
21 and DPL, with **few** changes observed in D00. S concentration was reduced in DPL, while little  
22 changes were observed in the other piles.

1 Total N **had** increasing trends in all the piles and increased by about 30% in DPL and more  
2 than 50% in the other piles, suggesting a benefit from co-composting materials in reducing loss  
3 or increasing mineralization of N during composting compared to the control. Previous research  
4 reported the recovery of N and other macro nutrients from wastes following composting (Rai and  
5 Suthar, 2020). Water extractable (available) **was** P enriched by 25, 30, 51, and 62% in the final  
6 composts for DCB, DMP, D00, and DMS, respectively (Table 3). On the other hand, DPL had  
7 little effect on P release, despite it **having** the highest amount of available P. Previous studies  
8 reported increases in available P release with composting, although the rate varies depending on  
9 the composted materials and the composting time (Sharma et al., 2018).

10 In the current study, S content in the DPL pile was the greatest, although it gradually  
11 decreased over the composting period, suggesting the main source of S was poultry litter. The  
12 decreased in S content in all the piles over the composting period may be linked to the loss in the  
13 form of H<sub>2</sub>S (Blazy et al., 2014). S breakdown during composting (and subsequent odor release)  
14 is sought to be mitigated to reduce S loss and pollution risk.

15

#### 16 3.4. Composting effect on trace elements release

17 Water extractable contents of Ba, Cu, Fe, Mn, Ni, and Zn showed a decreasing trend in all the  
18 piles (Fig. 3), indicating immobilization with increased OM stability during composting. More  
19 than 60% of Al, Ba, Fe, Mn, and Zn was immobilized with composting across the piles (except  
20 Al for DPL), while it was >40% for Ni; Cu showed the maximum immobilization (57%) in D00.  
21 In contrast, Pb showed little change, and Cd was enriched for all treatments except DPL (Table  
22 3). Concentrations of water extractable Al, Ba, Cu, Fe, Mn, Ni, and Zn were reduced by 16-93,  
23 74-95, 25-55, 75-89, 60-74, 42-70, and 66-85%, respectively. DPL was the least effective in

1 reducing Al, Cu, and Fe release compared to the other treatments. This may be ascribed to the  
2 lower C/N ratio compared to the other piles (Wu et al., 2017). Cd level increased after the first  
3 week of composting in all the piles, except DPL, and remained constant **until** the maturity period.  
4 Pb mineralization or immobilization was little in all piles, suggesting composting digestate did  
5 not influence the release or immobilization of Pb. This could also be because of the relatively  
6 little concentration found in the co-composting and digestate materials.

7 Immobilization of trace elements is one of the benefits of composting. However, like the case  
8 of Cd in the current study, mineralization of trace metals could also be possible with post-  
9 digestate composting (Miaomiao et al., 2009). A previous study by Awasthi et al. (2020)  
10 reported that composting was effective in immobilizing Cu and Zn with biochar as a bulking  
11 agent, indicating the importance of co-composting materials. Our study also gives an insight that  
12 stabilizing digestate with post-composting potentially stabilizes the material and reduces the  
13 release of trace elements, but also suggested careful selection of co-composting materials for  
14 optimization of nutrient digestate nutrient levels.

15

### 16 3.5. Effect of composting on enzyme activities

17 Among the 14 enzyme activities evaluated, arylsulfatase and xylosidase activities were either  
18 absent or very low in all piles over the composting period (Fig. 4). In contrast, acid  
19 phosphomonoesterase, alkaline phosphomonoesterase, chitinase, and  
20 pyrophosphate/phosphodiesterase showed clear increasing trends in all piles during the 90 days  
21 of composting, with the greatest values achieved during the maturity phase, thus implying their  
22 role in depolymerizing the most complex polysaccharides of the composting materials. The  
23 greatest activity was found for leucine aminopeptidase ( $11886 \text{ nmol kg}^{-1} \text{ h}^{-1}$ ) in DMS during the

1 second week of composting, followed by lipase nonanoate-esterase ( $9379 \text{ nmol kg}^{-1} \text{ h}^{-1}$ ) and  
2 alkaline phosphomonoesterase ( $7250 \text{ nmol kg}^{-1} \text{ h}^{-1}$ ) in DCB during the maturity phase. Acid  
3 phosphomonoesterase,  $\alpha$ -glucosidase,  $\beta$ -galactosidase,  $\beta$ -glucosidase, glucuronidase, cellulase,  
4 and chitinase reached their peak values during the thermophilic phase, whereas pyrophosphate  
5 and alkaline phosphomonoesterase, reached their peak during the maturity phase. Increased  
6 levels of activities of these latter two enzymes, indicates depletion of major unstable organic  
7 matter components (Herrmann and Shann, 1993), and may be indicators for stability of  
8 composts.

9 Some of our results agreed with the findings of Karwal and Kaushik (2020), Tiquia (2002),  
10 and Herrmann and Shann (1993), who conducted composting experiments with mixtures of  
11 buffalo dung and fly ash, manure, and municipal solid waste for 90, 154, and 90 days,  
12 respectively. In contrast, Castaldi et al. (2008) found a decreasing trend for dehydrogenase,  
13 urease, protease, and cellulase at maturity phase during municipal solid waste composting that  
14 lasted for 40 days. Ge et al. (2020) reported reduced activities of both alkaline and acid  
15 phosphatase during 60 days of cattle manure composting, but the trend of cellulase activity was  
16 consistent with our findings. This shows that enzymatic activity may serve as an indicator of  
17 compost maturity and stability, but with extended length of composting period ( $\geq 90$  days) and  
18 when the most degradable organic materials are exhausted.

19 There were variations among the different co-composting materials. Interestingly, enzymatic  
20 activities in DPL changed little over the composting period, indicating the possible presence of  
21 inhibitory substances in the poultry litter. Increases in the activities of different enzymes like  
22 acid phosphomonoesterase, alkaline phosphomonoesterase,  $\beta$ -glucosidase,  $\beta$ -galactosidase,  
23 chitinase, and pyrophosphate/phosphodiesterase activities indicated maturity of composts, but

1 their functional roles are different and, thus, they can be taken **indicators** of different chemical  
2 processes taking place during the composting process. For instance, cellulase are important  
3 **indicators** for cellulose degradation, which is expected to rise during the maturity stage of  
4 composting as easily degradable organic substances have already been degraded by microbes at  
5 the early stage of composting (Li et al., 2020). Cellulase had the highest activity in D00 and  
6 maintained **levels until** the end of composting, **thus implying low availability** for microbes.

### 7 3.6. **Correlations** and RDA analysis

8 **Alkaline phosphomonoesterase, Chitinase, Glucuronidase, Leucine aminopeptidase, and**  
9 **Pyrophosphate/phosphodiesterase**, and activities had strong positive correlation ( $p < 0.05$ ) with  
10 available P. Most of the enzyme activities had a negative correlation with C/N ratio, S, and most  
11 of the trace elements except Cd. **E-supplementary data of this work can be found in online**  
12 **version of the paper**. The correlation was **moderately** strong with Al, Ba, Cu, Fe, Ni, Mn, and  
13 Zn, indicating that an increase of trace elements release during the early period of composting  
14 might inhibit enzyme activities (Aponte et al., 2020), which further **suggests** increased enzyme  
15 activities at later composting stages could be indicators of stability and immobilization of trace  
16 elements. A similar strong negative correlation of trace elements with enzyme activities was  
17 reported to occur during a cow manure vermicomposting by Malley et al. (2006) . However, in  
18 this study, positive correlations were found between Cd and all enzyme activities, implying that  
19 it could be rather stimulatory. Unlike all the other trace elements, Pb showed weak and positive  
20 correlations, possibly because Pb release was **not** affected during composting. RDA analysis of  
21 the enzyme activities as response variables and pH, OM, C/N ratio, and macro-nutrients as  
22 environmental factors showed a variance of 70.8 explained by RDA1 and RDA2 (Fig. 5B). The  
23 PERMANOVA test conducted showed the contribution of variance explained by RDA1 was

1 significant ( $p < 0.01$ ), but RDA2 was not. In Fig. 5C, trace elements explained 69.5% of total  
2 enzyme activity variation with RDA1 and RDA2. The overall results showed the significance of  
3 organic matter dynamics and the release or immobilization of elements in predicting the enzyme  
4 activities.

#### 5 4. Conclusions

6  
7 Post-digestate composting and co-composting reduced C/N ratio to 11-17. Phytotoxicity was  
8 reduced and greatest GI (76%) was obtained by co-composting with food processing waste. Up  
9 to 90% trace elements immobilization was found, and better results were obtained by co-  
10 composting with maize silage and food processing waste. Many of the 14 enzyme activities  
11 were low or absent in digestate but increased in the final composts. Alkaline  
12 phosphomonoesterase, and Pyrophosphate/phosphodiesterase had strong negative correlations  
13 ( $p < 0.05$ ) with Ba, Cu, Fe, Mn, Ni, and Zn. The overall findings suggest the significance of post-  
14 digestate co-composting in improving quality and enzyme activities as compost maturity index.

#### 16 Acknowledgements

17 Authors are thankful to Dr. Eleonora Pettinari for the facilitation to get access to sampling the  
18 digestate biomass and co-composting materials. This research was funded through scholarship  
19 provided to the first author by the Università Politecnica delle Marche.

20  
21  
22  
23  
24  
25  
26

## 1 5. References

- 2 Alburquerque, J.A., de la Fuente, C., Ferrer-Costa, A., Carrasco, L., Cegarra, J., Abad, M.,  
3 Bernal, M.P., 2012a. Assessment of the fertiliser potential of digestates from farm and  
4 agroindustrial residues. *Biomass and Bioenergy* 40, 181–189.  
5 <https://doi.org/10.1016/j.biombioe.2012.02.018>
- 6 Alburquerque, J.A., Fuente, C. De, Bernal, M.P., 2012b. Chemical properties of anaerobic  
7 digestates affecting C and N dynamics in amended soils. *Agriculture, Ecosyst. Environ.*  
8 *160*, 15–22. <https://doi.org/10.1016/j.agee.2011.03.007>
- 9 Aponte, H., Meli, P., Butler, B., Paolini, J., Matus, F., Merino, C., Cornejo, P., Kuzyakov, Y.,  
10 2020. Meta-analysis of heavy metal effects on soil enzyme activities. *Sci. Total Environ.*  
11 *737*, 139744. <https://doi.org/10.1016/j.scitotenv.2020.139744>
- 12 Beggio, G., Schievano, A., Bonato, T., Hennebert, P., Pivato, A., 2019. Statistical analysis for  
13 the quality assessment of digestates from separately collected organic fraction of municipal  
14 solid waste ( OFMSW ) and agro-industrial feedstock . Should input feedstock to anaerobic  
15 digestion determine the legal status of digesta. *Waste Manag.* 87, 546–558.  
16 <https://doi.org/10.1016/j.wasman.2019.02.040>
- 17 Bernal, M.P., Alburquerque, J.A., Moral, R., 2009. Composting of animal manures and chemical  
18 criteria for compost maturity assessment. A review. *Bioresour. Technol.* 100, 5444–5453.  
19 <https://doi.org/10.1016/j.biortech.2008.11.027>
- 20 Blazy, V., de Guardia, A., Benoist, J.C., Daumoin, M., Lemasle, M., Wolbert, D., Barrington, S.,  
21 2014. Odorous gaseous emissions as influence by process condition for the forced aeration  
22 composting of pig slaughterhouse sludge. *Waste Manag.* 34, 1125–1138.  
23 <https://doi.org/10.1016/j.wasman.2014.03.012>



1 Bustamante, M.A., Albuquerque, J.A., Restrepo, A.P., de la Fuente, C., Paredes, C., Moral, R.,  
2 Bernal, M.P., 2012. Co-composting of the solid fraction of anaerobic digestates, to obtain  
3 added-value materials for use in agriculture. *Biomass and Bioenergy* 43, 26–35.  
4 <https://doi.org/10.1016/j.biombioe.2012.04.010>

5 Cao, Y., Wang, X., Liu, L., Velthof, G.L., Misselbrook, T., Bai, Z., Ma, L., 2020. Acidification  
6 of manure reduces gaseous emissions and nutrient losses from subsequent composting  
7 process. *J. Environ. Manage.* 264, 110454. <https://doi.org/10.1016/j.jenvman.2020.110454>

8 Cardelli, V., De Feudis, M., Fornasier, F., Massaccesi, L., Cocco, S., Agnelli, A., Weindorf,  
9 D.C., Corti, G., 2019. Changes of topsoil under *Fagus sylvatica* along a small latitudinal-  
10 altitudinal gradient. *Geoderma* 344, 164–178.  
11 <https://doi.org/10.1016/j.geoderma.2019.01.043>

12 Castaldi, P., Garau, G., Melis, P., 2008. Maturity assessment of compost from municipal solid  
13 waste through the study of enzyme activities and water-soluble fractions. *Waste Manag.* 28,  
14 534–540. <https://doi.org/10.1016/j.wasman.2007.02.002>

15 Ge, M., Zhou, H., Shen, Y., Meng, H., Li, R., Zhou, J., Cheng, H., Zhang, X., Ding, J., Wang,  
16 Jian, Wang, Jiarui, 2020. Effect of aeration rates on enzymatic activity and bacterial  
17 community succession during cattle manure composting. *Bioresour. Technol.* 304, 122928.  
18 <https://doi.org/10.1016/j.biortech.2020.122928>

19 Gurmessa, B., Pedretti, E.F., Cocco, S., Cardelli, V., Corti, G., 2020. Manure anaerobic digestion  
20 effects and the role of pre- and post-treatments on veterinary antibiotics and antibiotic  
21 resistance genes removal efficiency. *Sci. Total Environ.* 721, 137532.  
22 <https://doi.org/10.1016/j.scitotenv.2020.137532>

23 Heiri, O., Lotter, A., Lemcke, G., 2001. Loss on ignition as a method for estimating organic and

1 carbonate content in sediments. *J. Paleolimnol.* 25, 101–110.

2 Herrmann, R.F., Shann, J.R., 1993. Enzyme activities as indicators of municipal solid waste  
3 compost maturity. *Compost Sci. Util.* 1, 54–63.  
4 <https://doi.org/10.1080/1065657X.1993.10757904>

5 Karwal, M., Kaushik, A., 2020. Co-composting and vermicomposting of coal fly-ash with press  
6 mud: Changes in nutrients, micro-nutrients and enzyme activities. *Environ. Technol. Innov.*  
7 18, 100708. <https://doi.org/10.1016/j.eti.2020.100708>

8 Kupper, T., Bürge, D., Bachmann, H.J., Güsewell, S., Mayer, J., 2014. Heavy metals in source-  
9 separated compost and digestates. *Waste Manag.* 34, 867–874.  
10 <https://doi.org/10.1016/j.wasman.2014.02.007>

11 Leconte, M.C., Mazzarino, M.J., Satti, P., Crego, M.P., 2011. Nitrogen and phosphorus release  
12 from poultry manure composts: The role of carbonaceous bulking agents and compost  
13 particle sizes. *Biol. Fertil. Soils* 47, 897–906. <https://doi.org/10.1007/s00374-011-0591-z>

14 Lencioni, G., Imperiale, D., Cavirani, N., Marmiroli, N., Marmiroli, M., 2016. Environmental  
15 application and phytotoxicity of anaerobic digestate from pig farming by in vitro and in  
16 vivo trials. *Int. J. Environ. Sci. Technol.* 13, 2549–2560. [https://doi.org/10.1007/s13762-](https://doi.org/10.1007/s13762-016-1088-y)  
17 [016-1088-y](https://doi.org/10.1007/s13762-016-1088-y)

18 Li, C., Li, H., Yao, T., Su, M., Li, J., Liu, Z., Xin, Y., Wang, L., Chen, J., Gun, S., 2020. Effects  
19 of microbial inoculation on enzyme activity, available nitrogen content, and bacterial  
20 succession during pig manure composting. *Bioresour. Technol.* 306, 123167.  
21 <https://doi.org/10.1016/j.biortech.2020.123167>

22 Li, Yun, Liu, H., Li, G., Luo, W., Sun, Y., 2018. Manure digestate storage under different  
23 conditions: Chemical characteristics and contaminant residuals. *Sci. Total Environ.* 639,

1 19–25. <https://doi.org/10.1016/j.scitotenv.2018.05.128>

2 Li, Yangyang, Luo, W., Lu, J., Zhang, X., Li, S., Wu, Y., Li, G., 2018. Effects of digestion time  
3 in anaerobic digestion on subsequent digestate composting. *Bioresour. Technol.* 267, 117–  
4 125. <https://doi.org/10.1016/J.BIORTECH.2018.04.098>

5 Lopes, I.G., Braos, L.B., Cruz, M.C.P., Vidotti, R.M., 2021. Valorization of animal waste from  
6 aquaculture through composting: Nutrient recovery and nitrogen mineralization.  
7 *Aquaculture* 531, 735859. <https://doi.org/10.1016/j.aquaculture.2020.735859>

8 Luo, Y., Liang, J., Zeng, G., Chen, M., Mo, D., Li, G., Zhang, D., 2018. Seed germination test  
9 for toxicity evaluation of compost: Its roles, problems and prospects. *Waste Manag.* 71,  
10 109–114. <https://doi.org/10.1016/j.wasman.2017.09.023>

11 Malley, C., Nair, J., Ho, G., 2006. Impact of heavy metals on enzymatic activity of substrate and  
12 on composting worms *Eisenia fetida*. *Bioresour. Technol.* 97, 1498–1502.  
13 <https://doi.org/10.1016/j.biortech.2005.06.012>

14 Miaomiao, H., Wenhong, L., Xinqiang, L., Donglei, W., Guangming, T., 2009. Effect of  
15 composting process on phytotoxicity and speciation of copper, zinc and lead in sewage  
16 sludge and swine manure. *Waste Manag.* 29, 590–597.  
17 <https://doi.org/10.1016/j.wasman.2008.07.005>

18 Möller, K., Müller, T., 2012. Effects of anaerobic digestion on digestate nutrient availability and  
19 crop growth : A review. *Eng. Life Sci.* 12, 242–257. <https://doi.org/10.1002/elsc.201100085>

20 Mondini, C., Fornasier, F., Sinicco, T., 2004. Enzymatic activity as a parameter for the  
21 characterization of the composting process. *Soil Biol. Biochem.* 36, 1587–1594.  
22 <https://doi.org/10.1016/j.soilbio.2004.07.008>

23 **Nkoa, R., 2014. Agricultural benefits and environmental risks of soil fertilization with anaerobic**

1        [digestates: A review. Agron. Sustain. Dev. 34, 473–492. https://doi.org/10.1007/s13593-](https://doi.org/10.1007/s13593-013-0196-z)  
2        [013-0196-z](https://doi.org/10.1007/s13593-013-0196-z)

3        Oksanen, A.J., Blanchet, F.G., Friendly, M., Kindt, R., Legendre, P., Mcglinn, D., Minchin, P.R.,  
4        Hara, R.B.O., Simpson, G.L., Solymos, P., Stevens, M.H.H., Szoecs, E., 2019. vegan:  
5        Community Ecology Package. R package version 2.5-6. [https://CRAN.R-](https://CRAN.R-project.org/package=vegan)  
6        [project.org/package=vegan](https://CRAN.R-project.org/package=vegan).

7        Rai, R., Suthar, S., 2020. Composting of toxic weed Parthenium hysterophorus: Nutrient  
8        changes, the fate of faecal coliforms, and biopesticide property assessment. *Bioresour.*  
9        *Technol.* 311, 123523. <https://doi.org/10.1016/j.biortech.2020.123523>

10       Rehl, T., Müller, J., 2011. Life cycle assessment of biogas digestate processing technologies.  
11       *Resour. Conserv. Recycl.* 56, 92–104. <https://doi.org/10.1016/j.resconrec.2011.08.007>

12       Saveyn, H., Eder, P., 2014. End-of-waste criteria for biodegradable waste subjected to biological  
13       treatment ( compost & digestate ): Technical proposals. <https://doi.org/10.2791/6295>

14       Sharma, D., Yadav, K.D., Kumar, S., 2018. Role of sawdust and cow dung on compost maturity  
15       during rotary drum composting of flower waste. *Bioresour. Technol.* 264, 285–289.  
16       <https://doi.org/10.1016/j.biortech.2018.05.091>

17       [Świątczak, P., Cydzik-Kwiatkowska, A., 2018. Treatment of Ammonium-Rich Digestate from](https://doi.org/10.1007/s11270-018-3887-x)  
18       [Methane Fermentation Using Aerobic Granular Sludge. \*Water. Air. Soil Pollut.\* 229, 229:](https://doi.org/10.1007/s11270-018-3887-x)  
19       [247. https://doi.org/10.1007/s11270-018-3887-x](https://doi.org/10.1007/s11270-018-3887-x)

20       Tambone, F., Terruzzi, L., Scaglia, B., Adani, F., 2015. Composting of the solid fraction of  
21       digestate derived from pig slurry: Biological processes and compost properties. *Waste*  
22       *Manag.* 35, 55–61. <https://doi.org/10.1016/j.wasman.2014.10.014>

23       Tang, J., Zhang, L., Zhang, J., Ren, L., Zhou, Y., Zheng, Y., Luo, L., Yang, Y., Huang, H., Chen,

1 A., 2020. Physicochemical features, metal availability and enzyme activity in heavy metal-  
2 polluted soil remediated by biochar and compost. *Sci. Total Environ.* 701, 134751.  
3 <https://doi.org/10.1016/j.scitotenv.2019.134751>

4 Teglia, C., Tremier, A., Martel, J.L., 2011. Characterization of solid digestates: Part 2,  
5 assessment of the quality and suitability for composting of six digested products. *Waste and*  
6 *Biomass Valorization* 2, 113–126. <https://doi.org/10.1007/s12649-010-9059-x>

7 Tigni, V., Franchino, M., Bona, F., Varese, G.C., 2016. Is digestate safe? A study on its  
8 ecotoxicity and environmental risk on a pig manure. *Sci. Total Environ.* 551–552, 127–132.  
9 <https://doi.org/10.1016/j.scitotenv.2016.02.004>

10 Tiquia, S.M., 2002. Evolution of extracellular enzyme activities during manure composting. *J.*  
11 *Appl. Microbiol.* 92, 764–775. <https://doi.org/10.1046/j.1365-2672.2002.01582.x>

12 Torres-Climent, A., Martin-Mata, J., Marhuenda-Egea, F., Moral, R., Barber, X., Perez-Murcia,  
13 M.D., Paredes, C., 2015. Composting of the Solid Phase of Digestate from Biogas  
14 Production: Optimization of the Moisture, C/N Ratio, and pH Conditions. *Commun. Soil*  
15 *Sci. Plant Anal.* 46, 197–207. <https://doi.org/10.1080/00103624.2014.988591>

16 Wan, W., Wang, Y., Tan, J., Qin, Y., Zuo, W., Wu, H., He, H., He, D., 2020. Alkaline  
17 phosphatase-harboring bacterial community and multiple enzyme activity contribute to  
18 phosphorus transformation during vegetable waste and chicken manure composting.  
19 *Bioresour. Technol.* 297, 122406. <https://doi.org/10.1016/j.biortech.2019.122406>

20 Wu, S., Shen, Z., Yang, C., Zhou, Y., Li, X., Zeng, G., Ai, S., He, H., 2017. Effects of C/N ratio  
21 and bulking agent on speciation of Zn and Cu and enzymatic activity during pig manure  
22 composting. *Int. Biodeterior. Biodegrad.* 119, 429–436.  
23 <https://doi.org/10.1016/j.ibiod.2016.09.016>

1 Zeng, Y., De Guardia, A., Dabert, P., 2016. Improving composting as a post-treatment of  
2 anaerobic digestate. *Bioresour. Technol.* 201, 293–303.

3 <https://doi.org/10.1016/j.biortech.2015.11.013>

4

Table 1. Composition of the compost piles.

| Pile | Composition  | Mix ratio<br>(w/w) |
|------|--|--------------------|
| D00  | Solid digestate only   | -                  |
| DCB  | Solid digestate + food processing waste                      | 4:1                |
| DMS  | Solid digestate + Maize silage                               | 4:1                |
| DPL  | Solid digestate + Poultry litter <sup>‡</sup>                | 4:1                |
| DMP  | Solid digestate + Maize silage + Poultry litter <sup>‡</sup> | 8:1:1              |

<sup>‡</sup>The poultry litter was 85:15 chicken manure: wheat straw ratio; moisture  $\approx$ 58%.

Table 2. Physicochemical properties of digestate and the co-composting materials (Mean  $\pm$  SD)

(n = 3). For each parameter, mean values with different letters significantly differ ( $p < 0.05$ ).

| Variable                      | Digestate          | Food processing waste | Maize silage     | Poultry litter    | P-value |
|-------------------------------|--------------------|-----------------------|------------------|-------------------|---------|
| pH                            | 8.1 $\pm$ 0.0a     | 5.8 $\pm$ 0.1d        | 6.2 $\pm$ 0.0c   | 7.2 $\pm$ 0.1b    | <0.001  |
| TS (g kg <sup>-1</sup> )      | 384 $\pm$ 42c      | 941 $\pm$ 45a         | 413 $\pm$ 17c    | 565 $\pm$ 44b     | <0.001  |
| OM (g kg <sup>-1</sup> )      | 837 $\pm$ 2b       | 956 $\pm$ 1a          | 958 $\pm$ 0a     | 828 $\pm$ 0c      | <0.001  |
| Total C (g kg <sup>-1</sup> ) | 390 $\pm$ 35       | 416 $\pm$ 3           | 415 $\pm$ 6      | 387 $\pm$ 2       | 0.15    |
| Total N (g kg <sup>-1</sup> ) | 15.1 $\pm$ 1.0b    | 18.2 $\pm$ 1.1b       | 16.1 $\pm$ 1.6b  | 35.3 $\pm$ 3.4a   | <0.001  |
| C/N ratio                     | 25.8 $\pm$ 2.7a    | 22.9 $\pm$ 1.3a       | 25.8 $\pm$ 2.2a  | 11.0 $\pm$ 1.1b   | <0.001  |
| Ca (g kg <sup>-1</sup> )      | 3.0 $\pm$ 0.3c     | 0.8 $\pm$ 0.0d        | 6.0 $\pm$ 0.3b   | 12.6 $\pm$ 0.2a   | <0.001  |
| K (g kg <sup>-1</sup> )       | 141.0 $\pm$ 6.4b   | 36.2 $\pm$ 0.9d       | 83.5 $\pm$ 4.3c  | 259.6 $\pm$ 5.3a  | <0.001  |
| Mg (g kg <sup>-1</sup> )      | 2.9 $\pm$ 0.1d     | 4.2 $\pm$ 0.1c        | 7.7 $\pm$ 0.2b   | 15.3 $\pm$ 0.1a   | <0.001  |
| P (g kg <sup>-1</sup> )       | 10.2 $\pm$ 0.2c    | 17.9 $\pm$ 0.5a       | 10.2 $\pm$ 0.1c  | 14.4 $\pm$ 0.2b   | <0.001  |
| S (g kg <sup>-1</sup> )       | 11.7 $\pm$ 0.3b    | 2.0 $\pm$ 0.1d        | 4.4 $\pm$ 0.1c   | 57.5 $\pm$ 0.3a   | <0.001  |
| Al (mg kg <sup>-1</sup> )     | 108.2 $\pm$ 4.0b   | 145.9 $\pm$ 10.5a     | 113.1 $\pm$ 6.0b | 143.1 $\pm$ 4.3a  | <0.001  |
| Ba (mg kg <sup>-1</sup> )     | 0.1 $\pm$ 0.0c     | 1.7 $\pm$ 0.5b        | 1.3 $\pm$ 0.3b   | 11.5 $\pm$ 0.3a   | <0.001  |
| Cd (mg kg <sup>-1</sup> )     | 0.8 $\pm$ 0.1b     | 1.2 $\pm$ 0.1ab       | 0.7 $\pm$ 0.1b   | 1.6 $\pm$ 0.3a    | 0.001   |
| Cu (mg kg <sup>-1</sup> )     | 39.2 $\pm$ 1.5b    | 10.2 $\pm$ 0.3c       | 9.1 $\pm$ 0.2c   | 130.1 $\pm$ 0.4a  | <0.001  |
| Fe (mg kg <sup>-1</sup> )     | 1525.0 $\pm$ 65.9a | 93.0 $\pm$ 4.9c       | 60.6 $\pm$ 3.1c  | 552.2 $\pm$ 1.5b  | <0.001  |
| Mn (mg kg <sup>-1</sup> )     | 28.8 $\pm$ 1.0b    | 20.7 $\pm$ 0.4b       | 25.9 $\pm$ 2.5b  | 141.9 $\pm$ 5.8a  | <0.001  |
| Ni (mg kg <sup>-1</sup> )     | 11.6 $\pm$ 0.1b    | 1.5 $\pm$ 0.2c        | 0.3 $\pm$ 0.3d   | 44.1 $\pm$ 0.4a   | <0.001  |
| Pb (mg kg <sup>-1</sup> )     | 36.7 $\pm$ 3.4     | 34.4 $\pm$ 2.4        | 35.4 $\pm$ 1.8   | 36.6 $\pm$ 2.6    | 0.672   |
| Zn (mg kg <sup>-1</sup> )     | 95.3 $\pm$ 14.6b   | 41.1 $\pm$ 0.3c       | 4.3 $\pm$ 0.1d   | 258.0 $\pm$ 20.6a | <0.001  |

TS = total solid; OM = organic matter on a dry mass basis.



Table 3. Immobilization and release of trace elements and nutrients expressed as a percentage of the amount immobilized or released over the composting period from the concentration of the mix (mean  $\pm$  SD) (n = 3). Negative values indicate immobilization, positive values indicate release.

| Pile    | D00              | DCB              | DMS              | DPL             | DMP              |
|---------|------------------|------------------|------------------|-----------------|------------------|
| Ca      | -9.1 $\pm$ 4.5   | -53.5 $\pm$ 3.0  | -22.3 $\pm$ 0.6  | -50.9 $\pm$ 2.2 | -33.0 $\pm$ 6.2  |
| K       | 28.3 $\pm$ 3.9   | 51.3 $\pm$ 4.8   | 28.4 $\pm$ 2.8   | -1.0 $\pm$ 1.5  | -3.3 $\pm$ 1.2   |
| Mg      | 124.2 $\pm$ 5.7  | -38.5 $\pm$ 0.2  | 7.2 $\pm$ 1.5    | -19.0 $\pm$ 2.4 | -22.9 $\pm$ 3.6  |
| P       | 50.7 $\pm$ 2.7   | 25.4 $\pm$ 3.0   | 61.8 $\pm$ 3.1   | -1.5 $\pm$ 0.4  | 29.8 $\pm$ 6.7   |
| S       | -52.9 $\pm$ 1.6  | 2.6 $\pm$ 2.3    | -47.7 $\pm$ 1.1  | -18.1 $\pm$ 0.5 | -32.4 $\pm$ 2.3  |
| Total N | 26.1 $\pm$ 16.2  | 89.8 $\pm$ 16.4  | 92.9 $\pm$ 14.2  | 36.7 $\pm$ 6.8  | 64.0 $\pm$ 15.6  |
| Al      | -93.4 $\pm$ 0.3  | -94.6 $\pm$ 0.3  | -93.6 $\pm$ 0.4  | -16.2 $\pm$ 6.6 | -93.2 $\pm$ 1.5  |
| Ba      | -74.8 $\pm$ 1.0  | -95.3 $\pm$ 0.8  | -91.3 $\pm$ 1.4  | -97.7 $\pm$ 0.0 | -93.8 $\pm$ 0.1  |
| Cd      | 586.0 $\pm$ 76.1 | 475.9 $\pm$ 23.7 | 593.2 $\pm$ 83.9 | -34.3 $\pm$ 8.6 | 528.9 $\pm$ 77.9 |
| Cu      | -55.7 $\pm$ 2.2  | -47.9 $\pm$ 4.1  | -43.2 $\pm$ 2.4  | -25.7 $\pm$ 1.7 | -49.4 $\pm$ 0.2  |
| Fe      | -89.3 $\pm$ 0.5  | -82.3 $\pm$ 0.6  | -86.5 $\pm$ 0.4  | -75.3 $\pm$ 0.5 | -85.9 $\pm$ 0.5  |
| Mn      | -74.6 $\pm$ 0.9  | -60.6 $\pm$ 1.9  | -68.6 $\pm$ 0.5  | -72.6 $\pm$ 1.2 | -72.3 $\pm$ 0.4  |
| Ni      | -42.5 $\pm$ 4.9  | -42.6 $\pm$ 2.3  | -40.1 $\pm$ 5.2  | -70.4 $\pm$ 0.9 | -71.0 $\pm$ 5.7  |
| Pb      | 3.8 $\pm$ 16.8   | -0.6 $\pm$ 4.7   | -1.7 $\pm$ 8.4   | -3.2 $\pm$ 10.2 | -10.8 $\pm$ 1.8  |
| Zn      | -85.5 $\pm$ 2.2  | -70.9 $\pm$ 3.9  | -66.3 $\pm$ 4.1  | -93.9 $\pm$ 1.0 | -73.8 $\pm$ 2.5  |

D00 = solid digestate only; DCB = solid digestate + food processing waste; DMS = solid digestate + maize silage; DPL = solid digestate + poultry litter; DMP = solid digestate + maize silage + poultry litter.

## Figure captions

**Fig. 1.** Trend of temperature, pH, total C, C/N ratio, organic matter (OM), and germination index (GI) during the composting period. D00 = solid digestate only, DCB = solid digestate + food processing waste, DMS = solid digestate + maize silage, DPL = solid digestate + poultry litter, DMP = solid digestate + maize silage + poultry litter.

**Fig. 2.** Dynamics of macro-nutrients during the composting period. D00 = solid digestate only, DCB = solid digestate + food processing waste, DMS = solid digestate + maize silage, DPL = solid digestate + poultry litter, DMP = solid digestate + maize silage + poultry litter.

**Fig. 3.** Changes in the content of trace elements during the composting period. D00 = solid digestate only, DCB = solid digestate + food processing waste, DMS = solid digestate + maize silage, DPL = solid digestate + poultry litter, DMP = solid digestate + maize silage + poultry litter.

**Fig. 4.** Enzyme activity dynamics over the composting period. acP: acid phosphomonoesterase; alkP: alkaline phosphomonoesterase; alfaG:  $\alpha$ -glucosidase; aryS: arylsulfatase; betaG:  $\beta$ -glucosidase; betaGAL:  $\beta$ -galactosidase; bisP: phosphodiesterase; cell: cellulase, chit: chitinase; leu: leucine aminopeptidase; nona: lipase nonanoate-esterase; piroP: pyrophosphate/phosphodiesterase; uroni: glucuronidase; xilo: xylosidase. D00 = solid digestate only, DCB = solid digestate + food processing waste, DMS = solid digestate + maize silage, DPL = solid digestate + poultry litter, DMP = solid digestate + maize silage + poultry litter.

**Fig. 5.** Redundancy analysis of enzyme activities in relation to environmental factors (A: pH, C/N ratio, OM, and macro-nutrients; B: trace elements). acP: acid phosphomonoesterase; alkP: alkaline phosphomonoesterase; alfaG:  $\alpha$ -glucosidase; aryS: arylsulfatase; betaG:  $\beta$ -glucosidase; betaGAL:  $\beta$ -galactosidase; bisP: phosphodiesterase; cell: cellulase, chit: chitinase; leu: leucine aminopeptidase; nona: lipase nonanoateesterase; piroP: pyrophosphate/phosphodiesterase; uroni: glucuronidase; xilo: xylosidase. D00 = solid digestate only, DCB = solid digestate + food processing waste, DMS = solid digestate + maize silage, DPL = solid digestate + poultry litter, DMP = solid digestate + maize silage + poultry litter.




Fig. 1.




Fig. 2.




Fig. 3.




Fig. 4.








Fig. 5.





## Post-digestate composting benefits and the role of enzyme activity to predict trace element immobilization and compost maturity

Biyensa Gurmessa<sup>1\*</sup>, Stefania Cocco<sup>1</sup>, Amanda J. Ashworth<sup>2</sup>, Ester Foppa Pedretti<sup>1</sup>, Alessio Ilari<sup>1</sup>, Valeria Cardelli<sup>1</sup>, Flavio Fornasier<sup>3</sup>, Maria Letizia Ruello<sup>4</sup>, Giuseppe Corti<sup>1</sup>

<sup>1</sup>Department of Agricultural, Food and Environmental Sciences, Università Politecnica delle Marche, 60131 Ancona, Italy

<sup>2</sup>USDA-ARS, Poultry Production and Product Safety Research Unit, 1260 W. Maple St. Fayetteville, AR 72701, USA

<sup>3</sup>Consiglio per la Ricerca e la Sperimentazione in Agricoltura, Centro di Ricerca per lo Studio delle Relazioni tra Pianta e Suolo (C.R.E.A.-R.P.S.), 34170 Gorizia, Italy

<sup>4</sup>Department of Materials, Environmental Sciences and Urban Planning, Università Politecnica delle Marche, 60131 Ancona, Italy

\*Corresponding author: [biyensa@gmail.com](mailto:biyensa@gmail.com)




Fig. Pearson's correlation between enzyme activities and water extractable elements. Correlation coefficients indicated with \*, \*\*, and \*\*\* were statistically significant ( $p < 0.05$ ,  $p < 0.001$ , and  $p < 0.001$ , respectively).

**Biyensa Gurmessa:** Conceptualization, Methodology, Software, Data curation, Visualization, Writing original draft preparation. **Stefania Cocco:** Writing-Reviewing, **Amanda J. Ashworth:** Writing-Reviewing, **Ester Foppa Pedretti:** Supervision, Design, Conceptualization. **Alessio Ilari:** Design, conceptualization. **Valeria Cardelli:** Analysis, Investigation, **Flavio Fornasier:** Analysis, Investigation. **Maria Letizia Ruello:** Analysis, Writing-reviewing. **Giuseppe Corti:** Writing-Reviewing, supervision

**Declaration of interests**

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests: