



UNIVERSITÀ POLITECNICA DELLE MARCHE
Repository ISTITUZIONALE

Effect of coppice conversion into high forest on soil organic C and nutrients stock in a Turkey oak (*Quercus cerris* L.) forest in Italy

This is the peer reviewed version of the following article:

Original

Effect of coppice conversion into high forest on soil organic C and nutrients stock in a Turkey oak (*Quercus cerris* L.) forest in Italy / Camponi, Lorenzo; Cardelli, Valeria; Cocco, Stefania; Serrani, Dominique; Salvucci, Andrea; Cutini, Andrea; Agnelli, Alberto; Fabbio, Gianfranco; Bertini, Giada; Roggero, Pier Paolo; Corti, Giuseppe. - In: JOURNAL OF ENVIRONMENTAL MANAGEMENT. - ISSN 0301-4797. - 312:(2022). [10.1016/j.jenvman.2022.114935]

Availability:

This version is available at: 11566/300864 since: 2024-10-24T11:24:59Z

Publisher:

Published

DOI:10.1016/j.jenvman.2022.114935

Terms of use:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. The use of copyrighted works requires the consent of the rights' holder (author or publisher). Works made available under a Creative Commons license or a Publisher's custom-made license can be used according to the terms and conditions contained therein. See editor's website for further information and terms and conditions.

This item was downloaded from IRIS Università Politecnica delle Marche (<https://iris.univpm.it>). When citing, please refer to the published version.

(Article begins on next page)

17 **Abstract**

18 In forest ecosystems, a variety of abiotic and biotic soil forming factors drives soil organic matter
19 (SOM) and nutrients cycling with a profitable outcome on climate change mitigation. As a
20 consequence, type and intensity of forest management, through its impact on carbon (C) and
21 nutrient soil stocks, can be considered as an additional soil forming force. In this study, we
22 investigated the influence of the coppice conversion into high forest on pedogenesis and on soil C
23 and nutrient (N, P, Ca, Mg, and K) stocks, fifty years later the beginning of the conversion-cycle.
24 The trial was established in a Turkey oak forest historically managed under the coppice system in
25 central Italy. Specifically, we considered tree population density (natural evolution – control,
26 moderate thinning, heavy thinning) where soil samples were collected according to genetic horizon
27 to estimate C, N, and P stocks both in the forest floor and at fixed depth intervals (0-30, 30-50 and
28 50-75 cm). Further, the stocks of exchangeable Ca, Mg, and K were also assessed for the mineral
29 layers. The results showed that litter and the upper layer of mineral soil (0-30 cm) contained a
30 similar quantity of C (about 74-83 Mg ha^{-1}), independently of the trials and no differences were
31 observed also in the whole soil stocks (about 192-213 Mg ha^{-1}). The comparison of the mean stocks
32 calculated per 1-cm of thickness of organic (O), organo-mineral (OM), and mineral (M) layers,
33 although it did not display any difference among trials (excepted for P and Mg), showed a similar
34 capability of the organo-mineral horizons to store C and nutrients compared with the organic ones
35 (e.g., about 6-12 Mg ha^{-1} , 0.3-0.5 Mg ha^{-1} and 0.5-1.5 kg ha^{-1} for C, N and P, respectively). Our
36 findings showed that thinning operated on Turkey oak coppice did not affect soil capacity to store C
37 and nutrients. These results suggested that the forest ecosystem itself is the main soil forming force
38 and this is consistent with the target of adopting forest management able to control the global C
39 cycle through the storage of SOM in the mineral soil rather than in forest floor, where SOM
40 turnover is faster.

41 **Keywords:** forest soil, organic matter, rock fragments, pedogenetic horizons, coppice conversion
42 into high forest, sustainable forest management

43

44 **1. Introduction**

45 Soil organic matter (SOM) plays key roles in terrestrial ecosystems, where it is involved in many
46 processes of soil conservation. SOM is fundamental in stabilizing soil structures and reducing soil
47 erosion, improving water-holding capacity, and releasing nutrients to plants, microorganisms, and
48 soil fauna (Bot and Benites, 2005; Canedoli et al., 2020). In forest soils, the input of organic matter
49 depends on litter production, mortality of fine roots, roots exudates, and shoots residues (Lehmann
50 et al., 2015; Diao et al., 2020; Wu et al., 2020). Thus, depending on the interaction among the main
51 soil forming forces (parent material, climate, living organisms, relief, and time; Jenny, 1941), a
52 vastity of physicochemical and biological processes affects the transformation of plant-derived
53 organic materials in SOM. During this transformation, SOM is stabilized by the formation of
54 organo-metallic complexes with di- and trivalent cations (Kaiser et al., 2016), the formation of
55 organo-mineral complexes with clay minerals (Kögel-Knabner et al., 2008, Barré et al., 2014;
56 Gartzia-Bengoetxea et al., 2020), and the occlusion within aggregates (Schrumpf et al., 2013),
57 favouring its preservation in the soil. Therefore, plant species, soil properties, and their interactions
58 play a key role in determining the soil organic C (SOC) stock and, due to SOM elemental content,
59 also in the biogeochemical cycles of nutrients like N, P, Ca, Mg, and K (Tiessen et al., 1994). For
60 instance, estimations indicate that the mean world soil content to 1-m depth is 1462-1548 Pg for
61 organic C and 133-140 Pg for total N (Batjes, 1996), more than the global content obtained
62 combining vegetation and atmosphere (Lehmann et al., 2015; Mayer et al., 2020).

63 In forest ecosystems, forest management may impact on SOC and nutrient stock. In their review,
64 Mayer et al. (2020) reported that management practices like site preparation, harvesting operations,
65 removal of harvest residues, and removal of litter and biomass for fodder, fuel, or animal bedding
66 have a negative impact on SOC stock capacity. Conversely, N addition, introduction of N-fixing
67 plants, and herbivory regulation have a positive impact on SOC storage. Other practices like
68 management of tree species diversity and periodical thinnings over the whole stand lifespan that are
69 used to manage tree population density in high forest [which consists in a stand of trees, generally
70 originated from seed, that develop a high, closed canopy (SAF, 2008)] and in the conversion into
71 high forest of coppice systems seem not to interfere with the soil capacity to stock organic C under
72 both broadleaves and conifers (Bravo-Oviedo et al., 2015; Prasad Dangal et al., 2017; Zhang et al.,
73 2018; Mayer et al., 2020).

74 Coppicing represents the oldest form of systematic and sustainable use of forests. It is a very
75 flexible system that requires a low energy input and has been adapted and modified according to the
76 needs of rural societies, to whom coppice forests deliver small size wood primarily for energy
77 (firewood and charcoal), agriculture, and small scale businesses. As a matter of fact, coppice forests
78 characterize the European landscapes, especially in mountainous areas of central, east and southern

79 Europe. Due to rural migration and technical and economic restrictions, most of the coppice forests
80 are today neglected or abandoned, representing a significantly underused natural resource (Unrau et
81 al., 2018). In Italy, coppice forests cover 3.663 million hectares (Mairota et al., 2018)) and both
82 evergreen and deciduous *Quercus* spp. make a significant share of the total cover (nearly 1.6
83 million ha). Following the crisis of the firewood and charcoal system, the conversion of coppices
84 into high-forests is considered a sustainable forest management in many countries (Fabbio, 2016;
85 Fabbio and Cutini, 2017; Cutini et al., 2021) due to the low-frequency soil disturbance that would
86 favour the storage of SOC (Hölscher et al., 2001; Marchi et al., 2016). Therefore, the link between
87 forest management and soil properties, with its specific capacity to determine SOC stock and
88 climate change mitigation, has fostered a number of scientific researches (e.g., Caddeo et al., 2019;
89 Zhang et al., 2019; Zhao et al., 2019; Lee et al., 2020), but scarce has been the interest on the effect
90 of forest management on the soil stock of nutrients like N, P, Ca, Mg, and K, whose abundance and
91 availability is key to soil fertility and biomass production.

92 The aim of this work was to assess the role of thinning performed for the conversion of a coppice
93 forest into high forest on soil C, N, available P, and exchangeable Ca, Mg, and K stocks. The effect
94 of periodical thinning vs no silvicultural intervention (namely, natural evolution following the
95 suspension of periodical harvestings) was investigated in a Turkey oak (*Quercus cerris* L.) stand
96 under conversion into high forest and managed as coppice up to 1949 (last coppicing).

97 To test the hypothesis that different forest managements can affect soil C and nutrients stocks, and
98 to investigate on the contribution of each horizon to the whole soil stocks, we estimated: i) C, N,
99 and P stored in the genetic horizons (ranked in organic, organo-mineral and mineral horizons) and
100 at fixed depth intervals (0-30, 30-50, and 50-75 cm); and ii) exchangeable Ca, Mg, and K stored in
101 the mineral soil (organo-mineral and mineral horizons) and at fixed depth intervals (0-30, 30-50,
102 and 50-75 cm).

103

104 **2. Materials and Methods**

105 *2.1. Environmental and historical background*

106 The study was conducted in the Natural Reserve of Monterufoli-Caselli forest, Tuscany, Italy (Fig.
107 1), a Natura 2000 Site (SPA-SAC IT5170008 Complesso di Monterufoli). The whole reserve covers
108 a gentle hilly environment and extends for 4,828 ha within altitudes spanning between 100 and 560
109 m. The mean annual precipitation is 750 mm and the mean annual air temperature is 13.5 °C.
110 Geology of the area is rather complex being dominated by serpentinite and polygenic breccias
111 (Paleocene), followed by calcareous sandstone interbedded with limestone (Cretaceous), pelitic
112 marine sediments (Pliocene), silty-clay schists (Cretaceous), and quartzous sandstone interbedded

113 with arenaceous limestone (Paleocene). The area was heavily influenced by human activities since
114 ancient times. Central Italy, and especially Tuscany, was subject to an intense mining activity
115 during Bronze and Iron ages. As reported by Cartocci et al. (2007), Tuscany can be considered one
116 of the most important ancient metallurgical districts of Italy, with several active mining centres
117 since the Iron Age. In the study area, considerable was the production of iron, pyrite, base metals,
118 silver, antimony, mercury, and gold for millennia (Chiarantini et al., 2018). Because of this activity,
119 there was the need of fuel for metal smelting, with the consequent exploitation of forests (especially
120 oak forests of *Quercus cerris* L., *Quercus pubescens* Willd., and others) for charcoal production
121 (Carrari et al., 2017).

122

123 2.2. Study area

124 The study was run in a long-term monitoring area located within the Caselli Forest. Forest stands
125 consisted of Turkey oak (*Quercus cerris* L.) for about 90%, with broadleaves like *Fraxinus* spp.,
126 *Ulmus* spp., *Ostrya* spp., and *Quercus ilex* L. as subsidiary species. Under the coppice system, in
127 Italy Turkey oak cover $\approx 675,000$ ha, i.e. the 18.4% of coppice forests (Manetti et al., 2020). Leaf
128 area index (LAI) ranged from 4.3 to 5.2 (Cutini, 1996). Here, last coppicing was performed in 1949;
129 then, in 1969, a long-term experiment aimed at comparing the periodical thinning of standing crop
130 vs its natural evolutive pattern to achieve coppice conversion into high forest was established. The
131 main goal of the experiment was to verify stand dynamics as for its structural-compositional
132 arrangement and functional traits of tree biomass. The treatments on the ground were the full
133 release of the dominated layer, moderate thinning (MT) and heavy thinning (HT) with an average
134 release of 1500 and 1100 stems ha^{-1} in the dominant layer, respectively. The coppice under natural
135 evolution, in absence of any practice, was considered as control (CTR); here, the average full tree
136 density was 4269 ha^{-1} (Fabbio and Amorini, 2006; Manetti and Gugliotta, 2006). Each trial was
137 repeated four times according to a randomized blocks design. Within plots of several thousands of
138 m^2 , we selected a survey area of 900 m^2 all within a NNE-NNW exposure on slopes roughly
139 ranging from 10 to 20% (Table S1, Supplementary Materials). A second thinning was implemented
140 in 1989 releasing 715 and 1036 shoots ha^{-1} in HT and MT, respectively. Average stem density
141 decreased to 3589 ha^{-1} in CTR. The arrangement of stand structure following the applied
142 silviculture was a two-storied stand: the dominant layer mainly made of Turkey oak, and the
143 dominated layer made the set of subsidiary broadleaved species.

144 Other inventories were performed in 1998 and 2004. In 2004, there were 578 shoots ha^{-1} in the
145 dominant layer of HT and 869 shoots ha^{-1} in MT. Average full stem density was 3417 ha^{-1} in CTR.
146 Main stand parameters are summarized in Table 1.

147

148 *2.3. Sampling sites and soil sampling*

149 The study site spanned from 337 to 345 m above sea level, on soil formed on calcareous sandstone
150 interbedded with limestone. In each of the four plots of the three trials, in 2017 a survey was run to
151 evaluate the spatial variability of surface stoniness, rock outcrops, slope, micro-topography,
152 dominant vegetation, and understorey to select the location where to dig a soil profile. Then, a total
153 of 12 profiles (1 profile • 4 plots • 3 trials) representative of each plot conditions were opened
154 within locations with 12-15% slope and 90-95% soil cover. In each plot the soil profile was dug at
155 ≈ 1 m from the stem (downslope position) of one of the oldest trees and until the depth of ≈ 1 m,
156 except for lithic contact. For each profile, the organic horizons forming the forest floor were
157 morphologically described per Baize et al. (2008) and sampled in an area of about 3 m² around the
158 profile. The mineral soil was morphologically described per Schoeneberger et al. (2012) and
159 sampled by genetic horizons. Soil morphologies provided of understorey composition (Fрати et al.,
160 2021) are reported in Table S1 of Supplementary Material. During the field operations, the collected
161 samples were stored in a refrigerated bag and, once in the laboratory, they were allowed to air-dry.
162 Thus, the mineral samples were sieved at 2 mm to separate the fine earth (< 2-mm fraction) from
163 the skeleton (> 2-mm fraction).

164

165 *2.4. Laboratory analysis*

166 The bulk density of both fine earth and skeleton of each horizon was determined by soil cylinders.
167 Specifically, two horizontal soil cores were collected from each mineral horizon by using cylinders
168 of 503 cm³ (height: 10.8 cm; diameter: 7.7 cm). In the laboratory, the collected sample was sieved
169 at 2 mm and the volume of the skeletal particles was determined by water displacement after the
170 particles were water-saturated (Corti et al., 1998). The volume of the fine earth was obtained by
171 subtracting that of the skeletal particles from the total volume of the cylinder. Both fine earth and
172 skeleton were then heated at 105°C and weighed. The content of large cobbles was estimated by the
173 “percent of area covered” figure reported in Schoeneberger et al. (2012), and their bulk density
174 determined as mentioned above. For the organic horizons, the bulk density was estimated by
175 pedotransfer functions (De Nicola et al., 2014), which have been tested by other researcher in
176 various Italian contexts (Brenna et al., 2010; Garlato et al., 2009a, b; Guermandi et al., 2013). These
177 equations provide bulk density as a function of the percentage of estimated organic matter (OM = 2
178 • organic C) as follows:

179 1. For OM > 30%: bulk density (g cm⁻³) = 0.00589 • organic C + 0.554;

180 2. For OM = 30–15%: bulk density (g cm⁻³) = 0.00745 • organic C + 0.593;

181 3. For $OM < 15\%$: bulk density ($g\ cm^{-3}$) = $0.00797 \cdot \text{organic C} + 0.553$.

182 Aliquots of 20 g of fine earth were used to determine the particle-size analysis after they were
183 maintained submerged in deionised water for 24 h; sand was retrieved by wet sieving at 0.053 mm,
184 while silt and clay were obtained by sedimentation. All the following analyses were performed on
185 both fine earth and skeleton. The pH values were determined potentiometrically in water after one
186 night of solid:liquid contact at 1:2.5 w:v ratio for the mineral samples and 1:8 w:v ratio for the
187 organic samples (Cardelli et al., 2019). Total organic carbon (TOC) was estimated by K-dichromate
188 digestion, heating the suspension at 180 °C for 30 minutes (Nelson and Sommers, 1996). Water-
189 extractable organic matter (WEOM) was extracted after one night of the 1:10 solid:liquid
190 suspension in an orbital shaker at 140 rpm and filtered through a Whatman 42 filters (Agnelli et al.,
191 2014). The organic C content of the extract (WEOC, water-extractable organic carbon) was
192 determined by titration (Nelson and Sommers, 1996). Total N (TN) was measured by a dry
193 combustion analyser (EA-1110, Carlo Erba Instruments, Milan, Italy), while available P (P_{av}) was
194 determined following the Olsen et al. (1954) method. Exchangeable Ca, Mg, and K were displaced
195 by a 0.2 M $BaCl_2$ solution (solid:liquid ratio 1:10) and extracted after 10 min of shaking (Corti et
196 al., 1997). The obtained suspensions were centrifuged and filtered through Whatman 42 filters.
197 Elements were determined by atomic absorption with a Shimadzu AA-6300 spectrophotometer
198 (Tokyo, Japan). For the skeletal fraction, pH, P_{av} , and exchangeable Ca, Mg, and K were
199 determined on unground fragments, while TOC, WEOC, and TN were measured on ground aliquots
200 (Ugolini et al. 1996; Corti et al., 1997; Corti et al., 2002).

201

202 *2.5. Stock calculation*

203 Soil rock fragments can contain considerable amounts of nutrients (Ugolini et al., 1996). In
204 particular, as pointed out by Corti et al. (2002) and Cuniglio et al. (2009), calcareous skeleton may
205 represent a large reservoir of C, N, and nutrient cations. Thus, considering the soil as made of fine
206 earth only may result in significant overestimations of the soil nutrient budget. Therefore, C and
207 nutrients (N, P, Ca, Mg, K) stocks were calculated for each genetic horizon taking into
208 consideration both fine earth and skeleton contributions.

209 The amount of element stored in the fine earth and skeleton was calculated as following (De Nicola
210 et al., 2014):

211

212

$$213 \quad ES = EC \bullet BD \bullet TH \bullet CC$$

(1)

214

215 where ES is the element stock (in Mg ha⁻¹ for C, N, and exchangeable Ca, Mg, and K; in kg ha⁻¹ for
216 P_{av}), EC is the element concentration (g kg⁻¹ for C and N; mg kg⁻¹ for P_{av} and exchangeable Ca, Mg,
217 and K), BD is the bulk density (kg dm⁻³), TH is the horizon thickness (cm), and CC is the
218 coefficient applied to normalize the units of measure (10⁻¹ for C, N and P_{av}; 10⁻⁴ for exchangeable
219 Ca, Mg, and K).

220 Thus, the total C and nutrient stored in each genetic horizon were determined as the weighed mean
221 for the fine earth and skeleton contents:

222

$$223 \text{ ESTOT} = [(ES_{fe} \bullet FE\%) + (ES_{sk} \bullet SK\%)] / 100 \quad (2)$$

224

225 where ES_{TOT} is the total amount of element stored in the genetic horizon (in Mg ha⁻¹ for C, N, and
226 exchangeable Ca, Mg, and K; in kg ha⁻¹ for P_{av}), ES_{fe} is the amount of element contained in the fine
227 earth, FE% is the percentage of fine earth content in the horizon, ES_{sk} is the amount of element
228 contained in the skeleton, SK% is the percentage of skeleton content in the horizon.

229 For each element, the amount stored by 1-cm thickness of the organic, organo-mineral, and mineral
230 horizons was also calculated.

231

232 2.6. Statistical analysis

233 Because of the soil variability, profiles showed slight differences in the sequence of horizons. Thus,
234 genetic horizons were grouped into soil layers based on their nature: forest floor (OL_n, OL_v, OF_r,
235 and OH horizons), organo-mineral (A and AB horizons), and mineral (B_w, B_g, BC, and Cr
236 horizons). Properties of the soil layers were obtained by calculating the weighed mean of each
237 property based upon the thickness of each horizon. The element stocks for the 0-30, 30-50, and 50-
238 75 cm of soil were calculated considering the thicknesses of the organo-mineral and mineral soil
239 horizons. To highlight differences in C and nutrient stocks, one-way ANOVA was performed along
240 the soil layers and among soils under different forest managements. Prior to ANOVA, normality
241 and homoscedasticity of the dataset were assessed using Shapiro-Wilk statistical test and by
242 Levene's test at 5% significance level, respectively. Assumptions were not violated and Tukey's
243 Honest Significant Difference (HSD) test with $P \leq 0.05$ was used to compare differences among
244 means. Results of ANOVA (*F value* and *significance level*), showing the influence of management
245 and depth on physical and chemical properties and elements stock in the surveyed soils are reported
246 in Table S5 a/b of Supplementary Materials.

247

248 3. Results

249 *3.1. Soil morphology, and physical and chemical properties in the three forest trials*

250 Properties of the experimental site were similar in the three forest trials.

251 All the soils were classified as Humustepts (Soil Survey Staff, 2014). Soil morphology organized in
252 soil layers is reported in Table 2 and Table S1 of Supplementary Material. The litter layer was on
253 average 2 to 5 cm thick and was mainly made by Turkey oak leaves and branch fragments. Organo-
254 mineral horizons showed a thickness spanning from 2 to 6 cm thick, and the soil structure was
255 moderately to well-developed, in form of crumbs or subangular blocks; in the area massively
256 frequented by wild boars (MT), the structure was platy. Mineral horizons showed poorly to well-
257 developed structure mainly made of subangular and angular blocks. Gley B horizons (Bg) indicate
258 periodical soil water saturation (Soil Survey Staff, 2014). The skeleton content in the three trials
259 ranged from 0 to 50-60%, with the greatest contents in depth (Table 2).

260 The soil pH was sub-acid (ranging between 5.69 and 6.15), with no significant difference among
261 layers and trials (Table 3). The particle-size distribution showed a coarser texture in the organo-
262 mineral horizons (loam to sandy-loam textures) than in the mineral ones (silty clay and clay loam).
263 No statistically significant difference ($P>0.05$) among the trials was observed. As expected, the
264 largest contents of TOC, WEOC, TN, and P_{av} were in the litter and showed a decreasing trend with
265 depth. Among the trials, no difference occurred for TOC, WEOC, and TN, whereas the organo-
266 mineral horizons of HT displayed the highest P_{av} concentrations. The WEOC/TOC ratio showed
267 very high values with respect to other reports (Corvasce et al., 2006; De Feudis et al., 2017), with
268 statistically significant differences only in the MT trial, where the mineral horizons displayed the
269 highest value. The C/N ratio showed a significantly decreasing trend with depth in all the trials,
270 with no significant difference among them (Table 3, S2 and S3 of Supplementary Material for fine
271 earth and skeleton data, respectively).

272

273 *3.2. C and nutrient stocks in the three forest trials*

274 In the three trials, litter and the upper 0-30 cm mineral layer contained similar quantity of C, which
275 was higher than in the 30-50 and 50-75 cm mineral layers (Table 4). Other differences among litter
276 and mineral layers were observed for TN in CTR, exchangeable Ca in HT, and K in HT and MT,
277 always with the highest stock in the 0-30 cm layer. However, the stock of all elements showed no
278 statistical difference among the trials. The contribution of the skeleton to the element stocks was
279 negligible or null for C, TN, and P_{av} , but ranged from ≈ 1.5 to $\approx 11\%$ for exchangeable K, and from
280 ≈ 27 to $\approx 63\%$ for exchangeable Ca and Mg (Table 4 and Table S4 of Supplementary Material).

281 The amount of the elements stored by 1-cm thickness of the organic, organo-mineral, and mineral
282 horizons in the three trials is reported in Fig. 2. The quantity of C, TN, and P_{av} stored in 1 cm of

283 litter was often similar to that of the organo-mineral horizons (except for P_{av} in CTR), and from
284 three- to ten-fold higher than that of the mineral horizons (except for TN in CTR, where no
285 significant difference was observed). For the exchangeable Ca, Mg, and K, the stock capacity of 1
286 cm of organo-mineral horizons was generally greater than in the mineral horizons but because the
287 samples were small (Webster, 2001), the differences were not statistically significant except for Ca
288 in MT, Mg in HT, and K in both HT and MT, where the variability was proportionally less than in
289 the other cases. Contrasting the stock capacity per 1-cm thickness among the different trials, only
290 P_{av} and exchangeable Mg showed significant differences, with the highest contents of the two
291 elements in the organo-mineral horizons of HT (Table 3).

292

293 **4. Discussion**

294 *4.1. Effect of thinning on soil morphology and physicochemical properties*

295 The effect of thinning on the main pedological features appeared negligible in the studied forest. In
296 the topsoil, which is the soil portion most sensitive to disturbances and management practices (Song
297 et al., 2005), the effect of thinning could have been masked by the wild boar activity. In our case,
298 the topsoil mixing due to wild boars seemed to have not substantially affected the soil morphology,
299 probably because all trials have been characterized over time by their presence, albeit with different
300 intensities.

301 Although the soils developed from calcareous parent material, the soil profiles displayed sub-acid
302 pH values, indicating that soils have been subjected to a heavy decarbonation induced by several
303 acidification processes (Haynes, 1990; Richter et al., 2007; Lemanceau et al., 2009; Chapin et al.,
304 2011; Cocco et al., 2013; Corti et al., 2019). Because of this, and the long time needed to dissolve
305 all carbonates (Cocco et al., 2013), these soils can be considered as highly weathered (e.g.,
306 Sundquist and Visser, 2003) and, consequently, it was not expected that the forest thinning could
307 induce marked changes on soil pH in 50 years. Moreover, even when carbonates have been
308 dissolved, acidification is buffered by clay and organic matter (Brady and Weil, 2017), which
309 contribute to reduce pH changes. The soil texture is a parameter not responding quickly to
310 environmental changes; in fact, it is similar for all the trials. Along the profiles, the texture was finer
311 at depth than at the surface probably for the occurrence of lessivage, a process that requires long
312 time to produce differences in terms of soil texture and drainage (Buurman et al., 1998; Quénard et
313 al., 2011; Calabrese et al., 2018).

314 The decreasing of TOC, TN, and P_{av} with depth is a common trend in soil and especially in forest
315 soils, where the majority of the biomass produced is added in form of litter (Mason and Zanner,
316 2005). While TOC and TN were not affected by thinning, as they respond slowly to changes (Bai et
317 al., 2017), the different P_{av} content in the organo-mineral and mineral horizons of the three trials

318 were considered an effect of the forest thinning. In fact, working on soils under *Fagus sylvatica*
319 forests, Cardelli et al. (2019) reported that P liberation and activity of enzymes involved in the P
320 cycle are higher in the organo-mineral (A) than in the organic (O) horizons because of the major
321 content of decaying SOM and the consequently greater availability of P-bearing substances like
322 nucleic acids, carbohydrates, proteins, and fatty acids. The major reduction of the canopy density in
323 HT might have enhanced SOM degradation through the increased solar radiation and temperature
324 (e.g., Gressel et al., 1996; Scharenbroch and Bockeim, 2007; Cheng et al., 2021)), with the
325 subsequent higher release of P.

326 The WEOM content and the WEOM/TOC ratio did not change among the trials. This behaviour
327 was unexpected because WEOM, which is composed of easily degradable molecules that represent
328 the main C and energy source for the soil microbial community (De Feudis et al., 2019), is
329 considered as an indicator of microbial activity (Gutiérrez-Girón et al., 2015), very sensitive to
330 disturbances and management (Chantigny, 2003). However, since the WEOM is released following
331 SOM mineralization (Bartos et al., 2020) and the two thinning intensities did not produce different
332 litter thicknesses (Table 2) and soil TOC and TN concentrations (Table 3) in respect to the control,
333 it would justify the similar WEOC contents and WEOC/TOC ratios found among the trials.
334 Throughout the profiles, the decreasing content of WEOC and the parallel increase of WEOC/TOC
335 ratio (statistically significant only in MT) confirmed the importance of this soluble fraction as
336 energetic substrate for the organisms harbouring the deeper soil horizons. The increase of the
337 WEOC/TOC ratio in the mineral layers, where the clay content is the highest, could be also due to
338 adsorption of organics on the clay mineral lattices, with the formation of mobile organo-mineral
339 complexes (Corvasce et al., 2006).

340 The magnitude of the values of the C/N ratio for the three trials agreed with those reported in other
341 studies conducted on Mediterranean forests (Corral-Fernández et al., 2013; Cools et al., 2014). The
342 decreasing trend of the C/N ratio with depth is a common trend in forest soils, where the litter is
343 made by less degraded (and with higher C/N ratio) biomass than the organic molecules translocated
344 into the deeper soil horizons after SOM decaying (Marinari et al., 2021).

345

346 4.2. *Effect of thinning on the stocks of C and nutrients*

347 In all the trials, SOC stock is similar in both litter (with a general thickness of 6-10 cm) and 0-30
348 cm layer, while the SOC stored below 30 cm depth amounted to 54-69% of that in the upper layer.
349 The large amount of C stored in the sub-superficial mineral soil has a basic ecological relevance due
350 to the role of forest soils as C sink and, hence, in the climate change mitigation. In the mineral
351 layers, SOM is stabilized and protected from degradation (e.g., Ono et al., 2013; Yao et al., 2019)

352 mostly because of clay minerals associations and oxygen limitation (e.g., Wattel-Koekkoek et al.,
353 2003; Kleber, 2010). As a consequence, with increasing depth the C turnover rate slows down and
354 the mean residence time of SOM (e.g., Trumbore, 2000; Wang and Chang, 2001) and decaying
355 roots (e.g., Agnelli et al., 2014) tend to increase.

356 Regarding the different stocks observed along the depth for TN (in CTR) and exchangeable Ca (in
357 HT) and K (in HT and MT), they appeared not related with the amount of roots or with the presence
358 of leguminous species in the understorey that could have enriched the soil of N (Table S1 of
359 Supplementary Materials). The knowledge on the influence of thinning on the stock of elements in
360 the different layers along the soil depth is scarce but, working on a multi-centennial holm oak
361 (*Quercus ilex* L.) forest in a pedoclimatic condition similar to that of our trials, Agnelli et al. (2016)
362 found that many soil features, especially those not directly linked to the microbial activity, were
363 rather homogeneous for each soil depth because of the long lasting pedogenesis. Since our trials
364 were established under a forest cover as old as at least three millennia, we believe that, especially
365 for exchangeable Ca and K, differences derived from spatial differences of parent material and
366 skeleton content rather than to thinning experimentation started ≈ 50 years before this study.

367 The three trials showed no significant effect for none of the four (in case of C, N, and P_{av} stocks) or
368 three (for exchangeable Ca, Mg, and K) layers considered. Thinning has been reported not to be
369 able to produce changes on the organic C (and N) stock in naturally settled broadleaves and conifers
370 stands (e.g., Bravo-Oviedo et al., 2015; Bai et al., 2017; Prasad Dangal et al., 2017; Zhang et al.,
371 2018; Mayer et al., 2020). Opposite results were found in planted forests. For example, in their
372 review Gong et al. (2021) took into consideration 77 articles on the effect of forest thinning on SOC
373 stocks in the 0–30 cm mineral soil thickness across planted forests in China and concluded that a
374 moderate thinning significantly increased SOC stocks with respect to both no-thinning and heavy
375 thinning. Instead, working in a *Picea crassifolia* Kom. plantations, He et al. (2018) observed a
376 decrease of the C stock with increasing thinning intensity, with a parallel increase of soil water
377 storage. These reports reinforced the hypothesis that thinning cannot affect the stock of C and other
378 elements in soils with long forest cover history, where pedogenesis has heavily homogenized the
379 soil profile.

380 The 1-cm stock values confirmed that the organic and the organo-mineral horizons are able to stock
381 similar amounts of C, TN, and P_{av} , while more in depth this ability is minor. While the
382 concentration of C and TN in the organic and organo-mineral horizons was ascribed to their
383 richness of SOM, the large concentration of P_{av} was attributed to the degradation of SOM, which
384 released P from the organic structures (e.g., Pistocchi et al., 2018; Ni et al., 2021). With respect to 1
385 cm of mineral horizons, the ability of 1 cm of organo-mineral horizons to stock nutrient cations was

386 statistically significant only for Ca in MT, Mg in HT, and K in HT and MT, but a generalized trend
387 was observed in all cases, even though the differences were not significant because the samples
388 were relatively small. In all these cases, the differences were attributed to the relatively fast SOM
389 degradation occurring in the organo-mineral horizons (e.g., Pistocchi et al., 2018; Wang et al.,
390 2019).

391 When contrasting the 1-cm stock capacity among the trials, thinning appeared to have an effect only
392 on P_{av} and exchangeable Mg, which assumed the highest values in the organo-mineral horizons of
393 HT. Also in this case, although not significant, the same appeared true at least for the potentially
394 available Ca, while for the exchangeable K differences were probably disturbed by spatial
395 variability of parent material and skeleton contribution. However, since a more intense thinning is
396 expected to induce a diffuse higher soil water storage because of the resulted lower canopy density
397 (He et al., 2018), it is probable that a larger water availability in the organic and organo-mineral
398 horizons favoured a greater SOM mineralization with consequent release of nutrients (e.g.,
399 Vesterdal et al., 1995; Prescott, 2002; Chiti et al., 2015; Gross et al., 2018).

400

401 **5. Conclusions**

402 The soil physicochemical parameters and the stock of C and nutrients in the litter and in the 0-30,
403 30-50, and 50-75 cm layers under a multi-millennial Turkey oak forest cover, appeared slightly
404 influenced by thinnings operated along the last 50 years. This result, which contradicts our research
405 hypothesis, was achieved considering the contribution of the skeletal fraction that, especially in
406 depth, was present in a considerable amount. The only parameters that appeared to be more affected
407 by thinning were P_{av} and exchangeable Mg. The more intense thinning was able to increase the 1-
408 cm storage of the organo-mineral horizons via a major SOM mineralization. Our results contrast
409 with those reported for recently (decades) planted forests, especially if plantation occurs in former
410 cultivated fields, where thinning has tangible effects on element storage. This is equivalent to
411 saying that, after about three millennia of Turkey oak forest use, both forest cover and human
412 activity are the main soil forming forces.

413 Our study also showed that organo-mineral and mineral horizons under the Turkey oak forest are
414 able to store an amount of SOM similar to the litter layers. Since the SOM contained into the
415 organo-mineral and mineral horizons has higher recalcitrance and, consequently, is less involved in
416 the C turnover processes than that of the forest floor, it is mandatory to adopt forest managements
417 strategies able to increase SOM in depth rather than in the superficial organic horizons, to affect
418 positively the global C cycle.

419 Finally, considering that *i*) coppice stands under conversion into high forest via natural evolution
420 and by means of periodical thinnings appeared to be equal as for soil ecosystem properties 50 years
421 later, and that *ii*) the latter option is more profitable for environmental, socio-economic issues and
422 recreational purposes, thinning implementation can be considered as a valuable solution, among the
423 different and complementary strategies on the floor, to manage nowadays the original coppice area.

424

425 **Acknowledgements**

426 We are indebted to Leonardo Tonveronachi for his help during the field activities.

427

428 **Funding:** This work was supported by European project FutureForCoppices LIFE14
429 ENV/IT/000514.

430

431 **References**

432 Agnelli, A., Massaccesi, L., De Feudis, M., Cocco, S., Courchesne, F., Corti, G., 2016. Holm oak
433 (*Quercus ilex* L.) rhizosphere affects limestone-derived soil under a multi-centennial forest. *Plant*
434 *and soil*. 400, 297-314. <https://doi.org/10.1007/s11104-015-2732-x>

435 Agnelli, A., Bol, R., Trumbore, S.E., Dixon, L., Cocco, S., Corti, G., 2014. Carbon and nitrogen in
436 soil and vine roots in harrowed and grass-covered vineyards. *Agric. Ecosyst. Environ.* 193, 70–
437 82. <https://doi.org/10.1016/j.agee.2014.04.023>

438 Bai, S.H., Dempsey, R., Reverchon, F., Blumfield, T. J., Ryan, S., Cernusak, L. A., 2017. Effects of
439 forest thinning on soil-plant carbon and nitrogen dynamics. *Plant Soil* 411, 437–449.
440 <https://doi.org/10.1007/s11104-016-3052-5>

441 Baize, D., Girard, M.C., 2008. *Référentiel Pedologique*, Edition Quæ. Association française pour
442 l'étude du sol (Afes), Versailles, France.

443 Barré, P., Fernandez-Ugalde, O., Virto, I., Velde, B., Chenu, C., 2014. Impact of phyllosilicate
444 mineralogy on organic carbon stabilization in soils: Incomplete knowledge and exciting
445 prospects. *Geoderma* 235–236, 382–395. <https://doi.org/10.1016/j.geoderma.2014.07.029>

446 Bartos, A., Szymański, W., Klimek, M., 2020. Impact of conventional agriculture on the
447 concentration and quality of water-extractable organic matter (WEOM) in the surface horizons
448 of Retisols—A case study from the Carpathian Foothills in Poland. *Soil and Tillage Research*,
449 204. <https://doi.org/10.1016/j.still.2020.104750>.

450 Batjes, N.H., 1996. Total carbon and nitrogen in the soils of the world. *Eur. J. Soil Sci.*
451 https://doi.org/10.1111/ejss.12114_2

452 Bot, A., Benites, J., 2005. The Importance of Soil Organic Matter: Key to Drought-resistant Soil

453 and Sustained Food Production, ed. Food and Agriculture Organization of the United Nation
454 (FAO), Rome.

455 Brady, N.C., Weil, R.R., 2017. The Nature and Properties of Soils. Fifteenth ed. Upper Saddle
456 River, New York. <https://doi.org/10.2136/sssaj2016.0005br>

457 Bravo-Oviedo, A., Ruiz-Peinado, R., Modrego, P., Alonso, R., Montero, G., 2015. Forest thinning
458 impact on carbon stock and soil condition in Southern European populations of *P. sylvestris* L.
459 For. Ecol. Manag. 357, 259–267. <https://doi.org/10.1016/j.foreco.2015.08.005>

460 Brenna, S., Rocca, A., Sciacaluga, M., Valagussa, M., 2010. Sistema di Monitoraggio della Qualità
461 dei Suoli di Lombardia, Sperimentazione condotta nell'ambito del progetto di ricerca n. 1032
462 “Sistema di monitoraggio della qualità dei suoli di Lombardia”. “Foto a cura di: Marco
463 Sciacaluga, Alberto Rocca. - SOILQUALIMON” (d.g.r. 22 marzo 2006 n. VIII/2182 –
464 Programma di attività 2006). Quaderni della Ricerca n. 110 - maggio 2010. Regione Lombardia -
465 Direzione Generale Agricoltura U.O. – Ente regionale per i servizi all'agricoltura e alle foreste.

466 Buurman, P., Jongmans, A.G., PiPujol, M.D., 1998. Clay illuviation and mechanical clay
467 infiltration — Is there a difference? Quat. Int. 51–52, 66–69. [https://doi.org/10.1016/S1040-
468 6182\(98\)90225-7](https://doi.org/10.1016/S1040-6182(98)90225-7)

469 Caddeo, A., Marras, S., Sallustio, L., Spano, D., Sirca, C., 2019. Soil organic carbon in Italian
470 forests and agroecosystems: Estimating current stock and future changes with a spatial modelling
471 approach. Agric. For. Meteorol. 278, 107654.
472 <https://doi.org/10.1016/J.AGRFORMET.2019.107654>

473 Calabrese, S., Richter, D. D., & Porporato, A. M., 2018. The formation of clay-enriched horizons by
474 leaching. Geophysical Research Letters, 45, 7588– 7595.
475 <https://doi.org/10.1029/2018GL078778>

476 Canedoli, C., Ferrè, C., Abu El Khair, D., Comolli, R., Liga, C., Mazzucchelli, F., Proietto, A.,
477 Rota, N., Colombo, G., Bassano, B., Viterbi, R., Padoa-Schioppa, E., 2020. Evaluation of
478 ecosystem services in a protected mountain area: Soil organic carbon stock and biodiversity in
479 alpine forests and grasslands. Ecosyst. Serv. 44, 101135.
480 <https://doi.org/10.1016/j.ecoser.2020.101135>

481 Cardelli, V., De Feudis, M., Fornasier, F., Massaccesi, L., Cocco, S., Agnelli, A., Weindorf, D.C.,
482 Corti, G., 2019. Changes of topsoil under *Fagus sylvatica* along a small latitudinal-altitudinal
483 gradient. Geoderma 344, 164–178. <https://doi.org/10.1016/J.GEODERMA.2019.01.043>

484 Carrari, E., Ampoorter, E., Bottalico, F., Chirici, G., Coppi, A., Travaglini, D., Verheyen, K., Selvi,
485 F., 2017. The old charcoal kiln sites in Central Italian forest landscapes. Quat. Int. 458, 214–223.
486 <https://doi.org/10.1016/j.quaint.2016.10.027>

487 Cartocci, A., Fedi, M.E., Taccetti, F., Benvenuti, M., Chiarantini, L., Guideri, S., 2007. Study of a
488 metallurgical site in Tuscany (Italy) by radiocarbon dating. Nucl. Instruments Methods Phys.
489 Res. Sect. B Beam Interact. with Mater. Atoms 259, 384–387.
490 <https://doi.org/10.1016/j.nimb.2007.01.183>

491 Chantigny, M. H., 2003. Dissolved and water-extractable organic matter in soils: a review on the
492 influence of land use and management practices, Geoderma 113, issues 3–4, 357-380.
493 [https://doi.org/10.1016/S0016-7061\(02\)00370-1](https://doi.org/10.1016/S0016-7061(02)00370-1).

494 Chapin, F.S. III; Matson, P.A., Mooney, H.A., 2011. Principles of Terrestrial Ecosystem Ecology.
495 Springer, Heidelberg, Germany.

496 Cheng, X., Han, H., Zhu, J., Peng, X., Li, B., Liu, H., Epstein, H.E., 2021. Forest thinning and
497 organic matter manipulation drives changes in soil respiration in a *Larix principis-rupprechtii*
498 plantation in China. Soil Tillage Res. 211, 104996. <https://doi.org/10.1016/j.still.2021.104996>

499 Chiarantini, L., Benvenuti, M., Costagliola, P., Dini, A., Firmati, M., Guideri, S., Villa, I.M.,
500 Corretti, A., 2018. Copper metallurgy in ancient Etruria (southern Tuscany, Italy) at the Bronze-
501 Iron Age transition: a lead isotope provenance study. J. Archaeol. Sci. Reports 19, 11–23.
502 <https://doi.org/10.1016/j.jasrep.2018.02.005>

503 Chiti T., Sirca C., Rodeghiero M., Spano D., Valentini R. (2015) Soil Carbon Stocks and Fluxes. In:
504 Valentini R., Miglietta F. (eds) The Greenhouse Gas Balance of Italy. Environmental Science
505 and Engineering. Springer, Berlin, Heidelberg. https://doi.org/10.1007/978-3-642-32424-6_8

506 Cocco, S., Agnelli, A., Gobran, G.R., Corti, G., 2013. Changes induced by the roots of *Erica*
507 *arborea* L. to create a suitable environment in a soil developed from alkaline and fine-textured
508 marine sediments. Plant Soil 368, 297–313. <https://doi.org/10.1007/s11104-012-1501-3>

509 Cools, N., Vesterdal, L., De Vos, B., Vanguelova, E., Hansen, K., 2014. Tree species is the major
510 factor explaining C: N ratios in European forest soils. For. Ecol. Manage. 311, 3–16.
511 <https://doi.org/10.1016/j.foreco.2013.06.047>

512 Corral-Fernández, R., Parras-Alcántara, L., Lozano-García, B., 2013. Stratification ratio of soil
513 organic C, N and C:N in Mediterranean evergreen oak woodland with conventional and organic
514 tillage. Agric. Ecosyst. Environ. 164, 252–259. <https://doi.org/10.1016/J.AGEE.2012.11.002>

515 Corti, G., Agnelli, A., Cocco, S., Cardelli, V., Masse, J., Courchesne, F., 2019. Soil affects
516 throughfall and stemflow under Turkey oak (*Quercus cerris* L.). Geoderma 333, 43–56.
517 <https://doi.org/10.1016/J.GEODERMA.2018.07.010>

518 Corti, G., Agnelli, A., Ugolini, F.C., 1997. Release of Al by hydroxy-interlayered vermiculite and
519 hydroxy-interlayered smectite during determination of cation exchange capacity in fine earth and
520 rock fragments fractions. Eur. J. Soil Sci. 48, 249–262. <https://doi.org/10.1111/j.1365->

521 2389.1997.tb00545.x

522 Corti, G., Ugolini, F.C., Agnelli, A., 1998. Classing the soil skeleton (greater than two millimeters):
523 Proposed ppproach and procedure. *Soil Sci. Soc. Am. J.* 62, 1620–1629.
524 <https://doi.org/10.2136/sssaj1998.03615995006200060020x>

525 Corti, G., Ugolini, F.C., Agnelli, A., Certini, G., Cuniglio, R., Berna, F., Fernández Sanjurjo, M.J.,
526 2002. The soil skeleton, a forgotten pool of carbon and nitrogen in soil. *Eur. J. Soil Sci.* 53, 283–
527 298. <https://doi.org/10.1046/j.1365-2389.2002.00442.x>

528 Corvasce, M., Zsolnay, A., D’Orazio, V., Lopez, R., Miano, T.M., 2006. Characterization of water
529 extractable organic matter in a deep soil profile. *Chemosphere* 62, 1583–1590.
530 <https://doi.org/10.1016/J.CHEMOSPHERE.2005.07.065>

531 Cuniglio, R., Corti, G., Agnelli, A., 2009. Rock fragments evolution and nutrients release in
532 vineyard soils developed on a thinly layered limestone (Tuscany, Italy). *Geoderma* 148, 375–
533 383. <https://doi.org/10.1016/J.GEODERMA.2008.11.005>

534 Cutini A., 1996. The influence of drought and thinning on leaf area index estimates from canopy
535 transmittance method. *Annales des Sciences Forestieres* 53 (2-3): 595 – 603.

536 Cutini, A., Ferretti, M., Bertini, G., Brunialti, G., Bagella, S., Chianucci, F., Fabbio, G., Fratini, R.,
537 Riccioli, F., Caddeo, C., Calderisi, M., Chiucci, B., Corradini, S., Cristofolini, F., Di Salvatore,
538 U., Ferrara, C., Frati, L., Landi, S., Marchino, L., Patteri, G., Piovosi, M., Roggero, P.P.,
539 Seddaiu, G., Gottardini, E., 2021. Testing an expanded set of sustainable forest management
540 indicators in Mediterranean coppice area. *Ecological Indicators.* 130, 108040.
541 <https://doi.org/10.1016/j.ecolind.2021.108040>

542 De Feudis, M., Cardelli, V., Massaccesi, L., Hofmann, D., Berns, A.E., Bol, R., Cocco, S., Corti,
543 G., Agnelli, A., 2017. Altitude affects the quality of the water-extractable organic matter
544 (WEOM) from rhizosphere and bulk soil in European beech forests. *Geoderma* 302, 6–13.
545 <https://doi.org/10.1016/j.geoderma.2017.04.015>

546 De Feudis, M., Cardelli, V., Massaccesi, L., Trumbore, S.E., Vittori Antisari, L., Cocco, S., Corti,
547 G., Agnelli, A., 2019. Small altitudinal change and rhizosphere affect the SOM light fractions
548 but not the heavy fraction in European beech forest soil. *Catena* 181, 104091.
549 <https://doi.org/10.1016/j.catena.2019.104091>

550 De Nicola, C., Zanella, A., Testi, A., Fanelli, G., Pignatti, S., 2014. Humus forms in a
551 Mediterranean area (Castelporziano Reserve, Rome, Italy): classification, functioning and
552 organic carbon storage. *Geoderma* 235–236, 90–99.
553 <https://doi.org/10.1016/j.geoderma.2014.06.033>.

554 Diao, M., Yang, K., Zhu, J., Li, M., Xu, S., 2020. Native broad-leaved tree species play key roles on

555 maintaining soil chemical and microbial properties in a temperate secondary forest, Northeast
556 China. *Forest Ecology and Management*. 462, 117971.
557 <https://doi.org/10.1016/j.foreco.2020.117971>

558 Fabbio, G., 2016. Coppice forests, or the changeable aspect of things, a review. *Annals of*
559 *Silvicultural Research*. 40, 108-132. <http://dx.doi.org/10.12899/asr-1286>

560 Fabbio, G., Amorini, E., 2006. Avviamento ad altofusto e dinamica naturale nei cedui a prevalenza
561 di cerro. Risultati di una prova sperimentale a 35 anni dalla sua impostazione. Il protocollo di
562 Caselli (Pisa). *Ann. Ist. Sper. di Selvic.* 33, 79–104. [https://journals-](https://journals-crea.4science.it/index.php/asr/issue/viewIssue/154/8)
563 [crea.4science.it/index.php/asr/issue/viewIssue/154/8](https://journals-crea.4science.it/index.php/asr/issue/viewIssue/154/8)

564 Fabbio, G, Cutini, A, 2017. Il ceduo oggi: quale gestione oltre le definizioni? *Forest@ - Journal of*
565 *Silviculture and Forest Ecology*, Volume 14, Pages 257-274. [https://doi.org/10.3832/efor2562-](https://doi.org/10.3832/efor2562-014)
566 014

567 Frati, L., Brunialti, G., Landi, S., Filigheddu, R., Bagella, S., 2021. Exploring the biodiversity of
568 key groups in coppice forests (Central Italy): the relationship among vascular plants, epiphytic
569 lichens, and wood-decaying fungi. *Plant Biosystems*, 1-12.

570 Garlato, A., Obber, S., Vinci, I., Mancabelli, A., Parisi, A., Sartori, G., 2009a. La determinazione
571 dello stock di carbonio nei suoli del Trentino a partire dalla banca dati della carta dei suoli alla
572 scala 1:250.000. *Museo Tridentino di Scienze Naturali. Trento Studi Trent Sci Nat* 85, 157–160.

573 Garlato, A., Obber, S., Vinci, I., Sartori, G., Manni, G., 2009b. Stock attuale di carbonio organico
574 nei suoli di montagna del Veneto. *Museo Tridentino di Scienze Naturali. Trento Studi Trent Sci*
575 *Nat* 85, 69–8.

576 Gartzia-Bengoetxea, N., Virto, I., Arias-González, A., Enrique, A., Fernández-Ugalde, O., Barré,
577 P., 2020. Mineral control of organic carbon storage in acid temperate forest soils in the Basque
578 Country. *Geoderma* 358, 113998. <https://doi.org/10.1016/j.geoderma.2019.113998>

579 Gong, C., Tan, Q., Liu, G., Xu, M., 2021. Forest thinning increases soil carbon stocks in China.
580 *Forest Ecology and Management* 482, 118812. <https://doi.org/10.1016/j.foreco.2020.118812>

581 Gressel, N., McColl, J.G., Preston, C.M., Newman, R.H., Powers, R.F., 1996. Linkages between
582 phosphorus transformations and carbon decomposition in a forest soil. *Biogeochemistry* 33, 97–
583 123. <https://doi.org/10.1007/BF02181034>

584 Gross, C.D., James, J.N., Turnblom, E.C., Harrison, R.B., 2018. Thinning treatments reduce deep
585 soil carbon and nitrogen stocks in a Coastal Pacific Northwest forest. *Forests* 9, 238.
586 <https://doi.org/10.3390/f9050238>

587 Guermandi, M., Marchi, N., Tarocco, P., Calzolari, C., Ungaro, F., Villani, I., 2013. Siti locali
588 rappresentativi dei suoli della pianura e della collina Emiliano-Romagnola. In: *Emilia-Romagna,*

589 Regione (Ed.), Servizio Geologico, Sismico e dei Suoli. IRPI CNR, Provincia di Ferrara (41 pp.).

590 Gutiérrez-Girón, A., Díaz-Pinés, E., Rubio, A., Gavilán, R.G., 2015. Both altitude and vegetation
591 affect temperature sensitivity of soil organic matter decomposition in Mediterranean high
592 mountain soils. *Geoderma* 237–238, 1–8. <https://doi.org/10.1016/j.geoderma.2014.08.005>

593 Haynes, R.J. 1990. Active ion uptake and maintenance of cation-anion balance: A critical
594 examination of their role in regulating rhizosphere pH. *Plant Soil*, 126: 247–264.

595 He, Z., Fang, S., Chen, L., Du, J., Zhu, X., Lin, P., 2018. Spatial patterns in natural *Picea*
596 *crassifolia* forests of northwestern China, as basis for close-to-nature forestry. *Journal of*
597 *Mountain Science* 15, 1909-1919. <https://doi.org/10.1007/s11629-016-3998-z>

598 Hölscher, D., Schade, E., Leuschner, C., 2001. Effects of coppicing in temperate deciduous forests
599 on ecosystem nutrient pools and soil fertility. *Basic Appl. Ecol.* 2, 155–164.
600 <https://doi.org/10.1078/1439-1791-00046>

601 Jenny, H., 1941. *Factors of Soil Formation*. McGraw-Hill.

602 Kaiser, M., Zederer, D.P., Ellerbrock, R.H., Sommer, M., Ludwig, B., 2016. Effects of mineral
603 characteristics on content, composition, and stability of organic matter fractions separated from
604 seven forest topsoils of different pedogenesis. *Geoderma* 263, 1–7.
605 <https://doi.org/10.1016/j.geoderma.2015.08.029>

606 Kleber M., 2010. What is recalcitrant soil organic matter?. *Environmental Chemistry* 7, 320-332.
607 <https://doi.org/10.1071/EN10006>

608 Kögel-Knabner, I., Ekschmitt, K., Flessa, H., Guggenberger, G., Matzner, E., Marschner, B. and
609 von Lütow, M., 2008. An integrative approach of organic matter stabilization in temperate soils:
610 Linking chemistry, physics, and biology. *Z. Pflanzenernähr. Bodenk.*, 171, 5-13.
611 <https://doi.org/10.1002/jpln.200700215>

612 Lee, J.H., Lee, J.G., Jeong, S.T., Gwon, H.S., Kim, P.J., Kim, G.W., 2020. Straw recycling in rice
613 paddy: Trade-off between greenhouse gas emission and soil carbon stock increase. *Soil Tillage*
614 *Res.* 199, 104598. <https://doi.org/10.1016/J.STILL.2020.104598>

615 Lehmann, J., Kleber, M., 2015. The contentious nature of soil organic matter. *Nature* 528, 60-68.
616 <https://doi.org/10.1038/nature16069>

617 Lemanceau, P., Bauer, P., Kraemer, S., Briat, J.-F., 2009. Iron dynamics in the rhizosphere as a case
618 study for analyzing interactions between soils, plants and microbes. *Plant Soil* 321, 513–535.
619 <https://doi.org/10.1007/s11104-009-0039-5>

620 Mairota, P., Neri, F., Travaglini, D., Picchio, R., Terzuolo, P. G., Piussi P., Marchi, M., 2018.
621 Chapter 6 “Thirty-Five Countries”. In A. Unrau, G. Becker, R. Spinelli, D. Lazdina, N.
622 Magagnotti, V.N. Nicolescu, P. Buckley, D. Bartlett, P.D. Kofman (Eds.), *Coppice Forests in*

623 Europe (pp. 269-282). Freiburg i. Br., Germany: Albert Ludwig University of Freiburg.

624 Manetti, M.C., Becagli, C., Bertini, G., Cantiani, P., Marchi, M., Pelleri, F., Sansone, D., Fabbio,
625 G., 2020. The conversion into high forest of Turkey oak coppice stands: methods, silviculture
626 and perspectives. *iForest* 13: 309-317. - doi: 10.3832/ifor3483-013

627 Manetti, M.C., Gugliotta, O.I., 2006. Effetto del trattamento di avviamento ad altofusto sulla
628 diversità specifica e strutturale delle specie legnose in un ceduo di cerro. *Ann. Ist. Sper. di*
629 *Selvic.* 33, 105-114. <https://journals-crea.4science.it/index.php/asr/issue/viewIssue/154/8>

630 Marchi, E., Picchio, R., Mederski, P.S., Vusić, D., Perugini, M., Venanzi, R., 2016. Impact of
631 silvicultural treatment and forest operation on soil and regeneration in Mediterranean Turkey oak
632 (*Quercus cerris* L.) coppice with standards. *Ecol. Eng.* 95, 475–484.
633 <https://doi.org/10.1016/j.ecoleng.2016.06.084>

634 Marinari, S., Marabottinia, R., Falsone, G., Vianello, G., Vittori Antisari, L., Agnelli, A.,
635 Massaccesi, L., Cocco, S., Cardelli, V., Serrani, D., Corti, G., 2021. Mineral weathering and
636 leaching affect microbial community and enzyme activity in mountain soils. *Appl. Soil Ecol.*
637 167, 104024. <https://doi.org/10.1016/j.apsoil.2021.104024>

638 Mason, J.A., Zanner, C.W., 2005. Grassland Soils. *Encycl. Soils Environ.* 4, 138–145.
639 <https://doi.org/10.1016/B0-12-348530-4/00028-X>

640 Mayer, M., Prescott, C.E., Abaker, W.E.A., Augusto, L., Cécillon, L., Ferreira, G.W.D., James, J.,
641 Jandl, R., Katzensteiner, K., Laclau, J.P., Laganière, J., Nouvellon, Y., Paré, D., Stanturf, J.A.,
642 Vanguelova, E.I., Vesterdal, L., 2020. Tamm review: Influence of forest management activities
643 on soil organic carbon stocks: A knowledge synthesis. *For. Ecol. Manage.* 466, 118127.
644 <https://doi.org/https://doi.org/10.1016/j.foreco.2020.118127>

645 Nelson, D.W., Sommers, L.E., 1996. Total Carbon, Organic Carbon, and Organic Matter, in Sparks,
646 D., Page, A., Helmke, P., Loeppert, R.H., Soltanpour, P.N. Tabatabai, M.A., Johnston, C.T.,
647 Sumner, M.E. (Eds.), *Methods of Soil Analysis. SSSA Book Series.*
648 <https://doi.org/10.2136/sssabookser5.3.c34>

649 Ni, X., Lin, C., Chen, G., Xie, J., Yang, Z., Liu, X., Xiong, D., Xu, C., Yue, K., Wu, F., Yang, Y.,
650 2021. Decline in nutrient inputs from litterfall following forest plantation in subtropical China.
651 *Forest Ecol. and Manag.*, 496, 119445. <https://doi.org/10.1016/j.foreco.2021.119445>.

652 Olsen, S.R., Cole, C.V., Watanabe, F.S. and Dean, L.A. 1954. Estimation of available phosphorus
653 in soils by extraction with sodium bicarbonate, USDA. Circular 939

654 Ono, K., Hiradate, S., Morita S., Hirai, K., 2013. Fate of organic carbon during decomposition of
655 different litter types in Japan. *Biogeochem.* 112, 7-12. <https://doi.org/10.1007/s10533-011-9682->
656 z

657 Pistocchi, C., Mészáros, É., Tamburini, F., Frossard, E., Bünemann, E.K., 2018. Biological
658 processes dominate phosphorus dynamics under low phosphorus availability in organic horizons
659 of temperate forest soils. *Soil Biol. Biochem.* 126, 64–75.
660 <https://doi.org/10.1016/j.soilbio.2018.08.013>

661 Prasad Dangal, S., Kumar Das, A., Shyam Krishna, P., 2017. Effectiveness of management
662 interventions on forest carbon stock in planted forests in Nepal. *J. Environ. Manage.*
663 <https://doi.org/10.1016/j.jenvman.2017.03.056>

664 Prescott, C.E., 2002. The influence of the forest canopy on nutrient cycling. *Tree Physiol.* 22, 1193–
665 1200. <https://doi.org/10.1093/treephys/22.15-16.1193>

666 Quénard, L., Samouëlian, A., Laroche, B., Cornu, S., 2011. Lessivage as a major process of soil
667 formation: A revisit of existing data. *Geoderma* 167–168, 135–147.
668 <https://doi.org/10.1016/J.GEODERMA.2011.07.031>

669 Richter, D. de B., Oh, N.H., Fimmen, R., Jackson, J., 2007. The Rhizosphere and Soil Formation.
670 *Rhizosph.* 179-IN2. <https://doi.org/10.1016/B978-012088775-0/50010-0>

671 Scharenbroch, B.C., Bockheim, J.G., 2007. Impacts of forest gaps on soil properties and processes
672 in old growth northern hardwood-hemlock forests. *Plant and Soil.* 294, 219–233.
673 <https://doi.org/10.1007/s11104-007-9248-y>

674 Schoeneberger, P.J., Wysocki, D.A., Benham, E.C., Soil Survey Staff., 2012. *Field Book for*
675 *Describing and Sampling Soils, Version 3.0.* <https://doi.org/10.1038/258254a0>

676 Schrumpf, M., Kaiser, K., Guggenberger, G., Persson, T., Kögel-Knabner, I., Schulze, E.D., 2013.
677 Storage and stability of organic carbon in soils as related to depth, occlusion within aggregates,
678 and attachment to minerals. *Biogeosciences* 10, 1675–1691. <https://doi.org/10.5194/bg-10-1675->

679 Society of American Foresters (SAF), 2008. *Dictionary of Forestry.* Retrieved 1 February 2015.
680 http://dictionaryofforestry.org/dict/term/forest_management

681 Soil Survey Staff, 2014. *Keys to Soil Taxonomy, 12th edition.* USDA–Natural Resources
682 Conservation Service, Washington.

683 Song, G., Li, L., Pan, G., Zhang, Q., 2005. Topsoil organic carbon storage of China and its loss by
684 cultivation. *Biogeochemistry* 74, 47–62. <https://doi.org/10.1007/s10533-004-2222-3>

685 Sundquist, E.T., Visser, K., 2003. 8.09 - The Geologic History of the Carbon Cycle. *Treatise on*
686 *Geochemistry.* Pergamon (Ed.), 8-9, 425–472. US Geological Survey, Woods Hole, MA, United
687 States. <https://doi.org/10.1021/j100244a025>

688 Tiessen, H., Cuevas, E., Chacon, P., 1994. The role of soil organic matter in sustaining soil fertility.
689 *Nature* 371, 783–785. <https://doi.org/10.1038/371783a0>

690 Trumbore, S. 2000. Age of soil organic matter and soil respiration: radiocarbon constraints on

691 belowground C dynamics. *Belowground Processes and Global Change* 10, 399-411.

692 Ugolini, F.C., Corti, G., Agnelli, A., Piccardi, F., 1996. Mineralogical, physical, and chemical
693 properties of rock fragments in soil. *Soil Sci.* 161, 521-542. [https://doi.org/10.1097/00010694-](https://doi.org/10.1097/00010694-199608000-00007)
694 [199608000-00007](https://doi.org/10.1097/00010694-199608000-00007)

695 Unrau, A., Becker, G., Spinelli, R., Lazdina, D., Magagnotti, N., Nicolescu, V.-N., Buckley P.,
696 Bartlett D, Kofman, P. D.. (Eds.). 2018. *Coppice Forests in Europe*. EuroCoppice COST Action.
697 Albert Ludwig University of Freiburg, Germany. Retrieved from
698 <https://www.eurocoppice.unifreiburg.de/coppice-forests-in-europe>.

699 Vesterdal, L., Dalsgaard, M., Felby, C., Raulund-Rasmussen, K., Jørgensen, B.B., 1995. Effects of
700 thinning and soil properties on accumulation of carbon, nitrogen and phosphorus in the forest
701 floor of Norway spruce stands. *For. Ecol. Manage.* 77, 1–10. [https://doi.org/10.1016/0378-](https://doi.org/10.1016/0378-1127(95)03579-Y)
702 [1127\(95\)03579-Y](https://doi.org/10.1016/0378-1127(95)03579-Y)

703 Wang, L., Amelung, W., Prietzel, J., Willbold, S., 2019. Transformation of organic phosphorus
704 compounds during 1500 years of organic soil formation in Bavarian Alpine forests – A ³¹P
705 NMR study. *Geoderma*, 340, 192-205. <https://doi.org/10.1016/j.geoderma.2019.01.029>

706 Wang, M.C., Chang, S.H., 2001. Mean residence times and characteristics of humic substances
707 extracted from a Taiwan soil. *Can. J. Soil Sci.* 81, 299-307.

708 Wattel-Koekkoek, E.J.W., Buurman, P., van der Plicht, J., Wattel, E., van Breemen, N., 2003. Mean
709 residence time of soil organic matter associated with kaolinite and smectite. *Eur. J. Soil Sci.* 54,
710 1-10.

711 Webster, R., 2001. Statistics to support soil research and their presentation. *European Journal of*
712 *Soil Science.* 52, 331-340. <https://doi.org/10.1046/j.1365-2389.2001.00383.x>

713 Wu, J., Zeng, H., Zhao, F., Chen, C., Liu, W., Yang, B., Zhang, W., 2020. Recognizing the role of
714 plant species composition in the modification of soil nutrients and water in rubber agroforestry
715 systems. *Sci. Total Environ.* 723, 138042. <https://doi.org/10.1016/j.scitotenv.2020.138042>

716 Yao, S., Zhang, Y.L., Han, Y., Han, X.Z., Mao, J.D., Zhang, B., 2019. Labile and recalcitrant
717 components of organic matter of a Mollisol changed with land use and plant litter management:
718 An advanced ¹³C NMR study. *Sci. of The Total Environ.*, 660, 1-10.
719 <https://doi.org/10.1016/j.scitotenv.2018.12.403>

720 Zhang, H., Deng, Q., Hui, D., Wu, J., Xiong, X., Zhao, J., Zhao, M., Chu, G., Zhou, G., Zhang, D.,
721 2019. Recovery in soil carbon stock but reduction in carbon stabilization after 56-year forest
722 restoration in degraded tropical lands. *For. Ecol. Manage.* 441, 1–8.
723 <https://doi.org/10.1016/J.FORECO.2019.03.037>

724 Zhang, X., Guan, D., Li, W., Sun, D., Jin, C., Yuan, F., Wang, A., Wu, J., 2018. The effects of

725 forest thinning on soil carbon stocks and dynamics: A meta-analysis. *For. Ecol. Manage.* 429,
726 36–43. <https://doi.org/10.1016/j.foreco.2018.06.027>

727 Zhao, Q., Bai, J., Wang, X., Zhang, W., Huang, Y., Wang, L., Gao, Y., 2019. Soil organic carbon
728 content and stock in wetlands with different hydrologic conditions in the Yellow River Delta,
729 China. *Ecohydrol. Hydrobiol.* 20, 537-547. <https://doi.org/10.1016/J.ECOHYD.2019.10.008>

730