

UNIVERSITÀ POLITECNICA DELLE MARCHE Repository ISTITUZIONALE

Current trends and future perspectives towards sustainable and economically viable peach training systems

This is the peer reviewd version of the followng article:

Original

Current trends and future perspectives towards sustainable and economically viable peach training systems / Neri, D.; Crescenzi, S.; Massetani, F.; Manganaris, G. A.; Giorgi, V.. - In: SCIENTIA HORTICULTURAE. - ISSN 0304-4238. - ELETTRONICO. - 305:(2022). [10.1016/j.scienta.2022.111348]

Availability:

This version is available at: 11566/307021 since: 2024-12-06T12:58:42Z

Publisher:

Published DOI:10.1016/j.scienta.2022.111348

Terms of use:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. The use of copyrighted works requires the consent of the rights' holder (author or publisher). Works made available under a Creative Commons license or a Publisher's custom-made license can be used according to the terms and conditions contained therein. See editor's website for further information and terms and conditions. This item was downloaded from IRIS Università Politecnica delle Marche (https://iris.univpm.it). When citing, please refer to the published version.

1	Current trends and future perspectives towards sustainable and
2	economically viable peach training systems
3	
4	Davide Neri ¹ , Samuele Crescenzi ¹ , Francesca Massetani ² , George A. Manganaris ³ ,
5	Veronica Giorgi ^{1,*}
6	
7	¹ Department of agricultural, food and environmental sciences, Università Politecnica
8	delle Marche, Ancona, Italy
9	² HORT Soc. Coop.–Via Cardeto n.70, 60121 Ancona, Italy
10	³ Cyprus University of Technology, Department of Agricultural Sciences, Biotechnology
11	& Food Science, 3603 Lemesos, Cyprus
12	
13	
14	
15	
16	
17	
18	
19	
20	
21	
22	
23	*Author to whom correspondence should be addressed: Veronica Giorgi, D3A
24	Università Politecnica delle Marche, via Brecce Bianche 60131 Ancona, Italy. Email:
25	v.giorgi@staff.univpm.it

27 Abstract

Considering the limited areas suitable for peach cultivation, the short life cycle of the 28 29 orchards, as well as aspects regarding appropriate rootstock availability and soil 30 properties due to replant conditions, the sustainable intensification became increasingly 31 necessary on peach production systems. Based on the local environment and labor 32 availability, two new training systems have been introduced and are being widely 33 adopted towards efficient small canopies for high- and medium-density orchards, 34 respectively. The so-called two-dimensional (2-D) fruiting walls is an intensive, highly-35 mechanized/high-density training system and is commonly accompanied by 36 multifunctional net protection in order to counteract the adverse effects of climate 37 change. On the other hand, the medium-density three dimensional (3-D) small open 38 vases is suitable for low frost-risk areas and farms with fully available manpower. In 39 both cases, the decisions taken during the orchard establishment and the first years 40 regarding soil fertility and orchard mechanization have strategic importance to 41 guarantee a sustainable peach production system on both quantitative and qualitative 42 terms. The employment of both spring and summer pruning increases work efficiency 43 and improves fruit quality, as well as fruit thinning, management efficiency and 44 mechanization. Sustainable intensification not only reshapes the use of chemicals and 45 irrigation, but also weed control and soil amendment with organic matter to support the 46 complexity and heterogeneity of the agroecosystem with the circular economy.

47

48 Keywords: sustainable orchard management, intensification, high density planting,
49 pruning, labor efficiency, netting systems, *Prunus persica*

- 50
- 51

52 **1. Introduction**

The increase in consumption of horticultural commodities as high-quality fresh produce 53 54 food is now a major trend in the world (Cerdà et al., 2021). Agricultural production 55 for human consumption has implications for the environmental sustainability, therefore 56 modern agronomic techniques need to be developed to support the new planting 57 systems that must balance the increase of the demand and their negative environmental 58 impact. Nowadays, China is the major producer of peaches worldwide with ca. 15 59 million metric tons (Mt) per year, followed by Spain (1.8 Mt), Italy (1.3 Mt), Greece 60 (0.9 Mt) and the United States (0.8 Mt). Production volumes may vary from year to 61 year depending on the climatic conditions that may occasionally cause reduced 62 production due to frost damage, yet peach is still ranked as the most important temperate fruit crop in terms of production volumes, besides apple. 63

64 Notably, fresh peach consumption has registered over the recent years a 65 descending trend in several countries due to inferior fruit quality in the market that has 66 turned away consumers from fresh peaches (Cirilli et al., 2016; Crisosto, 2002; 67 Iglesias and Echeverría, 2009). Thus, optimizing consumer quality is necessary to promote fresh peach consumption. Sensory and nutritional quality of peach fruits are 68 69 the output of a system of interaction between different factors. These include rootstock-70 cultivar interactions, but also the training system, and the cultivation techniques 71 adopted (Minas et al., 2018).

Moreover, the globalisation and competition among flesh fruits from different parts of the world renders it necessary to engage in continuous innovation and refinement of orchard management techniques, in order to ensure the profit margins that are indispensable for the success of peach fruit industry. Over the recent years, the main challenge towards enhanced peach production was related to the selection of elite 77 cultivars, as a result of several efficient breeding programs worldwide. In one of the most active breeding periods, from 1997 to 2006, almost 1000 new cultivars were 78 79 registered in the world (516 peaches, 419 nectarines, 50 rachis peaches), in 18 different 80 countries (Fideghelli and Della Strada, 2008). Nowadays the cultivar selection is a 81 key aspect for farmers who have to meet the needs of a rapidly evolving supply chain, 82 and in the meantime, they have to cultivate the new orchard to be productive with 83 marketable fruit, knowing that the fruits of a given cultivar can be appreciated on the 84 market for a limited period, that can be eventually even shorter than the return on 85 investment. It is worth considering that, to pursue an environmentally sustainable 86 production, the choice of the production location is fundamental, especially considering 87 climate changes, for a production that requires lower inputs. Another great challenge is 88 to choose the appropriate cultivation techniques, which have to be efficient, adapted to 89 the cultivar characteristics, economically viable and environmentally-friendly towards 90 more sustainable production systems.

91 The goal to pursue in peach production is to meet the final consumer needs 92 within a global context, with a high-quality end product, and at the same time rationalize 93 management costs and logistics to guarantee sustainable production and 94 commercialization. The major cost in peach production is labor for tree pruning and training, fruit thinning and harvest operations. Preliminary works of our group showed 95 96 that, in Italian peach orchards, harvesting requires from 20 to 30% of the total cost per 97 hectare, pruning around 14% and thinning from 10 to 16% (data not shown). Such results are in accordance with those presented by Iglesias and Echeverria (2022) in 98 99 Spain. However, during the recent years the lack of local labor in the main producing 100 countries in Europe and the United States has led to modifications on planting systems 101 to counteract the lack of specialized workers. Therefore, simplified operations dealing

102 with pruning and thinning can be additionally conducted mechanically (Foschi et al.,

103 **2012; Anthony and Minas, 2021**).

104

105 The decisions taken at orchard establishment and in the first years became of 106 prime importance towards enhancement of fruit production at the lowest production 107 costs (**Loreti and Massai, 2006**). Environmental and social impact of the decisions 108 need to be considered to guarantee a sustainable peach production, while the limited 109 areas suitable for peach production and the short life cycle of the orchards create the 110 need for replantation with several problems regarding appropriate rootstock availability 111 and soil sickness.

112

113 **2.** Sustainable peach orchard intensification

114 Intensive fruit production systems are characterized by increasing planting density, 115 early fruit bearing, small tree size, high crop loads, short orchard life-span, easy 116 mechanical management, efficiency in the use of inputs and frequent replanting 117 (Musacchi et al 2021). The achievement of constant high fruit quality depends on the 118 efficient management of canopy architecture from the nursery to the orchard. 119 Environmental sustainability requires soil management practices to increase and 120 maintain soil fertility such as minimum tillage, multispecies ground cover (Mia et al., 121 2020a,b), supply of amendments and regulated deficit drip irrigation. Innovation in 122 crop-management regimes need strategies to control plant and root development, able to optimize and, where possible, to simplify orchard management. 123

124 The achievement of these goals requires the active participation of farmers and 125 accurate extension services (**Neri et al. 2020**). Integrated fruit production strategies can 126 provide many different ecosystem services (ES), defined as the benefits of nature to 127 human well-being. The ES conceptual framework assumes a dynamic interaction between people and ecosystems and requires a multiscale approach. Many biophysical and ecological processes in agriculture do not occur at the farm level, but at the landscape scale, while European Rural Development Programmes (RDP) typically neither require nor encourage landscape coordination.

132 The integration of knowledge from different stakeholders (e.g., farmers, 133 scientists, technicians, extension specialists) is thus a precondition for successful 134 sustainable land management (Neri et al., 2020). For this reason, it is important a spatial 135 scale match between the RDP and the ecological processes controlling the target agri-136 environmental issues. As an example, peach production should be carried out towards 137 increment of carbon soil content, reduction of soil erosion and augmentation of carbon 138 sequestration. Moreover, the use of plastic netting systems and drip irrigation with 139 plastic tubes should be organized with circular economy criteria which include plastic 140 recycling at the end of their life cycle in the field.

141 The competitiveness of peach industry is highly based on the efficient use of 142 labor and other inputs, in particular irrigation and use of agrochemicals, such as 143 fertilizers and pesticides. All of them require small and accessible trees according to 144 social and environmental conditions. Flatted canopies, commonly also named as planar 145 or bidimensional canopies, tend to be more efficient for enhanced fruit quality and use 146 of external inputs, being more accessible to workers and machines or robots compared 147 to volume, or 3D canopies. On the other hand, the latter are more autonomous, resilient 148 to climate change, but they can be considered as a profitable approach only if the trees 149 are of small and compact size, fully manageable from the ground. All these benefits 150 about the efficiency of labor, combined with environmental concerns, can be achieved 151 through a sustainable intensification (Willett et al., 2019). Most of the options provided in the section "Training systems" are based on flatted canopies or small open vase 152 153 systems, which are fully in line with the objectives of the "Green Deal" and the strategy

154 "From Farm to Fork" of the European Union concerning the sustainable fruit production

155 (EC, 2020; Musacchi et al. 2021).

156 It is worth noting that the Next Generation EU program (EC, 2020) requires to 157 dramatically reduce the use of the fertilizers and pesticides per hectare, and therefore 158 any training and pruning systems, which will be proposed from now onwards, should 159 be efficient not only for the production cost and the labor use but also for the easiness 160 in nutritional and soil fertility control and for precise fruit and pest management.

161

162 **3. Training systems for peach innovation**

163 The choice of the training system is depending on several factors interacting each other 164 (Figure 1Errore. L'origine riferimento non è stata trovata.); it is therefore imperative to 165 demonstrate which is the best solution, but this should be considered on a case by case basis. Biological characters of a given cultivar, labor requirements, mechanization and 166 167 protection systems need to be considered. In addition, the socio-economic conditions 168 which determine the efficiency of cash flow (farming and territory organization, food 169 chain and type of commercialization) and the level of ecologization or eco-170 sustainability of the orchard system must be taken into account.



Figure 1. Factors affecting the choice of a training system -

172 Technical and varietal renovation are necessary to cope with a market demand 173 of standardized quality products, but the answer to this renovation must be flexible, 174 capable to adapt to different techniques and planting systems available for the farmers 175 and able to face different socio-economic situations. As a result, in recent years there 176 are different solutions with a common intensification tendency. Intensification of the 177 planting systems aims not only to reduce unproductive period and rapidly reach full 178 production but, also, must facilitate mechanization and reduce labor cost per yield unit (Neri, 2015). 179

180 Historically, peach has been planted into low-density orchards, which were 181 characterized by wide inter and intra-row spacings, thus the trees had three-dimensional 182 canopy and were tall and robust, resulting very autonomous, productive and long lasting 183 but with a very high labor requirement and with difficult pest control. As in several 184 different fruit orchards, in peach the tendency nowadays is to increase planting density 185 and reduce the tree dimension. The aim is pursued with genetic and horticultural studies 186 to select cultivars and rootstocks that are suited for high density orchards, training and 187 pruning techniques that maintain the plant highly productive in a smaller volume with 188 high quality fruit. In particular, the new training systems manipulate the canopy 189 architecture to achieve various goals. In short, an ideal training system maintains 190 optimal levels of light interception, uniform light distribution and facilitates high yields 191 of premium quality fruit. Light interception and distribution are considered optimal if 192 all canopy parts are receiving more than 30% of incident light, meaning that shading is 193 not excessive in the lower and central part of the canopy to reduce efficiency (Anthony 194 and Minas, 2021). In this respect, the breeding programs try to select highly productive 195 cultivars with well ramified habitus and rootstocks able to grow well in vigor also in 196 replant conditions and able to support the growth of peach trees in a compact form

(canopy and root system). In every case, the rootstocks are not fully dwarfing because
most peach cultivars need yearly renewal of strong brindle and mixed-shoot to sustain
a continuous production along with fruit quality.

200 These goals can be reached with a wide variety of training systems due to the 201 fact that peach tree is characterized of high plasticity. Peach training systems varies 202 from more traditional 3D canopy architectures, with multiple leaders per tree that are 203 adapted to low-density plantings, to modern planar or flatted systems (mostly 2D 204 designs) with single or multiple leaders per tree adapted to high-density. Modern 3D 205 training systems that are now adapted to medium density orchards are the "delayed 206 open vase" in Northern Italy and the 'Catalan bush vase' (Spanish goblet) in Spain 207 which is wide spreading in all the Mediterranean climatic conditions (Mazzoni et al., 208 2022).

The "delayed open vase" is a technical variant of the traditional open vase with empty center, which is widely managed with winter pruning from the ground. It is trained in a free globe shaped canopy to induce early bearing but reducing the central leader vigor with summer pruning. Finally, to open the center at the 4 year (with very high vigor it can be anticipated at the 3^{rd} year) the central axis is drastically pruned (**Figure 2**). Therefore, the final vase form with stable production is delayed and can be reached at the 4^{th} - 5^{th} year with head back of the primary branches.



Figure 2. Pruning sequence during training of the "delayed open vase" from planting to the end of 4th year (trees before and after winter pruning in the first three years, and after pruning in the 4th year)

216

The "Catalan bush vase" is planned to be mechanically pruned and managed from the ground with repeated summer pruning (**Figure 3 and Figure S1**) and completed with a limited winter pruning according to the yield and quality control. It may have a significant production at the 3rd year and full at 4th year.



Figure 3. Catalan vase during the first (1, 2 and 3) and second growing seasons (4 and 5). (1) First manual topping when the shoots exceed 100 cm from the soil. (2) Second topping (manual or mechanical) when the shoots exceed 150 cm from the soil. (3) The plant in the winter at the end of the first year. (4) First mechanical topping when the shoots exceed 200 cm from the soil. (5) Second mechanical topping when the shoots exceed 250 cm from the soil and manual thinning of the main branches to open the center in very late summer, in case of high-yielding varieties (**Neri and Massetani, 2011**).

221

The branches with a curvilinear behavior are more efficient in producing mixed shoots, therefore the amount of pruning is reduced per unit of fruit production (**Figures S2, S3**). We can assume that modern peach orchard training systems are moving from a regular geometrical approach to a more functional approach, which favors the natural growth habit of the most common peach cultivars. The geometric shapes require intensive labor and structure, and thus are less used, even if they are very efficient in
collecting light energy and in improving light distribution inside the canopy, such as
the Y system (e.g., Tatura) and the V system.

The most common and traditional flattened hedgerow systems (so-called 2D) in Italian conditions were the palmette with several variants in the North. They require support structures (stakes, wires, etc.), training of the branches by bending and intensive and time-consuming winter pruning for training. Therefore, the growers are choosing to train peach trees with simpler and less labor requiring systems.

Among the new hedgerow or flatted systems for peach, the central leader has substituted the palmette, with a planting density up to 1000-1667 trees per hectare (4-5 m x 1.5-2 m, **Figures 4 and S4**). Higher planting density requires more summer pruning to enable light penetration to the basal portion of the plants. This form requires support (e.g., post and wires) and is composed of one permanent central axis and only 3-5 branches (less than 1 m long, according to the intra row distance between trees) inserted at 60-90 cm and oriented in all directions.



Figure 4. Description of the pruning for central leader during winter, at the end of the second growing season. The picture on the left shows a plant grown as central leader at the end of the 2^{nd} year summer season in medium density plantations.

The objective of this system, also called spindle or fuse, is to keep the one-yearold wood emerging from the central leader and/or from short pruned spurs. If the planting density is higher, up to 2500-3300 trees per hectare (3.5-4 m x 1 m), the central leader is well feathered and the branches is spur pruned in a columnar shape and can be assimilate to a fruiting wall (**Figure 5**).



Figure 5. Central leader in high density planting (4x1m), at the end of the fourth growing season and before winter pruning (Cuneo, Italy).

248

249 For training high density peach orchards, well feathered scions from the nursery 250 must be used to obtain fruit production in the second year. Above the basal scaffold, 251 the axis is almost empty for 35-45 cm and above that bears short renewable fruiting 252 branches oriented in all the directions with wide angles. The decreasing length of 253 branches from the scaffold to the top of the plant give the plants the typical spindle 254 shape. Even if summer pruning is essential for this system the demand for pruning is 255 low. After two years of almost free growth, precise pruning is applied to form a 256 hedgerow during the winter while minimum mechanical pruning can be applied in the

257 following summer. Training can include summer pruning to trim overcrowded branches 258 in the middle of the tree and to avoid the proliferation of vegetative shoots that shade 259 the tree's interior canopy and to retain only good fruiting shoots (Neri et al., 2015; 260 Figure 6). In the central leader when tree homeostasis is achieved, major part of net 261 photosynthesis during each growing cycle is used for the proper bud formation, 262 production and fruit quality of the subsequent cycle (Hoying et al., 2005), root 263 formation and functioning. In addition to presenting higher productivity per hectare, the 264 central leader system maintains the quality of the fruit, whether in size or sugar content 265 (Uberti et al., 2020) investing a little into new branches and trunk, but this reduces the 266 life span of the tree.



Figure 6. Summer mechanical pruning to improve shoot quality: top left – response to early summer pruning with sylleptic growth; top right – mechanical pruner (Rinieri, FC, Italy); bottom right – modern central axis training system; bottom left low part of the canopy with high quality fruits.

High-density peach orchards demonstrated significant horticultural and 268 269 economic benefit compared with the traditional systems. Increasing tree density may 270 overcome the loss of crop-bearing shoots as well as reduce pruning time (Glenn et al., 271 2011). An orchard trained with a narrow canopy has several advantages in management 272 (all manual operations are facilitated and can be done from ground, management of the 273 sub-row is easier), agronomic (more efficient light penetration in the canopy) and 274 environmental aspects (lower quantities of spraying products, easier to use 275 multifunctional nets) (Dorigoni, 2016).

276

277 4. Limitations towards intensification

278 The tendency to increase planting densities has been driven primarily by the need for 279 early production to pay back the initial investment cost and improve profitability. In 280 modern high-density peach orchards, production starts at the second growing season and reach a maximum at the 4th or 5th year. With higher tree planting densities, 281 282 cumulative fruit production over the first 10 years of an orchard's life has drastically 283 improved. Another reason for the intensification in orchard production systems has 284 been the need to reduce tree size to facilitate tree management. In addition, fruit color 285 is often poor in the center of the canopy of large trees. As the market standards for fruit 286 quality have increased, it has been difficult for fruit growers to achieve satisfactory pest control and fruit quality with large trees. The switch to smaller trees and higher tree 287 288 planting densities has allowed significant improvements on fruit quality (Robinson, 289 2007).

In species that bear fruits on short branches (apple, pear and cherry) it is possible to greatly intensify planting density up to 3000 trees/ha and optimize space occupation. With species that are fruiting in mixed shoots and/or in brindles (30-50 cm long) like peach, planting densities have a limit that is the space between trees that allows the easy

renewal of the mixed shoots. In this case it is possible to augment density up to 2000-2500 trees/ha with vertical axes distanced 1 m along the row, but the central leader is easier to be trained with 800-1,200 trees per hectare with a distance of 2 m along the row (Neri, 2015).

298 In both cases, the rootstock cannot be neither dwarfing because it excessively 299 reduces the renewal nor too vigorous as it creates excessive competition. Several 300 medium vigorous peach rootstocks have been developed by breeding programs in Italy, 301 Spain, France and the USA. However, they have not been adopted widely in Italy due 302 to the risk of reducing fruit size and too limited tree vigor that could excessively reduce 303 mixed shoot growth. Despite the lack of the ideal medium vigor rootstock, significant 304 increases in tree planting density have occurred in stone fruits, as improved canopy 305 management strategies were developed (Robinson, 2007). Moreover, the propagation 306 cycle of grafted peach trees helped in this direction producing small grafted scions in 307 short nursery cycle, reduced to less than 6 months using the mini chip-budding 308 (Musacchi and Neri, 2019). With this technique, in vitro propagated rootstocks in pot 309 are chip budded and 3 months later the bud is swelling. The plantlets with such fast-310 growing shoots are then transplanted directly in the orchard to originate the central axis 311 during the first growing season (Figure 7). This solution requires an efficient irrigation 312 system, but the roots are reactive and able to grow in very diverse conditions, while the 313 main growing shoot can be easily guided. Attention should be taken to control the weed 314 competition, protecting the young plant with a shelter.



Figure 7 Short nursery cycle for the production of scions by means of mini chipbudding on micropropagated rootstocks

316 5. Training system and economical sustainability of the orchard

317 Different training systems show a very different carbon balance during the growing 318 season, related to light interception (Monteith and Moss, 1977) that is dependent on 319 several features of the orchard e.g. plant density, canopy size, thickness, leaves density, 320 resulting a higher carbon accumulation of dry organic matter per hectare in high density 321 systems (Figure 8). Moreover, in structures as the central leader, with a limited 322 presence of secondary wood, only a small amount of plant energy is stocked in 323 permanent structures. It is important hence to adapt cultural practices with the aim of 324 addressing nutrients towards fruits, reproductive buds and roots, to increase yield 325 efficiency and economical sustainability of the orchard.



Training system: 1=Candelabra 2=Palmette 3=Vasette 4=Central leader

Figure 8 Cumulated dry matter produced during the growing season (0 means bud break) in the peach orchard according to the training system (**Silvestroni et al. 2004**)

327 The final aim of the orchard management is to obtain yield and fruit quality high 328 and constant in time, and to do so, the energy that is stocked in the orchard must be 329 used for the benefit of economically valuable plant organs, without dispersions. This 330 high efficiency can be reached only with a system approach which includes regulated 331 deficit irrigation, fertigation and eventually nutrient foliar application, precise 332 minimum winter pruning, mechanical summer pruning and accurate fruit thinning 333 (likely anticipated by mechanical flower thinning). In this way the plants will bear a 334 higher number of fruits per canopy unit. The fruits can be more exposed to light and 335 must be protected with shadowing nets from the excessive light insolation during the 336 summer periods when excessively high temperatures occur (Figure 9). Netting was 337 developed initially for anti-hail protection and later has been proven beneficial also for 338 other purposes. The use of photo-selective netting systems allows a reduction up to 10-339 20% of excess light, maintaining high photosynthetic level thanks to the selective light absorbance. The use of nets is becoming multifunctional compared to the original use
against hail, they help to reduce sunburn and cracking on the fruits while favoring
brilliant coloring and high sugar accumulation through supporting a high
photosynthetic capacity. The new net models are also able to block insects or protect
against rain (Neri et al., 2021).



Figure 9 Photo-selective nets on peach orchard, trained as Y-shape.

345

346 6. Environmental sustainability

A reduction in pesticide is nowadays an obligation for agriculture. Training systems that allow a more efficient foliar product distribution are preferable. Especially in cases where mechanical innovations are available as tower sprayers that direct air fluxes horizontally towards the canopy wall and need a training system in flattened systems so to efficiently spray all the canopy at once. Training systems can be designed to be adapted to spraying machines, like the recycling tunnel sprayers. This type of machines require training systems with compatible specific dimensions and currently are mainly being used in vineyards, since current fruit orchard architecture are not amenable (3Dtraining systems, narrow spaces between rows, canopy height).

356 Sustainability of an agro-ecosystem may increase if the system is efficient in terms of energy use (Mao et al., 2015). Primary energy in the orchard is light that leads 357 358 to CO₂ assimilation through photosynthesis. These processes are common to all plant 359 species in the orchard, including the grass cover, if present, that can increase the rate of 360 intercepted light and contribute to carbon balance of the system. A soil with a grass 361 cover has a reduced mineralization of organic matter in comparison with tilled soil and, 362 if the cover is adequately diversified, it can promote a high biodiversity level (Mia et 363 al., 2020). In such conditions, microbiological activity is high in the upper soil layers 364 where the root systems of the trees are developing. This biodiversity is of primary 365 importance for sustainability of modern orchards that are commonly composed of 366 cloned plants. The lack of biodiversification and low organic matter content in an 367 orchard can cause soil sickness and thus limitations during replantation (Polverigiani 368 et al., 2014). Elevated organic matter content and increased soil structure are effective 369 in maintaining the highest roots proliferation rate. All agronomical interventions aiming 370 to create the most suitable environment for root activity and proliferation have to be 371 studied as a tool in preventing the replant symptomatology. Conservation and enhancement of soil physical, chemical and biological fertility can minimize the 372 negative effects of replant disorders on root proliferation and functionality 373 374 (Polverigiani et al., 2014).

The organic matter of the grass cover residues is triggering a humification cycle in the soil that bring to an improvement of soil fertility during orchard lifespan. Ground cover with living vegetation can deliver several agroecosystem services by promoting functional agrobiodiversity in the orchard (**Canali et al., 2015**). Hence, adopting a sustainable orchard management strategy is vital for enhancing weed biodiversity,

which can provide ecological protection (Granatstein et al., 2010) by offering feed
and shelter to beneficial organisms, and improving soil fertility by hosting mycorrhizae,
and thereby promoting nutrient availability and resilience in the soil (Gangatharan
and Neri, 2012; Mia et al., 2020).

384 Inter-row ground is nowadays frequently managed with a temporary or 385 perennial grass cover, but the tree-row ground is often kept free from vegetable cover 386 (with chemical or mechanical means). Since it is vital to reduce the use of chemicals 387 and reduce the ecological impact of agriculture, while maintaining a production of high 388 quality and quantity (Palmer, 2011), research in weed management is therefore 389 focusing in optimizing the inter-row management with a plant cover. Tree-row 390 management involves the management of orchard weeds as they can compete 391 aggressively with fruit trees for available nutrients and water, essential for plant growth. 392 Therefore, proper weed management is vital in the fruit orchard to minimize weeds 393 competition against fruit trees, assuring fruit yield (Cavender et al., 2014; Steenwerth 394 and Guerra, 2012) and supporting weed biodiversity in the orchard (Mia et al., 2020). 395 Live-mulch is an option for the weed control in fact, sowing or planting a selected 396 species (or mix of species), able to efficiently cover the soil without competing with the 397 fruit trees, is an effective way to control the growth of spontaneous undesired weeds 398 (Neri et al., 2021). At the same time the presence of a plant cover of the soil is 399 improving soil biological and nutritional qualities.

- 400
- 401 **7. Conclusions and future perspectives**

402 Many training systems have been developed in Italy to maximize tree performance in 403 relation to rootstocks, cultivars, environment, and grower preferences. Different 404 training systems can offer an array of options towards productivity, improved quality, 405 and enhance labor efficiency or target market strategies. Until recently, the open center

406 vase (mainly "delayed vase" and "Catalan vase") has remained the system that is most 407 extensively planted worldwide because it is easy to manage and offers many training 408 alternatives. The Northern Italian peach industry has adopted the delayed vase and the 409 central axis depending on the availability of motorized platforms. The first one requires 410 reduced use of spring and summer pruning, the second one involves careful summer 411 pruning to manage shoot growth. The delayed removal of the central axis from the vase 412 creates strong open scaffolds without bending or spreading and provides early 413 production. The early production of a central axis is obtained with limited winter 414 pruning and high-density planting systems with vegetative rapid growth of small scions 415 from the nursery. This is important to help offset costs, considering that the average 416 useful commercial life of a peach cultivar in Europe is about 10 years.

The Catalan vase has been spreading in South climatic conditions with mild winter and it is able to organize tree pruning in a systematic and partially mechanized way, minimizing the management of the canopy and allowing the orchardist to easily train non-skilled workers with simple and programmed pruning and training operations. Nevertheless, new observations and extension research may lead to modification in the current tree training recommendations of the Catalan vase especially when adopted in different climatic growing conditions.

424 In all the training systems, spring and summer pruning increase the efficiency of 425 labor (both for the ease and speed of the work and for the capability of the tree to rapidly 426 compensate for errors and incorrect interventions) and improve fruit quality, and 427 eventually can be mechanized, so as the fruit thinning. Late summer pruning can 428 particularly improve management efficiency in modern peach orchards. It is necessary 429 to remodel the agronomic peach orchard management; that includes reduction in the 430 use of agrochemicals (in the soil and in foliar spraying), rationalization of irrigation and 431 soil management with a sustainable weed control, amendment with organic matter from 432 different sources to support the establishment of the complexity and heterogeneity of

433 the agroecosystem with circular economy.

434

435 Funding

This research did not receive any specific grant from funding agencies in the public,

- 437 commercial, or not-for-profit sectors.
- 438

439 **References**

- Anthony, B.M., Minas, I.S., 2021. Optimizing peach tree canopy architecture for
 efficient light use, increased productivity and improved fruit quality. *Agronomy* 11:1961.
- 443 Canali, S., Diacono, M., Campanelli, G., Montemurro, F., 2015. Organic no-till with
 444 roller crimpers: agro-ecosystem services and applications in organic
 445 Mediterranean vegetable productions. *Sustainable Agriculture Research* 4:70.
- Cavender, G., Liu, M., Hobbs, D., Frei, B., Strik, B., Zhao, Y., 2014. Effects of different
 organic weed management strategies on the physicochemical, sensory, and
 antioxidant properties of machine-harvested blackberry fruits. *Journal of Food Science* 79:S2107–S2116.
- 450 Cerdà, A., Daliakopoulos, I.N., Terol, E., Novara, A., Fatahi, Y., Moradi, E., Salvati,
 451 L., Pulido, M., 2021. Long-term monitoring of soil bulk density and erosion
 452 rates in two *Prunus persica* (L) plantations under flood irrigation and
 453 glyphosate herbicide treatment in La Ribera district, Spain. Journal of
 454 Environmental Management 282:111965
- 455 Cirilli, M., Bassi, D., Ciacciulli, A., 2016. Sugars in peach fruit: A breeding perspective.
- 456 Horticulture Research 3. https://doi.org/10.1038/HORTRES.2015.67/6447882

- 457 Crisosto, C.H., 2002. How Do We Increase Peach Consumption? Acta Horticolturae
 458 592, 601–605. https://doi.org/10.17660/ActaHortic.2002.592.82
- 459 Dorigoni, A., 2016. Innovative fruit tree architecture as a nexus to improve
 460 sustainability in orchards. Acta Hortic. 1137, 1–10.
- 461 European Commision (EC) 2020. Farm to Fork Strategy. For a Fair Healthy and
 462 Environmentally-Friendly Food System. Available online:
 463 https://ec.europa.eu/food/system/files/2020-05/f2f_actionplan_2020_strategy464 info en.pdf (accessed on 30 November 2021).
- Fideghelli, C., della Strada, G., 2008. The recent variety evolution of the international
 peach industry, in: VI Convegno Nazionale sulla Peschicoltura Meridionale.
 SOI, Firenze.
- Foschi, S., Neri, D., Massetani, F., 2012. Meccanizzare il pescheto per salvaguardare il
 reddito. Peach orchard mechanization to maintain profit. L'informatore
 Agrario 43–47.
- Gangatharan, R., Neri, D., 2012. Can biodiversity improve soil fertility resilience in
 agroecosystems? New Medit. 11:11–18.
- Glenn, D.M., Tworkoski, T., Scorza, R., Miller, S.S., 2011. Long-term effects of peach
 production systems for standard and pillar growth types on yield and economic
 parameters. *HortTechnology* 21:720–725.
- Granatstein, D., Wiman, M., Kirby, E., Mullinix, K., 2010. Sustainability trade-offs in
 organic orchard floor management. *Acta Hortic*. 873, 115–122.
- 478 Hoying, S.A., Terence, L., Andersen, R.L., Shane, B., Andersen, R., Freer, J., Bittner,
- 479 J., Robinson, T.L., 2005. Plant Higher Density Peach Orchards? *Fruit*480 *Quarterly- New York State Horticultural Society*. 13:1–34.

- 481 Iglesias, I., Echeverría, G., 2009. Differential effect of cultivar and harvest date on
 482 nectarine colour, quality and consumer acceptance. Scientia Horticulturae 120,
- 483 41–50. <u>https://doi.org/10.1016/j.scienta.2008.09.011</u>
- 484 Iglesias I., Echeverria G., 2022. Current situation, trends and challenges for efficient
 485 and sustainable peach production. Scientia Horticulturae 296,
 486 https://doi.org/10.1016/j.scienta.2022.110899.
- 487 Loreti, F., Massai, R., 2006. State of the art on peach rootstocks and orchard systems.
 488 *Acta Hortic.* 713, 253–268.
- Mao, L.L., Zhang, L.Z., Zhang, S.P., Evers, J.B., van der Werf, W., Wang, J.J., Sun,
 H.Q., Su, Z.C., Spiertz, H., 2015. Resource use efficiency, ecological
 intensification and sustainability of intercropping systems. Journal of
 Integrative Agriculture 14, 1542–1550. <u>https://doi.org/10.1016/S2095-</u>
 3119(15)61039-5
- Mazzoni, L., Medori, I., Balducci, F., Marcellini, M., Acciarri, P., Mezzetti, B.,
 Capocasa, F. Branch numbers and crop load combination effects on production
 and fruit quality of flat peach cultivars (*Prunus persica* (L.) Batsch) trained as
 Catalonian vase. Plants 2022, 11, 308.
- Mia, J., Monaci, E., Murri, G., Massetani, F., Facchi, J., Neri, D., 2020. Soil nitrogen
 and weed biodiversity: an assessment under two orchard floor management
 practices in a nitrogen vulnerable zone in Italy. *Horticulture* 6:96.
- 501 Mia, M.J., Massetani, F., Murri, G., Facchi, J., Monaci, E., Amadio, L., Neri, D., 2020.
 502 Integrated weed management in high density fruit orchards. Agronomy
 503 10:1492
- Minas, I.S., Tanou, G., Molassiotis, A., 2018. Environmental and orchard bases of
 peach fruit quality. Scientia Horticulturae 235, 307–322.
 https://doi.org/10.1016/J.SCIENTA.2018.01.028

- Monteith, J.L., Moss, C.J., 1977. Climate and the efficiency of crop production in
 Britain. Philosophical Transactions of the Royal Society of London. Series B,
 Biological Sciences 281, 277–294.
- Musacchi, S., Neri, D., 2019. Optimizing production of quality nursery plant for fruit
 tree cultivation, in: Lang, G. (Ed.), Achieving sustainable cultivation of
 temperate zone tree fruits and berries. Burleigh Dodds Science. Volume 1
 chapter 6, pp. 340.
- Musacchi, S., Iglesias, I., Neri, D., 2021. Training systems and sustainable orchard
 management for European pear (*Pyrus communis* L.) in the Mediterranean
 area: a review. Agronomy, 11:1765.
- 517 Neri, D., Bravetti, M., Murri, G., Nardini, G., Paroncini, M., 2021. Light spectrum
 518 modifications under photo-selective hail-nets. *Acta Hortic*. 1304: 191-200.
- 519 Neri, D., Massetani, F., 2011. Spring and summer pruning in apricot and peach
 520 orchards. *Advances Hortic. Sci.* 25: 170–178.
- 521 Neri, D., Massetani, F., Murri, G., 2015. Pruning and training systems: What is next?
 522 Acta Hortic. 1084:429–443.
- Neri, D., Polverigiani, S., Zucchini, M., Giorgi, V., Marchionni, F., Mia, J., 2021.
 Strawberry living mulch in organic vineyard. *Agronomy* 2021, 11:1643.
- 525 Neri, D., Silvestroni, S., Baldoni, N., et al., 2020. Sustainable crop production. In: The
- 526 First Outstanding 50 Years of "Università Politecnica delle Marche" Research
 527 Achievements in Life Sciences. Editors: Longhi, S., Monteriù, A., Freddi, A.,
- 528 Aquilanti, L., Ceravolo, M.G., Carnevali, O., Giordano, M., Moroncini, G.
- 529 (Eds.) Hardcover ISBN 978-3-030-33831-2, Springer CH: 583-600.
- 530 Palmer, J.W., 2011. Changing concepts of efficiency in orchard systems. *Acta Hortic*.

531 903:41-49

- 532 Polverigiani, S., Kelderer, M., Neri, D., 2014. Growth of 'M9' apple root in five central
 533 Europe replanted soils. *Plant Root* 8:55–63.
- Robinson, T.L., 2007. Recent advances and future directions in orchard planting
 systems. *Acta Hortic*. 732, 367–381.
- 536 Silvestroni, O., Mattioli, S., Manni, E., Neri, D., Sabbatini, P., Palliotti, A., 2004.
- 537 Seasonal dry matter production in field-grown Sangiovese and Montepulciano
 538 grapevines (*Vitis vinifera* L.). *Acta Hortic.* 640: 127-133.
- 539 Steenwerth, K., Guerra, B., 2012. Influence of floor management technique on
 540 grapevine growth, disease pressure, and juice and wine composition: A review.
 541 *American Journal of Enology and Viticulture* 63, 149-164.
- 542 Uberti, A., Santana, A.S., Lugaresi, A., Prado, J. do, Louis, B., Damis, R., Fischer, D.L.
 543 de O., Giacobbo, C.L., 2020. Initial productive development of peach trees
 544 under modern training systems. *Sci. Hortic*.272:10952
- 545 Willett, W., Rockström, J., Loken, B., Springmann, M., Lang, T., Vermeulen, S.,
- 546 Garnett, T., Tilman, D., DeClerck, F., Wood, A., Jonell, M., Clark, M., Gordon,
- 547 L.J., et al. 2019. Food in the Anthropocene: the EAT–Lancet Commission on
- healthy diets from sustainable food systems. The Lancet.
 https://doi.org/10.1016/S0140-6736(18)31788-4