UNIVERSITÀ POLITECNICA DELLE MARCHE
Repository ISTITUZIONALE

Contrasting land use legacy effects on forest landscape dynamics in the Italian Alps and the Apennines

This is the peer reviewd version of the followng article:

Original
Contrasting land use legacy effects on forest landscape dynamics in the Italian Alps and the Apennines / Garbarino, M., Morresi, D., Urbinati, C., Malandra, F., Motta, R., Sibona, E.M., Vitali, A., Weisberg, P.J.. - In: LANDSCAPE ECOLOGY. - ISSN 0921-2973. - ELETTRONICO. - 35:12(2020), pp. 2679-2694. [10.1007/s10980-020-01013-9]

Availability:
This version is available at: 11566/278409 since: 2024-12-10T13:43:42Z

Publisher:

Published
DOI:10.1007/s10980-020-01013-9

Terms of use:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. The use of copyrighted works requires the consent of the rights' holder (author or publisher). Works made available under a Creative Commons license or a Publisher's custom-made license can be used according to the terms and conditions contained therein. See editor's website for further information and terms and conditions.
This item was downloaded from IRIS Università Politecnica delle Marche (https://iris.univpm.it). When citing, please refer to the published version.

Publisher copyright:

Springer (article) - Postprint/Author's accepted Manuscript

This version of the article has been accepted for publication, after peer review (when applicable) and is subject to Springer Nature's AM terms of use https://www.springernature.com/gp/open-research/policies/accepted-manuscript-terms, but is not the Version of Record and does not reflect post-acceptance improvements, or any corrections. The Version of Record is available online at: 10.1007/s10980-020-01013-9.

(Article begins on next page)

05 October 2026

Landscape Ecology

Contrasting land use legacy effects on forest landscape dynamics in the Italian Alps and the Apennines

--Manuscript Draft--

Manuscript Number:	LAND-D-20-00035R1
Full Title:	Contrasting land use legacy effects on forest landscape dynamics in the Italian Alps and the Apennines
Article Type:	Original research
Keywords:	Forest expansion; Cultural landscape; historical ecology; aerial photographs; landscape structure; land abandonment
Corresponding Author:	Matteo Garbarino, Ph.D. University of Turin Grugliasco, Italy ITALY
Corresponding Author Secondary Information:	
Corresponding Author's Institution:	University of Turin
Corresponding Author's Secondary Institution:	
First Author:	Matteo Garbarino, Ph.D.
First Author Secondary Information:	
Order of Authors:	Matteo Garbarino, Ph.D.
	Donato Morresi, M.Sc.
	Carlo Urbinati, Ph.D.
	Francesco Malandra, Ph.D.
	Renzo Motta, Ph.D.
	Emanuele Sibona, M.Sc.
	Alessandro Vitali, Ph.D.
	Peter J Weisberg, Ph.D.
Order of Authors Secondary Information:	
Funding Information:	Regione Lombardia (PSR2005-n.857) — Prof. Matteo Garbarino Università Politecnica delle Marche (PA-123/2016) — Dr. Francesco Malandra Regione Piemonte (PSR 2000-2006-I.7) — Mr. Emanuele Sibona
Abstract:	Context Land use legacies of human activities and recent post-abandonment forest expansion have extensively modified numerous forest landscapes throughout the European mountain ranges. Drivers of forest expansion and the effects of changes on ecosystem services are currently debated. Objectives i) to compare landscape transition patterns of the Alps and the Apennines (Italy), ii) to quantify the dominant landscape transitions, and iii) to measure the influence of climatic, topographic and anthropogenic driving factors. Methods

Powered by Editorial Manager® and ProduXion Manager® from Aries Systems Corporation

	Land cover changes and landscape pattern modifications were investigated at the regional (over 28 years, Alps and Apennines, Corine Land Cover dataset) and landscape scale (over 58 years, 8 Alpine and 8 Apennine sites, aerial images). The main driving factors of post-abandonment forest landscape dynamics were assessed with a statistical modeling approach. Results Forest expansion was the dominant landscape transition at both Italian mountain ranges, with an annual overall rate of 0.6%. Forest expansion was more extensive at lower elevations in the Apennines where climate is less limiting and extensive abandoned croplands and pastures were available throughout the study period. Distance from pre-existing forest edges in the Alps and elevation in the Apennines emerged as the most important predictors. Conclusions Forest expansion is most rapid where areas of recent agricultural abandonment coincide with favorable climatic conditions. The prediction of forest landscape dynamics, in these mountain forests with a long history of cultural use, requires knowledge of how the magnitude and timing of land use changes intersect spatially and temporally with suitable conditions for tree establishment and growth.
Response to Reviewers:	Dear editors and reviewers, we would like to thank you for your efforts revising our manuscript that we believe greatly benefited from your comments and suggestions. A table containing the reply to reviewers related to the manuscript entitled “Contrasting land use legacy effects on forest landscape dynamics in the Italian Alps and the Apennines” has been uploaded and appended to the main text of the manuscript. The table precisely answer to all reviewers' comments and refers to all the changes we made within the manuscript. The latter are highlighted in yellow within the Word document named “LAND-D-20-00035_R1.docx”.

Powered by Editorial Manager® and ProduXion Manager® from Aries Systems Corporation

Manuscript R1

Click here to access/download;Manuscript;LAND-D-20-00035_R1.docx

Click here to view linked References

Contrasting land use legacy effects on forest landscape dynamics in the Italian Alps and the Apennines

Matteo Garbarino[1], https://orcid.org/0000-0002-9010-1731, matteo.garbarino@unito.it (Corr. author)
Donato Morresi[1], https://orcid.org/0000-0002-3545-872X
Carlo Urbinati[2], https://orcid.org/0000-0003-4879-1406
Francesco Malandra[2], https://orcid.org/0000-0001-9317-8446
Renzo Motta[1], https://orcid.org/0000-0002-1631-3840
Emanuele Marco Sibona, https://orcid.org/0000-0003-1594-3725
Alessandro Vitali[2], https://orcid.org/0000-0002-1713-2152
Peter J. Weisberg[3], https://orcid.org/0000-0003-4883-9297

1) Department of Agricultural, Forest and Food Sciences, University of Turin, Largo Braccini 2, IT 10095 Grugliasco (TO), Italy
2) Department of Agricultural, Food and Environmental Sciences, Marche Polytechnic University, Via Brecce Bianche 10, IT 60131 Ancona (AN), Italy
3) Department of Natural Resources and Environmental Science, University of Nevada, Reno, Mail Stop 186, Reno, NV 89557, USA

Abstract

Context: Land use legacies of human activities and recent post-abandonment forest expansion have extensively modified numerous forest landscapes throughout the European mountain ranges. Drivers of forest expansion and the effects of changes on ecosystem services are currently debated.

Objectives: i) to compare landscape transition patterns of the Alps and the Apennines (Italy), ii) to quantify the dominant landscape transitions, and iii) to measure the influence of climatic, topographic and anthropogenic driving factors.

Methods: Land cover changes and landscape pattern modifications were investigated at the regional (over 28 years, Alps and Apennines, Corine Land Cover dataset) and landscape scale (over 58 years, 8 Alpine and 8 Apennine sites, aerial images). The main driving factors of post-abandonment forest landscape dynamics were assessed with a statistical modeling approach.

Results: Forest expansion was the dominant landscape transition at both Italian mountain ranges, with an annual overall rate of 0.6%. Forest expansion was more extensive at lower elevations in the Apennines where climate is less limiting and extensive abandoned croplands and pastures were available throughout the study period. Distance from pre-existing forest edges in the Alps and elevation in the Apennines emerged as the most important predictors.

Conclusions: Forest expansion is most rapid where areas of recent agricultural abandonment coincide with favorable climatic conditions. Thus the prediction of forest landscape dynamics, in these mountain forests with a long history of cultural use, requires knowledge of how the magnitude and timing of land use changes intersect spatially and temporally with suitable conditions for tree establishment and growth.

Keywords Forest expansion, cultural landscape, historical ecology, aerial photographs, landscape structure, land abandonment

Acknowledgments

The authors would like to thank Nicolò Anselmetto, Michela Basso, Alessandra Bottero, Ariele Cinti, Francesca Lallo, Sara Martelletti, Fedele Maiorano, Fabio Meloni, Sara Saraceni, Laurent Perron and Marta Martinez Subirà for their help with image processing and GIS analyses.

Introduction

Land use history exerts a strong long term legacy on forest landscapes, affecting their structure, spatial pattern and associated ecosystem services (Bellemare et al. 2002; Ziter et al. 2017). At a landscape scale, forests after their removal can be replaced by other land cover types such as crops, pastures, and settlements, which in turn can lead to habitat fragmentation. Land abandonment has important effects on natural disturbance regimes especially in regions with a long history of intensive human influence (Mantero et al. submitted). After land abandonment, secondary succession can lead to forest expansion, a complex dynamic process influenced by different factors such as climate, topography, seed availability and anthropogenic variables. An important driver at local scales has proven to be the former land use intensity and type (Walker et al. 2010). For example, abandoned cropland, with compacted soil due to former plowing, may experience slower successional dynamics than former pastures (Dupouey et al. 2002). A pan-European scale study over the last 25 years revealed that the most important landscape transitions are urbanization and natural afforestation processes, both affecting landscape service provision (Van der Sluis et al. 2019). In forest landscapes post-abandonment processes generally cause increased wildfire frequency, extent and severity (e.g. Moreira 2001; Lloret 2002; Pausas et al. 2012) and a decrease in the frequency and intensity of rockfall (Lopez-Saez et al. 2016; Farvaque et al. 2019) and avalanches (Kulakowski et al. 2011; García-Hernández et al. 2017). Post-abandonment forest expansion can also cause changes to major bio-geochemical cycles, soil properties and eco-hydrological processes (Pellis et al. 2019), loss of biodiversity especially in semi-open areas with species-rich grasslands, and loss of cultural landscapes (Otero et al. 2015; Hermoso et al. 2018; Ridding et al. in press). Secondary succession processes can result in multiple pathways and abandoned lands may become more vulnerable to invasive species and fire (Munroe et al. 2013).

For thousands of years, anthropogenic pressure over the Mediterranean basin has shaped the numerous and diverse cultural landscapes (Naveh 1995). In many European rural areas abandonment after WW2 was a widespread socio-economic process, causing large human migrations toward urban areas (MacDonald et al. 2000; Poyatos 2003; Hochtl et al. 2005). The decline of traditional agro-pastoral activities was particularly intense in southern European mountains such as the Italian Alps and the Apennines. These two mountain ranges, covering approximately 35% of the entire country of Italy (Vacchiano et al. 2017), experienced a significant forest expansion after an extensive decrease of cultivated lands due to depopulation (Falcucci et al. 2007). Forest cover in Italy shifted from 6 million ha in 1936 (Forest Map of the Italian Kingdom) to 8.5 million ha in 1985 (First Italian Forest Inventory, IFNI85), and to 10.5 million in 2005 (Second Italian Forest Inventory, INFC05) and is currently estimated at about 11 million ha in 2015 with an increase of 20% in the last 30 years (Ferretti et al. 2018). These estimates rely on a wide range of sources, and studies using consistent datasets to quantify changes in land cover across broad areas prior to widespread availability of satellite imagery are lacking. In the Italian and French Alps, the depopulation of marginal lands started after the Industrial Revolution in approximately 1871 and, due to the two World Wars, lasted through the 1950s (Batzing et al. 1996). Before the 1950s, grazing was widely distributed in the Italian Alps where rangelands occupied 53% of the mountain areas (White 1950; Garbarino et al. 2013). This was primarily cattle grazing in unfenced pastures. In the Apennines a human migration from mountain areas toward the Adriatic and Tyrrhenian coastal areas occurred from 1951 to 1991, especially in the northern and southern Apennines (Malandra et al. 2018; Vitali et al. 2018). Many of these areas were subjected to former heavy exploitation for firewood and charcoal production, with wood pastures occurring frequently at higher elevations. A national reforestation program to reduce slope erosion, launched before WW2 and lasting until the 1970s, resulted in approximately 760,000 ha of new plantation forests composed mainly of coniferous tree species (Piermattei et al. 2016).

In both mountain regions, forest expansion has occurred as a gap-filling process at lower elevations and as an upward shift of treeline at higher elevations (Tasser et al. 2005). Forest expansion caused a direct

reduction of open areas, reduced the extent of forest-grassland ecotones, and led to decreases in species diversity as well as culturally important landscapes (Falcucci et al. 2007; Petanidou et al. 2008). However, there are fundamental differences in how forest expansion processes have unfolded in the two mountain regions. A gentler topography and a favorable climate led to more intense deforestation in the Apennines, creating open areas that were used as pastures and crops both at low and high elevations. However, landscape mosaic simplification due to forest gap-filling processes mainly occurred at lower elevations. At higher elevations, forest succession on abandoned croplands and grasslands led to a complex and fragmented landscape (Malandra et al. 2019; Vitali et al. 2019). Understanding the underlying drivers of forest landscape changes by comparing the different land use legacies of the Alps and the Apennines is a fundamental step for more ecologically based landscape planning and management oriented toward biodiversity conservation and other ecosystem services.

The Alps and the Apennines are large and highly representative areas for testing our three hypotheses on mountain forest landscape changes in Italy over the last 60 years: 1) forest cover is increasing everywhere, but with different patterns in the Alps and the Apennines; 2) pasture-to-forest is the dominant landscape transition at high elevation; 3) historical forest cover (i.e. land use legacy) is the most important environmental driver for predicting forest expansion today.

Methods

Study area and sampling design

Our multiscale research design was structured at two spatial scales (region and landscape) aimed at comparing the two mountain ranges of Italy, the Alps (AL) and the Apennines (AP). They have similar total length (1300 – 1350 km, respectively), but different geographic orientation (AL: from west-to-east across northern Italy; AP: northwest-to-southeast from Liguria to Calabria). The two mountain ranges differ in terms of climate, topography and land use history.

In AL, mean annual temperatures range from less than 0° to over 10° C, with very cold winters. Annual precipitation ranges from 400 to over 3000 mm and summer dry periods are very rare (Isotta et al. 2013). Metamorphic lithology with intrusive igneous rocks prevail in the inner sectors and sedimentary outcrops dominate in the outer ones. Oak forests dominate at lower elevations whereas beech-silver fir (*Fagus sylvatica* and *Abies alba)* forests prevail on mesic aspects of the montane zone, replaced by *Pinus sylvestris* on xeric slopes. Coniferous forests with *Picea abies* (L.) H.Karst., *Larix decidua* Mill. and *Pinus cembra* L. dominate the subalpine zone (Fauquette et al. 2018).

At AP, mean annual temperatures range from 6° to 10° C, and annual precipitation ranges from 730 to 877 mm, with a short and pronounced summer dry period at lower elevations (Blasi et al. 2014). The eastern Mediterranean side (Adriatic) is generally more continental and humid than the western (Tyrrhenian) one. The forest vegetation is largely dominated by broadleaf species of the Mediterranean and temperate biomes. Xeric oak forests of *Quercus ilex* L., *Quercus pubescens* Willd., *Quercus cerris* L. and *Ostrya carpinifolia* Scop. dominate at lower elevations and *Castanea sativa* Mill. the sub-montane zone. *Fagus sylvatica*, locally mixed with *Abies alba*, largely dominates the montane zone up to treeline except for a few locations in the central and southern sectors where natural pine forests occur, dominated by *Pinus mugo*, *P. heldreichii*, or *P. nigra laricio*.

For our regional scale analyses, the study area in each mountain chain included all contiguous land above 500 m a.s.l. excluding those island polygons separated from the main mountain chains (Fig. 1). We obtained two large areas of 52,002 km^2 (AL) and 44,615 km^2 (AP), where we assessed the land-cover changes (LCC) for the 1990 – 2018 period, based on the Corine Land Cover (CLC Level 3, Feranec et al. 2016) raster maps

(100-m spatial resolution) after merging the original CLC categories into five larger groups: forest (FO), grassland (GR), cropland (CR), urban (UR), and unvegetated (UV) (Table S1). We developed a transition matrix for both regions by assessing changes for the five selected land-cover categories, allowing us to compute the relative changes in AL and AP.

For our landscape-scale analyses, in each region (AL and AP) we selected 8 landscapes of variable extent ranging from 6.3 to 16 km^2. These landscapes were selected and harmonized from previous projects and unpublished data on land-use/land-cover changes in AL and AP (e.g. Garbarino et al. 2013; Malandra et al. 2019. The 16 study landscapes cover a total surface area of approximately 23,000 ha within an elevation range of 500 – 2,600 m a.s.l., including all vegetation zones (Table 1). We adopted the altitudinal threshold of 2,600 m a.s.l., as the potential alpine treeline (Caccianiga et al. 2008; Lingua et al. 2008; Garbarino et al. 2013) in order to limit the LUC analysis to the vegetated part of the 16 landscapes and to reduce the weight of the 'rock' land cover category, which is uninformative for our research. Historical aerial photographs for the years 1954-1962 (b/w, 1: 60,000 approximate cartographic scale, Italian Geographic Military Institute - IGMI) were scanned and digitized at 800 ppi, with a mean spatial resolution of 1 m. The IGMI images were georeferenced and orthorectified using PCI Geomatica 2012 software. Regularly distributed tie points were used to co-register IGMI images with 2012 orthophotos (RGB, 0.5 m cell size, National Agency for Funding in Agriculture - AGEA) resampled at 1 m. The average horizontal Root Mean Square Error (RMSE) was 23 m± 2SD. We used the TINITALY DEM at 10 m spatial resolution (Tarquini et al. 2012) for orthorectification.

We applied a semi-automated object-based classification by combining the automated image segmentation from eCognition software (scale factor 100, color factor 0.5) with on-screen photointerpretation of segmented polygons (Garbarino et al. 2013). We then performed a supervised classification of the objects based on an initial set of at least 10 training polygons for each category, selected through photointerpretation. This was followed by a manual classification of the previously unclassified polygons. Each polygon of the 32 land cover maps (16 landscapes × two time periods), was classified into five land cover classes that were used for the regional-scale analysis: forest (FO), grassland (GR), cropland (CR), urban (UR), unvegetated (UV). The UV category includes different land cover types such as rock, gravel, sand, bare soil and sparse vegetation areas. The latter is a mosaic of sparse grasslands and barren nonvegetated areas that are mostly located between 2000-3000 m a.s.l. A post-processing procedure on the resulting 32 land cover maps was performed in a GIS environment to enforce consistency among the two datasets. A minimum mapping unit (MMU) of 100 m^2 (Garbarino et al. 2011) was obtained by merging smaller polygons with neighboring larger ones by using the ArcGIS tool 'Eliminate' (Malandra et al. 2019). Merged polygons were rasterized at 1-m resolution and the resulting raster maps were smoothed using a moving window (3 × 3) majority filter. We obtained the level of accuracy by randomizing 16 polygons/ha on each map and classifying the objects visually using the same land cover categories adopted in the automatic segmentation (Radoux and Bogaert 2017). Overall, the classification accuracy ranged from 78% to 96% with a Cohen's Kappa coefficient between 0.67 and 0.93 (Table S2). The land cover change analysis at the landscape scale provided 16 transition matrices that were combined to detect overall transitions and differences between the AL and AP mountain regions. We converted the two transition matrices into two transition diagrams showing gain, loss, net change and persistence for each land cover category (Cousins 2001). With the same dataset (Garbarino et al. 2019), we computed the relative contribution (in hectares) of each land cover category to the transition to forest cover and we applied a Mann–Whitney test to compare the medians of the two mountain ranges (AL and AP). The Mann–Whitney test was performed for each of the five categories by using the 8 landscapes as sample size for each mountain range.

For our landscape-scale analyses, we assessed the main drivers affecting forest expansion in AL and AP using a Random Forest (RF) model (Rodman et al. 2019). Specifically, we modelled the occurrence of a transition to forest through a binary classification method using 'mlr' (Bischl et al. 2016) and 'ranger' (Wright & Ziegler 2017) R packages (R Core Team 2019). Given the unbalanced ratio between cells with a

transition to forest (minority class) and cells that remained unchanged (the majority class), we under-sampled unchanged cells using a spatially random selection within each landscape using the 'spatialEco' R package (Evans 2019). Before computing class transitions, we filtered out from the dataset the landscape portions that were already forests in the past, and we downscaled gridded land cover maps from 1 m to 30 and 60 m resolutions using the majority class within each coarser cell. The coarser resolutions allowed us to limit the influence of both co-registration and classification errors of aerial images on model predictions and to evaluate the dependence of model predictions on the spatial scale. For our final analysis, we used the 30 m resolution because RF models trained with data at 30 or 60 m resolutions produced very similar outputs in terms of variables importance and trends of partial dependences. At a coarser resolution prediction errors slightly increased (Table S3).



Figure 1. Location of the 16 landscapes (white circles) within the two Italian mountain regions: the Alps (light green) and the Apennines (light grey) with a minimum elevation of 500 m a.s.l. For landscape codes see Table 1.

Because there was a substantial area of *Pinus* plantations within the forest cover of 2012 images in certain landscapes of the Apennines, we removed plantation patches before computing class transitions in order to model only natural forest dynamics.

Table 1. Environmental descriptors (Area = total surface area; El = mean elevation; Sl = mean slope; Te = mean annual temperature; Pr = mean annual precipitations; BD = mean distance from buildings; RD = mean distance from roads) of 16 landscapes divided by mountain region (AL or AP).

Region	Landscape name	Landscape Code	Area (ha)	El (m a.s.l.)	Sl (°)	Te (°C)	Pr (mm)	BD (m)	RD (m)
AL	Bagni	BAN	1574.3	1788.1	36.0	3.9	1081.3	678.4	289.4
AL	Mello	MEL	1433.7	2045.8	34.1	1.7	983.7	935.0	359.8
AL	Sapè	SAP	959.4	1419.8	22.3	6.2	1001.9	343.8	178.4
AL	Pesio	PES	1599.2	1596.5	31.7	5.7	980.0	711.0	202.1
AL	Ventina	VEN	630.4	2248.6	30.1	0.1	900.8	1044.9	366.4
AL	Musella	MUS	921.4	2155.6	27.7	2.0	788.2	592.1	190.6
AL	Veglia	VEG	1407.2	2043.8	25.4	1.7	1045.1	815.3	205.1
AL	Devero	DEV	1570.0	2169.9	23.6	1.2	1138.6	1048.2	323.3
AP	Cimone	CIM	1593.4	1444.7	19.2	6.8	1352.5	349.3	111.6
AP	Sibillini	SIB	1601.3	1397.2	26.3	6.8	940.1	549.4	115.9
AP	Gran Sasso	GRS	1602.4	1577.5	27.2	6.3	872.3	1548.4	294.2
AP	Terminillo	TER	1600.7	1573.5	25.7	6.8	831.5	700.4	202.8
AP	Morrone	MOR	1603.7	1422.8	22.8	7.6	792.4	1447.3	276.9
AP	Genzana	GEN	1603.4	1234.7	22.4	8.2	789.1	664.7	228.5
AP	Monte Mare	MMA	1604.0	1149.7	24.1	9.0	780.9	1622.2	410.9
AP	Matese	MAT	1605.4	1112.5	22.5	9.2	694.7	481.4	163.4

We used several spatial predictors (Table S4) such as the distance from pre-existing forest edges, topographic variables (elevation, slope, heat load index *sensu* McCune and Grace 2002), climatic variables (mean annual temperature, annual precipitation), and anthropogenic variables (cost of movement, Euclidean distance to buildings, Euclidean distance to roads). We derived topographic variables from the 10 m DEM and climatic variables from the 'Climatologies at high resolution for the earth's land surface areas' (CHELSA) v1.2 datasets at 30 arcsec (~1 km) spatial resolution (Karger et al. 2017). We computed the accumulated cost of movement across the terrain through the Tobler's hiking function (on-path) implemented in the 'movecost' R package (Alberti 2019), using "buildings" in OpenStreetMap as starting locations. We applied two different approaches to obtain either the RF models predictions or the predictive performance estimates with a reduced bias. Specifically, we trained two RF models, one for AL and one for AP, using all the data and tuning hyper-parameters through an 8-fold spatial cross-validation (Brenning 2012, Schratz et al. 2019). We used bias-reduced predictive performance estimates using two common measures in binary classification, the Brier score (Brier 1950) and the area under the receiver operating characteristics curve (AUC). These measures were averaged over a total of 800 RF models obtained through an 8-fold spatial cross-validation repeated 100 times using a nested 5-fold spatial cross-validation for hyper-parameters tuning (Lovelace et al. 2019, Schratz et al. 2019). For both strategies, we used a sequential model-based optimization approach in 'mlrMBO' R package (Bischl et al. 2017) to search for the optimal RF hyper-parameters (mtry, sample fraction and minimum node size) using 50 steps. The spatial cross-validation resampling technique was based on k-means clustering of observation coordinates and allowed us to geographically partition the data, thus maintaining the assumption of independence among training and test sets which would be violated in the case of randomly sampled observations due to the presence of spatial autocorrelation. We assessed variable importance from each model using the permutation method (Breiman 2001) and we employed partial dependencies (Friedman 2001; Goldstein et al. 2015) to interpret the marginal effect of each variable on the predicted probability of forest expansion. Specifically, we computed the average and the standard deviation of individual marginal effects obtained using all the observations in the dataset through the 'generatePartialDependenceData' function in 'mlr' R package.

Results

Regional land cover changes – CORINE database

Forest expansion occurred in both areas, but was greater in AL (+2,951 ha, +9 %) than in AP (+1023 ha, +3.7 %) (Tab. 2). Cropland cover is generally stable whereas human infrastructures increased more at AP (+ 131 ha, +34 %) than at AL (+256 ha, +19 %). Grasslands greatly decreased at both mountain ranges, but more at AL (-2784 ha, -39 %) than at AP (-1160 ha, -27 %), a pattern also observed for unvegetated areas (AP -56 ha, -19 %; AL -503ha, -9 %). Regional scale results (1990-2018) were weakly in agreement with the landscape scale results (1954-2012) as shown in the supplementary material (Table S5).

Table 2. Land cover categories surface areas expressed as a percentage of the total mountain area derived from the Corine Land Cover in the Alps (AL) and the Apennines (AP) throughout the years. Changes across the entire period (1990-2018) are indicated in the last two columns as absolute and relative percent values.

AL	1990	2000	2006	2012	2018	1990-2018 Change	
						Absolute	*Relative*
Forest (FO)	62.9	62.4	62.5	68.6	68.6	+5.7	+9.0
Grassland (GR)	13.8	14.9	14.6	8.5	8.5	-5.4	-38.8
Cropland (CR)	10.1	10.1	10.4	10.2	10.2	+0.2	+1.5
Urban (UR)	2.6	2.7	2.8	3.0	3.1	+0.5	+19.3
Unvegetated (UV)	10.6	9.9	9.7	9.6	9.6	-1.0	-9.2
AP	1990	2000	2006	2012	2018	1990-2018 Change	
						Absolute	*Relative*
Forest (FO)	62.5	62.5	62.3	64.8	64.8	+2.3	+3.7
Grassland (GR)	9.8	9.8	9.8	7.2	7.2	-2.6	-26.6
Cropland (CR)	26.2	26.1	26.2	26.3	26.3	+0.1	+0.5
Urban (UR)	0.9	1.0	1.1	1.1	1.2	+0.3	+34.2
Unvegetated (UV)	0.7	0.6	0.5	0.5	0.5	-0.1	-19.1

Landscape transitions – aerial imagery

Forest expansion also occurred at the landscape scale in both ranges. For all study landscapes combined, the mean annual forest surface area increment was 60 ha/year (0.5 %/year), and was slightly greater at AP (41 ha yr^{-1} or 0.6% yr^{-1}) than at AL (19 ha yr^{-1} or 0.4% yr^{-1}). Grasslands and Crops decreased in both areas, but a larger reduction of crops (CR) occurred at AP (Fig. 2). Unvegetated lands (UV) decreased only at AL, and urban infrastructures (UR) increased more at AP. The relative weights of CR and UR were historically higher at AP, whereas UV values were historically higher at AL. Forest expansion was mostly related to the "GR to FO" transition (66.3 % overall), but was greater at AL than at AP (74.4 % and 62.4 % respectively, Fig. 3). The "CR to FO" transition was stronger at AP than AL (32.4 % and 1.4 %), whereas the opposite pattern was observed for the "UV to FO" transition (AL = 23.9 %, AP = 3.1 %). All land cover transitions to forest were significantly different between the two mountain ranges (Mann-Whitney test: GR, UR with $p < 0.05$; CR, UR with $p < 0.01$).



Figure 2. Area of land cover classes (ha) and land cover transitions from historic (1954-1962) to present time (2012) in the Alps (left panel) and the Apennines (right panel) study sites. Colored boxes refer to land cover categories with box size scaled to area: darker-colored inset boxes represent land cover class (LCC) persistence over time in the case of LCC increase (e.g. FO and UR categories); light-colored inset boxes represent persistence over time in the case of LCC decrease (e.g. GR, CR and UV). Transitions representing forest expansion are highlighted with arrows. Arrow thickness increases with magnitude of the land cover transition. The area converted to forest (ha) for each transition is reported in text above the arrow symbols.



Figure 3. Percent contribution of each land cover category (GR = grassland, CR = cropland, UR = urban, UV = unvegetated) to forest expansion over the studied period (1954-2012).

Drivers of forest expansion

RF models derived for the forest expansion portion of the landscape indicate that the best predictor was the distance from pre-existing forest edges (DF), particularly at AL (importance rate - *IR* = 0.13, Fig. 4). Forest expansion is predicted also by elevation (El), which is particularly important for AP mountain range (AL: *IR* = 0.08, AP: *IR* = 0.16). Climatic variables such as mean annual temperature (Te) and precipitation (Pr) were more important at AP (*IR* = 0.07-0.08). Anthropogenic impact proxy variables (MC, DB and DR) were less influential in the models (*IR* = 0.05-0.03), but DB was more important for AP than for AL (*IR* = 0.05 and 0.03 respectively).



Figure 4. Importance rate of variables in random forest (RF) models of the Alps (light green) and the Apennines (grey). Variables are: distance from pre-existing forest edges (DF), topographic (El = elevation, Sl = slope, HL = Heat load index), climatic (Te = mean annual temperature, Pr = annual precipitation) and anthropogenic (MC = cost of movement, DB = Euclidean distance to buildings, DR = Euclidean distance to roads).

At AL forest expansion probability was higher close to pre-existing forest edges, rapidly declining between 0 and 200 m and with a gradual decline between 200 and 800 m (Fig. 5). At AP the negative relationship between forest expansion probability and distance to pre-existing forest edges featured a rapid decline (from 0.6 to 0.4 of probability) between 0 and 150 m and a gradual decline between 150 and 900 m. The effect of distance from pre-existing forest edges (DF) observed at AP exhibited a higher heterogeneity compared to those observed at AL as highlighted by standard deviation computed at different values of the predictor variable (Fig. 5). Forest expansion at AP was more likely to occur at lower elevations (500 – 1,000 m a.s.l.) and the probability abruptly decreased between 1,000 and 1,500 m a.s.l. A similar pattern was observed at AL, but at higher elevations (1,000 – 2,000 m a.s.l.), with a clear decrease from 2,000-2,500 m a.s.l. Relationships between forest expansion probability and annual temperature were generally weak, although there was a slightly higher probability of forest expansion occurring between 2.5 - 7 °C at AL and 7.5 - 13 °C at AP.



Figure 5. Partial dependence plots showing relative influences of the five most important predictors on the probability of forest expansion, for the Alps (left) and the Apennines (right) across the respective input data ranges. We used all observations to train the model for computing the partial dependence function. Solid lines indicate the average over individual marginal effects of each selected explanatory variable (distance from pre-existing forest edges, elevation, temperature, precipitation and cost of movement) whereas dotted lines are the standard deviations of individual marginal effects.

Discussion

Forest expansion following land abandonment is a well-known process in mountain forests globally (Sitzia et al. 2010). For example, forest expansion in mountainous populated areas has been recently detected in South America (Nanni et al. 2019) and East Asia (Fang et al. 2014). Grazing decline and fire suppression favored forest expansion and forest infilling in California, USA (Lydersen and Collins 2018) and recently rain forest expansion into savanna has been detected in Australia (Ondei et al. 2017). Natural forest expansion is particularly evident and rapid in several mountain ranges of the World such as in the Mediterranean Basin (e.g. Roura-Pascual et al. 2005; Niedrist et al. 2009; Weisberg et al. 2013) where agro-silvo-pastoral traditional practices declined abruptly due to rural depopulation (Lasanta et al. 2017). However, there have been few studies comparing post-abandonment forest expansion patterns among different regions or mountain ranges (e.g. Tasser et al. 2007; Fontana et al. 2014).

By comparing the Italian Alps and the Apennines we found that environmental influences on forest expansion processes were similar between the two regions. Our results for the observed time span (1954-2012) indicate an overall forest area increase of 0.6% yr^{-1} in the Italian Alps and Apennines. These values match with the annual increments recently reported either for the entire Italian peninsula in 1985-2005 (0.3% yr^{-1}) and 2005-2015 (0.2% yr^{-1}) periods (RAF 2018) or for different sites of the Apennines (0.4 - 0.7% yr^{-1}, Brachetti et al. 2012; Malandra et al. 2019). Similar rates are reported for other Alpine regions such as Carnia (0.7% yr^{-1}), Tyrol (0.35% yr^{-1}) and South Tyrol (0.1% yr^{-1}) in the 1955-2000 period (Tasser et al. 2007).

Important differences between the two studied mountain ranges emerged from our analysis. Forest expansion was more intense in the Apennines during the 1954-2012 period. This outcome could be related to the different geographic layout of this mountain range and the wider latitudinal gradient that it encompasses, providing warmer climate conditions more favorable for forest regeneration. Another possible explanation for this pattern arises from to the large differences in elevation gradients between the two mountain ranges. Abandoned sites at lower elevations that were previously cultivated or grazed were more common in the Apennines than in the Alps, and such sites experienced more rapid and extensive forest expansion.

These differences highlight the importance of regional variation in climate and land use history for understanding and predicting forest landscape change following agricultural abandonment. Differences in land use history expressed by a mosaic of former croplands and pastures have important long-term implications for post-abandonment forest establishment (Zimmermann et al. 2010). At the regional scale, we found a greater reduction of grasslands in the Alps than in the Apennines, where we found a greater increase of anthropogenic land cover types (mostly UR). The landscape transitions from grasslands, croplands, and unvegetated lands to forests were by far the most relevant at our landscape scale of analysis. Grassland-to-forest was the dominant shift in both mountain ranges due to a general decline of traditional cattle grazing in mountain areas (e.g. Nagy et al. 2003). However, the two mountain ranges differ in that the widespread transition from unvegetated areas (e.g. rocks and bare soil) to forest occurred only in the Alps. Here, this transition is probably due to the tendency for coniferous treeline species (*Larix decidua* and *Pinus cembra*) to invade higher-elevation, shrub-dominated and alpine plant community types (Vittoz et al. 2008). On the other hand, *Fagus sylvatica* dominated treelines of the Apennines are less prone to upward migration; forest expansion here was mostly the outcome of gap infilling processes (Vitali et al. 2018, Malandra et al. 2019). High elevation forests in the Apennines are dominated by *Fagus sylvatica*, a strongly resprouting species, but with heavy seeds that disperse predominantly over short distances (Vitali et al. 2017). Conifer species with greater long-distance seed dispersal ability occur at a few sites of the central (*Pinus nigra* – Vitali et al. 2017; *Pinus mugo* – Dai et al. 2017) and southern (*Pinus heldreichii* – Vitali et al. 2018) Apennines. Here, croplands-to-forest was the second most important transition, given the possibility of growing a few rare food crops (e.g. potatoes, special cultivars of cereals, apples and chestnuts)

at higher elevations especially on southern or less exposed slopes (Bracchetti et al. 2012; Rovelli 2019). The abandonment of upland traditional farming systems in the Apennines is one of the most important socio-economic drivers of landscape degradation and biodiversity depletion (Farina 1995; Zimmermann et al. 2010).

Pre-existing forest edges emerged as a key land use legacy for future forest expansion both in the Alps (Abadie et al. 2017; Tasser et al. 2007) and the Apennines (Malandra et al. 2019) with a greater importance in the Alps. A general explanation for this is related to seed source availability and to the marginality of ecotones such as forest edges. These are the first pastoral zones to be abandoned when grazing pressure is reduced. The distance from pre-existing forest edges (years 1954-1962) appears a stronger driver in the Alps where, because of harsh conditions at higher elevations, favorable microsites are necessary for tree establishment. In the Apennines, where this variable was second in importance, high variability in effect size is likely caused by heterogeneity among individual observations belonging to different landscapes. On the other hand, in the Apennines, elevation was the most influential predictor variable with widespread forest expansion having occurred at lower altitudes on slopes severely exploited prior to the analyzed time span (1954-2012). The probability of forest expansion gradually decreased along an altitude gradient from 1,000 to 1,800 m a.s.l. in the Apennines, but decreased abruptly between 2,000 and 2,300 m a.s.l. in the Alps. Forest expansion on former pastures and croplands was also faster at lower elevations in mountains of southern Spain (Fernàndez et al. 2004).

The importance of land use legacy for forest landscape dynamics is emphasized by long-term studies demonstrating that legacies may persist for decades, affecting current and future land cover changes (Loran et al. 2017; Tasser et al. 2017). Human activities such as harvesting, grazing, fire and litter removal when practiced for long time periods, may greatly affect current forest dynamics (Gimmi et al. 2013). The role of land use legacies on past and future forest dynamics is typical of many southern European mountain ranges shaped by historical anthropogenic disturbance regimes such as an intensive land use (Albert et al. 2008; Ameztegui et al. 2016). Our study confirms the importance of the location of pre-existing forest edges as the legacy of centuries of human land use in mountain regions, as in other Mediterranean mountain ranges such as the Pyrenees and the French Alps (Mouillot et al. 2005; Gartzia et al. 2016; Abadie et al. 2017).

The differences observed in land cover change patterns between the Alps and the Apennines are not surprising because of strong regional differences in climate, geology, topography, vegetation (e.g. treeline species composition). Forest expansion by upward treeline rise or forest gap-filling processes has occurred primarily on warmer and gentler slopes (e.g. southern exposure), whether in the Alps (Tasser et al. 2007; Garbarino et al. 2013), the Apennines (Vitali et al. 2018) or the Pyrenees (Gartzia et al. 2016). More favorable climate conditions and the greater availability of abandoned open areas make south-exposed low elevation sites suitable areas for forest expansion. Ultimately, the more rapid rate of forest expansion in the Apennines was linked to the greater availability of open areas given the more intense previous land use. Tree encroachment on old pastures by secondary succession and on former unvegetated areas through primary succession prevail in the Alpine region where climate change appears to have a strong influence (Dirnböck et al. 2003; Gehrig-Fasel et al. 2007; Giorgi and Lionello 2008). Elevation plays a fundamental role on forest dynamics at both landscape and stand scales in the Alps (Garbarino et al. 2009; Kulakowski et al. 2011). Land use legacies such as conifer plantations and high elevation crop farming seem stronger in the Apennines where climate change effects on forest expansion at high elevation appear constrained by the unsuitability of *Fagus sylvatica* to migrate upwards due to its heavy seeds and its limited ability to invade adjacent plant community types (Vitali et al. 2018, 2019). However, the limited transitions from unvegetated areas to forests in the Apennines were also due to a combination of topographic influences and the previous land use. In particular, several Apennines peaks with sandy or marl-sandy soils

are less topographically limited (mountain mass effect) so that unvegetated areas are rare and in the absence of past human activities they are climatically suitable for forest dominance.

Forest expansion is an emerging and debated issue that requires accurate measurement and monitoring to allow for proper management of current and future dynamics (Otero et al. 2015). There are two main management strategies: i) passive management to support rewilding processes and limit human induced landscape fragmentation and ii) active management to control and limit the negative effects of re-vegetation processes (Lasanta et al. 2015). Negative effects of natural forest expansion include simplification of landscape structure, decline of species diversity, increased risk of fire and soil erosion, and the loss of cultural landscapes (Lasanta et al. 2015; Ferretti et al. 2019). A balance between conservation through monitoring and active management of secondary succession dynamics (new forests) should be attempted. A recent review, contrasting active management strategies with passive strategies allowing forest secondary succession, found that the most efficient technique seems to be a combination of clearing and extensive grazing (Lasanta 2019), maintaining high levels of landscape complexity and forest-meadow edge.

With this study, by means of a standardized aerial image processing protocol we provided a robust dataset that should be implemented with more comprehensive records (Garbarino et al. 2019). Quantitative historical ecology with data on land use legacies can provide excellent information for ecosystem modelling to predict forest landscape changes (Stürck and Verburg 2017).

We have shown that forest expansion in mountain ranges of Italy is controlled by land use legacies of pre-existing forest edges, interacting with topography and climate. The Alps and the Apennines showed similar landscape changes featuring grassland-to-forest transitions. However, the rate of forest expansion was faster in the Apennines for the larger occurrence at lower elevations of old-fields recolonized by secondary forests. In the Alps, climate and land use changes favored a widespread transition from unvegetated areas to forest at higher elevations. Our results could be biased by the stronger mass effect on the Alps, and the higher average elevation of alpine landscapes. A further limit in our approach is the spatial resolution mismatch between the 1-km climate data resolution and the 30 m unit of forest expansion analysis. Thus, the influence of climatic variables on forest expansion could be underestimated by our random forest models.

Despite these limitations, our results demonstrate that post-abandonment forest expansion is a widespread and ongoing process in Italian mountain forest landscapes. Future research should increase the number of surveyed sites for increased sensitivity in comparing regional differences. It would be informative to apply a land use change modeling approach. Predicting new landscape scenarios for Italian mountain forests should account for the possible changes to disturbance regimes linked to climatic changes (Vacchiano et al. 2017). The extensive forest cover that is blanketing large mountain areas has important implications for habitat and biodiversity. Forest expansion in these mountain landscapes additionally leads to an increase of fuel load continuity that increases the risk of wildfires, particularly in areas exposed to severe drought stress increased by recent climate changes. Large wildfires have recently increased in occurrence in mountain areas where they have historically been quite rare due to the prevalence of managed pastures and farmlands. Other implications regard snowfall and accumulation regimes in combination with soil erosion dynamics after forest fire. These issues are connected also to the naturalness of these processes triggered by man in human-shaped landscapes. A no-management approach of the successional processes would not guarantee, at least in the short term, a strictly natural forest encroachment. A comprehensive overview and assessment of the multiple ecosystem services provided by these complex and millenary landscapes is necessary in order to attempt a sustainable forest management (Schulze and Schulze, 2010).

Author contribution

Matteo Garbarino conceived the experiment and all authors contributed to the study design. Material preparation, data collection and analysis were performed by Matteo Garbarino, Donato Morresi, Emanuele Sibona, Francesco Malandra, Alessandro Vitali and Carlo Urbinati. The first draft of the manuscript was written by Matteo Garbarino and all authors commented on previous versions of the manuscript. All authors read and approved the final manuscript.

Data availability

The datasets generated during and analyzed during the current study are available in the Figshare repository, https://doi.org/10.6084/m9.figshare.11409444.v1

References

Abadie J, Dupouey J-L, Avon C, Rochel X, Tatoni T, Berges L (2017) Forest recovery since 1860 in a Mediterranean region: drivers and implications for land use and land cover spatial distribution. Landsc Ecol 33:289–305

Albert CH, Thuiller W, Lavorel S, Davies ID, and Garbolino E (2008) Land-use change and subalpine tree dynamics: colonization of Larix decidua in French subalpine grasslands. J Appl Ecol 45: 659–669

Alberti G (2019) movecost: An R package for calculating accumulated slope-dependent anisotropic cost-surfaces and least-cost paths. SoftwareX 10: 100331

Ameztegui A, Coll L, Brotons L, Ninot JM (2016) Land-use legacies rather than climate change are driving the recent upward shift of the mountain tree line in the Pyrenees. Glob Ecol Biog 25: 263-273

Bätzing W, Perlik M, Dekleva M (1996) Urbanization and depopulation in the Alps. Mt Res Dev 16: 335-350

Bellemare J, Motzkin G, Foster DR (2002) Legacies of the agricultural past in the forested present: an assessment of historical land-use effects on rich mesic forests. J Biog 29: 1401-1420

Bischl B, Lang M, Kotthoff L, Schiffner J, Richter J, Studerus E, Casalicchio G, Jones ZM (2016) Mlr: Machine Learning in R. J Mach Learn Res 17: 1–5

Blasi C, Capotorti G, Copiz R, Guida D, Mollo B, Smiraglia D, Zavattero L (2014) Classification and mapping of the ecoregions of Italy. Plant Biosyst 148:1255–1345

Bracchetti L, Carotenuto L, Catorci A (2012) Land-cover changes in a remote area of central Apennines (Italy) and management directions. Landscape Urban Plan 104:157–170

Breiman L (2001) Random Forests. Mach Learn 45: 5-32

Brenning A (2012) Spatial cross-validation and bootstrap for the assessment of prediction rules in remote sensing: The R package sperrorest. In: 2012 IEEE international geoscience and remote sensing symposium, IEEE, pp 5372-5375

Brier GW (1950) Verification of forecasts expressed in terms of probability. Mon Weather Rev 78: 1-3

Caccianiga M, Andreis C, Armiraglio S, Leonelli G, Pelfini M, Sala D (2008) Climate continentality and treeline species distribution in the Alps. Plant Bios 142: 66-78

487 Cousins SA (2001) Analysis of land-cover transitions based on 17th and 18th century cadastral maps
488 and aerial photographs. Landsc Ecol 16: 41-54

489 Dai L, Palombo C, Van Gils H, Rossiter DG, Tognetti R, Luo G (2017) Pinus mugo krummholz dynamics
490 during concomitant change in pastoralism and climate in the central Apennines. Mt Res Dev 2017: 37:
491 75–86

492 Dirnböck T, Dullinger S, Grabherr G (2003) A regional impact assessment of climate and land-use
493 change on alpine vegetation. J Biog 30: 401-417

494 Dupouey JL, Dambrine E, Laffite JD, Moares C (2002) Irreversible impact of past land use on forest soils
495 and biodiversity. Ecology 83: 2978-2984

496 Evans JS (2019). spatialEco. R package version 1.2-0, https://github.com/jeffreyevans/spatialEco.
497 Accessed October 2019

498 Falcucci A, Maiorano L, Boitani L (2007) Changes in land-use/land-cover patterns in Italy and their
499 implications for biodiversity conservation. Landsc Ecol 22: 617-631

500 Fang J, Guo Z, Hu H, Kato T, Muraoka H, Son Y (2014) Forest biomass carbon sinks in East Asia, with
501 special reference to the relative contributions of forest expansion and forest growth. Glob Change Biol
502 20: 2019-2030.

503 Farina A (1995) Upland farming systems of the Northern Apennines. Conserving Biodiversity Outside
504 Protected Areas The role of traditional agro-ecosystems Gland: Switzerland and Cambridge, 123-130

505 Farvacque M, Lopez-Saez J, Corona C, Toe D, Bourrier F, Eckert N (2019) How is rockfall risk impacted by
506 land-use and land-cover changes? Insights from the French Alps. Glob Planet Change 174: 138-152

507 Fauquette S, Suc JP, Médail F, Muller SD, Muller SD, Jiménez-Moreno G, Bertini A, Martinetto E,
508 Popescu MS, Zheng Z, De Beaulieu J-L (2018) The Alps: a geological, climatic and human perspective on
509 vegetation history and modern plant diversity. In Hoorn C, Perrigo A, Antonelli A (eds) Mountains,
510 climate and biodiversity. Wiley-Blackwell, Oxford, pp 413-428

511 Feranec J, Soukup T, Hazeu G, Jaffrain G (2016) European landscape dynamics: CORINE land cover data.
512 CRC Press, Boca Ranton

513 Fernández JBG, Mora MRG, Novo FG (2004) Vegetation dynamics of Mediterranean shrublands in
514 former cultural landscape at Grazalema Mountains, South Spain. Plant Ecol 172: 83-94

515 Ferretti F, Sboarina C, Tattoni C, Vitti A, Zatelli P, Geri F, Ciolli M (2018) The 1936 Italian Kingdom Forest
516 Map reviewed: a dataset for landscape and ecological research. Annals Silv Res 42: 3-19

517 Ferretti F, Alberti G, Badalamenti E, Campagnaro T, Corona P, Garbarino M, La Mantia T, Malandra F,
518 Maresi G, Morresi D, Piermattei A, Pividori M, Romano R, Salvadori C, Sibona E, da Silveira Bueno R,
519 Sitzia T, Urbinati C, Vitali A, Pelleri F (2019) Boschi di neoformazione in Italia: approfondimenti
520 conoscitivi e orientamenti gestionali. CREA, Roma, ISBN 978-88-3385-015-3

521 Fontana V, Radtke A, Walde J, Tasser E, Wilhalm T, Zerbe S, Tappeiner U (2014) What plant traits tell us:
522 consequences of land-use change of a traditional agro-forest system on biodiversity and ecosystem
523 service provision. Agr Ecosyst Environ 186: 44-53

524 Friedman J (2001) Greedy function approximation: a gradient boosting machine. Annals Stat 29: 1189-
525 1232

526 Garbarino M, Weisberg PJ, Motta R (2009) Interacting effects of physical environment and
527 anthropogenic disturbances on the structure of European larch (Larix decidua Mill) forests. For Eco Man
528 257: 1794-1802

529 Garbarino M, Lingua E, Subirà MM, Motta R (2011) The larch wood pasture: structure and dynamics of
530 a cultural landscape. Eur J For Res 130: 491-502

531 Garbarino M, Lingua E, Weisberg PJ, Bottero A, Meloni F, Motta R (2013) Land-use history and
532 topographic gradients as driving factors of subalpine Larix decidua forests. Landsc Ecol 28: 805-817

533 Garbarino M, Morresi D, Urbinati C, Vitali A (2019) Land cover raster data of the Italian mountain
534 ranges (Alps & Apennines) for the 1954 - 2012 period. figshare. Dataset.
535 https://doi.org/10.6084/m9.figshare.11409444.v1

536 García-Hernández C, Ruiz-Fernández J, Sánchez-Posada C, Pereira S, Oliva M, Vieira G (2017)
537 Reforestation and land use change as drivers for a decrease of avalanche damage in mid-latitude
538 mountains (NW Spain). Glob Planet Change 153: 35-50

539 Gartzia M, Pérez-Cabello F, Bueno CG, Alados CL (2016) Physiognomic and physiologic changes in
540 mountain grasslands in response to environmental and anthropogenic factors. Appl Geogr 66:1–11

541 Gehrig-Fasel J, Guisan A, Zimmermann NE (2007) Tree line shifts in the Swiss Alps: climate change or
542 land abandonment? J Veg Sci 18: 571-582

543 Gimmi U, Poulter B, Wolf A, Portner H, Weber P, Bürgi M (2013) Soil carbon pools in Swiss forests show
544 legacy effects from historic forest litter raking. Landsc Ecol 28: 835-846

545 Giorgi F, Lionello P (2008) Climate change projections for the Mediterranean region. Glob Planet
546 Change 63: 90-104

547 Goldstein A, Kapelner A, Bleich J, Pitkin E (2015) Peeking inside the black box: Visualizing statistical
548 learning with plots of individual conditional expectation. J Comput Graph Stat 24: 44-65

549 Hermoso V, Morán-Ordóñez A, Brotons L (2018) Assessing the role of Natura 2000 at maintaining
550 dynamic landscapes in Europe over the last two decades: implications for conservation. Landsc Ecol 33:
551 1447-1460

552 Höchtl F, Lehringer S, Konold W (2005) "Wilderness": what it means when it becomes a reality—a case
553 study from the southwestern Alps. Landsc Urb Pla 70: 85-95

554 Isotta FA, Frei C, Weilguni V, Perčec Tadić M, Lassegues P, Rudolf B, Pavan V, Cacciamani C, Antolini G,
555 Ratto SM, Munari M, Micheletti S, Bonati V, Lussana C, Ronchi C, Panettieri E, Marigo G, Vertacnik G
556 (2014) The climate of daily precipitation in the Alps: development and analysis of a high-resolution grid
557 dataset from pan-Alpine rain-gauge data. Int J Climatol 34: 1657-1675

558 Karger DN, Conrad O, Böhner J, Kawohl T, Kreft H, Soria-Auza RW, Zimmermann NE, Linder HP, Kessler
559 M (2017) Climatologies at high resolution for the earth's land surface areas. Scientific Data 4: 170122

560 Kulakowski D, Bebi P, Rixen C (2011) The interacting effects of land use change, climate change and
561 suppression of natural disturbances on landscape forest structure in the Swiss Alps. Oikos 120: 216-225

562 Lasanta T (2019) Active management against shrubland expansion: seeking a balance between
563 conservation and exploitation in the mountains. Cuadernos de Investigación Geográfica 45: 423-440

564 Lasanta T, Arnáez J, Pascual N, Ruiz-Flaño P, Errea MP, Lana-Renault N (2017) Space–time process and
565 drivers of land abandonment in Europe. Catena 149: 810-823

566 Lasanta T, Nadal-Romero E, Arnáez J (2015) Managing abandoned farmland to control the impact of re-
567 vegetation on the environment. The state of the art in Europe. Environ Sci Policy 52: 99-109

568 Lingua E, Cherubini P, Motta R, Nola P (2008) Spatial structure along an altitudinal gradient in the Italian
569 central Alps suggests competition and facilitation among coniferous species. J Veg Sci 19: 425-436

570 Lloret F, Calvo E, Pons X, Díaz-Delgado R (2002) Wildfires and landscape patterns in the Eastern Iberian
571 Peninsula. Landsc Ecol 17: 745-759

572 Loran C, Munteanu C, Verburg PH, Schmatz DR, Bürgi M, Zimmermann NE (2017) Long-term change in
573 drivers of forest cover expansion: an analysis for Switzerland (1850-2000). Reg Env Change 17: 2223-
574 2235

575 Lopez-Saez J, Corona C, Eckert N, Stoffel M, Bourrier F, Berger F (2016) Impacts of land-use and land-
576 cover changes on rockfall propagation: Insights from the Grenoble conurbation. Sci Tot Env 547: 345-
577 355

578 Lovelace R, Nowosad J, Muenchow J (2019) Geocomputation with R. Chapman and Hall/CRC, Boca
579 Raton, ISBN 978-11-3830-451-2

580 Lydersen JM, Collins BM (2018) Change in vegetation patterns over a large forested landscape based on
581 historical and contemporary aerial photography. Ecosystems 21: 1348–1363

582 MacDonald D, Crabtree JR, Wiesinger G, Dax T, Stamou N, Fleury P, Gutierrez Lazpita J, Gibon A (2000)
583 Agricultural abandonment in mountain areas of Europe: environmental consequences and policy
584 response. J Env Manag 59: 47-69

585 Malandra F, Vitali A, Urbinati C, Garbarino M (2018) 70 years of land use/land cover changes in the
586 Apennines (Italy): a meta-analysis. Forests 9: 551

587 Malandra F, Vitali A, Urbinati C, Weisberg PJ, Garbarino M (2019) Patterns and drivers of forest
588 landscape change in the Apennines range, Italy. Reg Env Change 19: 1973-1985

589 Mantero G, Morresi D, Marzano R, Mladenoff D, Motta R, Garbarino M (submitted) The influence of
590 land abandonment on forest disturbance regimes: a global review. Submitted to Landscape Ecology

591 McCune B, Grace JB, Urban DL (2002) Analysis of Ecological Communities. MjM Software Design,
592 Gleneden Beach, Oregon

593 Moreira F, Rego FC, Ferreira PG (2001) Temporal (1958–1995) pattern of change in a cultural landscape
594 of northwestern Portugal: implications for fire occurrence. Landsc Ecol 16: 557-567

595 Mouillot F, Ratte JP, Joffre R, Mouillot D, Rambal S (2005) Long-term forest dynamic after land
596 abandonment in a fire prone Mediterranean landscape (central Corsica, France). Landsc Ecol 20: 101–
597 112

598 Munroe DK, van Berkel DB, Verburg PH, Olson JL (2013) Alternative trajectories of land abandonment:
599 causes, consequences and research challenges. Curr Opin Environ Sustain 5: 471-476

600 Nagy L, Thompson D, Grabherr G, Körner C (2003) Alpine biodiversity in Europe: an introduction. Joint
601 Nature Conservation Comitee, Report: 1-10, Peterborough

602 Nanni AS, Sloan S, Aide TM, Graesser J, Edwards D, Grau HR (2019) The neotropical reforestation
603 hotspots: A biophysical and socioeconomic typology of contemporary forest expansion. Glob Environ
604 Change 54: 148-159

605 Naveh Z (1995) Interactions of landscapes and cultures. Landsc Urban Plan 32: 43-54

606 Niedrist G, Tasser E, Lüth C, Dalla Via J, Tappeiner U (2009) Plant diversity declines with recent land use
607 changes in European Alps. Plant Ecol 202: 195–210

608 Ondei S, Prior LD, Vigilante T, Bowman DM (2017) Fire and cattle disturbance affects vegetation
609 structure and rain forest expansion into savanna in the Australian monsoon tropics. J Biog 44: 2331-
610 2342

611 Otero I, Marull J, Tello E, Diana GL, Pons M, Coll F, Boada M (2015) Land abandonment, landscape, and
612 biodiversity: questioning the restorative character of the forest transition in the Mediterranean. Ecol
613 Soc 20: 7

614 Pausas JG, Fernández-Muñoz S (2012) Fire regime changes in the Western Mediterranean Basin: from
615 fuel-limited to drought-driven fire regime. Clim Change 110: 215-226

616 Pellis G, Chiti T, Rey A, Yuste JC, Trotta C, Papale D (2019) The ecosystem carbon sink implications of
617 mountain forest expansion into abandoned grazing land: The role of subsoil and climatic factors. Sci
618 Total Environ 672: 106-120

619 Petanidou T, Kizos T, Soulakellis N (2008) Socioeconomic dimensions of changes in the agricultural
620 landscape of the Mediterranean basin: a case study of the abandonment of cultivation terraces on
621 Nisyros Island, Greece. Environ Manag 41: 250–266

622 Piermattei A, Lingua E, Urbinati C, Garbarino M (2016) Pinus nigra anthropogenic treelines in the
623 central Apennines show common pattern of tree recruitment. Eur J For Res 135: 1119-1130

624 Poyatos R, Latron J, Llorens P (2003) Land use and land cover change after agricultural abandonment.
625 Mt Res Dev 23: 362-369

626 R Core Team (2019) R: a language and environment for statistical computing. R Foundation for
627 Statistical Computing, Vienna

628 Radoux J, Bogaert P (2017) Good practices for object-based accuracy assessment. Remote Sens 9: 646

629 RAF (2018) Rapporto sullo stato delle Foreste e del settore forestale in Italia. Compagnia delle Foreste,
630 Arezzo

631 Ridding LE, Watson SC, Newton AC, Rowland CS, Bullock JM (in press) Ongoing, but slowing, habitat loss
632 in a rural landscape over 85 years. Landsc Ecol doi: 10.1007/s10980-019-00944-2

633 Rodman KC, Veblen TT, Saraceni S, Chapman TB (2019) Wildfire activity and land use drove 20th-
634 century changes in forest cover in the Colorado front range. Ecosphere 10(2): e02594

635 Roura-Pascual N, Pons P, Etienne M, Lambert B (2005) Transformation of a rural landscape in the
636 eastern Pyrenees between 1953 and 2000. Mt Res Dev 25:252–261

637 Rovelli E (2019) Aspetti storici delle faggete appenniniche. http://www.mountainforest.org/aspetti-
638 storici-delle-faggete-appenniniche/ Accessed 1 Dec 2019

639 Schratz P, Muenchow J, Iturritxa E, Richter J, Brenning A (2019) Hyperparameter tuning and
640 performance assessment of statistical and machine-learning algorithms using spatial data. Ecol Model
641 406: 109-120

642 Schulze ED, Schulze I (2010) Sustainable foresting: easier said than done. Science 327: 957

643 Sitzia T, Semenzato P, Trentanovi G (2010) Natural reforestation is changing spatial patterns of rural
644 mountain and hill landscapes: a global overview. For Eco Man 259: 1354-1362

645 Stürck J, Verburg PH (2017) Multifunctionality at what scale? A landscape multifunctionality assessment
646 for the European Union under conditions of land use change. Landsc Ecol 32: 481-500

647 Tarquini S, Vinci S, Favalli M, Doumaz F, Fornaciai A, Nannipieri L (2012) Release of a 10-m-resolution
648 DEM for the Italian territory: Comparison with global-coverage DEMs and anaglyph-mode exploration
649 via the web. Comput Geosci 38: 168-170

650 Tasser E, Tappeiner U, Cernusca A (2005) Ecological effects of land use changes in the European Alps. In
651 Huber UM, Bugmann HKM, Reasoner MA (eds) Global Change and Mountain Regions. Advances in
652 Global Change Research. Springer, Dordrecht, pp 409–420

653 Tasser E, Walde J, Tappeiner U, Teutsch A, Noggler W (2007) Land-use changes and natural
654 reforestation in the Eastern Central Alps. Agric Ecosyst Environ 118: 115-129

655 Tasser E, Leitinger G, Tappeiner U (2017) Climate change versus land-use change - which affects the
656 landscape more? Land Use Policy 60: 60–72

657 Vacchiano G, Garbarino M, Lingua E, Motta R (2017) Forest dynamics and disturbance regimes in the
658 Italian Apennines. For Eco Man 388: 57-66

659 Van der Sluis T, Pedroli B, Frederiksen P, Kristensen SB, Busck AG, Pavlis V, Cosor GL (2019) The impact
660 of European landscape transitions on the provision of landscape services: an explorative study using six
661 cases of rural land change. Landsc Ecol 34: 307-323

662 Vitali A, Camarero JJ, Garbarino M, Piermattei A, Urbinati C (2017) Deconstructing human-shaped
663 treelines: Microsite topography and distance to seed source control Pinus nigra colonization of treeless
664 areas in the Italian Apennines. For Ecol Man 406: 37-45

665 Vitali A, Garbarino M, Camarero JJ, Malandra F, Toromani E, Spalevic V, Urbinati C (2019) Pine
666 recolonization dynamics in Mediterranean human-disturbed treeline ecotones. For Eco Man 435: 28-37

667 Vitali A, Urbinati C, Weisberg PJ, Urza AK, Garbarino M (2018) Effects of natural and anthropogenic
668 drivers on land-cover change and treeline dynamics in the Apennines (Italy). J Veg Sci 29: 189-199

669 Vittoz P, Rulence B, Largey T, Freléchoux F (2008) Effects of climate and land-use change on the
670 establishment and growth of cembran pine (Pinus cembra L) over the altitudinal treeline ecotone in the
671 Central Swiss Alps. Arct Antarct Alp Res 40: 225-232

672 Walker LR, Wardle DA, Bardgett RD, Clarkson BD (2010) The use of chronosequences in studies of
673 ecological succession and soil development. J Ecol 98: 725–736

674 Waser LT, Schwarz, M (2006) Comparison of large-area land cover products with national forest
675 inventories and CORINE land cover in the European Alps. Int J Appl Earth Obs Geoinf 8: 196-207

676 Weisberg PJ, Shandra O, Becker ME (2013) Landscape influences on recent timberline shifts in the
677 Carpathian Mountains: abiotic influences modulate effects of land-use change Arct Antarct Alp Res
678 45:404–414

679 White WT (1950) Pastures in the Italian Highlands. Rangeland Ecol Manag Arch 3: 22-28

680 Wright MN, Ziegler A (2017) ranger: A fast implementation of random forests for high dimensional data
681 in C++ and R. J Stat Softw 77: 1-17

Zimmermann PD, Tasser E, Leitinger G, Tappeiner U (2010) Effects of land-use and land-cover pattern on landscape-scale biodiversity in the European Alps. Agr Ecosyst Environ 139: 13–22

Ziter C, Graves RA, Turner MG (2017) How do land use legacies affect ecosystem services in United States cultural landscapes? Landsc Ecol 32: 2205-2218

687 **Supplemental material**

688 Table S1. Harmonization details of the Corine Land Cover legend in 5 land cover categories at AL and AP.

Corine Land Cover		Harmonization	
Code	**Description**	**Code**	**Description**
111	Continuous urban fabric	UR	Urban
112	Discontinuous urban fabric	UR	Urban
121	Industrial or commercial units	UR	Urban
122	Road and rail networks and associated land	UR	Urban
123	Port areas	UR	Urban
124	Airports	UR	Urban
131	Mineral extraction sites	UR	Urban
132	Dump sites	UR	Urban
133	Construction sites	UR	Urban
141	Green urban areas	UR	Urban
142	Sport and leisure facilities	UR	Urban
211	Non-irrigated arable land	CR	Cropland
212	Permanently irrigated land	CR	Cropland
213	Rice fields	CR	Cropland
221	Vineyards	CR	Cropland
222	Fruit trees and berry plantations	CR	Cropland
223	Olive groves	CR	Cropland
231	Pastures	GR	Grassland
241	Annual crops associated with permanent crops	CR	Cropland
242	Complex cultivation patterns	CR	Cropland
243	Land principally occupied by agriculture with … natural vegetation	CR	Cropland
244	Agro-forestry areas	CR	Cropland
311	Broad-leaved forest	FO	Forest
312	Coniferous forest	FO	Forest
313	Mixed forest	FO	Forest
321	Natural grasslands	GR	Grassland
322	Moors and heathland	FO	Forest
323	Sclerophyllous vegetation	FO	Forest
324	Transitional woodland-shrub	FO	Forest
331	Beaches dunes sands	UV	Unvegetated
332	Bare rocks	UV	Unvegetated
333	Sparsely vegetated areas	UV	Unvegetated
334	Burnt areas	FO	Forest
335	Glaciers and perpetual snow	UV	Unvegetated
411	Inland marshes	UV	Unvegetated
412	Peat bogs	UV	Unvegetated
421	Salt marshes	UV	Unvegetated
422	Salines	UV	Unvegetated
423	Intertidal flats	UV	Unvegetated
511	Water courses	UV	Unvegetated
512	Water bodies	UV	Unvegetated
521	Coastal lagoons	UV	Unvegetated
522	Estuaries	UV	Unvegetated
523	Sea and ocean	UV	Unvegetated
999	NODATA	999	No Data

689

690 Table S2. Classification accuracy (OA = overall accuracy, K = Cohen's Kappa coefficient) of 32 land cover
691 maps (16 landscapes x 2 periods).

Landscape	Year	*OA*	*K*
BAG	1962	79	0.69
	2012	89	0.83
CIM	1954	82	0.73
	2012	93	0.88
DEV	1954	82	0.71
	2012	86	0.77
GEN	1954	78	0.70
	2012	78	0.67
GRS	1954	80	0.67
	2012	86	0.76
MAT	1954	86	0.77
	2012	94	0.88
MEL	1962	82	0.73
	2012	88	0.82
MMA	1954	91	0.83
	2012	90	0.68
MOR	1954	82	0.69
	2012	90	0.83
MUS	1961	82	0.72
	2012	89	0.84
PES	1954	88	0.78
	2012	86	0.73
SAP	1954	84	0.73
	2012	91	0.84
SIB	1954	86	0.80
	2012	86	0.77
TER	1954	85	0.75
	2012	84	0.71
VEG	1954	87	0.81
	2012	91	0.87
VEN	1961	94	0.89
	2012	96	0.93

692
693
694
695
696
697
698
699

Table S3. Random forest model parameters used in this study (N transitions = number of forest expansion pixels, N total = number of total pixels in the model, OOB = Out-of-bag error or prediction error of RF models, Brier score = Brier predictive performance estimate, AUC = area under the receiver operating characteristics curve, K = Cohen's Kappa coefficient) computed for the AL and the AP datasets both at 30 and 60 m spatial resolutions.

RF models	AL 30 m	AP 30 m	AL 60 m	AP 60 m
N transitions	14420	21027	3555	5184
N total	28900	40575	7082	10010
OOB	0.108	0.091	0.126	0.098
Brier score	0.175	0.151	0.182	0.132
AUC	0.829	0.854	0.819	0.849
K	0.482	0.519	0.483	0.510
DF *IR*	0.127	0.088	0.114	0.072
El *IR*	0.077	0.156	0.065	0.168
MC *IR*	0.053	0.054	0.034	0.039
Te *IR*	0.049	0.076	0.033	0.051
Pr *IR*	0.048	0.073	0.028	0.060
DB *IR*	0.031	0.050	0.016	0.033
DR *IR*	0.026	0.025	0.014	0.012
Sl *IR*	0.022	0.035	0.013	0.030
HL *IR*	0.019	0.032	0.011	0.023
Total *IR*	0.452	0.589	0.329	0.489

Table S4. Description, data sources, spatial characteristics and usage rationale of explanatory and predictor variables used in the Random Forest models.

Variable	Code	Type	Resolution	Data source	Description
Forest expansion	FE	Binary	10 m	LC map	Transition to forest occurrence
Elevation	El	Topographic	10 m	DEM tinitaly	Gradient of site suitability
Slope	Sl	Topographic	10 m	DEM tinitaly	Proxy of human pressure
Heat Load Index	HL	Topographic	10 m	DEM tinitaly	Gradient of site suitability
Precipitation	Pr	Climatic	1 km	CHELSA	Average annual precipitation
Temperature	Te	Climatic	1 km	CHELSA	Average annual temperature
Distance to Forests	DF	LU Legacy	10 m	LU 1954	Distance to former forest borders
Distance to Roads	DR	Anthropic	10 m	OSM	Euclidean distance from roads
Distance to Buildings	DB	Anthropic	10 m	OSM	Euclidean distance from buildings
Moving cost	MC	Anthropic	10 m	DEM - OSM	Cost of movement across the terrain

Table S5. Land cover surface as a percentage of the total surface of AL and AP landscapes (10091 and 12830 ha, respectively) divided by regional (CORINE land cover) and landscape (aerial imagery) scales. For comparison purposes, the considered study area is included within the borders of the 16 selected landscapes.

Regional scale (CORINE LC)				Landscape (Aerial imagery)			
AL	1990	2018	Delta	AL	1954	2012	Delta
FO	51.58	53.02	1.44	FO	38.75	48.82	10.07
GR	16.16	4.43	-11.73	GR	28.35	24.51	-3.84
CR	0.81	0.55	-0.26	CR	0.37	0.10	-0.26
UR	0.04	0.04	0.00	UR	0.15	0.30	0.15
UV	31.40	41.96	10.55	UV	32.38	26.26	-6.11
AP	1990	2018	Delta	AP	1954	2012	Delta
FO	67.16	65.98	-1.18	FO	46.99	65.40	18.41
GR	16.03	10.82	-5.21	GR	36.11	26.67	-9.44
CR	3.52	3.66	0.14	CR	11.17	0.98	-10.19
UR	0.65	0.70	0.05	UR	0.95	1.47	0.52
UV	12.63	18.84	6.21	UV	4.78	5.49	0.70

Figure 1

Click here to access/download;Figure;Fig1.tif

Figure 3

Click here to access/download;Figure;Fig3.tif



Figure 2

Click here to access/download;Figure;Fig2_n.tif



Figure 4

Click here to access/download;Figure;Fig4_n.tif



Figure 5

Click here to access/download;Figure;Fig5_n.tiff



Reply to reviewers

Click here to access/download;Attachment to manuscript;LAND-D-20-00035_replytoreviewers.docx

Click here to view linked References

Reply to reviewers related to the manuscript entitled “Contrasting land use legacy effects on forest landscape dynamics in the Italian Alps and the Apennines”. All the changes are highlighted in yellow within the Word document named “LAND-D-20-00035_R1.docx”.

Reviewers	Comments	Replies
GENERAL COMMENTS		
R2	Overall, I felt that the analyses were sound and useful. However, clarity of certain portions of the methods section could be improved, and justification of some components (such as variable selection and selection of the focal landscapes) could be further detailed.	We better specified the rationale under the selection of landscapes and variables. See specific answers in the methods section.
R2	The paper is very well written, but would benefit from an additional review by the authors to catch several small grammatical errors (see specific comments below for some of these).	We fixed those grammatical errors following the specific comments.
R2	The two analyses that make up the core of this study (the CORINE land cover and air photos) felt a bit "detached" from one another and might be better integrated given that they convey similar ideas in similar study areas. In particular, I was surprised that the aerial imagery were discussed so much more than the CORINE land cover analyses. One suggestion below is to try to use the CORINE data to determine how representative the focal landscapes for the aerial imagery are of the broader Alp and Apennine ranges. Though there are probably other ideas that the authors may have for better integrating the two analyses.	To integrate CLC data and aerial imagery we added a table (S5) within the supplementary material to compare land cover proportions per year in the AL and AP regions. This table shows that at the landscape scale this data are not reliable because in some of our landscapes, due to CORINE misclassification, we observed a strong reduction of forest surface area, that is not the real situation. This is why we decided to discuss a lot more the aerial images derived data because we feel confident about the “quality” of our land cover maps, this is not the case of CORINE data. Another reason against the CORINE-aerial comparison is that this data analyze different periods of time (1990-2018 and 1954-2012, respectively).
R2	The effects of scale on the analyses and results could be further explored. One specific example is the selection of a 30 m spatial resolution in the Random Forest analysis.	The effect of scale was explored more thoroughly and methods/results sections are slightly modified. We added additional model results in the appendix section (Table S3).
R2	The mechanisms of forest expansion could be further elaborated in the discussion section to more strongly connect to fine-scale forest dynamics in these and similar systems.	We added some sentences in the discussion section to deepen the link between broad-scale forest expansion with species reproductive strategy.

R3	I have commented on some minor notes directly in the manuscript. Please note these. In addition, I have taken the liberty of including some literature citations specific to the Alps, which underline some points of discussion from the perspective of the Alps even more.	We report in this table all the comments by reviewer #3.
SPECIFIC COMMENTS		
R3	L 19: “dramatically” In some regions yes, in others much less.	That sentence was slightly modified.
R3	L 29: better “a-1”	Here and in the rest of the manuscript we used “ yr^{-1}”.
R2	L 46: "…and [associated] ecosystem services."	Modified accordingly.
R2	L 53: "…abandoned [cropland]…"	Modified accordingly.
R2	L 76: The wording "not directly comparable" here seems to undercut your point. Perhaps it could be restated? Or used to further justify the study by saying something like, "These estimates rely on a range of sources, and studies using consistent datasets to quantify changes in land cover across broad areas prior to widespread availability of satellite imagery are lacking."	We substituted that sentence with the suggested one.
R3	L 114: Where do these numbers come from? From my point of view these sums are far too high! Normally there is precipitation of 600-1600 (max. 2000) mm in the Central and Southern Alps.	We modified the sentence citing a recent paper by Isotta et al. (2013) highlighting the high variability that can be found in the Alpine region.
R2	L 114: As "drought" is typically used in a relative sense (relative to long-term conditions on a given site), perhaps this (and L 121) could be changed to "with a pronounced dry period during the summer" or something similar.	Right, we modified those two sentences accordingly.
R2	L 137-138: This section is related to a couple of my general comments above. I think that a justification for why these specific landscapes were selected would be helpful (was it based solely on data availability? Were they randomly selected from a pool of available images?), as would a description of how well they represent the broader conditions in each mountain range. In other words, are they representative samples or case studies? To determine this, perhaps a brief section in the appendix could be added that compares CORINE changes 1990-2018 within the (i.e., sample) of the study landscapes to the broader regional (i.e., population) results from CORINE presented in the first section of the results. I don't think this is necessary to add to the body of the paper (unless the authors choose to do so), but would be helpful in the appendix with a brief reference in the results to situate the study landscapes within the broader regions. I also think that it might be helpful to tie the two analyses together a bit more strongly.	We added a sentence on landscapes selection: “These landscapes were selected and harmonized from previous projects and unpublished data on land-use/land-cover changes in AL and AP (e.g. Garbarino et al. 2013; Malandra et al. 2019”. The landscape selection was necessarily random because of the different purposes of previous projects. We also believe they are representative of the studied mountain ranges because of the absence of a bias of our landscape scale data. Regarding the CORINE land cover data we added a table (S5) as a supplementary material reporting a comparison between the two different approaches (CORINE and

		aerial images).
R2	L 151: Why were these specific parameters selected in segmentation? It might be an interesting addition to test multiple parameters in the segmentation and to show their influence on the outcome of the classification or RF analyses.	These specific parameters were selected after several tests performed on different aerial images during these and previous projects (e.g. Malandra et al. 2019, Garbarino et al. 2013, Garbarino et al. 2011).
R2	L 152-154: At first read, this description seemed to suggest that the authors manually classified all segments (following segmentation in eCognition) in each study landscape. However, I think that they actually performed a supervised classification of the objects based on an initial set of training polygons selected through photointerpretation. If this is the case, it could be more clearly stated here (including the general protocol and supervised classification method). If the analysts did in fact manually classify all objects, then additional information should be added to this section to describe the procedure in detail, including methods to reduce variability in interpretation among different analysts. If the classification was supervised and object-based, ensuring that accuracy assessment was performed at the object-level rather than the pixel-level would be important (see: Radoux, J., & Bogaert, P. (2017). Good Practices for Object-Based Accuracy Assessment. Remote Sensing, 9(7), 646. doi: 10.3390/rs9070646). Additional detail could be added to the description of the accuracy assessment for this classification.	Right, we performed a supervised classification of the objects based on an initial set of training polygons selected through photointerpretation and then we manually classified those polygons that were left unclassified by the supervised classification. We added a sentence clarifying these important methodological details, comprised the object-based accuracy assessment.
R2	L 158-159: Why were the maps smoothed with a majority filter? If I understood the previous section correctly, this is already an object-based image classification, so smoothing the classified map would just change a few pixels along object boundaries?	The 3x3 majority filter was used to avoid aberrant very small objects coming from the previous RS-GIS operations.
R2	L 161: Are the authors referring to Cohen's Kappa coefficient here with K? If so, it should be defined and explicitly stated as there are many K statistics in different fields for different purposes. Also, if this is Cohen's Kappa, it is typically presented as an index of -1 to 1 rather than as a percentage.	Right, we specified that we used the Cohen's Kappa coefficient and corrected percentage values to -1 to 1 values.
R2	L 166: I appreciate that the authors have already made these data publicly available. It was nice to be able to look at them while reviewing this manuscript.	That was the idea, thanks!
R2	L 167: For the Mann-Whitney test, I'm a little unclear about the unit of the analysis here. Was it the count of 30 m cells within each study landscape? Or was it the amount of change within each landscape? Specific descriptions of sample size and units of analysis would help improve clarity here.	Right, we added a sentence clarifying that we used a sample size of 8 landscapes to compare the contribution (in hectares) of each category (5 categories = 5 M-W tests) to the transition into forest.
R2	L 169: It is a little unclear to me if the authors looked	Right, that part needed more

	only at cells that were unforested in the c. 1960 imagery for this analysis. So, the response was, for every 30 m cell (that was not forest in the 1960 period), a binary condition of 1) forest gain or 2) no transition or transition to another cover type? What about cells that were forest initially? This could be further clarified.	detailed explanations. We in fact considered only unforested pixels (30m size) and filtered out from the dataset those pixels that were forested in the past. Among the former unforested pixels we gave 1 to forest gain and 0 to unchanged or other transitions.
R2	L 173: Wouldn't this more appropriately be described as "1 m cells were aggregated to a 30 m resolution using the majority class within each 30 m pixel" or something similar? Nearest neighbor is more of an interpolation tool than an aggregation tool, and would imply that each 30 m cell in the aggregated map took the class label of the closest 1 m cell in the original map.	Right, we changed this part and the following analyses by using a majority resampling technique instead of using the NN.
R2	L 175-177: Given the target journal (scale being one of the key concepts in Landscape Ecology), I was a little surprised that this point was not discussed in more detail. I think this is particularly important because the RMSE of co-registration error (i.e., 23 m) means that some 30 m cells may actually be compared to an adjacent cell during the change detection. If results of the Random Forest analyses were qualitatively similar for the 30 and 60 m resolution (similar variable importance and trends in partial dependence) then the authors might present this (briefly) in the text, or as a separate figure in the appendix. If the variable importance and partial dependence plots were substantially different, then I think the scale-dependence of these results should be discussed quite a bit more than it is currently. One point of particular importance is that with this offset and a 30 m cell size, some cells that "transitioned" to forest along the previous forest edge could actually have been stable through time, but only appeared as change due to the offset between the two images. However, if the results are similar between 30 and 60 m then this may not be an issue. Either way, discussing this in further detail would be useful.	We explained more in detail the choice of the 30 m resolution and we added results regarding the RF model at 60 m resolution in the supplementary materials (Table S3). Specifically, our results showed that RF models were scarcely dependent on the spatial scale of the data and that the influence of co-registration errors on model predictions at 30 m was limited given the similarity with results obtained using data at 60 m resolution.
R3	L 182: For landscape codes see Table 1	We added the suggested sentence to that figure caption
R2	L 189-192: This section could benefit from a table, either in an appendix or in the main text, that lists all of the specific predictor variables, why they were included, and the authors' ideas of what they represent related to forests in these regions.	We agree and added a table (S4) in the supplementary material.
R2	L 199-201: I really like this approach of spatially stratified cross-validation in tuning hyperparameters. Any particular reason that eight clusters were selected? Given the nature of the data, I think that it might also make sense to use this spatially-stratified	The reason why we chose 8 partitions is that it corresponds to the number of landscapes in each mountain range, thus it should guarantee an adequate spatial

	method when calculating the predictive ability of the RF models themselves rather than using the default out-of-bag error. This would help to better assess the ability of these models to generalize to new areas.	partitioning. We ascertained this by plotting the spatial partitions of points created through repeated clustering of points coordinates based on k-means. Moreover, this number falls within the optimal number of partitions for cross-validation, ranging from 5 to 10. Now we assessed the predictive performances of the models using the Brier score and the AUC evaluation measures. We used a nested spatial cross-validation strategy in order to perform both repeated cross-validation and hyper-parameters tuning as described in the manuscript.
R2	L 211: For clarity, subsection titles could be modified to refer to specific data being used. For example, this subsection could be titled "Regional land cover changes - CORINE" and the following subsection could be "Landscape transitions - aerial imagery." Or the dataset could be referred to explicitly in the first sentence of text within each subsection. Here, something like: "Based on our analysis of CORINE land cover data from 1990-2018 in AL and AP, we found…" As it reads now, I had to glance back to the methods to remind myself what was being compared in each section.	We like the suggestion and modified the subsection titles.
R3	L 215: Realistic in such short time?	We believe that it is realistic at regional scale
R2	L 217: Nice. I think that the addition of this simple analysis with CORINE makes this more broadly interesting.	We agree with this.
R3	L 219: In my opinion, the change in area could also be expressed as a percentage of the total mountain area, allowing a direct comparison between mountain regions and a relativisation of the individual land cover changes.	We agree and modified the table accordingly.
R3	Table 2: I am surprised at the high proportion of this class. It is possible that Alpine grasslands was also included in the class. On unvegetated land a forest occurrence in such a short time is more than unrealistic. Normally there is no suitable soil. Please check.	With the new version of table 2 it is probably clearer that the high amount of UV category in the Alps (10%) is not a change, but the proportion of this category on the total surface of the alpine region that is rich of rocks, glaciers, gravel, etc… The % absolute change for this category is around -1, not so much.
R3	L 226: crops (CR)	Modified accordingly.
R2	Figure 2: I really like this figure but have a couple of suggestions. First, a different color palette might	We modified the colors in this figure to increase the contrast

	make things more visually appealing as the colors used here come off as a little "harsh" to the eye, particularly the light green and light blue. For example, "Set2" and "Dark2" in RColorBrewer might be nice choices for light and dark discrete palettes, or the "paired" palette that has paired light and dark colors. Second, I am a little bit unclear on the descriptions and use of "transitioning" vs "persistent" colors in this figure. It is hard for me to tell for sure, but in some cases (e.g., UV class in AL; GR class in AL and AP) the light-colored box actually represents persistence? Whereas in the FO class, the darker boxes appear to be persistence. Also, in at least one instance (e.g., CR to FO in AP), the number on the arrow is greater than the initial cover. Perhaps this is a typo in the label of the area of the CR class in AP historic?	between persistent and transitioning surfaces. We also improved the figure caption in order to make it clearer the difference between "transitioning" vs "persistent" land cover categories. We also fixed the typo in the label of CR category in AP.
R2	L 248: The numbers in parentheses here and later should be defined. Perhaps "importance =" with first use and "imp. =" after that?	We modified it by using the symbol “IR” meaning importance rate.
R2	L 249-251: Presumably, the climatic variables used in this analysis are strongly related to elevation and terrain. In the methods section, the authors might add a stronger justification for why they were all included instead of just climate or just topography. For example, climate could vary latitudinally (particularly in AP) or with distance to coastline (in AL) in ways that make these variables represent slightly different things than elevational trends in temp and precip. Alternatively, the authors could use a metric like climatic water deficit or actual evapotranspiration (e.g., Lutz et al. 2010; Journal of Biogeography) to combine the effects of temperature, precipitation, and heat load into a single variable at a higher resolution. At 1 km (which is a fairly high-resolution climate surface), the climatic predictors are still somewhat coarse compared to the 30 m unit of analysis, and there may only be a few unique climate cells within some of the smaller landscapes. This coarse spatial resolution seems like it has the potential to influence results of variable importance and OOB error (though using spatially-stratified cross-validation rather than OOB error might help to account for this component). I totally understand if the authors choose to use climate variables in the final analyses, but if they do, some discussion of the limitations of data resolution and its potential influence on the results would be valuable.	We understand the criticism raised by the reviewer, but the climatic dataset that was used (CHELSA) is less strongly influenced by topographic gradients compared to other climatic datasets (e.g. WorldClim). Anyway we added a sentence at the bottom of the discussion section reporting the limits of this approach linked to the spatial resolution mismatch between topographic-vegetation variables and climatic ones.
R2	Figure 5: Based on Fig. 4, it seems like the authors picked variables 1-3 and 5 for plotting. Why only these? Excluding the fourth ranked variable seems a little bit like cherry-picking what they thought were	Figure 4 and figure 5 were replaced by new versions following the new analyses made with resampled landscapes. In addition, we added

	the more the interesting results. I realize that there may not be enough space to show all of the partial plots, but including all of them in an appendix might be useful, and then specifically saying "We plotted the top 5 variables from each region and the remainder are in the appendix" or something along those lines. Also, for plotting, shouldn't the SD lines of partial dependence be included above and below the predicted effect as a sort of confidence interval? Rather than just below it?	the variable precipitation, in this way we plot the 5 most important predictors for both mountain ranges. We plotted SD lines only below the averaged predicted effect because they were presented in this way by the authors of the Individual Conditional Expectation plots (see Goldstein et al. 2015)
R2	L 281-282: The Norman et al. (2005) citation is a little bit dated for this kind of research in the western U.S. More recent citations showing afforestation in California -Lydersen, J. M., & Collins, B. M. (2018). Change in Vegetation Patterns Over a Large Forested Landscape Based on Historical and Contemporary Aerial Photography. Ecosystems, 21(7), 1348-1363. doi: 10.1007/s10021-018-0225-5 and Colorado - Rodman et al. (2019; already cited in methods section)	We modified that citation accordingly.
R3	L 286-287: Some examples: Tasser, E., Teutsch, A., Noggler, W. & Tappeiner U. (2007) Land-use changes and natural reforestation in the Eastern Central Alps. Agriculture, Ecosystems and Environment 118: 115–129. Fontana, V., Radtke, A., Walde, J., Tasser, E., Wilhalm, T., Zerbe, S., Tappeiner, U. (2014) What plant traits tell us: consequences of land-use change of a traditional agro-forest system on biodiversity and ecosystem service provision. Agriculture, Ecosystems and Environment 186, 44-53. Seidl, R., Albrich, K., Erb, K.-H., Formayer, H., Leidinger, D., Leitinger, G., Tappeiner, U., Tasser, E., Rammer, W. (2019) What drives the future supply of regulating ecosystem services in a mountain forest landscape? Forest Ecology and Management 445, 37-47. Wallentin G., Tappeiner U., Strobl J., Tasser E. (2008) Understanding alpine tree line dynamics: an individual based model. Ecological Modelling 218/3-4: 235-246. Tasser, E., Leitinger, G, Tappeiner, U. (2017) Climate Change versus Land-Use Change - which affects the landscape more? Land Use Policy 60: 60–72.	These papers are all very interesting, but we decided to cite the first two because they compare different landscapes in alpine regions characterized by post-abandonment forest expansion.
R3	L 293: Similar rates are also documented for the Alpine regions Carnia (0.7% a-1), Tyrol (0.35% a-1) and South Tyrol (0.1% a-1) in the period 1955 to 2000 (Tasser et al. 2007).	We added the whole sentence that gives an international breath to our paper.
R3	L 304-305: Some information in Zimmermann, P.D., Tasser, E., Leitinger, G., Tappeiner, U. (2010) Effects of land-use and land-cover pattern on landscape-scale biodiversity in the European Alps. Agriculture,	Right, we added that citation too.

	Ecosystem and Environment 139: 13–22.	
R2	L 306: As written, this kind of undersells the authors' analyses with CORINE. If I understand the issues with CORINE correctly (based on a quick look at the referenced paper) it seems like the primary issue is differences in mapping accuracy among different nations. Given that the study area is within a single country, this may not be an issue? So, perhaps this citation and portion of this sentence could be removed. Classification accuracy may be an issue, but this (and inaccuracy in classification of the air photos) could perhaps be briefly discussed in a limitations section instead. Similarly, classification accuracy of the CORINE datasets is not broadly quantified, but this could be completed pretty easily for the authors' study area (at least for a recent time period and the five aggregated classes) and it seems as if this would be a valuable contribution to the literature.	We removed part of that sentence accordingly. Anyway, the lack of speculative discussion on the CORINE analysis within our manuscript is due to the fact that even on a large study area such as the hole Italian Alps and the Apennines we found a sinusoidal behavior of forest surface trend from 1990 to subsequent study years.
R3	L 312: Hard to do in such a short time. I think there are classification problems.	The transition from unvegetated areas to forest is not so hard to believe. This is due to the fact that within this category we grouped rocks, gravel, sands, bare soil and sparse grasslands or sparse vegetation. The latter is a subcategory, mostly located at high elevation (between 2000 and 3000 m a.s.l.), composed by a mosaic of grasses and stones/sands.. Here the competition of encroaching trees with grasses and shrubs is less limiting. We added a sentence to methods section to explain the variability of this land cover category.
R2	L 316: If there is relevant literature, perhaps the authors could briefly discuss the methods of afforestation. Is it entirely from seed? Are there any places in which resprouting species had belowground tissue and are quickly growing back? This might be particularly important in rangeland/pastoral areas that were not intensive cropland - where grazing could allow resprouting species to persist but only at very low densities. A bit more about the ecology of these forest systems and mechanisms of expansion and infilling would be interesting in this paragraph or the next.	We added a couple of sentences and citations describing the main reproductive strategies of European beech and Mediterranean mountain pines.
R3	L 319-321: For the Alps: Zimmermann, P.D., Tasser, E., Leitinger, G., Tappeiner, U. (2010) Effects of land-use and land-cover pattern on landscape-scale biodiversity in the European Alps. Agriculture, Ecosystem and Environment 139: 13–22.	We added that citation for the Alps.

R3	L 322-328: See: Tasser, E., Teutsch, A., Noggler, W. & Tappeiner U. (2007) Land-use changes and natural reforestation in the Eastern Central Alps. Agriculture, Ecosystems and Environment 118: 115–129. Wallentin G., Tappeiner U., Strobl J., Tasser E. (2008) Understanding alpine tree line dynamics: an individual based model. Ecological Modelling 218/3-4: 235-246.	We added that citation for the Alps.
R3	L 336-338: See: Tasser, E., Leitinger, G, Tappeiner, U. (2017) Climate Change versus Land-Use Change - which affects the landscape more? Land Use Policy 60: 60–72.	We added that citation for the Alps.
R2	L 340: Shouldn't "legacy" be plural in this case? As in "disturbance legacies" or "biological legacies"?	Right, we modified accordingly.
R2	L 349-351: The effect of aspect (or heat load) is contingent upon elevation, correct? South-facing slopes at low elevations are fundamentally different than those at high elevations. It seems like they should be discussed together here to bring out some of that nuance. This same comment applies to the abstract. Climate is probably not universally more suitable for forests at low elevations?	We modified that sentence, accordingly.
R2	L 350: Perhaps "greater" rather than "larger"	We modified that sentence, accordingly.
R2	L 358-360: This sentence and the final one in the paragraph could use citations to support them.	We added those citations.
R2	L 368: The negative effects of afforestation in Italy could be listed here explicitly to help folks that are unfamiliar with management concerns in these regions. It is discussed in the intro but could be brought up again here.	We added a sentence to expand a bit more these concepts.
R2	L 378: This should be "legacies of land use" or "land use legacies", correct?	We used “legacies of land use”.
R2	L 379: Perhaps use: "…forest edges in combination with environmental factors such as…"	We modified that sentence, accordingly.
R3	L 378-385: This part contains only a summary and repetitions of the most important results. From my point of view this part should be deleted.	We decided to maintain this part because is a sort of concluding paragraph where we also highlight the limits of our research approach.
R2	L 387-388: I finished the article expecting more of a connection to the "big picture context" in the last paragraph. The authors do a nice job of this in the intro and in the start of the discussion, but this last paragraph could be adjusted slightly by connecting to some broader theoretical concepts or global issues. A couple of potential suggestions: 1) global patterns of afforestation vs deforestation and the carbon balance, or 2) linking recent change again with past patterns of forest cover and paleoecological literature (as is done effectively throughout the intro and reviewed in Vacchiano et al. 2017, Forest Ecology and	We added few sentences at the end of the discussion section, following the suggestion #2 of the reviewer #2.

	Management). Idea #2 could be situated in an interesting way to bring up the tricky question of "what is natural", "what is the desired condition", and are these convergent or achievable goals in a landscape heavily influenced by human impacts over many millenia.	
R3	Table S1: or grassland? (instead of “Sparsely vegetated areas”)	No, this is the official legend of CORINE land cover maps.

Authorship change form

Click here to access/download;Authorship change form;AuthorshipChangeSigned_LAND-D-20-00035R1.pdf

Change of authorship request form (pre-acceptance)

Please read the important information on page 4 before you begin

This form should be used by authors to request any change in authorship including changes in corresponding authors. Please fully complete all sections. Use black ink and block capitals and provide each author's full name with the given name first followed by the family name.

Please note: In author collaborations where there is formal agreement for representing the collaboration, it is sufficient for the representative or legal guarantor (usually the corresponding author) to complete and sign the Authorship Change Form on behalf of all authors.

Section 1: Please provide the current title of manuscript
(For journals: Please provide the manuscript ID, title and/or DOI if available.)
(For books: Please provide the title, ISBN and/or DOI if available.)

Manuscript ID: LAND-D-20-00035R1
DOI in case of published manuscript:
ISBN (for books):

Title: Contrasting land use legacy effects on forest landscape dynamics in the Italian Alps and the Apennines

Section 2: Please provide the previous authorship, in the order shown on the manuscript before the changes were introduced. Please indicate the corresponding author by adding (CA) behind the name.

	First name(s)	Family name	ORCID or SCOPUS id, if available
1st author	Matteo	Garbarino	0000-0002-9010-1731
2nd author	Donato	Morresi	0000-0002-3545-872X
3rd author	Francesco	Malandra	0000-0001-9317-8446
4th author	Renzo	Motta	0000-0002-1631-3840
5th author	Emanuele Marco	Sibona	0000-0003-1594-3725
6th author	Carlo	Urbinati	0000-0003-4879-1406
7th author	Alessandro	Vitali	0000-0002-1713-2152
8th author	Peter J	Weisberg	0000-0003-4883-9297

SPRINGER NATURE Springer Nature is one of the world's leading global research, educational and professional publishers, created in May 2015 through the combination of Nature Publishing Group, Palgrave Macmillan, Macmillan Education and Springer Science+Business Media.

Change of authorship request form (pre-acceptance)

Section 3: Please provide a justification for change. Please use this section to explain your reasons for changing the authorship of your manuscript, e.g. what necessitated the change in authorship? Please refer to the (journal) policy pages for more information about authorship. Please explain why omitted authors were not originally included and/or why authors were removed on the submitted manuscript.

The order of authors of our manuscript was slightly modified to better reflect the level of contribution of authors. **In particular, the "position" of Prof. Carlo Urbinati changed from 6th to 3rd, because while making the revisions** **suggested by the reviewers, the contributions by CU became greater, increasing his level of contribution to the overall paper.**

Section 4: Proposed new authorship. Please provide your new authorship list in the order you would like it to appear on the manuscript. Please indicate the corresponding author by adding (CA) behind the name. <u>If the corresponding author has changed, please indicate the reason under section 3.</u>

	First name(s)	Family name (this name will appear in full on the final publication and will be searchable in various abstract and indexing databases)
1st author	Matteo	Garbarino (CA)
2nd author	Donato	Morresi
3rd author	Carlo	Urbinati
4th author	Francesco	Malandra
5th author	Renzo	Motta
6th author	Emanuele Marco	Sibona
7th author	Alessandro	Vitali
8th author	Peter J	Weisberg

SPRINGER NATURE Springer Nature is one of the world's leading global research, educational and professional publishers, created in May 2015 through the combination of Nature Publishing Group, Palgrave Macmillan, Macmillan Education and Springer Science+Business Media.

Change of authorship request form (pre-acceptance)

Section 5: Author contribution, Acknowledgement and Disclosures. Please use this section to provide a new disclosure statement and, if appropriate, acknowledge any contributors who have been removed as authors and ensure you state what contribution any new authors made (if applicable per the journal or book (series) policy). Please ensure these are updated in your manuscript - after approval of the change(s) - as our production department will not transfer the information in this form to your manuscript.

New acknowledgements:
Not applicable

New Disclosures (financial and non-financial interests, funding):
Not applicable

New Author Contributions statement (if applicable per the journal policy):
Matteo Garbarino conceived the experiment and all authors contributed to the study design. Material preparation, data collection and analysis were performed by Matteo Garbarino, Donato Morresi, Emanuele Sibona, Francesco Malandra, Alessandro Vitali and Carlo Urbinati. The first draft of the manuscript was written by Matteo Garbarino and all authors commented on previous versions of the manuscript. All authors read and approved the final manuscript.

State 'Not applicable' if there are no new authors.

SPRINGER NATURE Springer Nature is one of the world's leading global research, educational and professional publishers, created in May 2015 through the combination of Nature Publishing Group, Palgrave Macmillan, Macmillan Education and Springer Science+Business Media.

Change of authorship request form (pre-acceptance)

Section 6: Declaration of agreement. *All* authors, unchanged, new and removed *must* sign this declaration.
(NB: Please print the form, sign and return a scanned copy. Please note that signatures that have been inserted as an image file are acceptable as long as it is handwritten. Typed names in the signature box are unacceptable.) *** Please delete as appropriate. Delete all of the bold if you were on the original authorship list and are remaining as an author.**

	First name	Family name		Signature	Affiliated institute	Date
1st author	Matteo	Garbarino	I agree to the proposed new authorship shown in section 4		University of Torino (IT)	April 4, 2020
2nd author	Donato	Morresi	I agree to the proposed new authorship shown in section 4		University of Torino (IT)	April 4, 2020
3rd author	Carlo	Urbinati	I agree to the proposed new authorship shown in section 4		Marche Polytechnic University (IT)	April 4, 2020
4th author	Francesco	Malandra	I agree to the proposed new authorship shown in section 4		Marche Polytechnic University (IT)	April 4, 2020
5th author	Renzo	Motta	I agree to the proposed new authorship shown in section 4		University of Torino (IT)	April 4, 2020
6th author	Emanuele Marco	Sibona	I agree to the proposed new authorship shown in section 4		University of Torino (IT)	April 4, 2020
7th author	Alessandro	Vitali	I agree to the proposed new authorship shown in section 4		Marche Polytechnic University (IT)	April 4, 2020
8th author	Peter J	Weisberg	I agree to the proposed new authorship shown in section 4		University of Nevada, Reno (USA)	April 4, 2020

Please use an additional sheet if there are more than 7 authors.

SPRINGER NATURE Springer Nature is one of the world's leading global research, educational and professional publishers, created in May 2015 through the combination of Nature Publishing Group, Palgrave Macmillan, Macmillan Education and Springer Science+Business Media.

Change of authorship request form (pre-acceptance)

Important information. Please read.

- Please return this form, fully completed, to Springer Nature. We will consider the information you have provided to decide whether to approve the proposed change in authorship. We may choose to contact your institution for more information or undertake a further investigation, if appropriate, before making a final decision.

- By signing this declaration, all authors guarantee that the order of the authors are in accordance with their scientific contribution, if applicable as different conventions apply per discipline, and that only authors have been added who made a meaningful contribution to the work.

- Please note, we cannot investigate or mediate any authorship disputes. If you are unable to obtained agreement from all authors (including those who you wish to be removed) you must refer the matter to your institution(s) for investigation. Please inform us if you need to do this.

- If you are not able to return a fully completed form within **30 days** of the date that it was sent to the author requesting the change, we may have to withdraw your manuscript. We cannot publish manuscripts where authorship has not been agreed by all authors (including those who have been removed).

- Incomplete forms will be rejected.

SPRINGER NATURE Springer Nature is one of the world's leading global research, educational and professional publishers, created in May 2015 through the combination of Nature Publishing Group, Palgrave Macmillan, Macmillan Education and Springer Science+Business Media.