

UNIVERSITÀ POLITECNICA DELLE MARCHE Repository ISTITUZIONALE

Mobilizing Crop Biodiversity

This is the peer reviewd version of the followng article:

Original

Mobilizing Crop Biodiversity / Mccouch, S.; Navabi, Z. K.; Abberton, M.; Anglin, N. L.; Barbieri, R. L.; Baum, M.; Bett, K.; Booker, H.; Brown, G. L.; Bryan, G. J.; Cattivelli, L.; Charest, D.; Eversole, K.; Freitas, M.; Ghamkhar, K.; Grattipaglia, D.; Henry, R.; Valadares Inglis, M. C.; Islam, T.; Kehel, Z.; Kersey, P. J.; King, G. J.; Kresovich, S.; Marden, E.; Mayes, S.; Ndjiondjiop, M. N.; Nguyen, H. T.; Paiva, S. R.; Papa, R.; Phillips, P. W. B.; Rasheed, A.; Richards, C.; Rouard, M.; Amstalden Sampaio, M. J.; Scholz, U.; Shaw, P. D.; Sherman, B.; Staton, S. E.; Stein, N.; Svensson, J.; Tester, M.; Montenegro Valls, J. F.; Varshney, R.; Visscher, S.; von Wettberg, E.; Waugh, R.; Wenzl, P.; Bieseberg, L. H.; MOLECULAR PLANT. - ISSN 1674-2052. - 19:10(2020), pp. 1341-1344. [10:1010/j.molp.2020.08.011]

Publisher:

Published DOI:10.1016/j.molp.2020.08.011

Terms of use:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. The use of copyrighted works requires the consent of the rights' holder (author or publisher). Works made available under a Creative Commons license or a Publisher's custom-made license can be used according to the terms and conditions contained therein. See editor's website for further information and terms and conditions. This item was downloaded from IRIS Università Politecnica delle Marche (https://iris.univpm.it). When citing, please refer to the published version.

note finali coverpage

(Article begins on next page)

Title: Mobilizing Crop Biodiversity

Authors: Susan McCouch¹, Katy Navabi^{2, 34}, Michael Abberton³, Noelle L Anglin⁴, Rosa Lia Barbieri⁵, Michael Baum⁶, Kirsten Bett⁷, Helen Booker⁸, Gerald L Brown⁹, Glenn J Bryan¹⁰, Luigi Cattivelli¹¹, David Charest¹², Kellye Eversole¹³, Marcelo Freitas⁵, Kioumars Ghamkhar¹⁴, Dario Grattapaglia⁵, Robert Henry¹⁵, Maria Cleria Valadares Inglis⁵, Tofazzal Islam¹⁶, Zakaria Kehel⁷, Paul J Kersey¹⁷, Stephen Kresovich¹⁸, Emily Marden¹⁹, Sean Mayes²⁰, Marie Noelle Ndjiondjop²¹, Henry T Nguyen²², Samuel Paiva⁵, Roberto Papa²³, Peter W B Phillips²⁴, Awais Rasheed²⁵, Christopher Richards²⁶, Mathieu Rouard²⁷, Maria Jose Amstalden Sampaio⁵, Uwe Scholz²⁸, Paul D Shaw¹⁰, Brad Sherman²⁹, S Evan Staton³⁰, Nils Stein²⁸, Jan Svensson³¹, Mark Tester³², Jose Francisco Montenegro Valls⁵, Rajeev Varshney³³, Stephen Visscher³⁴, Eric von Wettberg³⁵, Robbie Waugh¹⁰, Peter W B Wenzl³⁶, Loren H. Rieseberg*³⁰

Affiliations:

1 Plant Breeding and Genetics, School of Integrated Plant Sciences, Cornell University, Ithaca, NY, 14853, USA

2 DivSeek, Global Institute for Food Security, 110 Gymnasium Place, University of Saskatchewan, Saskatoon, SK, S7N 0W9, Canada

3 International Institute of Tropical Agriculture (IITA), PMB 5320, Oyo Rd, Ibadan, Nigeria

4 International Potato Center (CIP) 1895 Avenida La Molina, Lima Peru 12, Lima 15023, Peru

5 Embrapa Genetic Resources and Biotechnology, Parque Estação Biológica, Final Av W5 Norte, Caixa Postal 02372, 70770-917 - Brasília DF, Brazil

6 International Center for Agricultural Research in the Dry Areas (ICARDA), Station Exp. INRA-Quich. Rue Hafiane Cherkaoui. Agdal. Rabat – Instituts, 10111, Rabat, Morocco

7 University of Saskatchewan, 51 Campus Dr., Saskatoon, SK S7N 5A8, Canada

8 University of Guelph, Rm 316, Crop Science Bldg, 50 Stone Rd E, Guelph, ON, N1G 2W1, Canada

9 Genome Prairie, 111 Research Drive, Suite 101, Saskatoon, SK, S7N 3R2, Canada

10 The James Hutton Institute, Errol Road, Invergowrie, Dundee, DD2 5DA, UK

11 CREA, Research Centre for Genomics and Bioinformatics, via San Protaso 302, Fiorenzuola d'Arda, 29017, Italy

12 Genome British Columbia, 400-575 West 8th Avenue, Vancouver, BC, V5Z 0C4, Canada

13 International Wheat Genome Sequencing Consortium 2841 NE Marywood Ct, Lee's Summit, MO, 64086, USA

14 AgResearch Forage, Science, Grasslands Research Centre, AgResearch, Palmerston North, 4410, New Zealand

15 Queensland Alliance for Agriculture and Food Innovation, Queensland Alliance for Agriculture and Food Innovation, University of Queensland, Brisbane, QLD, 4072, Australia

16 Institute of Biotechnology and Genetic Engineering (IBGE), Bangabandhu Sheikh Mujibur Rahman Agricultural University, Gazipur 1706, Bangladesh

17 Royal Botanic Gardens, Kew, Richmond, Surrey, TW9 3AE, UK

18 Feed the Future Lab for Crop Improvement, 431 Weill Hall, Cornell University, Ithaca, NY, 14853, USA

19 University of British Columbia, 3083 Alma St, Vancouver, BC, V6R 3S9, Canada

20 Crops For the Future (UK) CIC 76-80 Baddow Road, Chelmsford, Essex, CM2 7PJ, UK

21 Africa Rice Center (AfricaRice), Mbe Research Station, Bouaké, 01 BP 2511 Bouaké, Côte d'Ivoire

22 University of Missouri, Division of Plant Sciences, 25 Agriculture Lab Bldg, College of Agriculture,

Food and Natural Resources, University of Missouri, Columbia, MO 65211, USA

23 Università Politecnica delle Marche, Via Brecce Bianche, 60131, Ancona, Italy

24 Johnson Shoyama Graduate School of Public Policy, University of Saskatchewan, 101 Diefenbaker Place, Saskatoon, S7N 5B8, Canada

25 CIMMYT-China office, Beijing 100081, Beijing, P.R. China

26 USDA-ARS National Laboratory for Genetic Resources Preservation, 1111 South Mason St, Fort Collins, CO, 80521, USA

27 Bioversity International, Parc Scientifique Agropolis II, 34397, Montpellier, Cedex 5, France

28 Leibniz Institute of Plant Genetics and Crop Plant Research (IPK), Gatersleben, Corrensstr. 3, D-06466 Stadt Seeland, Germany

29 Law School, University of Queensland, St Lucia, QLD, 4072, Australia

30 Department of Botany and Biodiversity Research Centre, University of British Columbia, Vancouver, BC, V6R 2A5, Canada

31 NordGen, Smedjevagen 3, 230 53 Alnarp, Sweden

32 King Abdullah University of Science & Technology (KAUST), Thuwal, 23955-6900, Saudi Arabia 33 International Crops Research Institute for the Semi-Arid Tropics (ICRISAT), Patancheru - 502 324, Telangana State, India

34 Global Institute for Food Security, 110 Gymnasium Place, University of Saskatchewan, Saskatoon, SK, S7N 4J8, Canada

35 University of Vermont, 63 Carrigan Drive, Jeffords Hall, Burlington, VT, 05405, USA

36 Centro Internacional de Agricultura Tropical (CIAT), Km 17 Recta Cali-Palmira, 763537 Cali, Colombia

* Correspondence: Loren H. Rieseberg,

Contact Information: https://www.icea.com https://www.icea.com

Journal Pre-proof

1 Over the past 70 years, the world has witnessed extraordinary growth in crop productivity, enabled by a 2 suite of technological advances, including higher yielding crop varieties, improved farm management, synthetic agrochemicals, and agricultural mechanization. While this "Green Revolution" intensified crop 3 production, and is credited with reducing famine and malnutrition, its benefits were accompanied by 4 5 several undesirable collateral effects (Pingali, 2012). These include a narrowing of agricultural 6 biodiversity, stemming from increased monoculture and greater reliance on a smaller number of crops and 7 crop varieties for the majority of our calories. This reduction in diversity has created vulnerabilities to 8 pest and disease epidemics, climate variation, and ultimately to human health (Harlan, 1972). The value of crop diversity has long been recognized (Vavilov, 1992). A global system of genebanks (e.g. 9 10 www.genebanks.org/genebanks/) was established in the 1970s to preserve the abundant genetic variation found in traditional "landrace" varieties of crops and in crop wild relatives (Harlan, 1972). While 11 preserving crop variation is a critical first step, the time has come to make use of this variation to breed 12 more resilient crops. The DivSeek International Network (https://divseekintl.org/) is a scientific, not-for-13 14 profit organization that aims to accelerate such efforts. 15

16 Crop diversity: value, barriers to use, and mitigation strategies

There are >1750 national and international genebanks worldwide. They house ~7 million crop germplasm
accessions (<u>http://www.fao.org/3/i1500e/i1500e00.htm</u>), including samples of diverse natural
populations, with many more managed *in situ*. These accessions arguably represent one of humanity's
greatest treasures, as they contain genetic variation that can be harnessed to create better tasting, higher
yielding, disease/pest resistant, and climate resilient cultivars that require fewer agricultural inputs (Figure
1).

23 Unfortunately, most genebank accessions are poorly characterized, and few have been utilized in 24 breeding. Yet when a serious effort has been made to search genebanks for traits of interest, the effort has 25 been highly rewarded. Examples include the discovery of a submergence-tolerant landrace used to breed new, high-yielding, submergence-tolerant rice varieties currently grown on tens of millions of acres 26 27 (Mackill et al., 2012) and durable resistance to late blight, a devastating pathogen of potato, derived from a wild relative (Bernal-Galeano, 2020). Given the high value of the genetic diversity found in crop wild 28 29 relatives and traditional landraces, why are these genetic resources not more widely employed in breeding 30 programs?

31 One reason for the limited use of genebank holdings is the paucity of information about them, which 32 increases the time, expense, and risk associated with mining genebank diversity. To address this

Journal Pre-proof

deficiency, we support the development of digital catalogs that provide essential information about the genetic composition, phenotypic diversity and phylogenetic relationships of genebank holdings, along with traditional passport data, images of whole plant morphology, growth habit, physiological data showing response to biotic/abiotic stress, nutrient profiles, and other information where available. Some genebanks have already begun building catalogs of their collections to improve the efficiency of genebank management, as well as to permit users to pre-screen for traits of interest, thereby facilitating variety development (König et al., 2020).

Another challenge to widespread use of genebank materials is the nature of genetic variation itself. Exotic
 germplasm often contains valuable cryptic variation, which is revealed only after crosses have been made

42 with cultivated and elite breeding lines (Tanksley and McCouch, 1997). For example, wild populations

43 frequently carry alleles that increase seed/ fruit/ tuber size or disease resistance when introduced into

44 cultivars, but these are often masked by genes with opposing effects. Also, some traits that look

45 promising in wild or landrace populations may not be expressed in adapted genetic backgrounds due to

- 46 quantitative inheritance.
- 47 In addition, strategies are needed to overcome crossing barriers and to ameliorate the impacts of genetic
- 48 material that is inadvertently introduced into cultivars along with traits/alleles of interest (i.e. linkage

drag). Even when crosses are successful, specific chromosomal segments may fail to introgress if they

50 underlie hybrid incompatibilities or experience reduced recombination, further exacerbating linkage drag

51 (Canady et al., 2006). Lastly, traits and alleles introgressed from wild germplasm may exhibit incomplete

- penetrance or unexpected epistatic interactions, forfeiting expected gains from introgressions (Lippman et al., 2007).
- 54 Sorting through the myriad combinations of alleles generated in wild x elite crosses requires a systematic
- approach if it is to be productive. The use of structured populations, appropriate experimental designs,
- and effective use of reference varieties in combination with cost-effective genotyping, high throughput
- 57 phenotyping, automated data capture and appropriate analyses make it possible to link genotype with
- 58 phenotype, identify valuable haplotypes, drive recombination, and make predictions about offspring
- 59 phenotypes. Techniques that enhance recombination and mitigate crossability barriers offer additional
- 60 means for accessing diversity from divergent wild relatives while reducing linkage drag (Fernandes et al.,
- **61** 2018).
- 62 To address these challenges, we encourage communities of researchers to undertake systematic pre-
- 63 breeding efforts to generate recombinant populations of introgressed lines in adapted cultivated
- backgrounds, evaluate them in diverse environments, and share the lines and associated information with
- breeders, farmers, researchers, and policy makers. The long time horizon and uncertainties associated

- 66 with "pre-breeding" often impede investment from private breeding programs. Therefore, we urge
- 67 increased investment from foundations and the public sector to support such efforts across major crop
- 68 families, expanding on recent efforts by the Global Crop Diversity Trust, CGIAR, and other
- 69 organizations. As products of pre-competitive research, pre-bred lines could be deposited into genetic
- stock centers and made available to both public and private breeding programs with explicit procedures to
- 71 fulfill access and benefit sharing obligations (see below).
- An alternative approach to pre-breeding involves the use of genome editing, which can be used to re-
- 73 introduce favorable alleles from wild and exotic relatives into crop plants, purge deleterious alleles, break
- 74 linkage drag, or create new alleles designed to enhance plant performance and resilience (Johnsson et al.,
- 75 2019; Zsögön et al., 2018). This approach allows researchers to explore natural variation as a key to
- resilience, and its application rests on a deep knowledge of the genetics and evolution of key traits and
- alleles. The use of genome editing also introduces a need for community discussion about regulatory
- requirements and international agreements to address the complex political, social, legal and economic
- 79 concerns surrounding the use of this technology (Lassoued et al., 2019).
- 80 Lastly, national and international policies related to benefit sharing derived from the use of plant genetic
- 81 resources impact how such resources and associated information are collected, stored, shared, studied,
- 82 and used, creating additional obstacles to research and the utilization of crop diversity (Marden 2018;
- 83 McCouch et al., 2013). The International Treaty for Plant Genetic Resources in Food and Agriculture
- 84 facilitates multi-lateral access to plant genetic resources under mutually agreed-upon terms. It currently
- 85 covers 64 crops, but ambiguity regarding benefit-sharing requirements impedes the use of genebank
- 86 holdings by many plant breeders, researchers, and farmers (Sherman and Henry, 2020). There also are
- 87 concerns that the benefit sharing provisions of the Treaty conflict with the long-accepted practice of
- 88 providing open access to genetic sequence data (Marden 2018). In our view, open sharing of information
- 89 about plant genetic resources, represents an essential form of benefit-sharing and provides a critical
- 90 foundation for capacity building strategies that help address UN sustainable development goals. It is
- 91 important that those employing genomic and phenomic information for crop research and breeding are
- 92 fully aware of international treaties and comply with their requirements.

93 Mission of DivSeek International Network

- 94 The DivSeek International Network is a global, community-driven organization that facilitates the
- 95 generation, integration and sharing of information related to plant genetic resources, thereby empowering
- 96 genebank managers, researchers, breeders, and farmers to more effectively utilize genetic variation for
- 97 research, accelerated crop improvement, and sustainable production. DivSeek comprises ~65 members
- 98 from >30 countries, and includes a broad array of academic and research institutions, government

Journal Pre-proo

- 99 agencies, and inter-governmental organizations (<u>https://divseekintl.org/members/</u>). To help achieve its
- 100 goals, DivSeek has established several Working Groups to engage members and assist them in addressing
- 101 issues of importance to the DivSeek Community. The activities undertaken by DivSeek's three current
- 102 Working Groups are summarized below.

103 *Genomics for Plant Genetic Resources:* DivSeek supports open-source genomic-assisted germplasm

- 104 management and breeding, which represents a decentralized form of empowerment for genebanks and
- 105 national breeding programs (Santantonio et al., 2020). Similar to the revolution in information technology
- that invented the internet and put cell phones in the hands of people throughout the world, open-source
- 107 genomics tools, strategies and datasets are being developed and shared internationally. The tools provide
- 108 data and information to support decisions about germplasm management and variety development, and
- 109 the use and iterative improvement of these tools by communities of practice has the power to accelerate
- the deployment of crop diversity in farmers' fields, helping to address several of the UN's Sustainable
- 111 Development Goals.
- 112 *Phenomics, ontologies and standards:* DivSeek promotes the use of new technologies for quantitative
- 113 phenotypic evaluation of plant genetic resources across a network of test environments, and the
- application of community-based standards, ontologies and data management practices that help make data
- findable, accessible, interoperable and reusable (FAIR) (Pommier et al., 2019). Utilizing efficient and
- affordable technologies will be key to engaging genebanks and plant breeders in modern phenomics-
- 117 based screening (Mir et al., 2019). Integration of diverse datasets boosts the power of global efforts to
- 118 document phenotypic variation found in both genebank accessions and in breeding populations (Roitsch
- et al., 2019), and can greatly improve the accuracy of predictions about plant performance across
- 120 environments. This is especially critical for accelerating the breeding of climate-resilient varieties in
- 121 vulnerable environments.
- 122 International Policies: DivSeek aims to help members of the international plant community to understand
- 123 the legal and policy framework for sharing information about plant genetic resources, lead discussions
- about the technological requirements for data-sharing across constituencies, and share perspectives on
- 125 benefit sharing practices that are aligned with international treaties. All international agreements
- 126 governing the utilization of plant genetic resources share the same basic objectives: conservation and
- 127 sustainable use of resources, ease of access to them, and fair and equitable sharing of benefits derived
- 128 from their use (<u>https://www.cbd.int/; http://www.fao.org/plant-treaty/en/</u>). However, rapid technological
- 129 developments are changing the way scientists explore, utilize, and exchange information about plant
- 130 genetic resources, creating new value for the information itself, and new opportunities for access and
- 131 benefit-sharing, while at the same time challenging existing agreements (Marden 2018). In particular, new

- 132 breeding techniques that can take advantage of genomic and phenotypic data without accessing physical
- 133 germplasm have led to a debate about open access to sequence data and the best ways to implement
- 134 benefit-sharing requirements (Laird et al., 2020).
- 135

136 Call for Global Participation

- 137 International collaborative partnerships are essential for addressing global challenges, ranging from
- 138 climate change to the control of pests and diseases to the conservation of biodiversity. DivSeek represents
- 139 one such global partnership, focusing on the characterization and utilization of agricultural biodiversity
- 140 and its impact on food and nutritional security.
- 141 The success of DivSeek will depend on attracting a broad coalition of members, observers, and
- stakeholders dedicated to discussion and constructive exchange of ideas, perspectives, and expertise. We
- 143 are pleased to invite the global agricultural science community to join the DivSeek International Network,
- 144 either as members or observers (<u>https://divseekintl.org/apply-to-join/</u>). By joining forces, we can mobilize
- the value of crop diversity to sustainably improve yields in farmers' fields and ensure that the benefits of
- 146 our efforts are equitably distributed across the globe.

References

Bernal-Galeano V, N.G., Ellis D, Anglin NL, Hareau G, Smale M, Jamora N, Alwang J, Pradel W. (2020). The effect of Andean potato diversity conserved in the International Potato Center genebank helps develop agriculture in Uganda: the example of Victoria. Food Security Journal, https://doi.org/10.1007/s12571-020-01037-8

Canady, M.A., Ji, Y., and Chetelat, R.T. (2006). Homeologous recombination in Solanum lycopersicoides introgression lines of cultivated tomato. Genetics 174:1775–1788.

Fernandes, J.B., Séguéla-Arnaud, M., Larchevêque, C., Lloyd, A.H., and Mercier, R. (2018). Unleashing meiotic crossovers in hybrid plants. Proceedings of the National Academy of Sciences USA 115:2431-2436.

Harlan, J.R. (1972). Genetics of Disaster. Journal of Environmental Quality 1:212-215.

Johnsson, M., Gaynor, R.C., Jenko, J., Gorjanc, G., de Koning, D.-J., and Hickey, J.M. (2019). Removal of alleles by genome editing (RAGE) against deleterious load. Genetics Selection Evolution 51:14.

König, P., Beier, S., Basterrechea, M., Schüler, D., Arend, D., Mascher, M., Stein, N., Scholz, U., and Lange, M. (2020). BRIDGE - A visual analytics web tool for barley genebank genomics. Frontiers in Plant Science 11:701-701.

Laird, S., Wynberg, R., Rourke, M., Humphries, F., Muller, M.R., and Lawson, C. (2020). Rethink the expansion of access and benefit sharing. Science 367:1200.

Lassoued, R., Macall, D.M., Smyth, S.J., Phillips, P.W.B., and Hesseln, H. (2019). Risk and safety considerations of genome edited crops: Expert opinion. Current Research in Biotechnology 1:11-21.

Lippman, Z.B., Semel, Y., and Zamir, D. (2007). An integrated view of quantitative trait variation using tomato interspecific introgression lines. Current Opinion in Genetics & Development 17:545-552.

Mackill, D.J., Ismail, A.M., Singh, U.S., Labios, R.V., and Paris, T.R. (2012). Development and rapid adoption of submergence-tolerant (Sub1) rice varieties. Advances in Agronomy 115:299-352.

Marden, E. (2018). International agreements may impact genomic technologies. Nature Plants 4:2-4.

McCouch, S., Baute, G.J., Bradeen, J., Bramel, P., Bretting, P.K., Buckler, E., Burke, J.M., Charest, D., Cloutier, S., Cole, G., et al. (2013). Feeding the future. Nature 499:23-24.

Mir, R.R., Reynolds, M., Pinto, F., Khan, M.A., and Bhat, M.A. (2019). High-throughput phenotyping for crop improvement in the genomics era. Plant Science 282:60-72.

Pingali, P.L. (2012). Green Revolution: Impacts, limits, and the path ahead. Proceedings of the National Academy of Sciences USA 109:12302-12308.

Pommier, C., Michotey, C., Cornut, G., Roumet, P., Duchêne, E., Flores, R., Lebreton, A., Alaux, M., Durand, S., Kimmel, E., et al. (2019). Applying FAIR principles to plant phenotypic data management in GnpIS. Plant Phenomics 2019:1671403.

Sherman, B., and Henry, R.J. (2020). The Nagoya Protocol and historical collections of plants. Nature Plants 6:430.

Tanksley, S.D., and McCouch, S.R. (1997). Seed banks and molecular maps: unlocking genetic potential from the wild. Science 277:1063-1066.

Vavilov, N.I. (1992). Origin and geography of cultivated plants. New York, NY, USA: Cambridge University Press.

Zsögön, A., Cermak, T., Naves, E., Notini, M., Edel, K., Weinl, S., Freschi, L., Voytas, D., Kudla, J., and Peres, L. (2018). *De novo* domestication of wild tomato using genome editing. Nature Biotechnology 36:1211–1216.

Journal Pre-proof

Figure 1. Sunflower pre-bred line containing introgressions from wild *Helianthus annuus* performing well in drought stress trial in Uganda. Pre-bred lines developed by Greg Baute and Loren Rieseberg at the University of British Columbia. Drought stress trial performed by Walter Anyanga, National Semi-Arid Resources Research Institute, Uganda. Photo Credit: Walter Anyanga.



