

Age and onset timing of Raynaud's phenomenon and first non-Raynaud symptom as prognostic factors in systemic sclerosis: a retrospective analysis from the Italian national multicenter Systemic Sclerosis Progression INvestiGation registry of the Italian Society for Rheumatology (SPRING-SIR)

Silvia Peretti* , Cosimo Bruni* , Francesco Bonomi ,
Rossella De Angelis, Gianluigi Bajocchi, Dilia Giuggioli, Martina Orlandi,
Giovanni Zanframundo , Roberta Foti, Elisa Visalli, Giovanna Cuomo, Alarico Ariani,
Edoardo Rosato, Gemma Lepri, Francesco Girelli, Valeria Riccieri, Elisabetta Zanatta,
Silvia Laura Bosello, Ilaria Cavazzana, Francesca Ingegnoli, Maria De Santis,
Fabio Cacciapaglia , Giuseppe Murdaca, Giuseppina Abignano, Giorgio Pettiti,
Alessandra Della Rossa, Maurizio Caminiti, Annamaria Iuliano, Giovanni Ciano,
Lorenzo Beretta, Gianluca Bagnato, Ennio Lubrano, Ilenia De Andres, Luca Idolazzi ,
Marta Saracco, Cecilia Agnes, Corrado Campochiaro, Giacomo De Luca,
Edoardo Cipolletta, Marco Fornaro, Federica Lumetti, Amelia Spinella,
Luca Magnani, Veronica Codullo, Carlo Iandoli, Antonietta Gigante, Greta Pellegrino,
Erika Pigatto, Maria Grazia Lazzaroni, Enrico De Lorenzis, Gianna Angela Mennillo,
Marco Di Battista, Giuseppa Pagano Mariano, Federica Furini, Licia Vultaggio,
Simone Parisi, Clara Lisa Peroni, Gerolamo Bianchi, Enrico Fusaro,
Gian Domenico Sebastiani, Marcello Govoni, Salvatore D'Angelo ,
Franco Cozzi, Franco Franceschini, Serena Guiducci, Lorenzo Dagna, Andrea Doria ,
Carlo Salvarani, Maria Antonietta D'Agostino, Florenzo Iannone ,
Marco Matucci-Cerinic, Clodoveo Ferri and Silvia Bellando Randone; on behalf of
SPRING-SIR (Systemic Sclerosis PROgression INvestiGation group of the Italian
Society for Rheumatology)

Abstract

Background: The sequence and temporal relationship between Raynaud's phenomenon (RP) and the first non-Raynaud's sign/symptom (NRP) in systemic sclerosis (SSc) have been partially investigated.

Objectives: To evaluate whether the mode and ages of clinical onset are associated with disease endotype and survival in SSc.

Design: We included SSc patients from the Systemic sclerosis Progression INvestiGation registry of the Italian Society of Rheumatology (SPRING-SIR) registry in a cohort study, with post hoc cross-sectional and longitudinal analysis.

Methods: Patients were grouped based on age-RP and age-NRP quartiles. Additionally, categories were defined based on mode of onset: RP group—RP onset at least 1 year before NRP; Simultaneous group—RP onset within the same year of NRP; NRP group—RP onset after at least 1 year after NRP. Comparisons were made using Chi-square and ANOVA tests.

Ther Adv Musculoskelet Dis

2026, Vol. 18: 1–19

DOI: 10.1177/
1759720X251410243

© The Author(s), 2026.
Article reuse guidelines:
sagepub.com/journals-
permissions

Correspondence to:

Cosimo Bruni
Department of
Rheumatology, University
Hospital Zurich, University
of Zurich, Zurich 8091,
Switzerland

Department of
Experimental and Clinical
Medicine, University of
Florence, Florence, Italy
cosimo.bruni@usz.ch

Silvia Peretti
Francesco Bonomi
Serena Guiducci
Gemma Lepri
Silvia Bellando Randone
Department of
Experimental and Clinical
Medicine, University of
Florence, Florence, Italy

Division of Rheumatology,
Department of Medical
and Geriatric specialties,
Careggi University
Hospital, University of
Florence, Florence, Italy

Rossella De Angelis
Edoardo Cipolletta
Rheumatology Unit,
Department of Clinical
and Molecular Sciences,
Polytechnic University of
Marche, Ancona, Italy

Gianluigi Bajocchi
Luca Magnani
Rheumatology Unit,
S. Maria Hospital-USL,
IRCCS Institute, Reggio
Emilia, Italy

Dilia Giuggioli
Martina Orlandi
Federica Lumetti
Amelia Spinella
Carlo Salvarani
Department of Medical
and Surgical Sciences
for Children and Adults,
University Hospital of
Modena and Reggio
Emilia School of Medicine,
Modena, Italy

Giovanni Zanframundo
Veronica Codullo
Department of
Internal Medicine and
Therapeutics, Università di
Pavia, Pavia, Italy

Division of Rheumatology,
Fondazione IRCCS
Policlinico San Matteo,
Pavia, Italy

Roberta Foti

Elisa Visalli

Rheumatology Unit, AOU
Policlinico San Marco,
Catania, Italy

Giovanna Cuomo

Carlo Iandoli

Department of Precision
Medicine, University
of Campania—Luigi
Vanvitelli University,
Naples, Italy

Alarico Ariani

Department of Medicine,
Internal Medicine
and Rheumatology,
Azienda Ospedaliero—
Universitaria di Parma,
Parma, Italy

Edoardo Rosato

Antonietta Gigante

Department of
Translational and
Precision Medicine,
Sapienza University of
Rome, Rome, Italy

Francesco Girelli

Department of
Medicine, Rheumatology
Unit, Ospedale GB
Morgagni—L. Pierantoni,
Forlì, Italy

Valeria Riccieri

Department of Internal
Medicine, Anesthesiology
and Cardiovascular
Sciences, Sapienza
University of Rome,
Rome, Italy

Elisabetta Zanatta

Andrea Doria

Department of
Rheumatology, University
of Padua, Padua, Italy

Silvia Laura Bosello

Enrico De Lorenzis

Maria Antonietta

D'Agostino

Rheumatology Division,
Catholic University of the
Sacred Heart, Fondazione
Policlinico Universitario
A. Gemelli—IRCCS,
Rome, Italy

Ilaria Cavazzana

Maria Grazia Lazzaroni

Franco Franceschini

Rheumatology and
Clinical Immunology,
ASST Spedali Civili of
Brescia, Brescia, Italy

Department of Clinical
and Experimental
Sciences, University of
Brescia, Brescia, Italy

Logistic, linear, and multinomial regression models were applied to assess associations, while Kaplan–Meier curves and Cox regression were used to assess mortality.

Results: A total of 1748 patients were eligible: 682 (39.0%) in the RP group, 1026 (58.8%) in the simultaneous group, and 39 (2.2%) in the NRP group. A higher prevalence of anti-centromere antibodies was found in the RP group, while the simultaneous group had more diffuse cutaneous SSc (dcSSc), anti-topoisomerase-I antibodies, and higher Rodnan's skin score (mRSS). The NRP group presented higher prevalence of pulmonary arterial hypertension. On logistic regression, the simultaneous group was associated with a higher prevalence of dcSSc compared to the RP group (odds ratio, 1.491, 95% confidence interval (CI): 1.032–2.154). Younger age at RP onset was associated with lower systolic pulmonary artery pressure and mRSS. In 943 patients with available follow-up (median 24 months), the simultaneous group had higher mortality compared to the RP group (hazard ratio, 1.975, 95% CI: 1.002–3.893).

Conclusion: The timing of RP and NRP onset may help define SSc endotype and survival. Patients with simultaneous RP-NRP onset have more severe disease features and higher mortality risk, emphasizing the relevance of onset timing in disease stratification.

Plain language summary

How systemic sclerosis starts impacts how the disease will behave

Systemic sclerosis (SSc) is a rare autoimmune disease that affects the skin and internal organs. In most cases the disease starts with Raynaud's phenomenon (RP), a discolouring of the fingers in reaction to cold temperatures; some other patients may experience different symptoms first, such as joint pain, skin changes or shortness of breath. Understanding which symptom appears first, and at what age, may help doctors identify patients who are more likely to develop severe disease. This study included 1,748 SSc patients enrolled in the Italian SPRING-SIR registry, divided patients into three groups: those who first developed RP (RP group), those who developed RP and other symptoms in the same year (Simultaneous), and those who developed non-RP symptoms first (NRP group). We found that patients in the RP group were younger at disease onset and often had milder disease. Patients in the Simultaneous group were more likely to develop diffuse skin disease, lung complications, and had a higher risk of death. Patients in the NRP group were rare but often had aggressive involvement of heart and lung. Overall, our results show that the timing of RP and non-RP symptoms can help predict disease course and survival in SSc. These findings may help doctors recognize higher-risk patients earlier and provide closer monitoring and tailored treatments.

Keywords: disease onset, non-Raynaud's symptoms, Raynaud's phenomenon, risk assessment, systemic sclerosis

Received: 28 May 2025; revised manuscript accepted: 8 December 2025.

Introduction

Systemic sclerosis (SSc) is a rare autoimmune disease characterized by widespread vascular damage, immune dysregulation, and fibrotic changes affecting the skin and internal organs.^{1,2} The disease primarily affects females, with a female-to-male ratio ranging from 3:1 to 8:1, and

shows two incidence peaks in the age groups 45–54 and 65–74 years based on race and sex.³

Among its diverse clinical presentations, Raynaud's phenomenon (RP) occurs in over 95% of SSc patients. It is the commonest initial symptom in SSc patients and generally precedes the onset of the

disease by several years,⁴⁻⁷ especially in the limited cutaneous SSc subset (lcSSc). Conversely, patients exhibit other non-Raynaud's signs or symptoms (NRP) soon after the development of RP in the diffuse cutaneous subset (dcSSc).^{8,9} A short interval (≤ 1 year) between RP and the appearance of NRP symptoms has been previously associated with more aggressive disease endotype, including a higher prevalence of anti-topoisomerase I antibodies (ATA), dcSSc, and a lower 10-year survival rate. This association was demonstrated in two large Italian cohort studies conducted by Ferri *et al.*,^{10,11} which highlighted the prognostic significance of symptom onset timing in SSc. More recently, De Angelis *et al.*¹² analyzed the SPRING registry focusing on the sine scleroderma subset (ssSSc), and demonstrated that the timing from RP onset to diagnosis significantly differed across SSc skin subsets, being longer in ssSSc compared to both lcSSc and dcSSc.¹² These findings underline the importance of onset timing as a prognostic factor, but evidence from large cohorts stratifying patients by RP- or NRP-onset is lacking.

Furthermore, other studies have identified factors associated with worse outcomes in SSc, such as male sex, dcSSc, interstitial lung disease (ILD), and older age at disease onset.^{10,11,13-19} The relationship between age at RP onset, the age of onset of the first non-Raynaud's phenomenon sign or symptom (NRP), and their timing (i.e., whether RP occurs before, after, or simultaneously with NRP) remains partially underexplored, although initial evidence suggests that NRP symptoms/signs at onset are independent predictors of more aggressive disease features such as visceral involvement, and worse survival compared to RP at onset.²⁰

The primary aim of our investigation was to assess whether the mode of SSc onset (RP vs NRP) along with the age at their onset are associated with the disease endotype and the prognosis in SSc patients. This data would further underline the importance of stratifying SSc patients based on RP and NRP patterns of onset, aiming to improve risk assessment and support the development of personalized management strategies.²⁰

Methods

Study design

We performed a cohort study, including a post hoc, cross-sectional analysis to evaluate the association between the mode of disease onset and

clinical endotype in patients with SSc. In addition, we conducted an exploratory longitudinal analysis to compare survival rates across patients presenting with different onset patterns. The reporting of this study conforms to the "Strengthening the Reporting of Observational Studies in Epidemiology (STROBE)" statement.²¹

Data source

We extracted relevant data from the Systemic sclerosis Progression INvestiGation registry of the Italian Society of Rheumatology (SPRING-SIR). SPRING-SIR is a multicenter, national cohort study promoted by the Italian Society of Rheumatology (SIR) which involved 38 referral centers across Italy, with expertise in the diagnosis and management of SSc. The study was conducted in accordance with the Declaration of Helsinki. The protocol was approved by the coordinating ethics committee of AOU Careggi, Florence (reference number OSS 15.010 AOU Careggi, Firenze, Italy), and by the ethics committees of all participating centers. All participants had previously provided written informed consent at the time of enrollment in the SPRING-SIR registry; therefore, no additional consent was required for this retrospective analysis.

For each patient, demographic, clinical, and immunological data were collected in line with the nature of the registry, with follow-up up to 5 years.^{22,23}

Inclusion and exclusion criteria

Patients were eligible for inclusion in our analysis if they:

- were enrolled in the SPRING-SIR registry;
- were aged ≥ 18 years at the time of registry enrollment;
- fulfilled the 2013 American College of Rheumatology/European League Against Rheumatism (EULAR) classification criteria for systemic sclerosis;
- had available data on the year of onset for both RP and the first NRP sign or symptom;

otherwise excluded from our study.

Exposure

Patients were categorized into four groups both for age-RP and age-NRP, based on the quartiles of the

Francesca Ingegnoli
Clinica Reumatologica, Dipartimento di Reumatologia e Scienze Mediche, ASST Gaetano Pini-CTO, Dipartimento di Scienze Cliniche e di Comunità, Dipartimento di Eccellenza, Università degli Studi di Milano, Milan, Italy

Maria De Santis
Department of Biomedical Sciences, Humanitas University, Pieve Emanuele—Milan and Research Hospital, Milan, Italy

Fabio Cacciapaglia
Rheumatology Unit, Department of Precision and Regenerative Medicine—Ionian Area, University of Bari "Aldo Moro," Bari, Italy

Rheumatology Service, Internal Medicine Unit "F. Miulli" General Hospital, Acquaviva delle Fonti, Bari, Italy

Department of Medicine and Surgery, LUM "G. De Gennaro," Casamassima, Bari, Italy

Giuseppe Murdaca
Department of Internal Medicine, University of Genoa and Allergy, and Clinical Immunology Unit, Ospedale San Bartolomeo, Sarzana, Italy

Giuseppina Abignano
Gianna Angela Mennillo
Salvatore D'Angelo
Rheumatology Unit, Department of Health Science, San Carlo Hospital, University of Basilicata, Potenza, Italy

Giorgio Pettiti
Rheumatology Unit ASO S., Croce e Carle Hospital, Cuneo, Italy

Alessandra Della Rossa
Marco Di Battista
Department of Rheumatology, University of Pisa, Pisa, Italy

Maurizio Caminiti
Giuseppe Pagano Mariano
Departmental Rheumatology Unit, Grande Ospedale Metropolitano, Reggio Calabria, Italy

Annamaria Iuliano
Gian Domenico Sebastiani
Rheumatology Unit, San Camillo—Forlanini Hospital, Rome, Italy

Giovanni Ciano

Local Health
Department, Hospital
of Ariano Irpino, Ariano
Irpino, Italy

Lorenzo Beretta

Referral Center for
Systemic Autoimmune
Diseases, Fondazione
IRCCS Ca' Granda,
Ospedale Maggiore
Policlinico di Milano,
Milan, Italy

Gianluca Bagnato

Department of Clinical
and Experimental
Medicine, University of
Messina, Messina, Italy

Ennio Lubrano

Department of
Rheumatology,
University of Molise,
Campobasso, Italy

Ilenia De Andres

Rheumatology Unit,
Azienda Ospedaliera
di Rilievo Nazionale ed
Alta Specializzazione
Garibaldi, Catania, Italy

Luca Idolazzi

Rheumatology Unit,
Department of Medicine,
University of Verona,
Verona, Italy

Marta Saracco

Rheumatology Unit,
Mauriziano-Umberto I
Hospital, Turin, Italy

Cecilia Agnes

Division of
Rehabilitation,
Department of Medicine,
Torino, ASL T05,
Carmagnola, Italy

Corrado Campochiaro

Giacomo De Luca

Lorenzo Dagna

Marco Matucci-Cerinic
IRCCS San Raffaele
Scientific Institute,
Unit of Immunology,
Rheumatology, Allergy
and Rare Diseases, &
Inflammation, Fibrosis
and Ageing initiative,
Milan, Italy

Vita-Salute San Raffaele
University, Milan, Italy

Marco Fornaro

Florenzo Iannone

Rheumatology Unit,
Department of Precision
and Regenerative
Medicine—Ionian Area,
University of Bari "Aldo
Moro," Bari, Italy

Greta Pellegrino

Department of
Rheumatology, IRCCS
Ospedale Galeazzi—
Sant'Ambrogio, Milan,
Italy

Dipartimento di Scienze
Biomediche e Cliniche,
Università degli Studi di
Milano, Milan, Italy

distribution in the total population. Since age at RP onset and age at NRP onset follow different distributions within the cohort, the resulting quartile groups differed in terms of age range. This approach ensured that each group contained a similar number of patients, allowing for balanced comparisons across categories. Additionally, patients were divided into three groups based on RP onset patterns: (1) RP group: RP onset at least 1 year before NRP sign/symptom; (2) Simultaneous group: RP onset within the same calendar year of NRP sign/symptom (therefore corresponding to a maximum interval of 12); and (3) NRP group: RP onset after at least 1 year after NRP sign/symptom.

Outcome

The primary outcome of this study was to assess the association between the mode of onset defined as the temporal sequence between RP and NRP and the clinical endotype of SSc. As an exploratory outcome, the association between mode of onset and mortality was additionally tested.

Covariates

For all patients, demographic and clinical data were collected at the baseline visit. While the age of onset of RP was patient-reported, the first NRP sign or symptom was documented by the physician, based on both patient report and reviewed clinical documentation. ILD was defined as the presence of interstitial pattern changes on high resolution computed tomography (HRCT) such as reticulation, traction bronchiectasis, pleural irregularities, lung volume loss, and honeycombing. Pulmonary arterial hypertension (PAH) was defined as a mean pulmonary artery pressure (PAPm) >20 mmHg on right heart catheterization.²⁴ Cardiopulmonary manifestations encompassed ILD, PAH, pericardial effusion, arrhythmias, and coronary artery disease of unknown origin. Esophageal involvement was defined by the presence of gastroesophageal reflux, hypomotility on esophageal manometry, or abnormalities on esophageal/intestinal barium radiography.²⁵ Gastric symptoms included delayed gastric emptying, bloating, early satiety, nausea, and vomiting. Intestinal symptoms were defined as abdominal distension, chronic diarrhea, fecal incontinence, fecal soilage, constipation, or malabsorption. The EUSTAR disease activity index was calculated using the established composite score incorporating the modified Rodnan skin score

(mRSS), tendon friction rubs, elevated C-reactive protein, reduced diffusion capacity of the lung for carbon oxide, and active digital ulcers (DU).²⁶ The Charlson Comorbidity Index (CCI) was used to assess the burden of comorbidities through a weighted scoring system covering conditions such as cardiovascular disease, malignancies, and diabetes.²³

Follow-up time

In the survival analysis, patients were followed from the enrollment in the SPRING-SIR Registry. Follow-up ended at the earliest occurrence among the date of death, last available clinical assessment, study end date (March 29, 2021), or withdrawal from the registry for any reason.

Statistical analysis

The data were analyzed using descriptive statistics to calculate the absolute (*n*) and relative (%) frequencies of categorical variables. Continuous variables were described in terms of mean and standard deviation or median and interquartile range, as appropriate following their distribution. All group comparisons were performed using Chi-square test (with Fisher's correction where applicable), otherwise with ANOVA test. A significance level of 0.05 was used for all tests. Variables with missing data were analyzed using a complete case approach; no data imputation was performed. Sensitivity analyses and multiple imputation were considered; however, given the retrospective nature of the registry, the level and pattern of missingness, and the exploratory aim of the study, we opted for a complete-case analysis. All eligible patients from the SPRING-SIR registry were included in the analysis to ensure representativeness and generalizability of the results.

We used logistic, linear, and multinomial regression models to assess the association between selected predictors (age-RP, age-NRP, and RP-NRP onset pattern) and dichotomous, continuous, and categorical outcomes, respectively.

For all analyses involving the RP-NRP pattern variable, the RP group was used as the reference category, against which the simultaneous and NRP groups were compared. Similarly, when stratifying by age quartiles, the youngest quartile served as the reference.

Kaplan–Meier survival curves and multivariable Cox proportional hazards models were used to investigate the association of the aforementioned predictors with overall survival. Proportional hazards assumptions were tested and multicollinearity between covariates was excluded ($r < |0.7$).

All models were adjusted for potential confounders selected based on prior literature and expert clinical judgment. These included age, sex, cutaneous subset, presence of ILD, CCI.^{13,23}

Results

This study included a total of 1748/2134 SSc patients from the SPRING-SIR registry (Figure 1); of those, 1549 (86.6%) were females, with a mean age of 59 ± 14 years. Additional clinical characteristics are presented in Table 1.

Population characteristics according to the RP-NRP patterns of onset

The RP group consisted of 682 patients (39%) with a mean age-RP of 39 ± 15 years and age-NRP of 49 ± 14 years. The simultaneous group included 1026 (58.8%) patients, in whom age-RP was 49 ± 15 years and age-NRP was 48 ± 15 years. Lastly, the NRP group comprised 39 patients (2.2%), with a mean age-RP of 53 ± 12 years and age-NRP of 48 ± 12 years.

When comparing the three groups, we observed specific differences in the distribution of SSc features (Table 1).

Patients in the RP group were characterized by a higher prevalence of anti-centromere antibodies (ACA, 41.5% vs 25.2% in the simultaneous group and 26.5% in NRP; $p < 0.001$) and earlier age at RP onset ($p < 0.001$). In line with this autoantibody profile, patients in the RP group presented a lower prevalence of ILD (35.3% vs 41.7% in simultaneous group and 41.0% in NRP; $p = 0.03$), dyspnea (35.3% vs 41.7% in simultaneous group and 41.0% in NRP, $p = 0.018$), and sclerodactyly (65.6% vs 73.6% in simultaneous group and 73.7% in NRP, $p = 0.002$), while intestinal symptoms were slightly more common (21.2% vs 20.2% in simultaneous group and 5.4% in NRP, $p = 0.048$).

Patients with simultaneous RP-NRP onset had a higher prevalence of ATA (41.1% vs 26.4% in RP




Figure 1. Patients selection process.

and 21.6% in NRP, $p < 0.001$). Consistently, they presented more frequently with dcSSc (25.2% vs 13.2% in RP and 5.2% in NRP, $p < 0.001$), a higher median mRSS, and more prevalent clinical manifestation such as esophageal involvement (54.1% vs 47.5% in RP and 37.8% in NRP, $p = 0.007$) and ILD (41.7% vs 35.3% in RP and 41% in NRP, $p = 0.03$).

Conversely, NRP patients showed no relevant positivity of a specific autoantibody pattern. Clinically, NRP patients had lower prevalence of dcSSc and esophageal symptoms, but a higher prevalence of cardiopulmonary involvement, specifically of PAH. Consistently, we observed a higher systolic pulmonary artery pressure (sPAP) on echocardiography, lower forced vital capacity (FVC), and total lung capacity, although the latter two fell close to the normal range.

Population characteristics according to the age of RP and the age of NRP

When dividing our population according to the quartiles of the distribution of age-RP, the first group included 436 patients whose RP appeared before 34 years of age, the second group consisted of 422 patients with RP onset between 35 and 45 years, the third group had 439 patients with RP onset between 46 and 56 years, while the fourth group included patients in whom RP appeared after aging 56 years (450 patients).

The most striking differences were observed when comparing the first and the fourth groups. Patients with RP onset in younger age had a higher prevalence of ATA (42.2% vs 31.5, $p = 0.003$) and

Erika Pigatto
Federica Furini
Franco Cozzi
Department of Medicine, Villa Salus Hospital, Mestre, Italy

Licia Vultaggio
Marcello Govoni
Rheumatology Unit, Department of Medical Sciences, University of Ferrara and Azienda Ospedaliera—Universitaria S. Anna, Ferrara, Italy

Simone Parisi
Clara Lisa Peroni
Enrico Fusaro
Rheumatology Unit, Azienda Ospedaliera Universitaria Città Della Salute e della Scienza di Torino, Turin, Italy

Gerolamo Bianchi
Rheumatology Unit, Department of Medical Specialities, Local Health Trust 3, Genoa, Italy

Clodoveo Ferri
Department of Medical and Surgical Sciences for Children and Adults, University Hospital of Modena and Reggio Emilia School of Medicine, Modena, Italy
Rheumatology Clinic "Madonna dello Scoglio" Crotoni, Crotoni, Italy

*These authors contributed equally

Table 1. General characteristics of the whole population and then stratified according to RP-NRP onset groups.

Clinical characteristics	Total, n (%)	Missing, n (%)	NRP onset	Simultaneous onset	RP onset	p-Value
Patients, n (%)	1748 (100)	0	39 (2.2)	1026 (58.8)	682 (39)	
Smoking habit, n (%)	530 (33.7)	175 (10)	12 (34.3)	322 (34.5)	196 (32.4)	0.689
Male sex, n (%)	199 (11.4)	4 (0.2)	8 (20.5)	133 (13)	58 (8.5)	0.004
Age years mean, y, mean ± SD	59 ± 14	1 (0.06)	60 ± 11	58 ± 13	59 ± 14	0.527
Age at RP onset, y, mean ± SD	45 ± 16	1 (0.06)	53 ± 12	49 ± 15	39 ± 15	<0.001
Age at NRP onset, y, mean ± SD	48 ± 14	1 (0.06)	48 ± 12	48 ± 15	49 ± 14	0.892
CCI (mean ± SD)	3 ± 2	0	3 ± 2	3 ± 2	3 ± 2	0.276
RP-NRP time, y, mean ± SD	3.94 ± 7.9	0	-4.74 ± 3.44	0.22 ± 0.46	10 ± 9.8	<0.001
Skin and vascular						
dcSSc, n (%)	351 (20.1)	0	2 (5.1)	259 (25.2)	90 (13.2)	<0.001
mRSS, median (IQR)	4 (2–9)	149 (8.5)	3 (0–6)	5 (2–10)	4 (1–8)	<0.001
Puffy fingers, n (%)	897 (51.9)	20 (1.1)	15 (40.5)	516 (50.8)	366 (54.1)	0.158
Digital ulcers, n (%)	368 (21.3)	17 (1)	3 (7.9)	220 (21.7)	145 (21.4)	0.113
Digital pitting scars, n (%)	830 (48.1)	21 (1.2)	16 (43.2)	506 (49.9)	308 (45.6)	0.179
Sclerodactyly, n (%)	1218 (70.4)	19 (1.1)	28 (73.7)	746 (73.6)	444 (65.6)	0.002
Telangiectasias, n (%)	1052 (60.8)	18 (1)	25 (65.8)	580 (57.1)	447 (66.1)	<0.001
Capillaroscopy pattern late, n (%)	409 (26)	176 (10.1)	6 (15.8)	253 (27.7)	150 (24.2)	0.109
Gastrointestinal						
Esophageal symptoms, n (%)	885 (51.2)	19 (1.1)	14 (37.8)	550 (54.1)	321 (47.5)	0.007
Gastric symptoms, n (%)	340 (19.7)	21 (1.2)	5 (13.5)	212 (20.9)	123 (18.2)	0.258
Intestinal symptoms, n (%)	350 (20.3)	20 (1.1)	2 (5.4)	205 (20.2)	143 (21.2)	0.048
Cardiopulmonary						
Cardiopulmonary manifestation, n (%)	487 (28.2)	22 (1.3)	17 (45.9)	314 (31)	156 (23.1)	<0.001
Dyspnea, n (%)	676 (39.3)	26 (1.5)	18 (48.6)	421 (41.6)	237 (35.3)	0.018
ILD, n (%)	685 (39.2)	0	16 (41)	428 (41.7)	241 (35.3)	0.030
PAH, n (%)	28 (3.9)	0	3 (15.8)	10 (2.3)	15 (5.7)	0.003
Conduction block, n (%)	126 (12.2)	716 (41)	3 (16.7)	87 (14.2)	36 (9)	0.027
Pericardial effusion, n (%)	100 (7)	319 (18.2)	2 (6.7)	63 (7.4)	35 (6.4)	0.761
Abnormal diastolic function, n (%)	331 (23.5)	339 (19.4)	9 (31)	187 (22.3)	135 (25)	0.316
EF (%)	61 ± 6	383 (21.9)	60 ± 9	61 ± 5	61 ± 5	0.652
sPAP (mmHg)	22 ± 16	321 (18.4)	29 ± 19	22 ± 16	23 ± 16	0.012

(Continued)

Table 1. (Continued)

Clinical characteristics	Total, <i>n</i> (%)	Missing, <i>n</i> (%)	NRP onset	Simultaneous onset	RP onset	<i>p</i> -Value
FVC (%)	101 ± 23	440(25.2)	97 ± 24	100 ± 22	104 ± 23	0.004
DLCO (%)	69 ± 21	503 (28.8)	65 ± 24	68 ± 20	69 ± 20	0.322
TLC (%)	97 ± 20	990 (56.7)	89 ± 16	96 ± 21	99 ± 20	0.025
Musculoskeletal						
Calcinosis, <i>n</i> (%)	201 (11.6)	21 (1.2)	5 (13.5)	120 (11.8)	76 (11.3)	0.883
Arthritis, <i>n</i> (%)	199 (11.6)	30 (1.7)	3 (8.1)	123 (12.2)	73 (10.9)	0.662
Other symptoms						
Renal crisis, <i>n</i> (%)	21 (1.2)	20 (1.1)	1 (2.7)	14 (1.4)	6 (0.9)	0.293
Autoantibodies						
ATA, <i>n</i> (%)	597 (34.9)	37 (2.1)	8 (21.6)	411 (41.1)	178 (26.4)	<0.001
ARA, <i>n</i> (%)	27 (2)	374 (21.4)	0	24 (2.9)	3 (0.6)	0.007
ACA, <i>n</i> (%)	508 (31.6)	141 (8.1)	9 (26.5)	239 (25.2)	260 (41.5)	<0.001

Bold indicates statistically significant result.
 ACA, anti-centromere antibodies; ARA, anti-RNA polymerase III antibodies; ATA, anti-topoisomerase antibodies; BMI, body mass index; CCI, Charlson comorbidity index; dcSSc, diffuse cutaneous systemic sclerosis; DLCO, diffusing capacity of the lung for carbon monoxide; EF, ejection fraction; FVC, forced vital capacity; ILD, interstitial lung disease; IQR, interquartile range; mRSS, modified Rodnan skin score; *n*, numbers; NRP, non-Raynaud's phenomenon; PAH, pulmonary arterial hypertension; RP, Raynaud's phenomenon; SD, standard deviation; sPAP, systolic pulmonary artery pressure; TLC, total lung capacity; *y*, years.

consistently showed a greater prevalence of dcSSc (23.2% vs 14.9%, $p=0.01$), DU (29.2% vs 14.6%, $p<0.001$), digital pitting scars (58.2% vs 35.7%, $p<0.001$), and joint contractures (17.1% vs 9.7%, $p=0.014$). Conversely, the group with RP onset after 56 years of age was characterized by a higher prevalence of cardiopulmonary manifestations (35.3% vs 21.1%, $p<0.001$) and a numerical increase in sPAP values (25 ± 18 vs 21 ± 14 mmHg, $p<0.001$). Other features are shown in Table S1.

When clustering our population according to the distribution of the age-NRP, the first group consisted of 311 (17.8%) patients whose NRP symptoms appeared before the age of 38. The second group included 663 (37.9%) patients with NRP onset between 39 and 49 years. The third group comprised 407 (23.3%) patients with NRP onset between 50 and 59 years, while the fourth group, with a total of 367 (21%) patients, included those with NRP sign on symptom onset after the age of 59.

Similarly, when comparing the groups, the most relevant differences were observed between the

first and fourth groups. Again, we observed a higher prevalence of ATA (50.7% vs 29.9%, $p<0.001$), dcSSc (30.2% vs 13.1%, $p<0.001$), and DU (34.4% vs 14%, $p<0.001$) in the group with NRP onset at younger age. In contrast, the group with age-NRP over 59 years showed higher prevalence of ACA (35.7% vs 22.8%, $p=0.004$) and more prominent cardiopulmonary manifestations (36.5% vs 22.4%, $p<0.001$). Other features are shown in Table S2.

Associating age-RP, age-NRP, and RP-NRP timing with SSc presentation

Logistic regression analysis revealed that the group with simultaneous RP-NRP onset was associated with a higher risk of presenting the dcSSc subset, compared to the RP group (odds ratio (OR), 1.491, 95% confidence interval (CI): 1.032–2.154, $p=0.033$), independently of ATA positivity and sex (Table 2). Additionally, we identified age at RP onset as a predictor of sPAP values ($\beta -0.149$, 95% CI: -0.297 to -0.001 ; $p=0.049$, $N=1426$) and mRSS ($\beta -0.53$, 95% CI: 0.094–0.011; $p=0.012$, $N=1598$; Table S3). No statistically significant association was

Table 2. Logistic multivariable regression models predicting clinical features of SSc.

Outcome	N patients included	Predictors	OR	95% CI lower bound	95% CI upper bound	p-Value
ILD	1705/1748 (97.5%)	RP onset	Reference			0.892
		NRP onset	1.033	0.486	2.197	0.933
		Simultaneous onset	0.944	0.719	1.240	0.678
		Age_RP	1.014	0.996	1.033	0.119
		Constant	0.093			<0.001
dcSSc	1705/1748 (97.5%)	RP onset	Reference			0.006
		NRP onset	0.263	0.057	1.206	0.086
		Simultaneous onset	1.491	1.032	2.154	0.033
		Age_RP	1.030	0.997	1.063	0.072
		Constant	0.373			0.002
DU	1557/1748 (89.1%)	RP onset	Reference			0.218
		NRP onset	0.352	0.099	1.250	0.106
		Simultaneous onset	1.041	0.740	1.465	0.816
		Age_RP	0.995	0.974	1.018	0.686
		Constant	0.243			<0.001
DPS	1553/1748 (88.8%)	RP onset	Reference			0.748
		NRP onset	0.907	0.419	1.961	0.804
		Simultaneous onset	1.091	0.820	1.452	0.550
		Age_RP	0.992	0.974	1.009	0.345
		Constant	0.730			0.266
Calcinosis	1701/1748 (97.3%)	RP onset	Reference			0.948
		NRP onset	1.197	0.406	3.534	0.744
		Simultaneous onset	1.019	0.677	1.534	0.927
		Age_RP	1.003	0.976	1.030	0.846
		Constant	0.098			<0.001
Esophageal symptoms	1431/1748 (81.9%)	RP onset	Reference			0.133
		NRP onset	0.653	0.278	1.535	0.329
		Simultaneous onset	1.215	0.920	1.605	0.170
		Age_RP	1.030	0.997	1.063	0.072
		Constant	0.630			0.238

(Continued)

Table 2. (Continued)

Outcome	N patients included	Predictors	OR	95% CI lower bound	95% CI upper bound	p-Value
Gastric symptoms		RP onset	Reference			0.532
	1415/1748 (80.9%)	NRP onset	0.484	0.134	1.742	0.267
		Simultaneous onset	0.921	0.644	1.318	0.654
		Age_RP	1.017	0.992	1.042	0.188
		Constant	0.082			<0.001
Intestinal symptoms		RP onset	Reference			0.171
	1416/1748 (81.0%)	NRP onset	0.186	0.024	1.431	0.106
		Simultaneous onset	1.132	0.802	1.599	0.480
		Age_RP	0.982	0.963	1.002	0.082
		Constant	0.173			<0.001
Diastolic dysfunction		RP onset	Reference			0.450
	1165/1748 (66.6%)	NRP onset	1.232	0.450	3.368	0.685
		Simultaneous onset	0.827	0.570	1.200	0.316
		Age_RP	0.997	0.976	1.019	0.805
		Constant	0.008			<0.001

Mode of onset and age at RP onset represented the predictors of interest. All analyses were adjusted for covariates sex autoantibodies and disease duration.
 Bold indicates statistically significant result.
 CI, confidence interval; dcSSc, diffuse cutaneous systemic sclerosis; DPS, digital pitting scars; DU, digital ulcers; ILD, interstitial lung disease; NRP, non-Raynaud's phenomenon; OR, odds ratio; RP, Raynaud's phenomenon; SSc, systemic sclerosis.

found when predicting other SSc-related manifestations.

Predicting the pattern of SSc onset

A multinomial regression analysis was performed to test which characteristics were associated with the risk of belonging to a specific group of RP-NRP onset (Table 3). In this analysis including 1710/1748 (97.8%) patients, we observed that female sex was associated with a higher probability of being included in the RP group (OR, 2.445; 95% CI: 1.052–5.681; $p=0.038$). Older age at RP onset increased the likelihood of being included in the NRP group (OR, 1.067; 95% CI: 1.042–1.092; $p<0.001$) or in the simultaneous group (OR, 1.048;

95% CI: 1.041–1.056; $p<0.001$). Additionally, ACA positivity was associated with a higher probability of being in the RP group (OR, 2.004; 95% CI: 1.550–2.597; $p<0.001$), while the presence of ATA and anti-RNA polymerase III (ARA) antibodies was linked to a higher likelihood of being included in the Simultaneous group (OR, 1.671; 95% CI: 1.295–2.157; $p<0.001$ and OR, 4.002; 95% CI: 1.151–13.914; $p=0.029$, respectively).

Survival analysis

Survival status was available for 943 patients (53.95%), with median follow-up duration of 24 (interquartile range: 12–48) months and 44 (4.7%) deaths recorded.

Table 3. Multinomial logistic regression predicting NRP and Simultaneous onset patterns compared to RP onset in patients with systemic sclerosis.

Group	Predictor	OR	95% CI lower bound	95% CI upper bound	p-value
RP ONSET	Reference				
NRP ONSET	Intercept				<0.001
	Age_RP	1.067	1.042	1.092	<0.001
	Female sex	0.409	0.176	0.950	0.038
	ACA	0.329	0.147	0.738	0.007
	ATA	0.511	0.212	1.230	0.134
	ARA	2.060E-8	2.060E-8	2.060E-8	NA
SIMULTANEOUS ONSET	Intercept				<0.001
	Age_RP	1.048	1.041	1.056	<0.001
	Female sex	0.726	0.513	1.027	0.071
	ACA	0.499	0.385	0.645	<0.001
	ATA	1.671	1.295	2.157	<0.001
	ARA	4.002	1.151	13.914	0.029

The analyses included age at RP onset, sex, autoantibodies as predictors. Bold indicates statistically significant result. ACA, anti-centromere antibodies; ARA, anti-RNA polymerase III antibodies; ATA, anti-topoisomerase I antibodies; CI, confidence interval; NA: not applicable; NRP, non-Raynaud's phenomenon; OR, odds ratio; RP, Raynaud's phenomenon.

The estimated survival rates were comparable among the RP (97%), NRP (95%), and simultaneous (94%) groups ($p=0.09$), as presented in Figure 2.

However, when testing the risk carried by the RP-NRP onset patterns in an exploratory, multi-variable Cox regression model, including 936/943 (99.3%) of the patients with follow-up data, we observed that the simultaneous group had a significant higher risk of mortality compared to RP group (hazard ratio (HR), 1.975; 95% CI: 1.002–3.893, $p=0.049$), which was independent from age at enrollment in the registry, sex, EUSTAR disease activity index, and CCI (Figure 3).

As expected, Kaplan–Meier survival analysis revealed lower survival rates for patients with older age at time of RP and NRP onset, respectively, 94% and 95.2% and significantly lower than younger ages on onset (data not shown).

Discussion

Our study provides a comprehensive analysis of SSc patients enrolled in the Italian SPRING-SIR registry, emphasizing the influence of the pattern of RP-NRP timing, the age of RP onset, and the age of the first NRP sign or symptom on the clinical presentation of SSc, as well as preliminary results on a possible impact on outcomes.

The mode of clinical onset in SSc remains a critical area of investigation, particularly regarding the differences between RP onset and NRP onset. Although RP is the expected initial manifestation in most SSc patients, studies have highlighted the distinct and aggressive nature of NRP onset.^{6,27} These studies confirm and expand on previous observations from the SSc Study Group of the Italian Society of Rheumatology, which reported a significant correlation between the mode of disease onset (RP or NRP) and adverse SSc features, including the dcSSc subset, ATA seropositivity,