

ORIGINAL RESEARCH

Inflammatory rheumatic diseases with onset after SARS-CoV-2 infection or COVID-19 vaccination: a report of 267 cases from the COVID-19 and ASD group

Francesco Ursini ^{1,2}, Piero Ruscitti ³, Olga Addimanda,⁴ Rosario Foti,⁵ Vincenzo Raimondo,⁶ Giuseppe Murdaca,⁷ Virginia Caira,⁸ Erika Pigatto,⁹ Giovanna Cuomo,¹⁰ Alberto Lo Gullo ⁵, Ilaria Cavazzana,¹¹ Corrado Campochiaro ¹², Caterina Naclerio,¹³ Rossella De Angelis,¹⁴ Jacopo Ciaffi,¹ Luana Mancarella,¹ Veronica Brusi,¹ Elena Marchetti,² Francesca Motta ^{15,16}, Marcella Visentini,¹⁷ Sebastiano Lorusso,^{18,19} Maria De Santis ^{15,16}, Giacomo De Luca,¹² Laura Massaro,²⁰ Domenico Olivo,²¹ Roberta Pellegrini,²² Francesca Francioso,²³ Jessica Luppino,⁶ Ilenia Di Cola,³ Roberta Foti,⁵ Giuseppe Varcasia,⁸ Francesco Caso,²⁴ Massimo Reta,⁴ Lorenzo Dagna ¹², Carlo Selmi ¹⁵, Annamaria Iagnocco,²⁵ Roberto Giacomelli,^{18,19} Florenzo Iannone,²⁶ Clodoveo Ferri ^{6,27}

To cite: Ursini F, Ruscitti P, Addimanda O, *et al.* Inflammatory rheumatic diseases with onset after SARS-CoV-2 infection or COVID-19 vaccination: a report of 267 cases from the COVID-19 and ASD group. *RMD Open* 2023;**9**:e003022. doi:10.1136/rmdopen-2023-003022

Received 23 January 2023
Accepted 22 May 2023



© Author(s) (or their employer(s)) 2023. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Francesco Ursini;
francesco.ursini2@unibo.it

ABSTRACT

Objectives To better define the spectrum of new-onset post-COVID-19 and post-COVID-19 vaccine inflammatory rheumatic diseases (IRD) from a large multicentric observational study.

Methods Consecutive cases of IRD encountered during a 12-month period and satisfying one of the following inclusion criteria: (a) onset of the rheumatic manifestations within 4 weeks from SARS-CoV-2 infection or (b) onset of the rheumatic manifestations within 4 weeks from the administration of one of the COVID-19 vaccines were recruited.

Results The final analysis cohort comprised 267 patients, of which 122 (45.2%) in the post-COVID-19 and 145 (54.8%) in the postvaccine cohort. Distribution of IRD categories differed between the two cohorts: the post-COVID-19 cohort had a higher percentage of patients classified as having inflammatory joint diseases (IJD, 52.5% vs 37.2%, $p=0.013$) while the post-vaccine cohort had a higher prevalence of patients classified as polymyalgia rheumatica (PMR, 33.1% vs 21.3%, $p=0.032$). No differences were detected in the percentage of patients diagnosed with connective tissue diseases (CTD 19.7% vs 20.7%, $p=0.837$) or vasculitis (6.6% vs 9.0%, $p=0.467$). Despite the short follow-up period, IJD and PMR patients' response to first-line therapy was favourable, with both groups achieving a drop in baseline disease activity scores of ~30% and ~70% respectively.

Conclusion Our article reports the largest cohort published to date of new-onset IRD following SARS-CoV-2 infection or COVID-19 vaccines. Although causality cannot be ascertained, the spectrum of possible clinical manifestations is broad and includes IJD, PMR, CTD and vasculitis.

WHAT IS ALREADY KNOWN ON THIS TOPIC

⇒ Since the beginning of the COVID-19 pandemic, isolated cases of inflammatory rheumatic diseases (IRD) with onset after SARS-CoV-2 infection or COVID-19 vaccine administration have been described by several groups. However, the anecdotal nature of the previous reports does not allow to catch the 'big picture' of the spectrum of IRD potentially associated with those exposures.

WHAT THIS STUDY ADDS

⇒ In this study, we collected a large cohort of IRD developed in close temporal association with SARS-CoV-2 infection or vaccine administration. According to our data, the spectrum of possible presentation is broad and includes inflammatory joint diseases (IJD), polymyalgia rheumatica (PMR), connective tissue diseases (CTD) and vasculitis with IJD prevailing in post-COVID-19 and PMR prevailing in postvaccine patients.

HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE OR POLICY

⇒ Our study provides a broad, descriptive analysis of the possible array of IRD associated with SARS-CoV-2 infection and COVID-19 vaccines. This may contribute to strengthen the awareness on similar events—although uncommon—among physicians involved in musculoskeletal care and stimulate further research on this topic.

INTRODUCTION

Following the rapid spread of the COVID-19 pandemic, the scientific community demonstrated an unprecedented resilience to tackle an unknown biological threat, obtaining rapid progress in understanding the pathogenesis of the disease¹ and the subsequent development of highly specific ‘countermeasures’ such as the novel mRNA vaccines² and antiviral medications.³ This allowed to scale down the burden of the disease and save tens of millions of lives globally.⁴ Indeed, most European countries achieved large vaccine coverage and Italy, in particular, attained one of the most successful campaigns with more than 50 million of people (93% of the population >12 years) getting at least one dose.⁵

Since the beginning, it was evident that the profound interaction between SARS-CoV-2 and the host immune system¹ drives the most severe acute manifestations of the disease (eg, acute respiratory distress syndrome and microvascular injury); on the other hand, in the subsequent phases, a long-term immune dysregulation was demonstrated in at least a proportion of patients⁶ that may bolster prolonged symptoms persisting for months or years after viral clearance.

Postacute COVID-19 syndrome (PACS)⁷ has now been defined as an extremely heterogeneous condition following COVID-19 and characterised by a multitude of signs and symptoms including complaints of rheumatological interest.⁸ Musculoskeletal pain, indeed, is reported in almost 10% of individuals infected by SARS-CoV-2 at some time during the first year after the infection⁸ and, as suggested in a previous study published by our group, most of these patients satisfy the classification criteria for fibromyalgia.⁹

Concurrently, a growing number of articles described new-onset serological evidence of autoimmunity^{10–11} or frank inflammatory rheumatic diseases (IRD)—including inflammatory joint diseases (IJD),¹² polymyalgia rheumatica (PMR),¹³ connective tissue diseases (CTD)^{14–15} and vasculitis¹⁶—in close temporal association with SARS-CoV-2 infection. However, this was not surprising as the potential role of infections in triggering rheumatic diseases has been hypothesised since decades on the basis of preclinical studies¹⁷ and is clearly recapitulated by reactive arthritis¹⁸ or virally induced arthritis (eg, parvovirus B19 or hepatitis C virus (HCV)).¹⁹ On the other hand, it was previously speculated that also the exposure to vaccines may elicit autoimmunity.²⁰ The controversial autoimmune/inflammatory syndrome induced by adjuvants (ASIA syndrome),²¹ in example, is believed to result from the exposure to common vaccine adjuvants. Also in this case, reports of IRD following vaccination with available anti-COVID-19 vaccines appeared in literature since the earliest phases of the vaccine campaign.^{22–25}

On this background, aim of the present study was to contribute to the knowledge on the spectrum of new-onset post-COVID-19 and post-COVID-19 vaccine IRD with a descriptive and comparative analysis from a large multicentric observational study.

METHODS

Study design and inclusion/exclusion criteria

The present study was conceived as an observational cohort study. To this purpose, we published a web-based data collection form and invited all members of the Italian *COVID-19 and autoimmune systemic diseases* (COVID-19 and ASD) collaborative research group²⁶ to submit consecutive cases of IRD or acrosyndromes encountered during routine clinical practice from 1 November 2021 to 31 October 2022 (12 months) and satisfying one of the following inclusion criteria: (a) onset of the rheumatic manifestations within 4 weeks from SARS-CoV-2 infection, demonstrated by RT-PCR and/or antigenic nasopharyngeal swab or (b) onset of the rheumatic manifestations within 4 weeks from the administration of one of the COVID-19 vaccines approved for administration in Italy (BNT162b2—Pfizer/BioNTech), mRNA-1273 (Moderna) during the collection period. Exclusion criteria were predefined as a past history of IRD or acrosyndromes or a diagnosis of PACS or ‘long-COVID’.

The COVID-19 and ASD group is a network of more than 40 physicians with senior experience in the management of patients with IRD, belonging to 11 public academic medical centres, 7 public general hospitals, 3 private academic medical centres and 2 private general hospitals, covering regions from northern to southern Italy and including rheumatologists, clinical immunologists and specialists in internal medicine.

Data were collected anonymously through an online form built using the Google Forms platform. Google Forms is a free and easy to use questionnaire administration tool that has been largely used in medical research.²⁷ Cases submitted by individuals centres were assigned an alphanumeric identifier code, including a centre-specific string and a progressive patient number; before the analysis, records were manually checked for recognising potential duplicate entries (eg, identical demographic data and/or other variables despite different identifier code).

Information gathered included demographic characteristics, past exposure to SARS-CoV-2 infection, vaccine status, clinical features of rheumatological interest, auto-antibodies status, erythrocyte sedimentation rate (ESR), C reactive protein (CRP), final diagnosis of the rheumatologist, first-line treatment.

For patients classified as having arthritis, disease activity before and after at least 4 weeks of treatment was recorded using the disease activity score including 28 joints (DAS28).²⁸ For patients classified as having PMR, disease activity before and after at least 4 weeks of treatment was recorded using the PMR activity score (PMR-AS) as previously published.²⁹

The research was conducted in compliance with the Declaration of Helsinki and its latest amendments³⁰ and approved by a central Ethics Committee (Comitato Etico Azienda Sanitaria Locale 1 Avezzano/Sulmona/L’Aquila, L’Aquila, Italy; approval number: 0204194/22). Written informed consent was obtained from all study subjects.



Statistical analysis

Data are expressed as mean±SD, median (25th–75th percentile) or number (percentage) as appropriate. Student's *t* test was used to compare differences in normally distributed continuous variables between the two groups; highly skewed variables were *ln*-transformed before the analysis. Fisher's exact test was used to compare categorical variables. A *p* value <0.05 was considered statistically significant. All analyses were performed using the SPSS software V.26.0 (IBM, Armonk, New York).

RESULTS

Clinical features of the overall study population

A total of 270 cases were entered in the database. Three records were excluded because identified as mechanical pain related to osteoarthritis (*n*=1) or fibromyalgia (*n*=2). No duplicate records were identified. The final analysis cohort comprised a total of 267 patients, of which 122 (45.2%) patients in the post-COVID-19 (age 54±17 years, 69.7% female subjects) and 145 (54.8%) patients (age 58±16 years, 66.2% female subjects) in the postvaccine cohort. Number of vaccine doses received did not differ between post-COVID-19 and postvaccine patients (2±1 vs 3±1, *p*=0.123).

In the post-COVID-19 cohort, four (3.3%) patients experienced asymptomatic infection, 104 (85.2%) had mildly symptomatic disease, while 14 (11.5%) had severe disease according to the WHO severity classification.³¹ Accordingly, 113 patients (92.6%) were treated at home and only 9 (7.4%) were hospitalised; of these, only one required subsequent admission to intensive care unit.

In the postvaccine cohort, most cases occurred after the booster (third) dose (*n*=66, 45.8%); the remaining were associated with a second dose (*n*=51, 35.4%), first dose (*n*=23, 16%) and only three (2.1%) after a fourth dose. In most cases, the specific vaccine associated with IRD onset was the Pfizer/BioNTech BNT162b2 (*n*=114, 78.6%) followed by Moderna mRNA-1273 (*n*=24, 16.6%),

mirroring the overall distribution of COVID-19 vaccines dispensed in Italy.⁵ In seven (4.8%) patients, the specific vaccine was unknown. A total of 15 (10.3%) patients reported a history of asymptomatic or mildly symptomatic COVID-19 before the vaccine dose associated with IRD onset.

Mean delay between COVID-19 diagnosis or vaccine administration and rheumatic manifestations development was 14.5±7.8 vs 13.9±8.5 days, respectively (*p*=0.59).

Distribution of various IRD categories differed between the two cohorts (figure 1): the post-COVID-19 cohort had a higher percentage of patients classified as having IJD (52.5% vs 37.2%, *p*=0.013), while the postvaccine cohort had a higher prevalence of patients classified as PMR (33.1% vs 21.3%, *p*=0.032). No differences were detected in the percentage of patients diagnosed with CTD (post-COVID-19: 19.7% vs postvaccine: 20.7%, *p*=0.837) or vasculitis (post-COVID-19: 6.6% vs postvaccine: 9.0%, *p*=0.467).

Inflammatory joint diseases

The most represented IRD category in both groups was IJD. The range of specific IJD was broad (table 1), with peripheral spondyloarthritis (pSpA), representing the most frequent final diagnosis. Although IJDs were relatively more frequent in post-COVID-19 patients, a significantly higher proportion of patients in the post-vaccine group was diagnosed as having PsA (29.6% vs 12.5%, *p*=0.021), while no significant differences were observed in other domains, including baseline inflammatory markers, disease activity, follow-up duration and first-line treatment used. The two patients in the post-COVID-19 vaccine cohort classified as having adult-onset Still's disease (AOSD) were included as part of the IJD subgroup because of a predominantly articular clinical picture. Notably, despite the short follow-up, patients' response to first-line therapy was favourable, with both




Figure 1 Relative frequencies of inflammatory rheumatic diseases categories in the two study cohorts. CTD, connective tissue diseases; IJD, inflammatory joint diseases; PMR, polymyalgia rheumatica; Vasc, vasculitis.

Table 1 Comparative analysis of clinical features in inflammatory joint diseases with onset after SARS-CoV-2 infection or COVID-19 vaccine administration

	Post-COVID-19 (n=64)	Post-vaccine (n=54)	P value
Age, years	49±14	51±13	0.277
Female sex, n (%)	47 (73.4)	37 (68.5)	0.557
N° vaccine doses, n	2±1	3±1	0.434
ESR (baseline), mm/h	32±22	36±25	0.364
ESR (follow-up), mm/h	19±12	20±14	0.689
CRP (baseline), mg/dL	1.29 (1.00–2.50)	1.50 (0.56–2.50)	0.639
CRP (follow-up), mg/dL	0.53 (0.40–0.90)	0.50 (0.30–0.80)	0.456
TJC (baseline), n	6±4	8±6	0.081
TJC (follow-up), n	3±3	3±4	0.895
SJC (baseline), n	2 (2–4)	3 (2–4)	0.754
SJC (follow-up), n	0	0	0.575
DAS28-CRP (baseline), score	4.3±1.0	4.4±1.1	0.501
DAS28-CRP (follow-up), score	3.0±1.0	3.1±1.1	0.887
Average follow-up, weeks	8 (6–12)	8 (7–12)	0.866
Specific diseases			
pSpA (other than PsA), n (%)	32 (50.0)	19 (35.2)	0.106
HLA-B27 present, n (%)	8 (25)	6 (31.6)	0.748
Enthesitis, n (%)	4 (12.5)	1 (5.3)	0.401
Dactylitis, n (%)	3 (9.4)	1 (5.3)	0.597
RA, n (%)	19 (29.7)	14 (25.9)	0.650
RF positive, n (%)	11 (57.9)	10 (71.4)	0.486
ACPA positive, n (%)	10 (52.6)	7 (50.0)	1.000
PsA, n (%)	8 (12.5)	16 (29.6)	0.021
Enthesitis, n (%)	3 (37.5)	8 (50.0)	0.562
Dactylitis, n (%)	1 (12.5)	4 (25.0)	0.477
AxSpA, n (%)	5 (7.8)	3 (5.6)	0.627
HLA-B27 present, n (%)	3 (60.0)	2 (66.7)	1.000
Enthesitis, n (%)	1 (20.0)	0 (0.0)	0.408
Dactylitis, n (%)	0 (0.0)	0 (0.0)	1.000
AOSD, n (%)	0 (0.0)	2 (3.7)	0.120
Treatment			
Paracetamol, n (%)	8 (12.5)	13 (24.1)	0.101
NSAIDs, n (%)	30 (46.9)	22 (40.7)	0.504
GCs, n (%)	33 (51.6)	33 (61.1)	0.298
Intra-articular GCs, n (%)	1 (1.6)	3 (5.6)	0.232
Colchicine, n (%)	2 (3.1)	0 (0.0)	0.190
MTX, n (%)	22 (34.4)	22 (40.7)	0.476
SSZ, n (%)	9 (14.1)	3 (5.6)	0.128
HCQ, n (%)	4 (6.3)	5 (9.3)	0.540
TNFi, n (%)	3 (4.7)	0 (0.0)	0.107
ANR, n (%)	0 (0.0)	1 (1.9)	0.274

ANR, anakinra; AOSD, adult-onset Still's disease; AxSpA, axial spondyloarthritis; CRP, C reactive protein; DAS28, disease activity score including 28 joints; ESR, erythrocyte sedimentation rate; GCs, glucocorticoids; HCQ, hydroxychloroquine; MTX, methotrexate; NSAIDs, non-steroidal anti-inflammatory drugs; PsA, psoriatic arthritis; pSpA, peripheral spondyloarthritis; RA, rheumatoid arthritis; SJC, swollen joint count; TJC, tender joint count; TNFi, TNF inhibitors.

Table 2 Comparative analysis of clinical features in polymyalgia rheumatica with onset after SARS-CoV-2 infection or COVID-19 vaccine administration

	Post-COVID-19 (n=28)	Post-vaccine (n=46)	P value
Age, years	58±20	62±17	0.632
Female sex, n (%)	5 (62.5)	9 (69.2)	0.751
N° vaccine doses, n	2±1	2±1	0.739
ESR (baseline), mm/h	50±22	59±32	0.150
ESR (follow-up), mm/h	22±18	22±15	0.954
CRP (baseline), mg/dL	2.15 (1.46–4.10)	1.99 (1.62–3.09)	0.630
CRP (follow-up), mg/dL	0.60 (0.50–1.00)	0.50 (0.50–0.95)	0.910
PMR-AS (baseline), score	28.3 (25.2–33.0)	24.8 (23.7–27.5)	0.392
PMR-AS (follow-up), score	8.44 (5.4–10.0)	6.43 (4.6–7.0)	0.268
Average follow-up, weeks	8±5	14±13	0.004
Treatment			
Paracetamol, n (%)	1 (3.6)	4 (8.7)	0.666
NSAIDs, n (%)	2 (7.1)	7 (15.2)	0.303
GCs, n (%)	28 (100.0)	43 (93.5)	0.168
MTX, n (%)	2 (7.1)	9 (19.6)	0.145

CRP, C reactive protein; ESR, erythrocyte sedimentation rate; GCs, glucocorticoids; MTX, methotrexate; NSAIDs, non-steroidal anti-inflammatory drugs; PMR-AS, polymyalgia rheumatica—activity score.

groups achieving a 30% drop from baseline in disease activity, as measured by DAS28-CRP.

Polymyalgia rheumatica

PMR was the second most common IRD category (table 2). None of the patients had clinical features reminiscent of concurrent giant cell arteritis (GCA) or required second-level imaging for suspected GCA on the basis of clinical judgement; those patients with prominent symptoms and/or supporting imaging suggestive of GCA with or without associated PMR are reported in the vasculitis subgroup.

No significant differences were observed between post-COVID-19 and postvaccine PMR patients, except for a shorter duration of follow-up in the post-COVID-19 patients (8±5 vs 14±13 weeks, p=0.004). Despite this, the patients responded well to first-line therapy, with reductions in PMR-AS of 70% and 74% from baseline, respectively. Of note, six patients in the postvaccine and one in the post-COVID-19 cohort were aged below 50 years (range: 33–48). However, all patients had increased CRP (range: 1.15–5.4 mg/dL) and/or ESR levels (range: 20–110 mm/hour), a clear PMR-like clinical presentation, and steroid responsiveness (overall PMR-AS: baseline 23.5±2.3 vs follow-up 7.7±2.8) and thus were attributed to the PMR subgroup by the treating rheumatologist.

Connective tissue diseases

CTDs were also frequently reported. The range of specific CTD presentation was broad (table 3) and included undifferentiated CTD, systemic sclerosis, idiopathic inflammatory myopathies, amyopathic dermatomyositis, systemic

lupus erythematosus, mixed CTD, Sjogren's syndrome and isolated new-onset acrosyndromes. Notably, when comparing the two cohorts, the only variable reaching statistical significance was the prevalence of acrosyndromes, which was greater in post-COVID-19 group (20.8 vs 3.3%, p=0.02). No information on the follow-up of CTD patients was collected due to the high heterogeneity between outcome measures in such a mixed category of patients.

Vasculitis

Vasculitis was the least common category in both groups. However, the range of specific presentation was broad (table 4) and included small-vessel vasculitis, large-vessel vasculitis, variable vessel vasculitis and isolated skin vasculitis. No significant differences were found when comparing post-COVID-19 and postvaccine patients. No information on the follow-up of vasculitis patients was collected due to the high heterogeneity between outcome measures in such a mixed category of patients.

DISCUSSION

Since the earliest phases of the pandemic, several isolated case reports or small case series reporting the suspicious onset of rheumatic diseases in association with SARS-CoV-2 infection or COVID-19 vaccine administration have been published. Even though the anecdotal nature of those reports did not shed sufficient light on this phenomenon, they represent a stimulating hypothesis-generating basis for encouraging additional research on this topic.

Table 3 Comparative analysis of clinical features in connective tissue diseases with onset after SARS-CoV-2 infection or COVID-19 vaccine administration

	Post-COVID-19 (n=24)	Post-vaccine (n=30)	P value
Age, years	47±18	55±19	0.134
Female sex, n (%)	19 (79.2)	22 (73.3)	0.618
N° vaccine doses, n (%)	2±1	3±1	0.263
ANA positive, n (%)	21 (87.5)	28 (93.3)	0.462
>1:160, n (%)	12 (57.1)	16 (57.1)	1.000
>1:640, n (%)	3 (14.3)	5 (17.8)	1.000
ESR, mm/h	19±16	28±22	0.728
CRP, mg/dL	0.75 (0.30–1.50)	0.56 (0.20–2.50)	0.551
Specific diseases			
UCTD, n (%)	6 (25.0)	11 (36.7)	0.359
Anti-SSA positive, n (%)	0 (0.0)	3 (27.3)	0.159
Anti-Scl-70 positive, n (%)	2 (33.3)	2 (18.2)	0.482
SSc, n (%)	6 (25.0)	6 (20.0)	0.661
Anti-Scl-70 positive, n (%)	2 (33.3)	2 (33.3)	1.000
ACA positive, n (%)	3 (50.0)	4 (66.7)	0.558
IIM, n (%)	4 (16.7)	6 (20.0)	0.754
Anti-Mi2 positive, n (%)	1 (25.0)	2 (33.3)	0.778
Anti-Pl7 positive, n (%)	1 (25.0)	0 (0.0)	0.197
Anti-TIF1-γ, n (%)	0 (0.0)	2 (33.3)	0.197
ADM, n (%)	0 (0.0)	2 (6.7)	0.197
Anti-Mi2 positive, n (%)	0 (0.0)	1 (50.0)	N/A*
Anti-MDA5 positive, n (%)	0 (0.0)	1 (50.0)	N/A*
SLE, n (%)	2 (8.3)	1 (3.3)	0.425
Anti-ds-DNA positive, n (%)	1 (50.0)	1 (100.0)	0.386
Anti-Sm positive, n (%)	1 (50.0)	0 (0.0)	0.386
MCTD, n (%)	0 (0.0)	3 (10.0)	0.111
Anti-RNP positive, n (%)	0 (0.0)	3 (100.0)	N/A*
SjS, n (%)	1 (4.2)	0 (0.0)	0.259
SSA positive, n (%)	1 (100.0)	0 (0.0)	N/A*
Isolated acrosyndrom, n (%)	5 (20.8)	1 (3.3)	0.042
Acrocyanosis, n (%)	2 (40.0)	0 (0.0)	0.439
Raynaud's phenomenon, n (%)	3 (60.0)	1 (100.0)	0.439
Treatment			
Paracetamol, n (%)	7 (29.2)	10 (33.3)	0.743
NSAIDs, n (%)	5 (20.8)	6 (20.0)	0.940
GCs, n (%)	9 (37.5)	17 (56.7)	0.161

Continued

Table 3 Continued

	Post-COVID-19 (n=24)	Post-vaccine (n=30)	P value
Intra-articular GCs, n (%)	2 (8.3)	0 (0.0)	0.107
Colchicine, n (%)	0 (0.0)	3 (10.0)	0.111
MTX, n (%)	3 (12.5)	3 (10.0)	0.771
HCQ, n (%)	4 (16.7)	6 (20.0)	0.754
AZA, n (%)	0 (0.0)	2 (6.7)	0.197
MMF, n (%)	1 (4.2)	1 (3.3)	0.872
CYC, n (%)	0 (0.0)	1 (3.3)	0.367
IVIg, n (%)	1 (4.2)	1 (3.3)	0.472
CCBs, n (%)	4 (16.7)	2 (6.7)	0.245

*Expected number in one category is equal to zero.

ACA, anti-centromere antibodies; ADM, amyopathic dermatomyositis; ANA, antinuclear antibodies; AZA, azathioprine; CCBs, calcium channel blockers; CRP, C reactive protein; CYC, cyclophosphamide; ESR, erythrocyte sedimentation rate; GCs, glucocorticoids; HCQ, hydroxychloroquine; IIM, idiopathic inflammatory myopathy; IVIg, intravenous immunoglobulin; MCTD, mixed connective tissue disease; MMF, mycophenolate mofetil; MTX, methotrexate; N/A, not applicable; NSAIDs, non-steroidal anti-inflammatory drugs; SjS, Sjögren's syndrome; SLE, systemic lupus erythematosus; SSc, systemic sclerosis; UCTD, undifferentiated connective tissue disease.

To contribute to the body of knowledge in this field, we systematically collected consecutive cases of IRD onset in close temporal association with COVID-19 or vaccine administration, in order to identify potential differences between the two cohorts and more accurately depict the spectrum of diseases.

According to our results, IRD may occur roughly equivalently both following SARS-CoV-2 infection or COVID-19 vaccination and the range of manifestations is broad, covering different forms of IJD, CTD or vasculitis. However, it is crucial to disclose that temporal association alone does not imply causality and may merely represent a coincidental association, and, thus, no pathophysiological explanation can be extrapolated from ours' and others' available data. Furthermore, although we report a similar number of cases in the two cohorts, it must be pointed out that the denominator of the overall population is not known, preventing any epidemiological consideration to be drawn. Anyway, from a merely speculative point of view and according to the Italian Ministry of Health official data, 25 millions of cases of COVID-19 have been registered to date while more than 143 million of doses of COVID-19 vaccines have been administered.⁵ This implies that the number of individuals potentially susceptible to vaccine-related events is at least fivefold higher than that exposed to SARS-CoV-2 infection. Consequently, the association is anticipated to occur more frequently by coincidence in the postvaccination cohort (ie, no more than the anticipated number of incident cases of the specific IRD in such a heavily immunised population).

Table 4 Comparative analysis of clinical features in vasculitis with onset after SARS-CoV-2 infection or COVID-19 vaccine administration

	Post-COVID-19 (n=8)	Post-vaccine (n=13)	P value
Age, years	58±20	62±17	0.632
Female sex, n (%)	5 (62.5)	9 (69.2)	0.751
N° vaccine doses, n	2±1	2±1	0.739
ANA positive, n (%)	2 (25)	4 (30.8)	0.776
ESR, mm/h	42±33	67±46	0.162
CRP, mg/dL	3.57±2.80	5.48±5.89	0.329
Specific diseases			
Large vessel vasculitis, n (%)	2 (25.0)	3 (23.1)	0.920
C-GCA, n (%)	1 (12.5)	2 (15.4)	0.854
LV-GCA, n (%)	1 (12.5)	1 (7.7)	0.716
Small vessel vasculitis, n (%)	3 (37.5)	3 (23.1)	0.477
ANCA positive, n (%)	2 (66.7)	1 (33.3)	0.414
EGPA, n (%)	2 (66.7)	2 (66.7)	1.000
GPA, n (%)	0 (0.0)	1 (33.3)	0.273
MPA, n (%)	1 (33.3)	0 (0.0)	0.273
Variable vessel vasculitis, n (%)	0 (0.0)	1 (7.7)	0.421
BD, n (%)	0 (0.0)	1 (7.7)	N/A*
Skin vasculitis, n (%)	3 (37.5)	6 (46.2)	0.697
Palpable purpura, n (%)	2 (66.7)	6 (100.0)	0.134
Urticarial vasculitis, n (%)	1 (33.3)	0 (0.0)	0.134
Treatment			
NSAIDs, n (%)	1 (12.5)	2 (15.4)	0.854
GCs, n (%)	6 (75.0)	12 (92.3)	0.271
Colchicine, n (%)	0 (0.0)	1 (7.7)	0.421
MTX, n (%)	1 (12.5)	0 (0.0)	0.191
SSZ, n (%)	1 (12.5)	0 (0.0)	0.191
AZA, n (%)	2 (25.0)	1 (7.7)	0.271
CYC, n (%)	0 (0.0)	1 (7.7)	0.421
ANR, n (%)	0 (0.0)	1 (7.7)	0.421

*Expected number in one category is equal to zero.
 ANA, antinuclear antibodies; ANCA, antineutrophil cytoplasmic antibodies; ANR, anakinra; AZA, azathioprine; BD, Behçet's disease; C-GCA, cranial giant cell arteritis; CRP, C reactive protein; CYC, cyclophosphamide; EGPA, eosinophilic granulomatosis with polyangiitis; ESR, erythrocyte sedimentation rate; GCs, glucocorticoids; GPA, granulomatosis with polyangiitis; LV-GCA, large vessel giant cell arteritis; MPA, microscopic polyangiitis; MTX, methotrexate; N/A, not applicable; NSAIDs, non-steroidal anti-inflammatory drugs; SSZ, sulfasalazine.

Additionally, increase in disease activity following SARS-CoV-2 infection has been reported in a significant portion of patients with pre-existent IRD,^{32 33} but it is difficult to uncouple the direct role of the virus and/or secondary immune system activation from the

consequences of background DMARD treatment disruption. Similarly, vaccine administration has been associated with worsening of disease activity in a minor, but still present, fraction of patients.^{34 35} In this context, although it is not possible to exclude that a small proportion of patients in our cohort may actually represent a SARS-CoV-2 or vaccine-induced exacerbation of latent or previously undiagnosed IRD, evidence of disease flare-up in previously diagnosed IRD patients may suggest common underlying pathophysiological mechanisms, leading to the development of a true, 'de novo', immunopathological event.

Despite these limitations, our data provide an interesting overview of the spectrum of the diseases possibly associated with either SARS-CoV-2 infection or COVID-19 vaccine and highlight some potential differences between the two cohorts. Indeed, we found a relatively higher percentage of patient developing IJD after COVID-19, while, on the other hand, a higher percentage of vaccine recipients developed PMR. Conversely, the distribution of other IRD categories did not differ between the two groups.

Viruses are well-known triggers for acute, subacute or chronic arthritis as commonly occur after parvovirus B19, HCV or chikungunya virus infection.³⁶ A previous report from our group²⁵ suggested that the delay between COVID-19 and arthritis onset is associated with the clinical phenotype of arthritis: while an RA-like pattern (polyarthritis mainly involving wrist and hand)—reminiscent of other prototypical viral arthritis—was more common in patients with early onset of rheumatic manifestations after infection (≤ 2 weeks), a late onset was more frequently associated with oligoarthritis of large joints suggestive of reactive arthritis. Viral infections have been evoked as a potential pathophysiological contributor to a number of other autoimmune diseases with different mechanisms including, but not limited to, molecular mimicry, bystander activation and epitope spreading.³⁷

Similarly to infections, vaccine administration has been associated with the development of autoimmunity.³⁸ The debated ASIA syndrome²¹ was defined in 2011 by Shoenfeld and collaborators as a condition in which the exposure to an adjuvant (eg, aluminium-based or squalene) leads to an abnormal, autoimmunogenic, immune response. However, it must be pointed out that novel mRNA vaccines do not contain traditional adjuvants because nucleic acids are sufficiently immunogenic to induce an effective immunisation.³⁹

Our findings regarding post-vaccine IRD are in line with available literature, demonstrating that PMR is the most reported IRD after COVID-19 vaccine administration.^{25 40 41} Similar data were previously reported for influenza vaccine^{42 43} and supposed to result from an external trigger leading to an aberrant response in the context of a senescent immune system. Similarly to what observed in the post-COVID-19 cohort, however, in the post-vaccine group we reported a number of other ARD belonging to all categories mirroring isolated case reports or small

series available in literature.^{22–44–46} Furthermore, a growing body of evidence support an association between mRNA vaccines and non-rheumatic autoimmune diseases such as myocarditis,^{47–48} while adenovirus-based vaccines have been associated with vaccine-induced immune thrombotic thrombocytopenia^{49–50} and Guillain-Barré syndrome.^{51–52}

Interestingly, when comparing clinical features of IJD between the two cohorts, we observed a higher relative percentage of PsA in the post-vaccine cohort. There are no obvious explanations for this finding although both new-onset psoriasis and psoriasis flares have been reported following COVID-19 vaccination.⁵³ A speculative interpretation may be that the presence of concurrent skin disease (either new-onset or exacerbation of previously unrecognised psoriasis) promoted by vaccine administration, may lead to classify as PsA some patients of the post-vaccine cohort that otherwise would be attributed to the pSpA category. Unfortunately, we did not collect information regarding the cutaneous domain and therefore no sub-analysis can contribute to interpret this finding.

Another interesting finding is the higher prevalence of new-onset acrosyndromes in post-COVID-19 patients. This finding is not unsurprising in the light of the well-described endothelial tropism of SARS-CoV-2⁵⁴; indeed, cutaneous microvascular changes have been described since the beginning of the pandemic as chilblain-like lesions or ‘COVID toes’.⁵⁵ In a Spanish observational study, acro-ischæmic lesions were observed in 0.6% outpatients and 2.9% hospitalised COVID-19 patients.⁵⁶ More recently, Sulli *et al*⁵⁷ reported that the mean nail-fold capillary number per linear millimetre was significantly lower in COVID-19 survivors when compared with primary Raynaud’s phenomenon patients and control individuals.

In conclusion, although the aforementioned limitations, our article reports the largest cohort published to date and supports the hypothesis that new-onset IRD may be triggered by SARS-CoV-2 infection or COVID-19 vaccines and that the spectrum of possible clinical manifestations is broad and includes IJD, PMR, CTD and vasculitis. However, it is highly plausible that the risk conferred by SARS-CoV-2 infection is higher and thus the usefulness and safety of vaccines, in our opinion, are not to be questioned. Larger epidemiological studies are needed to better clarify the causal role of each individual exposure on this association, quantify the true incidence and eventually explore the pathophysiological mechanism linking SARS-CoV-2 infection or vaccination to the development of IRD.

Author affiliations

¹Medicine & Rheumatology Unit, IRCCS Istituto Ortopedico Rizzoli, Bologna, Italy

²Department of Biomedical and Neuromotor Sciences (DIBINEM), Alma Mater Studiorum University of Bologna, Bologna, Italy

³Department of Biotechnological and Applied Clinical Sciences, University of L’Aquila, L’Aquila, Italy

⁴UO Interaziendale Medicina Interna ad Indirizzo Reumatologico, AUSL BO—IRCCS AOU BO, Bologna, Italy

⁵Unit of Rheumatology, Department of Medicine, ARNAS Garibaldi Hospital, Catania, Italy

⁶Rheumatology Unit, Rheumatology Hospital ‘Madonna dello Scoglio’, Cotronei, Italy

⁷Department of Internal Medicine, Ospedale Policlinico San Martino, Genoa, Italy

⁸UO Reumatologia, Ospedale di Castrovillari, Castrovillari, Italy

⁹Rheumatology Outpatient Clinic, Ospedale Villa Salus, Mestre, Italy

¹⁰Department of Precision Medicine, University of Campania ‘L. Vanvitelli’, Naples, Italy

¹¹Rheumatology and Clinical Immunology Unit, ASST Spedali Civili, Brescia, Italy

¹²Unit of Immunology, Rheumatology, Allergy and Rare Diseases (UnIRAR), IRCCS San Raffaele Hospital, Milan, Italy

¹³Rheumatology Unit, ‘Mauro Scarlato’ Hospital, Scafati, Italy

¹⁴Department of Clinical and Molecular Sciences, Polytechnic University of Marche, Ancona, Italy

¹⁵Division of Rheumatology and Clinical Immunology, IRCCS Humanitas Research Hospital, Rozzano, Italy

¹⁶Department of Biomedical Sciences, Humanitas University, Pieve Emanuele, Milan, Italy

¹⁷Department of Translational and Precision Medicine, Sapienza University of Rome, Rome, Italy

¹⁸Fondazione Policlinico Universitario Campus Bio-Medico, Rome, Italy

¹⁹Research and Clinical Unit of Immunorheumatology, Department of Medicine and Surgery, Università Campus Bio-Medico, Rome, Italy

²⁰Rheumatology Outpatient Clinic, ASP Cosenza, Scalea, Italy

²¹Rheumatology Outpatient Clinic, ASP Crotone, Crotone, Italy

²²UOC Medicina Interna ‘M.Valentini’, Presidio Ospedaliero ‘Annunziata’, Cosenza, Italy

²³Rheumatology Unit, Department of Clinical and Molecular Sciences, Polytechnic University of Marche, Ancona, Italy

²⁴Department of Clinical Medicine and Surgery, University of Naples Federico II, Naples, Italy

²⁵Academic Rheumatology Centre, Dipartimento Scienze Cliniche e Biologiche, Università degli Studi di Torino, Torino, Italy

²⁶Rheumatology Unit, Department of Precision and Regenerative Medicine-Ionian Area, University of Bari ‘Aldo Moro’, Bari, Italy

²⁷Rheumatology Unit, Policlinico di Modena, University of Modena and Reggio Emilia, Modena, Italy

Contributors FU, CF, PR, FI, AI, RG contributed to the conception and design of the work. FU analysed the data; FU, CF, PR, FI, AI, RG contributed to data interpretation. FU and CF prepared the first draft of the work. All the other authors contributed to data acquisition, revised the draft critically and approved the final version to be published. CF is responsible for the overall content as guarantor.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Not applicable.

Ethics approval This study involves human participants and was approved by Comitato Etico Azienda Sanitaria Locale 1 Avezzano/Sulmona/L’Aquila, L’Aquila, Italy. Participants gave informed consent to participate in the study before taking part.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available upon reasonable request.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Francesco Ursini <http://orcid.org/0000-0001-8194-8642>

Piero Ruscitti <http://orcid.org/0000-0003-3487-8551>

Alberto Lo Gullo <http://orcid.org/0000-0003-4383-0314>

Corrado Campochario <http://orcid.org/0000-0001-6806-3794>

Francesca Motta <http://orcid.org/0000-0002-8093-2734>

Maria De Santis <http://orcid.org/0000-0002-3196-1336>

Lorenzo Dagna <http://orcid.org/0000-0002-7428-315X>
 Carlo Selmi <http://orcid.org/0000-0002-0323-0376>
 Clodoveo Ferri <http://orcid.org/0000-0002-0840-5817>

REFERENCES

- Lamers MM, Haagmans BL. SARS-Cov-2 pathogenesis. *Nat Rev Microbiol* 2022;20:270–84.
- Li M, Wang H, Tian L, et al. COVID-19 vaccine development: milestones, lessons and prospects. *Sig Transduct Target Ther* 2022;7:146.
- Vangeel L, Chiu W, De Jonghe S, et al. Remdesivir, Molnupiravir and Nirmatrelvir remain active against SARS-Cov-2 Omicron and other variants of concern. *Antiviral Research* 2022;198:105252.
- Watson OJ, Barnsley G, Toor J, et al. Global impact of the first year of COVID-19 vaccination: a mathematical Modelling study. *Lancet Infect Dis* 2022;22:1293–302.
- Italian Ministry of Health. Report Vaccini anti COVID-19; Italian Ministry of health Available: <https://www.governo.it/it/cscovid19/report-vaccini/>
- Phetsouphanh C, Darley DR, Wilson DB, et al. Immunological dysfunction persists for 8 months following initial mild-to-moderate SARS-Cov-2 infection. *Nat Immunol* 2022;23:210–6.
- Nalbandian A, Sehgal K, Gupta A, et al. Post-acute COVID-19 syndrome. *Nat Med* 2021;27:601–15.
- Fernández-de-las-Peñas C, Navarro-Santana M, Plaza-Manzano G, et al. Time course prevalence of post-COVID pain systemic of musculoskeletal origin in patients who had survived severe acute respiratory syndrome Coronavirus 2 infection: a systematic review and meta-analysis. *Pain* 2022;163:1220–31.
- Ursini F, Ciaffi J, Mancarella L, et al. Fibromyalgia: a new facet of the post-COVID-19 syndrome spectrum? results from a web-based survey. *RMD Open* 2021;7:e001735.
- Rojas M, Rodríguez Y, Acosta-Ampudia Y, et al. Autoimmunity is a hallmark of post-COVID syndrome. *J Transl Med* 2022;20:129.
- Gracia-Ramos AE, Martín-Nares E, Hernández-Molina G. New onset of autoimmune diseases following COVID-19 diagnosis. *Cells* 2021;10:3592.
- Ursini F, Ruscitti P, D'Angelo S, et al. Broad clinical spectrum of SARS-Cov-2-associated inflammatory joint disease in adults: a report of 35 cases from the COVID-19 & autoimmune systemic disease Italian study group. *Ann Rheum Dis* 2021;80:1498–501.
- Metyas S, Chen C, Aung T, et al. Rheumatologic manifestations of post SARS-Cov-2 infection: A case series. *CRR* 2022;18:346–51.
- Shimizu H, Matsumoto H, Sasajima T, et al. New-onset dermatomyositis following COVID-19: A case report. *Front Immunol* 2022;13:1002329.
- Kazzi B, Fine D, Geetha D, et al. New-onset lupus nephritis associated with COVID-19 infection. *Lupus* 2022;31:1007–11.
- Allez M, Denis B, Bouaziz J-D, et al. COVID-19-related IgA vasculitis. *Arthritis Rheumatol* 2020;72:1952–3.
- Bach J-F. Infections and autoimmune diseases. *J Autoimmun* 2005;25 Suppl:74–80.
- Courcoul A, Brinster A, Decullier E, et al. A Bicentre retrospective study of features and outcomes of patients with reactive arthritis. *Joint Bone Spine* 2018;85:201–5.
- Vassilopoulos D, Calabrese LH. Virally associated arthritis 2008: clinical, epidemiologic, and pathophysiological considerations. *Arthritis Res Ther* 2008;10:215.
- Agmon-Levin N, Paz Z, Israeli E, et al. Vaccines and Autoimmunity. *Nat Rev Rheumatol* 2009;5:648–52.
- Watad A, Quaresma M, Bragazzi NL, et al. The autoimmune/inflammatory syndrome induced by Adjuvants (ASIA)/Shoenfeld's syndrome: descriptive analysis of 300 patients from the International ASIA syndrome Registry. *Clin Rheumatol* 2018;37:483–93.
- Chen Y, Xu Z, Wang P, et al. New-onset autoimmune phenomena post-COVID-19 vaccination. *Immunology* 2022;165:386–401.
- Al-Allaf A-W, Neethu M, Al-Allaf Y. A case series and literature review of the Association of COVID-19 vaccination with autoimmune diseases: causality or chance *Cureus* 2022.
- Ursini F, Ruscitti P, Raimondo V, et al. Systemic syndromes of Rheumatological interest with onset after COVID-19 vaccine administration: a report of 30 cases. *Clin Rheumatol* 2022;41:2261–7.
- Ursini F, Ruscitti P, Raimondo V, et al. Spectrum of short-term inflammatory musculoskeletal manifestations after COVID-19 vaccine administration: a report of 66 cases. *Ann Rheum Dis* 2022;81:440–1.
- Ferri C, Ursini F, Gagnani L, et al. Impaired Immunogenicity to COVID-19 vaccines in autoimmune systemic diseases. high prevalence of non-response in different patients' subgroups. *J Autoimmun* 2021;125:102744.
- Rayhan RU, Zheng Y, Uddin E, et al. Administer and collect medical questionnaires with Google documents: a simple, safe, and free system. *Appl Med Inform* 2013;33:12–21.
- Prevo ML, van 't Hof MA, Kuper HH, et al. Modified disease activity scores that include twenty-eight-joint counts. development and validation in a prospective longitudinal study of patients with rheumatoid arthritis. *Arthritis Rheum* 1995;38:44–8.
- Leeb BF, Bird HA. A disease activity score for Polymyalgia Rheumatica. *Ann Rheum Dis* 2004;63:1279–83.
- World Medical Association. World Medical Association Declaration of Helsinki: ethical principles for medical research involving human subjects. *JAMA* 2013;310:2191–4.
- World Health Organization (WHO). Living guidance for clinical management of COVID-19. 2021. Available: <https://www.who.int/publications/i/item/WHO-2019-nCoV-clinical-2021-2>
- Di Iorio M, Cook CE, Vanni KMM, et al. DMARD disruption, rheumatic disease flare, and prolonged COVID-19 symptom duration after acute COVID-19 among patients with rheumatic disease: A prospective study. *Semin Arthritis Rheum* 2022;55:152025.
- Dharia T, Venkatchalam S, Baker JF, et al. Medication interruptions and subsequent disease flares during the COVID-19 pandemic: A longitudinal Online study of patients with rheumatic disease. *Arthritis Care Res (Hoboken)* 2022;74:733–40.
- Álvarez-Gracia JM, Sánchez-Piedra C, Cuiqui D, et al. Effects of COVID-19 vaccination on disease activity in patients with rheumatoid arthritis and Psoriatic arthritis on targeted therapy in the COVIDSER study. *RMD Open* 2023;9:e002936.
- Machado PM, Lawson-Tovey S, Strangfeld A, et al. Safety of vaccination against SARS-Cov-2 in people with rheumatic and musculoskeletal diseases: results from the EULAR Coronavirus vaccine (COVAX) physician-reported Registry. *Ann Rheum Dis* 2022;81:695–709.
- Marks M, Marks JL. Viral arthritis. *Clin Med (Lond)* 2016;16:129–34.
- Smatti MK, Cyprian FS, Nasrallah GK, et al. Viruses and Autoimmunity: A review on the potential interaction and molecular mechanisms. *Viruses* 2019;11:762.
- Toussiot É, Bereau M. Vaccination and induction of autoimmune diseases. *Inflamm Allergy Drug Targets* 2015;14:94–8.
- Verbeke R, Hogan MJ, Loré K, et al. Innate immune mechanisms of mRNA vaccines. *Immunology* 2022;55:1993–2005.
- Mettler C, Jonville-Bera A-P, Grandvuillemin A, et al. Risk of giant cell arteritis and Polymyalgia Rheumatica following COVID-19 vaccination: a global Pharmacovigilance study. *Rheumatology (Oxford)* 2022;61:865–7.
- Ottaviani S, Juge P-A, Forien M, et al. Polymyalgia Rheumatica following COVID-19 vaccination: A case-series of ten patients. *Joint Bone Spine* 2022;89:105334.
- Falsetti P, Conticini E, Acciai C, et al. Polymyalgia Rheumatica following infective triggers or Vaccinations: a different subset of disease *Reumatologia* 2020;58:76–80.
- Liozon E, Parreau S, Filloux M, et al. Giant cell arteritis or Polymyalgia Rheumatica after influenza vaccination: A study of 12 patients and a literature review. *Autoimmun Rev* 2021;20:102732.
- Chen C-C, Chen C-J. New-onset inflammatory arthritis after COVID-19 vaccination: A systematic review. *Int J Rheum Dis* 2023;26:267–77.
- Khanna U, Oprea Y, Mir A, et al. New diagnosis of systemic lupus erythematosus after COVID-19 vaccination: A case report and review of literature. *JAAD Case Reports* 2022;30:30–4.
- Abdelmaksoud A, Wollina U, Temiz SA, et al. SARS-Cov-2 vaccination-induced cutaneous vasculitis: report of two new cases and literature review. *Dermatol Ther* 2022;35:e15458.
- Ling RR, Ramanathan K, Tan FL, et al. Myopericarditis following COVID-19 vaccination and non-COVID-19 vaccination: a systematic review and meta-analysis. *Lancet Respir Med* 2022;10:679–88.
- Heymans S, Cooper LT. Myocarditis after COVID-19 mRNA vaccination: clinical observations and potential mechanisms. *Nat Rev Cardiol* 2022;19:75–7.
- Kelton JG, Arnold DM, Nazy I. Lessons from vaccine-induced immune thrombotic thrombocytopenia. *Nat Rev Immunol* 2021;21:753–5.
- Elberry MH, Abdelgawad HAH, Hamdallah A, et al. A systematic review of vaccine-induced thrombotic thrombocytopenia in individuals who received COVID-19 adenoviral-vector-based vaccines. *J Thromb Thrombolysis* 2022;53:798–823.
- Abolmaali M, Reznania F, Behnagh AK, et al. Guillain-Barré syndrome in association with COVID-19 vaccination: a systematic review. *Immunol Res* 2022;70:752–64.

- 52 Hanson KE, Goddard K, Lewis N, *et al.* Incidence of Guillain-Barré syndrome after COVID-19 vaccination in the vaccine safety Datalink. *JAMA Netw Open* 2022;5:e228879.
- 53 Wu P-C, Huang I-H, Wang C-W, *et al.* New Onset and exacerbations of psoriasis following COVID-19 vaccines: A systematic review. *Am J Clin Dermatol* 2022;23:775–99.
- 54 Varga Z, Flammer AJ, Steiger P, *et al.* Endothelial cell infection and Endotheliitis in COVID-19. *Lancet* 2020;395:1417–8.
- 55 Kanitakis J, Lesort C, Danset M, *et al.* Chilblain-like Acral lesions during the COVID-19 pandemic ('COVID toes'): histologic, Immunofluorescence, and immunohistochemical study of 17 cases. *J Am Acad Dermatol* 2020;83:870–5.
- 56 Alonso MN, Mata-Forte T, García-León N, *et al.* Incidence, characteristics, laboratory findings and outcomes in Acro- ischemia in COVID-19 patients. *Vasc Health Risk Manag* 2020;16:467–78.
- 57 Sulli A, Gotelli E, Bica PF, *et al.* Detailed Videocapillaroscopic Microvascular changes detectable in adult COVID-19 survivors. *Microvasc Res* 2022;142:104361.