



UNIVERSITÀ POLITECNICA DELLE MARCHE
Repository ISTITUZIONALE

Susceptibility of polar cod (*Boreogadus saida*) to a model carcinogen

This is the peer reviewed version of the following article:

Original

Susceptibility of polar cod (*Boreogadus saida*) to a model carcinogen / Lerebours, A.; Murzina, S.; Song, Y.; Tollefsen, K. E.; Benedetti, M.; Regoli, F.; Rotchell, J. M.; Nahrgang, J.. - In: MARINE ENVIRONMENTAL RESEARCH. - ISSN 0141-1136. - ELETTRONICO. - 170:(2021). [10.1016/j.marenvres.2021.105434]

Availability:

This version is available at: 11566/295505 since: 2024-04-08T16:13:29Z

Publisher:

Published

DOI:10.1016/j.marenvres.2021.105434

Terms of use:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. The use of copyrighted works requires the consent of the rights' holder (author or publisher). Works made available under a Creative Commons license or a Publisher's custom-made license can be used according to the terms and conditions contained therein. See editor's website for further information and terms and conditions.

This item was downloaded from IRIS Università Politecnica delle Marche (<https://iris.univpm.it>). When citing, please refer to the published version.

(Article begins on next page)

Marine Environmental Research

Susceptibility of the key Arctic fish polar cod (*boreogadus saida*) to a model carcinogen

--Manuscript Draft--

Manuscript Number:	MERE-D-21-00104R1
Article Type:	Full Length Article
Keywords:	Carcinogenesis; polar cod; Benzo(a)pyrene; cancer genes; DNA repair genes
Corresponding Author:	Adelaide Lerebours FRANCE
First Author:	Adelaide Lerebours
Order of Authors:	Adelaide Lerebours Svetlana Murzina You Song Knut Erik Tollefsen Maura Benedetti Francesco Regoli Jeanette M. Rotchell Jasmine Nahrgang
Abstract:	<p>Studies that aim to characterise the susceptibility of the ecologically relevant and non-model fish polar cod (<i>Boreogadus saida</i>) to model carcinogens are required. Polar cod were exposed under laboratory conditions for six months to control, 0.03 µg BaP/ g fish/ week and 0.3 µg BaP/ g fish/ week dietary benzo(a)pyrene (BaP), a reference carcinogen. The concentrations of the 3-OH-BaP bile metabolite and transcriptional responses of genes involved in DNA adduct recognition (<i>xpc</i>), helicase activity (<i>xpd</i>), DNA repair (<i>xpf</i>, <i>rad51</i>) and tumour suppression (<i>tp53</i>) were assessed after 0, 1, 3 and 6 months of exposure, alongside body condition indexes (gonadosomatic index, hepatosomatic index and condition factor). Micronuclei and nuclear abnormalities in blood and spleen, and liver histopathological endpoints were assessed at the end of the experiment.</p> <p>Fish grew steadily over the whole experiment and no mortality was recorded. The concentrations of 3-OH-BaP increased significantly after 1 month of exposure to the highest BaP concentration and after 6 months of exposure to all BaP concentrations showing the biotransformation of the mother compound. Nevertheless, no significant induction of gene transcript involved in DNA damage repair or tumour suppression were observed at the selected sampling times. These results together with the absence of chromosomal damage in blood and spleen cells, the subtle increase in nuclear abnormalities observed in spleen cells and the low occurrence of foci of cellular alteration suggested that the exposure was below the threshold of observable effects. Taken together, the results showed that polar cod was not susceptible to carcinogenesis using the BaP exposure regime employed herein.</p>
Suggested Reviewers:	<p>Jim Smith jim.smith@port.ac.uk Jim Smith is an eminent scientist in the field of environmental pollution (fish).</p> <p>Lourdes Cruz lourdescruzg@gmail.com Lourdes Cruz is an excellent fish toxicologist working at MRC, UK.</p> <p>Angelique Lazartigues alazartigues@yahoo.fr Angelique Lazartigue is a great fish ecotoxicologist.</p> <p>Helene Thomas helene.thomas@univ-lr.fr</p>

	Helene Thomas has a great expertise in the area of aquatic toxicology.
Response to Reviewers:	

1 Susceptibility of polar cod (*Boreogadus saida*) to a model carcinogen

2
3
4
5 3 Adélaïde Lerebours^{1,2,*}, ³Svetlana Murzina, ⁴You Song, ⁴Knut Erik Tollefsen, ⁵Maura
6
7 4 Benedetti, ⁵Francesco Regoli, ⁶Jeanette M. Rotchell and ⁷Jasmine Nahrgang
8
9
10 5

11
12 6 ¹School of Biological Sciences, University of Portsmouth, Portsmouth, PO1 2DY, United
13
14 7 Kingdom
15

16
17 8 ²UMR CNRS LIENSs, Littoral Environnement et Sociétés, Université de La Rochelle, La
18
19 9 Rochelle, 17 000, France
20
21

22 10 ³Institute of Biology of the Karelian Research Centre of the Russian Academy of Sciences (IB
23
24 11 KarRC RAS) 185910 Petrozavodsk, Russia
25
26

27 12 ⁴Norwegian Institute for Water Research (NIVA), Section of Ecotoxicology and Risk
28
29 13 Assessment, Gaustadalléen 21, N-0349 Oslo, Norway
30
31

32 14 ⁵ Department of Life and Environmental Sciences, Polytechnic University of Marche, Ancona,
33
34 15 60131, Italy
35

36 16 ⁶Department of Biological and Marine Sciences, University of Hull, Cottingham Road, Hull,
37
38 17 HU6 7RX, United Kingdom
39
40

41 18 ⁷Department of Arctic and Marine Biology, Faculty of Biosciences, Fisheries and Economics
42
43 19 UiT The Arctic University of Norway, 9037 Tromsø, Norway
44
45

46 20
47
48 21 **Keywords:** Carcinogenesis, polar cod, benzo(a)pyrene, cancer genes, DNA repair genes
49
50

51 22
52
53 23 * Corresponding author: Adélaïde Lerebours, phone number: +33 (0) 5 46 50 76 23, email:
54
55 24 adelaiderebours@univ-lr.fr
56
57
58
59 25
60
61
62
63
64
65

26 **Abstract**

1
2 27 Studies that aim to characterise the susceptibility of the ecologically relevant and non-model
3
4 28 fish polar cod (*Boreogadus saida*) to model carcinogens are required. Polar cod were exposed
5
6 29 under laboratory conditions for six months to control, 0.03 µg BaP/ g fish/ week and 0.3 µg
7
8 30 BaP/ g fish/ week dietary benzo(a)pyrene (BaP), a reference carcinogen. The concentrations of
9
10 31 the 3-OH-BaP bile metabolite and transcriptional responses of genes involved in DNA adduct
11
12 32 recognition (*xpc*), helicase activity (*xpd*), DNA repair (*xpf*, *rad51*) and tumour suppression
13
14 33 (*tp53*) were assessed after 0, 1, 3 and 6 months of exposure, alongside body condition indexes
15
16 34 (gonadosomatic index, hepatosomatic index and condition factor). Micronuclei and nuclear
17
18 35 abnormalities in blood and spleen, and liver histopathological endpoints were assessed at the
19
20 36 end of the experiment.
21
22
23
24
25

26
27 37 Fish grew steadily over the whole experiment and no mortality was recorded. The
28
29 38 concentrations of 3-OH-BaP increased significantly after 1 month of exposure to the highest
30
31 39 BaP concentration and after 6 months of exposure to all BaP concentrations showing the
32
33 40 biotransformation of the mother compound. Nevertheless, no significant induction of gene
34
35 41 transcript involved in DNA damage repair or tumour suppression were observed at the selected
36
37 42 sampling times. These results together with the absence of chromosomal damage in blood and
38
39 43 spleen cells, the subtle increase in nuclear abnormalities observed in spleen cells and the low
40
41 44 occurrence of foci of cellular alteration suggested that the exposure was below the threshold of
42
43 45 observable effects. Taken together, the results showed that polar cod was not susceptible to
44
45 46 carcinogenesis using the BaP exposure regime employed herein.
46
47
48
49
50

51 47

52
53 48

54
55 49

56
57 50

51 Introduction

52 Polycyclic aromatic hydrocarbons (PAHs) are a very large group of ubiquitous organic
53 compounds that can originate from petrogenic, pyrogenic, biogenic and diagenic sources. A
54 number of PAHs are identified as having carcinogenic properties and have been associated to
55 an increased incidence of liver tumours in flatfish in highly polluted environments (Malins *et al.*
56 *al.*, 1985; Myers *et al.*, 1991; Harshbarger and Clark, 1990; Vogelbein *et al.*, 1990; Baumann
57 and Harshbarger, 1998). This pathology has been used to monitor the effects of exposure to
58 PAHs and the health of marine ecosystem since the 1980s (Malins *et al.*, 1985; Veethaak and
59 Ap Rheinallt, 1992) and its assessment recommended by the International Council for
60 Exploration of the Sea (ICES) and the Oslo and Paris Convention (OSPAR) Joint Assessments
61 and Monitoring Programme (JAMP) (Lyons *et al.*, 2010).

62 Benzo(a)pyrene (BaP) is a well-known pyrogenic carcinogen in a plethora of animals
63 such as marine mammals (Acevedo-Whitehouse *et al.*, 2018; Poirier *et al.*, 2019), fish (Wang
64 *et al.*, 2010; Wills *et al.*, 2010) and mice (Kasala *et al.*, 2015; Chen *et al.*, 2019). The reference
65 oral dose below which no effect is expected is $3 \cdot 10^{-4}$ $\mu\text{g BaP /g per day}$, based on animal and
66 human studies (reviewed in EPA/635/R-17/003). The metabolites generated by endogenous
67 metabolism (biotransformation) are highly genotoxic. Phase I biotransformation of BaP is
68 mediated by cytochrome P450 (CYP) enzymes and produces highly reactive metabolic
69 intermediates such as diol-epoxide, dihydrodiol and 3-hydroxybenzo(a)pyrene (3-OH-BaP)
70 (Karle *et al.*, 2004; Zhu *et al.*, 2008; Rey-Salgueiro *et al.*, 2011). Those metabolites form DNA
71 adducts that interfere with DNA repair and replication (Phillips and Arlt, 2007). This represents
72 a critical event in the initiation of tumorigenesis, potentially leading to mutations within
73 specific regions of DNA, such as proto-oncogenes and tumour suppressor genes (Rotchell *et al.*
74 *al.*, 2001, Du Corbier *et al.*, 2005, Lerebours *et al.*, 2014, 2016). The carcinogenicity of BaP
75 has been well studied in several temperate fish species where specific DNA adducts are used

76 as markers for exposure and potential genotoxic effects. Exposure to BaP specifically caused
77 DNA adducts in fish such as pale chub (*Zacco platypus*) (Lee *et al.*, 2014) and killifish species
78 (*Fundulus grandis* and *F. similis*) (Willett *et al.*, 1995; Rose *et al.*, 2000, 2001). Moreover, BaP
79 exposure was associated with neoplastic lesions in brown bullhead (*Ameiurus nebulosus*)
80 (Ploch *et al.*, 1998), English sole (*Parophrys vetulus*) (Reichert *et al.*, 1998) and rainbow trout
81 (Hendricks *et al.*, 1985). PAH-induced lesions have also recently been suggested in marine
82 mammals such as harbour porpoises (*Phocoena phocoena*) (Acevedo-Whitehouse *et al.*, 2018)
83 and beluga whales (Poirier *et al.*, 2019). Pollution induced cancer affects many aquatic species
84 and represents a growing concern for aquatic wildlife (for a review see Baines *et al.*, 2021).

85 Tumourigenesis is a progressive process characterised by different stages for which the
86 underlying molecular steps and the role of environmental exposure are not always well-known.
87 Nonetheless, liver tumourigenesis has been well studied in flatfish (Stentiford *et al.*, 2010;
88 Lerebours *et al.*, 2013; 2014; 2017) and in the model fish Japanese medaka (*Oryzias latipes*)
89 (Rotchell *et al.*, 2001) and zebrafish (*Danio rerio*) (Li *et al.*, 2017; 2019). DNA repair
90 mechanisms have been associated with tumour formation perturbing several steps of the
91 nucleotide excision repair (NER) pathway, which recognises and repairs DNA adducts induced
92 by numerous environmental mutagens, including PAHs (Gillet and Schärer, 2006; Rastogi *et*
93 *al.*, 2010). While such mechanisms involved in the development of tumours in certain fish
94 species are well characterised, a substantial knowledge gap exists for non-model and
95 ecologically important species inhabiting remote regions in particular. The Arctic is currently
96 experiencing a rapid decline in sea ice (Kumar *et al.*, 2021) that may lead to a significant
97 increase in marine shipping (Ho *et al.*, 2010), oil and gas exploration and operation (Elias,
98 2018), and tourism (Meier *et al.*, 2014) and associated release of potential carcinogenic
99 contaminants (Elias, 2018). The polar cod (*Boreogadus saida*) is a keystone fish species in the
100 arctic marine ecosystem due to its abundance, distribution and central role in the food web

101 (Welch *et al.*, 1992). Polar cod has been considered a model fish for arctic ecotoxicology
102 studies (Jonsson *et al.*, 2010; Nahrgang *et al.*, 2009, 2010a,b,c). The toxicity of petroleum
103 compounds on the physiology of polar cod has been well studied (Geraudie *et al.*, 2014; Bender
104 *et al.*, 2016; Nahrgang *et al.*, 2016; Vieweg *et al.*, 2018; Nahrgang *et al.*, 2019) but the
105 tumourigenic potential of a potent carcinogen remains unknown in that species. A few studies
106 however have reported a potential susceptibility to carcinogenic contaminants including BaP.
107 The hepatic metabolism of BaP is particularly efficient in polar cod and a significant increase
108 of covalently bound reactive intermediates of BaP in the bile of fish has been found after dietary
109 exposure to BaP (Ingebrigtsen *et al.*, 2000; Bakke *et al.*, 2016). These reactive intermediates
110 were found to induce the formation of DNA adducts in the liver of that species (Aas *et al.*,
111 2003). This genotoxic effect can in turn result in cellular abnormalities and cancer initiation.
112 Finally, a recent study showed that expression of genes involved in DNA repair and cell cycle
113 regulation processes was modified in liver of polar cod dietary exposed to BaP (Song *et al.*,
114 2019).

115 In order to evaluate the susceptibility of polar cod to a carcinogenic compound, adult
116 specimens were exposed under laboratory conditions for six months to control, 0.03 µg BaP/ g
117 fish/ week and 0.3 µg BaP/ g fish/ week dietary BaP. Selected body condition indexes, bile
118 metabolite concentrations and transcriptional responses of genes involved in DNA adduct
119 recognition (*xpc*), helicase activity (*xpd*), DNA repair (*xpf*, *rad51*) and tumour suppression
120 (*tp53*) were assessed after 0, 1, 3 and 6 months of exposure. Blood and spleen micronuclei,
121 nuclear abnormalities and liver histopathological endpoints were assessed at the end of the
122 experiment.

123 The sampling times were selected because carcinogenesis is a long-term process. They
124 were comparable to the exposure durations used in several studies interested in carcinogenesis
125 in European eel (Nogueira *et al.*, 2006), brown bullhead and channel catfish (Ploch *et al.*, 1985)

126 rainbow trout (Hendricks et al., 1985, Black et al., 1985) and coho salmon (Black *et al.*, 1985)
127 exposed to BaP. The BaP doses selected were lower than the concentrations frequently used in
128 previous studies. They were 10 and 100 times lower than the concentration of 3 µg BaP/g of
129 fish /week (Colli-Dula *et al.*, 2018) that induced a decrease of body indexes in Nile tilapia after
130 one month of exposure. In addition, our highest concentration was 4 times lower than the lowest
131 concentration used in the study of Song *et al.*, (2019) that found gene expression changes in
132 polar cod after two weeks of exposure, a twelve times shorter exposure duration.

134 **Methods**

135 *Fish collection and exposure*

136 Adult polar cod (4 years old) were collected along the west coast of the Svalbard archipelago
137 (Norway) onboard RV Helmer Hanssen in January 2014 using a Campelen bottom trawl (at
138 200m depth) and a fish-lift (Holst and McDonald, 2000). At the Tromsø aquaculture research
139 station (Havbrukstasjon i Tromsø), fish were kept in 3000 L acclimation tank under a natural
140 light and temperature (1.5 - 3 °C) regime of 79°N (based on mooring data in Wallace *et al.*,
141 2010). During this period, fish were fed until satiation with thawed *Calanus sp.* copepods
142 (*Calanus AS*, Tromsø). Ninety fish were selected based on similar length (15 ± 1 cm) and
143 weight (25 ± 7 g) for the experiment (June 2014).

144 Polar cod were dietarily exposed to 0, 0.03 and 0.3 µg BaP per gram fish per week, for
145 6 months (2nd of July 2014 to 31st January 2015). The experiment was conducted in compliance
146 with the policies of the Norwegian animal welfare authorities (application ID 6571). Briefly, a
147 BaP (Sigma Aldrich, St. Louis, USA) solution in acetone was mixed with *Calanus spp* (*Calanus*
148 *AS*) to yield 0.5 or 5 µg BaP per g feed or acetone alone (acetone control). The
149 acetone was volatilized by constant stirring on a magnetic stirrer for 2.5 hours at 30 °C. Small
150 pellets were then created with the addition of 0.5 mL gelatin per g feed. Fish were fed pellets

151 corresponding to 4% of their body wet weight (bw) 5 days a week. On the first, third and fifth
152 day of a week, fish were exposed to dietary BaP or a solvent control by receiving the 2% bw
153 exposed feed (or solvent control) and 2% bw of unexposed feed (no BaP, no acetone). Feeding
154 was done by distributing the pellets to the surface of the tank. Thus, feeding hierarchies may
155 have occurred resulting in some intra-tank individual exposure variations. On the remaining 2
156 days of a week, all fish were fed 4% bw of unexposed feed. The amount of food given to each
157 tank was adjusted at each sampling point to account for both growth and sampling of
158 specimens. With this feeding regime, the fish nominally received an average of 0, 0.03 and 0.3
159 μg BaP per gram of fish per week. After 1 (2nd August), 3 (3rd October) and 6 (31th of January)
160 months, 10 fish per condition were anaesthetized and killed by a sharp blow to the head. Total
161 body weight (g) and fork length (cm) were measured and the presence of parasites recorded.
162 Liver and gonads were removed and weighed. Bile was snap frozen in liquid nitrogen and
163 stored at -80°C until 3-OH BaP metabolite determination. A liver section was snap frozen in
164 liquid nitrogen and stored at -80°C for molecular analyses. During the final sampling (6 months
165 of exposure), a standardized liver cross-section was fixed for 24 hrs in neutral buffered
166 formaldehyde (4%) before being transferred to 70% ethanol for subsequent histological
167 assessment. Blood and spleen samples were preserved in Carnoy solution (3 methanol: 1 acetic
168 acid) and stored at +4°C for subsequent identification of nuclear abnormalities and
169 micronuclei. Finally, somatic weight (g) was determined as weight of eviscerated fish.
170 Gonadosomatic index (GSI) and hepatosomatic index (HSI) were calculated as follows:

$$\text{GSI} = (\text{gonad weight} / \text{somatic weight}) \times 100$$

$$\text{HSI} = (\text{liver weight} / \text{somatic weight}) \times 100.$$

173 *3-OH-benzo[a]pyrene measurement*

174 Biliary 3-OH-benzo[a]pyrene metabolite concentration was determined after 1, 3 and 6 months
175 of exposure following the procedure detailed in Song *et al.*, (2018). Preparation of hydrolysed

176 bile samples was performed as described in Krahn *et al.*, (1992). Briefly, bile (1–20 μ L) was
177 mixed with an internal standard (triphenylamine) and diluted with demineralised water (10–50
178 μ L) and hydrolysed with β -glucuronidasearylsulphatase (20 μ L, 1 h at 37 °C). Methanol (75–
179 200 μ L) was added and the sample was mixed thoroughly before centrifugation. The
supernatant was then transferred to vials and analysed. High pressure liquid chromatography
180 (Waters 2695 Separations Module) was used to separate 3-OH-BaP in a Waters PAH C18
181 column (4.6 \times 250 mm, 5 μ m particle size). The mobile phase consisted of a gradient from
182 40:60 acetonitrile:ammonium acetate (0.05 M, pH 4.1) to 100% acetonitrile at a flow rate of 1
183 mL/min, and the column was heated to 35 °C. A 2475 fluorescence detector measured
184 fluorescence at the optimum for each analyte (excitation/emissions: 380/430). A total of 25 μ
185 L extract was injected for each analysis. The results were calculated by use of the internal
186 standard method (Grung *et al.*, 2009). The calibration standards utilized were obtained from
187 Chiron AS, Trondheim, Norway, and were in the range 0.2–200 ng/g. Values below the limit
188 of detection were considered as equal to 0 ng/g in the analyses.

189 *Histopathological analyses*

190 Livers were processed in a vacuum infiltration processor (Shandon Citadel 1000) using
191 standard histological protocols (Feist *et al.*, 2004). The tissues were embedded in paraffin using
192 an STP-120 spin tissue processor (Thermo fisher scientific, USA). Sections of 4 μ m thickness
193 were cut using a microtome HM 450 (Thermo Fisher Scientific, USA) and subsequently stained
194 with haematoxylin and eosin (H&E). The liver sections were examined for microscopic pre-
195 tumour and tumour lesions according to BEQUALM and ICES criteria (Feist *et al.*, 2004). The
196 pre-tumour lesions sought were the vacuolated, basophilic and eosinophilic foci of cellular
197 alteration (FCA). Tumour lesions were the benign hepatocellular adenoma and the malignant
198 hepatocellular carcinoma (HCC). Lesions associated to nuclear and cellular polymorphism,
199

200 cell death, inflammation and regeneration were also examined. A total of 5, 6 and 4 fish were
201 assessed from control, low and high exposure condition, respectively.

202 *Micronucleus test and nuclear abnormalities*

203 The micronuclei and nuclear abnormalities frequencies were measured in blood and spleen of
204 polar cod tissues fixed in Carnoy's solution; subsequently separated cells were dispersed on
205 glass slides, and stained with the fluorescent dye 4',6-diamidino- 2-phenylindole at 100 ng/
206 mL. For each experimental condition, a range of 6 to 8 fish were investigated, and for each
207 specimen 2000 cells with preserved cytoplasm were scored to assess the presence of
208 micronuclei and nuclear abnormalities. Micronuclei are defined as round structures, smaller
209 than 1/3 of the main nucleus diameter, on the same optical plan and clearly separated from
210 nucleus; Nuclear abnormalities include (i) binucleated: cell with two nuclei, (ii) notch nuclei:
211 looks like nucleus but do not have nuclear materials, (iii) nuclear bud: evagination of bud-like
212 structure from the nucleus, and (iv) blebbed nuclei: small euchromatin evagination of the
213 nuclear membrane (Gorbi *et al.*, 2009; Islam *et al.*, 2021).

214 *Gene expression analyses*

215 Following 1, 3 and 6 months of exposure, a cross section of each liver, next to the one dedicated
216 to histological analyses at 6 months was used for gene transcriptional response analyses. Total
217 RNAs were extracted using the High Pure RNA Tissue kit (Roche Diagnostics Ltd, West
218 Sussex, U.K.) according to the supplier's instructions which included a DNase treatment. RNA
219 quality (integrity of 18S and 28S ribosomal bands) was evaluated by electrophoresis on a 1%
220 agarose-formaldehyde gel. RNA purity was assessed by measuring the ratios of absorbance:
221 A_{260}/A_{280} and A_{260}/A_{230} using a spectrophotometer (NanoDrop, ThermoFisher). All samples
222 were of high purity (ratios' values > 2.1).

223 First strand cDNAs were synthesized from 1 µg of total RNA using the AffinityScript
224 Multiple Temperature cDNA Synthesis Kit (Agilent Technologies, Stockport, U. K.) using

225 hexa primers and according to the supplier's instructions. Putative coding sequences (Figure
226 S1) were identified by nucleotide and protein BLAST searches on the NCBI database
227 (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>) and sequence homologies across fish species on the
228 EMBL-EBI platform (<https://www.ebi.ac.uk/Tools/msa/clustalo/>). The contigs produced in the
229 study of Song *et al.*, (2019) were also used. Primer pairs and FAMTM-TAMRATM dye probes
230 used to amplify the target sequences were designed using the Prime Express software (Applied
231 Biosystem) (Table 1). Ten ng of the reverse transcribed product measured by a qubit
232 fluorometer (Thermo Fisher Scientific) was used as a template for subsequent polymerase
233 chain reaction (PCR) in a 20 µL final volume using 1x of TaqMan® Fast Advanced Master
234 Mix (Life technologies, Paisley, U.K.), 900 nM primers and 250 nM probe (final
235 concentrations) according to the supplier's protocol. PCR reactions were performed in the
236 Applied BiosystemsTM ViiATM 7 Real-Time PCR System using the following programme: one
237 cycle at 95°C for 20 s and 40 amplification cycles at 95°C for 3 s and 60°C for 30 s. Primer
238 efficiencies were determined by 10 times dilution series of the cDNA template and were about
239 100%. The optimal normalization gene was selected by testing the expressions of 3 reference
240 genes (*β tubulin*, *hpri1* and *28S*) on all the samples using the NormFinder algorithm. The
241 expression of the *β tubulin* gene displayed the highest stability. The melting curves were
242 carefully checked after each qPCR run. The gene expression was calculated according to the
243 delta delta Ct method.

245 *Statistical analyses*

246 Statistical analyses were performed using R (version 3.1.2). The effect of the BaP exposure
247 concentration and time of exposure were assessed on all the biological parameters measured
248 using 1-way ANOVA. When the normality of the residuals was not verified by the Shapiro-
249 Wilk test, the non-parametric Kruskal-Wallis tests were used. Post-hoc comparisons were

250 performed using the least-square mean test for parametric test and the Wilcoxon rank test for
251 non-parametric test. The α error was adjusted using the Bonferroni correction for each post-
252 hoc test.

253 Results

254 Over the course of the six-month exposure, all specimens grew significantly in weight and
255 underwent gonadal maturation with mean GSI ranging from 1.5 ± 0.7 to 20.8 ± 3.9 % (Table
256 2). Endoparasites were commonly found across all treatments and sampling times. Nematodes
257 on the liver surface were the most common parasites with a frequency of occurrence of 26%.
258 Parasites of the phylum Platyhelminthes were less common (7%). No mortality was observed.

259 The dietary BaP exposure of polar cod led to a dose-dependent production of biliary 3-
260 OH BaP metabolites for the low ($0.03 \mu\text{g BaP/g}$ of fish/week) and high ($0.3 \mu\text{g BaP/g}$ of
261 fish/week) BaP exposure conditions after 1 and 6 months of exposure (Figure 1). The
262 concentrations of bile 3-OH BaP ranged from 20 to 40 ng/g of bile for the low exposure
263 condition and were approximately 10 times higher, from 132 to 390 ng/g of bile, for the highest
264 exposure condition.

265 The transcriptional responses related to DNA adduct recognition (*xpc*), helicase activity
266 (*xpd*), DNA repair (*xpf*, *rad51*) and tumour suppression (*tp53*) were not significantly changed
267 by any BaP dietary exposures as compared to controls ($p > 0.05$) (Figure 2).

268 The number of micronuclei recorded in the blood and spleen of polar cod ($p > 0.05$)
269 exposed to BaP did not significantly vary as compared to control (Figure 3 A, B). Nuclear
270 abnormalities in polar cod spleen were significantly increased in the high dose group ($p = 0.03$),
271 while close to significant in the low dose group ($p = 0.057$) (Figure 3 C, D). No significant
272 nuclear abnormalities were observed after BaP exposure in blood cells ($p > 0.05$).

273 Histopathological analyses revealed one basophilic foci of cellular alteration in liver of
274 two individuals exposed to the low exposure condition after six months of exposure (Figure

275 S2). No tumour-related lesions were observed in livers of control and highly exposed
276 individuals.

277 Discussion

278 The present study showed that polar cod grew steadily through the entire experiments for all
279 exposure conditions (0, 0.03 and 0.3 µg BaP/g of fish/ week). These results were expected
280 under chronic low dose exposure scenarios and are consistent with similar results obtained in
281 polar cod exposed to higher dietary BaP concentrations (1.2 and 60.9 µg BaP/ g of fish/ week)
282 but for a shorter period of two weeks (Song *et al.*, 2019). In Nile tilapia exposed to
283 intraperitoneal injections of 3 µg BaP/g of fish/ week for 4 weeks, K, GSI and GSI were slightly
284 decreased (Colli-Dula *et al.*, 2018). This suggests that there is a dose and time dependent
285 threshold above which exposure to BaP induce significant body condition indexes changes.
286 The fish species and the mode of BaP administration (injected intraperitoneally versus dietary)
287 may also be important factors to consider.

288 The bile concentration of 3-OH-BaP has been used as an indicator fo BaP exposure ad
289 biotransformation in many fish species including polar cod (Baake *et al.*, 2016; Baali *et al.*,
290 2016; Kammann *et al.*, 2017; Song *et al.*, 2019). The increase in biliary 3-OH-BaP metabolite
291 concentrations after 1 and 6 months exposure supported that of a similar, albeit shorter,
292 exposure study (Song *et al.*, 2019). When exposed to a four times higher exposure dose (1.2
293 µg BaP/ g of fish/ week) than the highest dose used in the present study, a 3-OH-BaP
294 concentration of 800 ng/ g of bile was found, which was two to six times higher than the
295 metabolite concentration range identified in our study. The reactive BaP intermediates have
296 been found to accumulate and covalently bind DNA in the biliary system of polar cod one
297 month after exposure to a single dietary concentration (Baake *et al.*, 2016) equivalent to the
298 cumulative dose received in the high BaP exposure condition during the first month of our
299 study. Those reactive BaP metabolites covalently bind to biological molecules such nucleic

300 acids and form DNA adducts that can lead to tumour formation. For instance, higher levels of
301 BaP-7,8-diol metabolites and DNA binding activity were found in bile of English sole
302 (*Parophrys vetulus*) a fish species more sensitive to carcinogenesis than the more resistant
303 starry flounder (*Platichthys stellatus*) (Varanasi *et al.*, 1986). In polar cod dietarily exposed to
304 higher BaP concentrations (from 5 µg BaP/ g of fish in a single injection) DNA adducts were
305 found (Aas *et al.*, 2003), revealing an increased risk of liver tumour formation later on. Indeed,
306 50% of rainbow trout displayed pre-tumour (basophilic FCA) and tumour (HCC) liver lesions
307 after six months of exposure to a similar dose injected intraperitoneally (Hendricks *et al.*,
308 1985). In their study, 25 % of the trouts displayed similar liver lesions after twelve months of
309 dietary exposure to a high dose of BaP (estimated to 1-2 mg BaP/ g fish/ week) (Hendricks *et*
310 *al.*, 1985). In the present study, the potential genotoxic damage generated by the BaP
311 metabolites produced did not cause significant tumour lesions. This could be the result of
312 several factors potentially in combination, including low dose, low exposure duration and
313 effective DNA repair mechanisms.

314 The DNA repair system and cell cycle regulators can prevent DNA adducts and the
315 onset of tumorous events. In the present study, the transcriptional response of genes involved
316 in the nucleotide excision repair (NER) process (*xpc*, *xpd*, *xpf*), DNA double strand breaks
317 repair (*rad51*) and cell division regulation (*tp53*) did not vary significantly after 1, 3, and 6
318 months of exposure to both BaP dietary concentrations. The exposure levels of BaP may have
319 been too low to cause significant accumulation of cell damage and trigger a significant gene
320 transcriptional response. Interestingly, a dose-specific transcriptional response of some genes
321 has been observed in liver of polar cod dietarily exposed to BaP (Song *et al.*, 2019). For instance,
322 some genes involved in apoptosis (*bax* and *casp9*), a process that eliminates damaged cells and
323 prevent the proliferation of abnormal cells in tumour formation, were upregulated in polar cod
324 exposed to the high exposure level (60.9 µg BaP/ g of fish/ week). The expression levels of

325 those genes were not modified in fish exposed to the low exposure dose (1.2 µg BaP/ g of
326 fish/week) suggesting a threshold above which gene transcription is modified (Song *et al.*,
327 2019). The basal gene expression level may also be sufficient to repair DNA and/or delay the
328 cell cycle to maintain the genetic integrity. Moreover, the DNA repair gene measured in our
329 study, *rad51*, may not be involved in the repair of specific DNA damage induced. Similarly to
330 the results herein, this gene was not differentially expressed in liver of polar cod dietary
331 exposed to BaP (Song *et al.*, 2019). *Rad51* is involved in the repair of DNA double strand
332 breaks, which belong to a different pathway than the NER. The mechanism of DNA damage
333 induced by BaP exposure is more likely to involve DNA adducts than double strand breaks.
334 Other genes involved in DNA repair processes and control of cell cycle have been found
335 induced at higher exposure regimes. For example, the gene encoding for the growth arrest and
336 DNA damage inducible beta gene (*gadd45b*) was induced in the liver of the tropical fish, Nile
337 Tilapia (*Oreochromis niloticus*) after one month of exposure to 3 µg BaP/ g of fish/week (Colli-
338 Dula *et al.*, 2018). In polar cod exposed to 1.2 and 60.9 µg BaP/g of fish/ week, genes involved
339 in the excision DNA repair process (such as *hmg2b* and *rad23a*) were differentially expressed
340 (Song *et al.*, 2019). Shorter timepoints may have also been necessary to observe a gene
341 expression modulation as an early response to stressors. In the liver of polar cod, transcriptional
342 responses of genes involved in DNA damage repair were changed after two weeks of dietary
343 exposure to BaP (Song *et al.*, 2019). Some studies using a reference genotoxic compound
344 showed that DNA damage was rapidly repaired with increased transcription of DNA repair
345 genes such as *rad51* in zebrafish larvae, as early as 6 hours (Reinardy *et al.* 2013). The addition
346 of early sampling times seems relevant to include in future studies.

347 Other biological processes such as detoxification mechanisms could have prevented
348 polar cod from the genotoxic effects of BaP exposure. Activation of detoxification events could
349 explain the resistance of polar cod to BaP exposure and the absence of liver tumours in the

350 present study. Variation in the expression of genes and proteins belonging to the cytochrome
351 P450 family involved in phase I of BaP detoxification process has been well described in liver
352 of fish (Nahrgang *et al.*, 2009; Lee *et al.*, 2014; Colli-Dula *et al.*, 2018). Interestingly, *cyp1a1*
353 and *cyp1b1* genes were upregulated in liver of polar cod dietary exposed to 60.9 µg BaP/g of
354 fish/ week but were not differentially expressed after exposure to a lower dose of BaP (1.2 µg
355 BaP/g of fish/ week) (Song *et al.*, 2019). This suggests a dose threshold for activating the
356 detoxification mechanisms during a chronic exposure. Activation of genes and proteins
357 involved in phase II detoxification process has been also described in liver of fish exposed to
358 BaP (Nahrgang *et al.*, 2009). For instance, *gstA1* gene expression was modified in the liver of
359 Nile tilapia exposed to 3 µg BaP/ g of fish/ week (Colli-Dula *et al.*, 2018). Interestingly, GST
360 activity was higher in starry flounder, a tumour resistant species, than in English sole, a tumour
361 sensitive species, after exposure to a BaP dose that induced carcinogenesis (Varanasi *et al.*,
362 1987). Finally, phase III detoxification process based on active efflux of chemicals by ATP-
363 binding cassette (ABC) transporters could be involved in BaP elimination. For example, a
364 rainbow trout ABCG2 transporter was found to interact with BaP (Zaja *et al.*, 2016).

365 In accordance with relatively low biliary BaP metabolite concentrations observed
366 throughout the exposure and limited responses of genes involved in DNA damage
367 identification and repair, no effect of dietary BaP on the micronuclei and nuclear abnormalities
368 was recorded in the present study except in spleen where nuclear abnormalities increased in
369 the high dose group. Micronuclei are formed during the anaphase stage of the cell division.
370 They are considered as a reliable index of chromosomal breakage, chromosomal loss and
371 cellular spindle malfunction (Bolognesi and Hayashi, 2011). Additionally, micronuclei
372 constitute an irreversible form of genotoxic damage compared to DNA strand breaks and their
373 induction are regulated by a large number of experimental carcinogens, including chlorinated
374 hydrocarbons, benzidine, aflatoxins, methylcholanthrene, and common carcinogenic

375 pollutants, such as PAHs, heavy metals, and pesticides (Bolognesi and Hayashi, 2011). Many
376 research studies reported the increased in micronuclei frequency in erythrocytes of different
377 fish species exposed to PAHs (Shirmohammadi *et al.*, 2018).

378 Contrary to micronuclei, nuclear abnormalities origin has not been clearly explained; some
379 suggest that nuclear abnormalities can be a primary response, prior to the micronuclei
380 formation, highlighting their relevance in the evaluation of genotoxic damage (Bolognesi and
381 Hayashi, 2011; Seriani *et al.*, 2011). An increase of erythrocytic nuclear abnormalities and
382 strand breaks was observed in eels (*Anguilla anguilla* L.) and juvenile sea bass (*Dicentrarchus*
383 *labrax*) exposed to a range of 0.3 to 2.7 μM of BaP and naphthalene (Maria *et al.*, 2002; Teles
384 *et al.*, 2003; Gravato and Santos, 2002), while on the contrary, lower concentrations of BaP
385 (0.1 μM) did not affect DNA integrity (Nogueira *et al.*, 2006). The induction of micronuclei
386 and other nuclear abnormalities were also caused by crude oil exposure in turbot (*Scophthalmus*
387 *maximus*) and Atlantic cod (*Gadus morua*) (Baršienė *et al.*, 2004; 2006). The exposure duration
388 and levels are extremely important in determining micronuclei and nuclear abnormalities
389 formation; long-term chemical exposures can cause genetic changes and consequently
390 physiological alterations or pathologies including cancer development (Depledge and Hopkin,
391 1995). In a study led on the European flounder, *Platichthys flesus*, Köhler and Ellesat, (2008),
392 first suggested that nuclear anomalies inside liver lesions of hepatocellular cancers were
393 correlated with micronuclei frequencies in fish blood and that the
394 histopathological grading of cancers from preneoplastic, benign to malignant types was clearly
395 associated with micronuclei increase.

396 The present study showed that polar cod were consistently exposed to dietary BaP
397 through the entire experiment and biotransformed the mother compound to intermediate
398 metabolites. However, this exposure did not lead to significant changes in the transcription of
399 selected genes, nor in chromosomal alterations and significant tissue lesions. Some early

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

400 responses to stress may have occurred prior to the first sampling time point at one month of
401 exposure, and basal expression of genes or potentially activated compensatory mechanisms
402 may have been sufficient to control the damage caused by the reactive metabolites. Moreover,
403 protective mechanisms such as detoxification and apoptosis could have prevented the cells
404 from the accumulation of cell damage caused by the reactive metabolites. Therefore, we deduce
405 that the BaP exposure concentrations were below the threshold of observable effects. As a
406 whole, our results showed that polar cod exposed to 0.03 and 0.3 µg BaP/ g fish/ week was not
407 sensitive to the model carcinogen and liver carcinogenesis. The present work encourages the
408 addition of earlier sampling points and indicators of detoxification mechanisms in future
409 studies.

410 411 412 413 414 415 416 417 418 419 **References**

420
421 Aas E, Liewenborg B, Grøsvik BE, Camus L, Jonsson G, Børseth JF, Balk L. 2003. DNA
422 adduct levels in fish from pristine areas are not detectable or low when analysed using the
423 nuclease P1 version of the ³²P-postlabelling technique. *Biomarkers*, 8: 445-460.

425 Acevedo-Whitehouse K, Cole KJ, Philips DH, Jepson PD, Deaville R, Arlt VM. 2018. Hepatic
1
2 426 DNA Damage in Harbour Porpoises (*Phocoena phocoena*) Stranded along the English and
3
4
5 427 Welsh Coastlines. *Environmental and Molecular Mutagenesis*, 59: 613-624.

6
7 428
8
9
10 429 Baali A, Kammann U, Hanel R, El Qoraychy I, Yahyaoui A. 2016. Bile metabolites of
11
12 430 polycyclic aromatic hydrocarbons (PAHs) in three species of fish from Morocco.
13
14 431 *Environmental Sciences Europe*. 28, 25.

15
16
17 432
18
19 433 Baines C, Lerebours A, Thomas F, Fort J, Kreitsberg R, Gentes S, Meitern R, Saks L, Ujvari
20
21
22 434 B, Giraudeau M, Stepp T. 2021. Linking pollution and cancer in aquatic environments: A
23
24 435 review. *Environment International* 149: 106391.

25
26
27 436
28
29 437 Bakke MJ, Nahrgang J, Ingebrigtsen K. 2016. Comparative absorption and tissue distribution
30
31 438 of ¹⁴C-benzo(a)pyrene and ¹⁴C-phenanthrene in the polar cod (*Boreogadus saida*) following
32
33
34 439 oral administration. *Polar biology*, 39: 165–1173.

35
36
37 440
38
39 441 Baršienė J, Dedonytė V, Rybakovas A, Andreikėnaitė L, Andersen OK. 2006. Investigation of
40
41 442 micronuclei and other nuclear abnormalities in peripheral blood and kidney of marine fish
42
43
44 443 treated with crude oil. *Aquatic Toxicology*, 78: S99-S104

45
46 444
47
48
49 445 Baršienė J, Lazutka J, Šyvokienė J, Dedonytė V, Rybakovas A Bagdonas E, Bjornstad A,
50
51 446 Andersen OK. 2004. Analysis of micronuclei in blue mussels and fish from the Baltic and
52
53
54 447 North Seas. *Environmental Toxicology*, 19: 365-371.

55
56 448
57
58
59
60
61
62
63
64
65

449 Baumann, PC, Harshbarger JC. 1998. Long term trends in liver neoplasm epizootics of brown
1 bullhead in the Black River, Ohio. Environmental Monitoring and Assessment 53: 213-223.
2
3
4
5 451
6
7 452 Bender ML, Frantzen M, Vieweg I, Falk-Petersen IB, Johnsen HK, Rudolfsen G, Tollefsen
8
9 453 KE, Dubourg P, Nahrgang J. 2016. Effects of chronic dietary petroleum exposure on
10 reproductive development in polar cod (*Boreogadus saida*). Aquatic Toxicology, 180: 196–
11 454 208.
12
13
14 455
15
16
17 456
18
19 457 Black JJ, Maccubbin AE and Martin M. 1985. A Reliable, Efficient, Microinjection Apparatus
20 and Methodology for the In Vivo Exposure of Rainbow Trout and Salmon Embryos to
21 458 Chemical Carcinogen. Journal of the National Cancer Institute, 75: 1123-8.
22
23
24 459
25
26
27 460
28
29 461 Bolognesi C, Hayashi M. 2011. Micronucleus assay in aquatic animals. Mutagenesis, 26: 205–
30 462 213.
31
32
33
34 463
35
36 464 Chen W, Zhang H and Liu Y. 2019. Anti-Inflammatory and Apoptotic Signaling Effect of
37 Fucoxanthin on Benzo(A)Pyrene-Induced Lung Cancer in Mice. Journal of Environmental
38 465 Pathology, Toxicology and Oncology, 38: 239-251.
39
40
41 466
42
43
44 467
45
46 468 Colli-Dula RC, Fang X, Moraga-Amador D, Albornoz-Abud N, Zamora-Bustillos R, Conesa
47 A, Zapata-Perez O, Moreno D, Hernandez-Nuñez E. 2018. Transcriptome analysis reveals
48 469 novel insights into the response of low-dose benzo(a)pyrene exposure in male tilapia. Aquatic
49 470 Toxicology, 162-173.
50
51
52
53 471
54
55
56 472
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

473 Depledge M.H., Hopkin S.P. 1995. Methods to assess effects on brackish, estuarine, and near-
474 coastal water organisms. Wiley (Ed.), SCOPE 53-Methods to Assess the Effects of Chemicals
475 on Ecosystems, the Scientific Committee On Problems of the Environment (SCOPE).

476
477 Du Corbier, FA, Stentiford GD, Lyons BP, Rotchell JM. 2005. Isolation of the retinoblastoma
478 cDNA from the marine flatfish dab (*Limanda limanda*) and evidence of mutational alterations
479 in liver tumors. *Environmental Science and Technology*, 39: 9785-9790.

480
481 Elias, 2018. Conservation issues: Tundra ecosystems. *Encyclopedia of the Anthropocene*,
482 2, 2018,165-175

483
484 EPA/635/R-17/003Fa. Toxicological Review of Benzo[a]pyrene. 2017. U.S. Environmental
485 Protection Agency Washington, DC.

486
487 Feist SW. 2004. Biological effects of contaminants: use of liver pathology of the European
488 flatfish dab (*Limanda limanda L.*) and flounder (*Platichthys flesus L.*) for monitoring
489 (International Council for the Exploration of the Sea, Copenhagen).

490
491 Geraudie P, Nahrgang J, Forget-Leray J, Minier C, Camus L. 2014. In vivo effects of en
492 vironmental concentrations of produced water on the reproductive function of polar cod
493 (*Boreogadus saida*). *Journal of Toxicology and Environmental Health A*, 77: 557–573.

494
495 Gillet LCJ and Scharer OD. 2006. Molecular mechanisms of mammalian global genome
496 nucleotide excision repair. *Chemical Reviews*, 106: 253–276.

497 Gorbi S., Benedetti M., Virno Lamberti C., Pisanelli B., Moltedo G., Regoli F. 2009. Biological
1 effects of diethylene glycol (DEG) and produced waters (PWs) released from offshore
2 498 activities: A multi-biomarker approach with the sea bass *Dicentrarchus labrax*. Environmental
3
4 499 Pollution, 157: 3166-3173.
5
6 500
7
8
9 501
10
11 502 Gravato C, Santos MA. 2002. Juvenile sea bass liver P450, EROD induction, and erythrocytic
12
13 503 genotoxic responses to PAH and PAH-like compounds. Ecotoxicology and Environmental
14
15 504 Safety, 51: 115-127
16
17 505
18
19 506 Harshbarger JC, Clark JB. 1990. Epizootiology of neoplasms in bony fish of North America.
20
21 507 Science of the Total Environment, 94: 1-32.
22
23
24 508
25
26 509 Hendricks JD, Meyers TR, Shelton DW, Casteel JL, Bailey GS. 1985. Hepatocarcinogenicity
27
28 510 of benzo[a]pyrene to rainbow trout by dietary exposure and intraperitoneal injection. Journal
29
30 511 of national cancer institute, 74 : 839-851.
31
32
33 512
34
35 513 Ho, J. 2010. The implications of Arctic sea ice decline on shipping. Marine Policy, 34 : 713–
36
37 514 715.
38
39
40 515
41
42 516 Holst, JC, McDonald, A. 2000. Fish-lift: a device for sampling live fish with trawls. Fish.
43
44 517 Research, 48 : 87–91.
45
46
47 518
48
49 519 Ingebrigtsen K, Christiansen JS, Lindhe Ö, Brandt I. 2000. Disposition and cellular binding
50
51 520 of ³H-benzo(a)pyrene at subzero temperatures: studies in an agglomerular arctic teleost fish -
52
53 521 the polar cod (*Boreogadus saida*). Polar Biology, 23: 503-509.
54
55
56
57
58
59
60
61
62
63
64
65

522

1
2 523 Islam MJ, Slater MJ, Thiele R, Kunzmann A. 2021. Influence of extreme ambient cold stress
3
4
5 524 on growth, hematological, antioxidants, and immune responses in European seabass,
6
7 525 *Dicentrarchus labrax* acclimatized at different salinities. Ecological Indicators, 122: 107280.

8
9
10 526

11 527 Jonsson H, Sundt RC, Aas E, Sanni S. 2010. The Arctic is no longer put on ice: evalua- tion of
12
13
14 528 polar cod (*Boreogadus saida*) as a monitoring species of oil pollution in cold waters. Marine
15
16
17 529 Pollution Bulletin, 60: 390–395.

18
19 530

20
21
22 531 Kammann U, Akcha F, Budzinski H, Burgeot T, Gubbins MJ, Lang T, Le Menach K, Vethaak
23
24 532 AD, Hylland K. 2017. PAH metabolites in fish bile: from the seine estuary to Iceland. Marine
25
26
27 533 Environmental Research, 124: 41–45.

28
29 534

30
31
32 535 Karle IL, Yagi H, Sayer JM, and Donald MJ. 2004. Crystal and molecular structure of a
33
34 536 benzo[a]pyrene 7,8-diol 9,10-epoxide N2-deoxyguanosine adduct: Absolute configuration and
35
36
37 537 conformation. Proceedings of the National Academy of Sciences of the United States of
38
39 538 America, 6: 1433–1438.

40
41 539

42
43
44 540 Kasala ER, Bodduluru LN, Barua CC, Sriram CS, Gogoi R. 2015. Benzo(a)pyrene induced
45
46 541 lung cancer: Role of dietary phytochemicals in chemoprevention. Pharmacological Reports.
47
48
49 542 67: 996-1009.

50
51 543

52
53 544 Köhler A, Ellesat K. 2008. Nuclear changes in blood, early liver anomalies and hepatocellular
54
55
56 545 cancers in flounder (*Platichthys flesus* L.) as prognostic indicator for a higher cancer risk?
57
58 546 Marine Environmental Research, 66 : 149-150
59
60
61
62
63
64
65

547

1
2 548 Krahn, MM, Burrows DG, Ylitalo GM, Brown DW, Wigren CA, Collier TK, Chan SL,

3
4
5 549 Varanasi U. 1992. Mass-spectrometric analysis for aromatic-compounds in bile of fish sampled

6
7 550 after the Exxon-Valdez oil-spill. *Environmental Science and Technology*, 26: 116–126.

8
9
10 551

11
12 552 Kumar A, Yadav J, Mohan R. 2021. Spatio-temporal change and variability of Barents-Kara

13
14 553 sea ice, in the Arctic: Ocean and atmospheric implications. *Science of the Total Environment*,

15
16 554 753: 142046.

17
18
19 555

20
21
22 556 Lee, JW, Kim, YH, Yoon, S, Lee SK. 2014. Cytochrome P450 System Expression and DNA

23
24 557 Adduct Formation in the Liver of *Zacco platypus* Following Waterborne Benzo(a)pyrene

25
26 558 Exposure: Implications for Biomarker Determination. *Environmental Toxicology*, 29: 1032-

27
28
29 559 1042.

30
31
32 560

33
34 561 Lerebours A, Bignell JP, Stentiford GD, Feist SW, Lyons BP, Rotchell JM. 2013. Advanced

35
36 562 diagnostics applied to fish liver tumours: relating pathology to underlying molecular aetiology.

37
38
39 563 *Marine Pollution Bulletin*, 72 : 94-98.

40
41
42 564

43
44 565 Lerebours A, Stentiford GD, Lyons BP, Bignell JP, Derocles SA, Rotchell JM. 2014. Genetic

45
46 566 alterations and cancer formation in a European flatfish at sites of different contaminant

47
48
49 567 burdens. *Environmental Science and Technology*, 48 :10448-10455.

50
51
52 568

53
54 569 Lerebours A, Chapman E, Lyons BP, Bignell JP, Stentiford GD, Rotchell JM. 2017.

55
56 570 Hepatocellular adenoma in a European flatfish (*Limanda limanda*): genetic alterations in laser-

571 capture micro-dissected tissue and global transcriptomic approach. Marine Pollution Bulletin,
1
2 572 119: 120-127.
3
4
5 573
6
7 574 Li Y, Li H, Spitsbergen JM, Gong Z. 2017. Males develop faster and more severe
8
9 575 hepatocellular carcinoma than females in krasV12 transgenic zebrafish. Scientific Reports, 7:
10
11 576 41280.
12
13
14 577
15
16 578 Li H, Li Y, Lu JW, Huo X, Gong Z. 2019. Liver-specific androgen receptor knockout attenuates
17
18 579 early liver tumor development in zebrafish. Scientific Reports, 9: 10645.
19
20
21 580
22
23 581 Lyons, BP, Thain, JE, Stentiford, GD, Hylland, K, Davies, IM, Vethaak, AD. 2010. Using
24
25 582 biological effects tools to define Good Environmental Status under the European Union Marine
26
27 583 Strategy Framework Directive. Marine Pollution Bulletin, 60 : 1647-1651.
28
29 584
30
31 585 Malins, DC, Krahn MM, Brown DW, Rhodes LD, Myers MS, McCain BB, Chan SL. 1985.
32
33 586 Toxic-Chemicals in Marine Sediment and Biota from Mukilteo, Washington - Relationships
34
35 587 with Hepatic Neoplasms and Other Hepatic-Lesions in English Sole (*Parophrys Vetulus*).
36
37 588 Journal of the National Cancer Institute, 74: 487-494.
38
39
40
41 589
42
43 590 Maria VL, Correia AC, Santos MA. 2002. *Anguilla anguilla* L. biochemical and genotoxic
44
45 591 responses to benzo[a]pyrene. Ecotoxicology and Environmental Safety, 53: 86-92.
46
47
48
49 592
50
51 593 Meier WN, Hovelsrud GK, van Oort BEH, Key JR, Kovacs KM, Michel C, Haas C, Mats A
52
53 594 G, Gerland S, Donald KP, Makshtas A, Reist JD. 2014. Arctic sea ice in transformation: A
54
55
56
57
58
59
60
61
62
63
64
65

595 review of recent observed changes and impacts on biology and human activity. Reviews of
1
2 596 Geophysics, 51: 185–217
3
4
5 597
6
7 598 Myers, MS, Landahl JT, Krahn MM, McCain BB. 1991. Relationships between hepatic
8
9 599 neoplasms and related lesions and exposure to toxic chemicals in marine fish from the U.S.
10
11
12 600 west coast. Environmental Health Perspectives, 90: 7-15.
13
14
15 601 Nahrgang J, Camus L, Gonzalez P, Goksøyr A, Christiansen J, Hop H, 2009. PAH biomarker
16
17 602 responses in polar cod (*Boreogadus saida*) exposed to benzo(a)pyrene. Aquatic Toxicology.
18
19 603 14, 309-319.
20
21
22
23 604 Nahrgang J, Camus L, Broms F, Christiansen JS, Hop H. 2010a. Seasonal baseline levels of
24
25 605 physiological and biochemical parameters in polar cod (*Boreogadus saida*): implications for
26
27 606 environmental monitoring. Marine Pollution Bulletin, 60: 1336–1345.
28
29
30
31
32 607 Nahrgang J, Camus L, Carls MG, Gonzalez P, Jonsson M, Taban IC, Bechmann RK,
33
34 608 Christiansen JS, Hop H. 2010b. Biomarker responses in polar cod (*Boreogadus saida*) exposed
35
36 609 to the water soluble fraction of crude oil. Aquatic Toxicology, 97: 234–242.
37
38
39
40
41 610 Nahrgang J, Camus L, Gonzalez P, Jonsson M, Christiansen JS, Hop H. 2010c. Biomarker
42
43 611 responses in polar cod (*Boreogadus saida*) exposed to dietary crude oil. Aquatic Toxicology,
44
45 612 96: 77–83.
46
47
48
49
50 613 Nahrgang J, Dubourg P, Frantzen M, Storch D, Dahlke F, Meador JP. 2016. Early life stages
51
52 614 of an arctic keystone species (*Boreogadus saida*) show high sensitivity to a water-soluble
53
54 615 fraction of crude oil. Environmental Pollution, 218: 605–614.
55
56
57
58
59
60
61
62
63
64
65

616 Nahrgang J, Bender ML, Meier S, Nechev J, Berge J, Frantzen M. 2019. Growth and
1 metabolism of adult polar cod (*Boreogadus saida*) in response to dietary crude oil.
2
3
4
5 618 Ecotoxicology and Environmental Safety, 180: 53–62.
6
7
8 619 Nogueira PR, Lourenço J, Mendo S, Rotchell LM. 2006. Mutation analysis of ras gene in the
9
10
11 620 liver of European eel (*Anguilla anguilla* L.) exposed to benzo[a]pyrene. Marine Pollution
12
13 621 Bulletin, 52: 1611-1616.
14
15
16 622
17
18 623 Phillips and Artlt. 2007. The ³²P-postlabeling assay for DNA adducts. Nature Protocols, 2:
19
20 624 2772–2781.
21
22
23 625
24
25 626 Ploch SA, King LC, Kohan MJ, Di Giulio RT.1998. Comparative *in Vitro* and *in Vivo*
26
27 627 Benzo[a]pyrene–DNA Adduct Formation and Its Relationship to CYP1A Activity in Two
28
29 628 Species of Ictalurid Catfish. Toxicology and applied pharmacology 149: 90–98.
30
31
32
33
34 629 Poirier et al 2019. Intestinal Polycyclic Aromatic Hydrocarbon-DNA Adducts in a Population
35
36 630 of Beluga Whales with High Levels of Gastrointestinal Cancers Environ Mol Mutagen, 60: 29–
37
38
39 631 41.
40
41 632
42
43 633 Rastogi A, Richa, Kumar A, Tyagi MB, Sinha RP. 2010. Molecular mechanisms of ultraviolet
44
45 634 radiation-induced DNA damage and repair. Journal of Nucleic Acids, 2010: 592980.
46
47
48 635
49
50
51 636
52
53 637 Reinardy KC, Dharamshi J, Jha AN, Henry TB. 2013. Changes in expression profiles of genes
54
55 638 associated with DNA repair following induction of DNA damage in larval zebrafish *Danio*
56
57 639 *rerio*. Mutagenesis, 28: 601-608.
58
59
60
61
62
63
64
65

640
1 Rei-Salgueiro L, Costa J, Ferreira M Reil-Henriques MA. 2011. Evaluation of 3-hydroxy-
2 641 benzo[*a*]pyrene levels in Nile tilapia (*Oreochromis niloticus*) after waterborne exposure to
3
4 benzo[*a*]pyrene. *Ecotoxicology*, 2040-2054.
5 642
6
7 643
8
9
10 644
11 Reichert WL, Myers MS, Peck-Miller K, French B, Anulacion BF, Collier TK, Stein JE,
12 645 Varanasi U. 1998. Molecular epizootiology of genotoxic events in marine fish: Linking
13
14 646 contaminant exposure, DNA damage, and tissue-level alterations. *Mutation Research*, 411:
15
16 647 215-225
17
18 648
19
20
21
22 649
23 Rose, WL, French, BL, Reichert, WL, Faisal, M. 2001. Persistence of benzo[*a*]pyrene–DNA
24 650 adducts in hematopoietic tissues and blood of the mummichog, *Fundulus heteroclitus*. *Aquatic*
25
26 651 *Toxicology*, 52: 319–328.
27
28 652
29
30
31 653
32 Rose, WL, French, BL, Reichert, WL, Faisal, M., 2000. DNA adducts in hematopoietic tissues
33 654 and blood of the mummichog (*Fundulus heteroclitus*) from a creosote-contaminated site in the
34
35 655 Elizabeth River, Virginia. *Marine Environmental Research*, 50: 581–589.
36
37 656
38
39
40
41 657
42 Rotchell, JM, Blair, JB, Shim, JK, Hawkins, WE, Ostrander, GK. 2001. Cloning of the
43 658 Retinoblastoma cDNA from the Japanese medaka (*Oryzias latipes*) and preliminary evidence
44
45 659 of mutational alterations in chemically-induced retinoblastomas. *Gene*, 263: 231-237.
46
47
48
49
50
51 661
52
53 662
54 Seriani R, Tavares Ranzani-Paiva M J., Silva- Souza A. T., Silva Roseli N. 2011. Hematology,
55 663 micronuclei and nuclear abnormalities in fishes from São Francisco River, Minas Gerais state,
56
57 664 Brazil. *Acta Scientiarum Biological Sciences*. 33: 107–112.
58
59
60
61
62
63
64
65

665

1
2 666 Shirmohammadi M., Mohammad N. S., Ronagh T., Movahedinia A., Hamidian G. 2018. Using
3
4
5 667 cell apoptosis, micronuclei and immune alternations as biomarkers of phenanthrene exposure
6
7 668 in yellowfin seabream (*Acanthopagrus latus*). Fish and Shellfish Immunology, 72: 37-47.
8
9

10 669

11
12 670 Song Y, Nahrgang Y, Tollefden KE. 2019. Transcriptomic analysis reveals dose-dependent
13
14 671 modes of action of benzo(a)pyrene in polar cod (*Boreogadus saida*). Science of the total
15
16 672 environment, 653: 176-189.
17
18

19 673

20
21
22 674 Stentiford, GD, Bignell, JP, Lyons, BP, Thain, JE, Feist, SW. 2010. Effect of age on liver
23
24 675 pathology and other diseases in flatfish: Implications for assessment of marine ecological
25
26 676 health status. Marine Ecology Progress Series, 411: 215-230.
27
28

29 677

30
31
32 678 Teles M, Pacheco M, Santos MA. 2003. *Anguilla anguilla* L. liver ethoxyresorufin O-
33
34 679 deethylation, Glutathione S-transferase, erythrocytic nuclear abnormalities, and endocrine
35
36 680 responses to naphthalene and β -naphthoflavone. Ecotoxicology and Environmental Safety, 55:
37
38 681 98-107
39

40 682

41
42
43
44 683 Varasini U, Nishimoto M, Reichert WL, Le Eberhart BT. 1986. Comparative Metabolism of
45
46 684 Benzo(a)pyrene and Covalent Binding to Hepatic DNA in English Sole, Starry Flounder, and
47
48 685 Rat. Cancer Research, 46: 3817-3824.
49

50 686

51
52
53 687 Varasini U, Stein JE, Nishimoto M, Reichert WL, Collier TK. 1987. Chemical Carcinogenesis
54
55 688 in Feral Fish: Uptake, Activation, and Detoxication of Organic Xenobiotics. Environmental
56
57
58 689 Health Perspectives, 71: 155-170.
59
60
61
62
63
64
65

690
1 Vethaak, AD, Rheinallt Tap. 1992. Fish disease as a monitor for marine pollution: the case of
2 691
3 the North Sea. *Reviews in Fish Biology and Fisheries*, 2: 1-32.
4
5 692
6
7 693
8 Vieweg I, Bilbao E, Meador JP, Cancio I, Bender ML, Cajaraville MP, Nahrgang J. 2018.
9 694
10 Effects of dietary crude oil exposure on molecular and physiological parameters related to lipid
11 695
12 homeostasis in polar cod (*Boreogadus saida*). *Comparative Biochemistry and Physiology, Part*
13 696
14 *C: Toxicology and Pharmacology*. 206-207, 54–64.
15
16 697
17
18 698
19 Vogelbein, WK, Fournie JW, Van Veld PA, Huggett RJ. 1990. Hepatic neoplasms in the
20 699
21 mummichog *Fundulus heteroclitus* from a creosote-contaminated site. *Cancer Research* 50:
22 700
23 5978-5986.
24
25 701
26
27 702
28 Wallace MI, Cottier FR, Berge J, Tarling GA, Griffiths C, et al. 2010. Comparison of
29 703
30 zooplankton vertical migration in an ice-free and a seasonally ice-covered Arctic fjord: An
31 704
32 insight into the influence of sea ice cover on zooplankton behavior. *Limnology and*
33 705
34 *Oceanography* 55: 831–845.
35
36 706
37
38 707
39 Wang L, Camusb AC, Donga W, Thorntona C, Willetta KL. 2010. Expression of CYP1C1 and
40 708
41 CYP1A in *Fundulus heteroclitus* during PAH-induced carcinogenesis. *Aquatic Toxicology*,
42 709
43 99: 439–447.
44
45 710
46
47 711
48
49 712
50 Welch HE, Bergmann MA, Siferd TD, Martin KA, Curtis MF, Crawford RE Conover RJ, Hop
51 713
52 H. 1992. Energy flow through the marine ecosystem of the Lancaster Sound region, Arctic
53 714
54 Canada. *Arctic* 45, 343-375.
55
56
57
58
59
60
61
62
63
64
65

715

716 Willett, K, Steinberg M, Thomsen J, Narasimhan TR, Safe S, McDonald S, Beatty K,

717 Kennicutt, MC. 1995. Exposure of killifish to benzo[a]pyrene: comparative metabolism, DNA

718 adduct formation and aryl hydrocarbon (Ah) receptor agonist activities. *Comparative*

719 *Biochemistry and Physiology. Part B: Biochemistry and Molecular Biology*, 112: 93–103.

720

721 Wills, LP, Jung D, Koehn K, Zhu S, Willett KL, Hinton DE, di Giulio RT. 2010. Comparative

722 chronic liver toxicity of benzo[a]pyrene in two populations of the atlantic killifish (*Fundulus*

723 *heteroclitus*) with different exposure histories. *Environmental Health Perspectives*, 118: 1376-

724 1381.

725 Zaja R, Popovic M, Loncar J, Smital T. 2016. Functional characterization of rainbow trout

726 (*Oncorhynchus mykiss*) Abcg2a (Bcrp) transporter. *Comparative Biochemistry and*

727 *Physiology, Part C*, 190: 15–2.

728 Zhu S, Li L, Thornton C, Carvalho P, Avery BA, Willett KL. 2008. Simultaneous

729 determination of benzo[a]pyrene and eight of its metabolites in *Fundulus heteroclitus* bile

730 using ultra-performance liquid chromatography with mass spectrometry. *Journal of*

731 *Chromatography B*, 863: 141-149.

732

733

734 **Figure and Table Legend.**

735

736 **Figure 1.** Concentrations of biliary 3-OH-benzo[a]pyrene (ng/g bile, $n = 4$ to 9) in fish exposed

737 to three treatments of BaP (acetone control, low and high) during 0, 1, 3 and 6 months. Plots

738 represent the median (line), 25–75% percentiles (box), non-outlier range (whisker), outliers

739 (circle) and extreme values (coloured triangle). The effect of the dose and time on the
740 metabolite concentrations were assessed using the Kruskal-Wallis rank test. When significant,
741 a Wilcoxon test and a Bonferroni correction were applied. Asterisks (*) show significant
742 difference from the control treatment ($p < 0.05$). Numbers above boxes represent the n .

743 **Figure 2.** Relative expression of genes (mean \pm SD, arbitrary units) in liver of polar cods ($n =$
744 10 per treatment and time) exposed to acetone control, low and high BaP treatments after 1, 3
745 and 6 months. Plots represent the median (line), 25–75% percentiles (box), non-outlier range
746 (whisker), outliers (circle) and extreme values (coloured triangle). The effect of the dose on the
747 gene expression levels was assessed using the Kruskal-Wallis rank test. No significant
748 differences ($p > 0.05$) among treatments were found. Numbers above boxes represent the n .


749 **Figure 3.** DNA damage in the form of micronuclei per thousand in the blood (A) and spleen
750 (B) and nuclear abnormalities in the blood (C) and spleen (D) of polar cod sampled after 6
751 month of exposure. Plots represent the median (line), 25–75% percentiles (box), non-outlier
752 range (whisker), outliers (circle) and extreme values (coloured triangle). The effect of the dose
753 on the number of micronuclei and nuclear abnormalities was assessed using the Kruskal-Wallis
754 rank test. When significant, a Wilcoxon test and a Bonferroni correction were applied.
755 Asterisks (*) show significant difference from the control treatment ($p < 0.05$). Numbers above
756 boxes represent the n .

757 **Table 1.** Sequences of primer pairs and FAM/TAMRA probes used in RT-qPCR reactions for
758 each of the target genes studied. *β tubulin* was used as the reference gene.

759 **Table 2.** Fulton condition (K), hepatosomatic index (HSI), gonadosomatic index (GSI), liver,
760 gonad and body weight (g), and fork length (cm) (mean \pm SD, $n = 10$) and sex ratio determined
761 after 0, 1, 3, and 6 months of exposure to different BaP treatments (acetone control, low and
762 high exposures).

763

764 Figure 1



765

766

767

768

769

770

771


772

773

774

775

776 Figure 2



777

778


779

780

781

782

783 Figure 3



784

785

786

787

788 Table 1

Gene name	Primers and Probe Sequences 5'-3'
<i>β tubulin</i>	F: GCCCGGCACCATGGA R: TGGCCGAAAACGAAGTTGTC P: TCCGGTGCTTTCGGTCAGATCTTCA
<i>XPC</i>	F: GCTTCGACTTCCATGGAGGAT R: CTTCGTGCTCCTCACACACAA P: CGCATGCTGTGACCGACGGCTAC
<i>XPD</i>	F: TCATGTTCCGGAGTCCCTTATGTT R: GGAAGTGGTCCCGGAGGTA P: ACACACAGAGCCGCATTCTGAAGGC
<i>XPF</i>	F: ATCTGGACCTGGCGAGGAA R: TCCTGCTTTGCGGGTGTT P: CTGGAGCCCGCCAACGCTACC
<i>Rad51</i>	F: AAGAAGCCGATTGGAGGAAAC R: CGCCCCTTCCTCAGGTACA P: TCATGGCCCACGCCTCCACC
<i>tp53</i>	F: CCTCTGAGGGGCATGTTCTC R: GGGGCTCTTTCTTTTTTTTGG P: TCCTGGGCGCGACCGCA

789

790

791

792

793

794

795

796

797

798

799

800

801 Table 2

	0 month	1 month			3 months			6 months		
		Control	Low	High	Control	Low	High	Control	Low	High
Fork l. (cm)	15.9 ± 1.6	16.4 ± 1	16.7 ± 1.6	16.9 ± 1.7	17.8 ± 1.6	16.8 ± 1.9	17.3 ± 1.5	16.4 ± 1.5	17.2 ± 0.7	17.3 ± 1.2
Total w. (g)	25.6 ± 7.1	29.3 ± 5.6	32.4 ± 8.8	33.2 ± 9.1	38.3 ± 11.6	32.1 ± 8.5	35.6 ± 10.4	32.7 ± 10	37.3 ± 4.5	36.7 ± 8
K	5.2 ± 0.5	5.2 ± 0.4	5.5 ± 0.5	5.4 ± 0.3	5.3 ± 0.7	5.4 ± 0.5	5.3 ± 0.4	5.2 ± 0.7	5.3 ± 0.4	5 ± 0.4
Liver w. (g)	2 ± 0.9	2.2 ± 0.7	2.5 ± 0.9	2.6 ± 0.7	2.7 ± 1.2	2.3 ± 0.6	2.8 ± 1	2.7 ± 0.9	2.6 ± 0.5	3 ± 0.9
HSI	9.6 ± 3.3	9.4 ± 2.1	9.4 ± 2.1	10.4 ± 2.8	8.6 ± 2.6	9.3 ± 2.3	9.8 ± 2.1	12 ± 3.9	9.5 ± 2	11.3 ± 2.3
Gonad w. (g)	0.3 ± 0.2	0.5 ± 0.2	0.5 ± 0.3	0.5 ± 0.3	1.4 ± 0.5	1.1 ± 0.6	1.6 ± 0.9	4.2 ± 2.3	5.6 ± 1.3	5 ± 1.8
GSI	1.5 ± 0.7	2 ± 0.5	2.2 ± 1	1.8 ± 0.6	4.6 ± 1.2	4.1 ± 1.8	5.6 ± 2.3	16.9 ± 6.8	20.8 ± 3.9	19.3 ± 6.4
Sex ratio	40	60	30	50	50	40	30	50	22	50

802

803

Supplementary material

Figure S1. Putative coding sequences used to design the primers and probe for the 6 genes studied. The pink marks indicate the location of the introns in the DNA sequence. Primers were designed to overlap the introns whenever possible to check the specificity of the qPCR reactions.

>*β tubulin*
TCTACTACAATGAGGCCTCAGGTAATTTAAAGCAATGTCAAATGGATGACGATACTTTCTCTCGTCTGTTTCAGCTCTAACTGTT
TCTAATCTATTAGGTGGCAAATACGTCCCCCGCGCTGTTCTGGTCGATCTTGAGCCCGGCACCATGGACTCTGTGAGGTCCGG
TGCTTTCCGGTCAGATCTTCAGGCCAGACAACTTCGTTTTTCGGCCAGAGTGGTGTGGCAACAACCTGGGCCAAGGGTCACTACA
CGGAAGGTGCCGAGCTGGTGGACTCTGTGCTCGACGTGGTGGAGAAAGAGGCAGAGAGCTGTGACTGCCTGCAGGGCTTCCAG
CTC

>*XPC*
ACGGGCTTCGACTTCCATGGAGGATACTCGCATGCTGTGACCGACGGCTACATTGTGTGTGAGGAGCACGAAGATTTCTCAG
AGCAGCCTGGGAGGAAGATCAAGCGCTCCAGAAACAGAAGGAGATTGAGAAGCGAGAGAAGCGGGCCACCACCAACTGGAAGC
TACTGGTGAAGGGCTTCTGATCAGGGAGAGGCTCCAGCTACGATACGGCAA

>*XPD*
CGGGGTGGTCCCTGACGGCATCGTGGCGTTCTTACCAGCTACATGTACATGGAGAACATCGTGGCGTCTGGTATGAACAGG
GAATCCTGGAGAACATCCAGAGGAACAAGCTGATCTTCATTGAGACGCCAGATGCTGCAGAGACCAGCATGGCTCTGGAGAAA
TACCAGGAGGCATGTGAGAACGGGAGAGGAGCCATCCTTCTGTCTGTGGCCCGAGGAAAAGTGTGCGAAGGAATCGATTTTGT
GCACCACTTTGGTCCGGCAGTGATCATGTTCCGGAGTCCCTTATGTTTACACACAGAGCCGATTCTGAAGCGCGCTCTGGAGT
ACCTCCGGGACCAAGTTCAGATCCGGGAGAACGACTTCCCTGACGTTTCGACGCCATGCGCCATGCGGCCAGTGCCTGGCCCGG
GTCATCAGGGGCAAGACGGACTACGGACTCATGATCTTCGCTGACAAGCGCTACGCCCGGGCGGACAAGCGGGGGAAGCTGCC
CCGCTGGATCCAGGACACATCAGCGACGGCAGCCTGAACCTCACGGTGGACGAGACGGTGCAGCTCTCCAAGCACTTCTTGA
GGCAGATGGCCAGCCCTTCCAGACAGGAGGACCAGCTGGGTCTGTCACTGCTGACGATAGAACAGCTGGAGTCAAGGAGATG
CTGAAGAAGATCAGCCAAATGGCTCACCAGGCCTGACCACAT

>*XPF*
GATCTACAAGGCCAACCGCCCCGGGAAGACGCTGCGGGTGTATTTCTGATCTATGGAGGATCCACAGAGGAACAGAAGTATC
TCACCGCCTCTCCAAGGAGAAGAAAGCCTTCGAACACCTCATCAGGGAGAAGGCGACCATGGTTGTGCCGGAGGAGCGAGAG
GGTCGAGAAGACACCAATCTGGACCTGGCGAGGAATCTGGAGCCCGCCAACGCTACCACCAACACCCGCAAGCAGGAGGCCA
GGACCAGCCAGGGAGCCCTCCCGGGTTCATCGTGGACATGCGGGAGTTCGCGAGCGAGCTGCCCTCCCTGCTGCACCGCCCGG
GGCTGGACATCGAGCCGGTCAACCTGGAGGTGGCGACTACATCCTGACGGCGGACACCTGCGTGGAGCGCAAGAGCGTGAGC
GACCTGATCGGCTCGCTGCAGAGCGGCCCTTACACGCAGTGCCTGTCCATGACGCGCTACTACAAGCGGGCCGCTGCTGCT
CATCGAGTTCGACCCGGCCAAGCCCTTCTCGCTGGTGGCGCGCTCCGAGTTCGCGCACGAGCTGTGCGCCAACGACGTCACGT
CCAAGCTGACGCTGCTCACCTGCACCTTCCCGCGCCTGCGCCTCCTTGGTGGCCCTCGCCCCACGCCACGGCCGAGCTCTTC
GAGGAGATGAAGCGGGGCCGCGGCGAGCCGCGCCCGCCGCGCAGGCCATCGCGGCCGAGTTCGGACGCCAGGACGACCGGC
GGAGCTGTACAACCCGGCGCCGTACGACTTCTGCTGAAGATGCCGGGGTCAACGCCAAGAAGCTGCGGGGCGCTGGTGAGCA
AGGCGGACAGCCTGGCCGCGCTGGCCGAGTTCAGCCAGGAGAGGCTGGCGCAGGTCTGGGGCACACCCGCAACGCCAAGATG
CTCTACGAGTTCCTGCACAACGTGGCCGACGTGCCCGGCCAGCTGCCAAGGGCAGACGGACGTGAAGGGAAAGACT


>*rad51*
ATGGCTATGAGGAGTGAAGTGCCTTTGGAGGAGGAGGTAGAGGTGGAGGAGAACCTCGGCCCTCAGCCATCAGTTCGACTGGA
GCAAAGCGGTGTGAGCAGCAGTGACCTGAAGAAGCTGGAGGAGGCGGGCTTCCACACCATCGAGGCCGTGGCTACACCCCCA
AGAAAGAGCTGCTCCACATCAAGGGCATCAGCGAGGCCAAGGCCGACAAGATCCTGGCGGAGGCAGCCAAGCTGGTGGCCATG
GGCTTACCACGGCAACCGAGTTCCACCAGCGGGCGGGCGGAGATCATCCAGATCTCCACCAGGCTCCAAGGAGCTGGACAAAAT
GTTAATGGGGGAATGGAGACGGGTTCCATCAGGAGATGTTTGGAGAGTTCGGACGGGGAAGACACAGCTGTGCCACACTC
TGGCTGTACCTGCCAGCTGCCATCGACCAGGGCGGGGAGAGGGCAAGGCCATGTACATCGACACCGAGGGAACTTCCGG
CCGGAGCGCCTGTGGCCGTGGCCGAGAGGTACGGGCTGGTGGCAGCGACGTCTGGACAACGTGGCGTACGCCCGGGCCTT
CAACACCGACCACCAGACCAGCTGCTGTACCAGGCTCCGCCATGATGGCAGAGTACGCTATGCCCTGCTCATCGTGGACA
GCGCCACCGCTCTGTACCGACGGACTACTCCGGTCCGGGGGAGCTGTGCGCCCGCAGGGCCACCTCGGCCCTTCTCCCG


ATGCTGCTGCGGCTGGCCGACGACTTGGCGTTGCCGTGGTGTATACCAACCAGGTGGTGGCCAGGTGGACGGGGCGGCCAT
GTTCTCGGCCGACCCCAAGAAGCCGATTGGAGGAAACATCATGCCCACGCCCTCCACCACGCGCTGTACCTGAGGAAGGGC
GGGGAGAGACCCGGATCTGTAAGATCTACGACTCCCCCTGCCCTCCCGGAGTCGGAGGCCATGTTCCGCCATCAACGCCGACGGC
GTGGCGACGCCAAGGACTGA






>p53

```
1 CGACGCTCCTTCTGAACTACATGTGCAACAGCTCCTGCATGGGAGGGATGAACCGGAGAGCCATCCTGACCATCCTGACCCCTG  
2 GAGTCCTCTGAGGGGCATGTTCTCGGGCGGGCGTTGCTTCGAGGTGCGCGTCTGTGCCTGTCTGGGGCGCGACCGCAAGACGGA  
3 GGAGGGCAACGTTGGAGAAAAAGACGGAGGGATCCAAGCCCACAAAAAAGAAAGAGCCCCCCTCAGGCCCCACGGCT  
4 CCGCCCAAGAGGGTCTGTCCGCTCCAGCGCTGAAGAGGAGGATAAAGGAGGTGTTTGTGCTACAGGTCGTTGGCCCGAAGAG  
5 ATTCGAGATCCTGAGGCAGATAAACGATGCACTCGCGCTGCAGGAGAGGATGACAGTCAAGCAGGAGGTCCAAGGAGGGCCGT  
6 CGCGGGGAAAGAGACGGCTGGGGGACCGGACAGACGAGGGGACCGACTGAGCGACCGTCCAACCGACCGTCCAACATAACT  
7 GCCACAGCGAACCCCGTTATTTTCTACTTTTCTTTTGTTCATTCCTATTTTATTTTTTTCATAGCTTTTTTTTCAGCATA  
8 TAGTTTTATATACAATGTATTTTTATTTTTTCATGTATTTTTTCTTTCTTGATACTAATTCTTTTTATTCGTTTTTATAAG  
9 AGGCCATG
```

10
11
12
13
14
15
16 **Figure S2.** Basophilic foci of cellular alteration diagnosed in liver of a polar cod exposed to
17
18 the low BaP concentration (magnification x40).
19







Exposure time (month)


Treatment

Control


Low

High


A - blood



B - spleen



C - blood



D - spleen

