

#### UNIVERSITÀ POLITECNICA DELLE MARCHE Repository ISTITUZIONALE

Polystyrene nanoplastics in the marine mussel Mytilus galloprovincialis

This is a pre print version of the following article:

Original

Polystyrene nanoplastics in the marine mussel Mytilus galloprovincialis / Gonçalves, J M; Benedetti, M; D'Errico, G; Regoli, F; Bebianno, M J. - In: ENVIRONMENTAL POLLUTION. - ISSN 0269-7491. - STAMPA. - 333:(2023). [10.1016/j.envpol.2023.122104]

Availability:

This version is available at: 11566/318652 since: 2024-04-08T15:34:49Z

Publisher:

Published DOI:10.1016/j.envpol.2023.122104

Terms of use:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. The use of copyrighted works requires the consent of the rights' holder (author or publisher). Works made available under a Creative Commons license or a Publisher's custom-made license can be used according to the terms and conditions contained therein. See editor's website for further information and terms and conditions. This item was downloaded from IRIS Università Politecnica delle Marche (https://iris.univpm.it). When citing, please refer to the published version.

note finali coverpage

(Article begins on next page)

Polystyrene nanoplastics in the marine mussel Mytilus 1 2 galloprovincialis. J.M. Gonçalves<sup>a</sup>, M. Benedetti<sup>b,c</sup>, G. d'Errico<sup>b</sup>, F. Regoli<sup>b,c</sup>, and M.J. Bebianno<sup>a\*</sup> 3 4 <sup>a</sup>CIMA, Centre of Marine and Environmental Research\ARNET - Infrastructure Network in 5 Aquatic Research, University of Algarve, Campus de Gambelas, 8000-139 Faro, Portugal. 6 <sup>b</sup>Dipartimento di Scienze della Vita e dell'Ambiente, Università Politecnica delle Marche, Via Brecce Bianche, 7 60131 Ancona, Italy 8 <sup>°</sup>National Future Biodiversity Centre (NFBC), Palermo, Italy 9 \*Corresponding author: mbebian@ualg.pt / +351 289 244 434 10 11 Abstract

12 Concerns about plastic pollution and its toxicity towards animals and people are growing. 13 Polystyrene (PS) is a plastic polymer highly produced in Europe for packaging purposes and 14 building insulation amongst others. The marine environment is inevitably the fate of PS products, 15 whether originating from illegal dumping, waste mismanagement, or lack of treatment for the 16 removal of plastic debris from wastewater treatment plants. Nanoplastics (< 1000 nm) is the new 17 focus for plastic pollution gaining a wide array of interest. Whether primary or secondary, their 18 relatively small size permits nanoparticles to cross cellular boundaries, and consequently lead to 19 adverse toxic effects. An in vitro assay of Mytilus galloprovincialis haemocytes exposed to 10 µg/L 20 of polystyrene nanoplastics (nPS; 50 nm) for 24 h was used to test cellular viability along with the 21 median lethal concentration (LC<sub>50</sub>) of Aliivibrio fischeri bacteria to evaluate acute toxicity. Cellular 22 viability of mussel haemocytes decreased significantly after a 24 h exposure and nPS LC<sub>50</sub> range 23 from 180 and 217, µg/L. Moreover, a 28-day exposure of the marine mussel M. galloprovincialis 24 to nPS (10 µg/L; 50 nm) was carried out to evaluate neurotoxic effects and the ingestion of these 25 plastic particles in three mussel tissues (gills, digestive gland, and gonads). The ingestion of nPS 26 was time and tissue-specific, suggesting that nPS are ingested through the gills and then 27 translocated through the mussel bloodstream, to the digestive gland and gonads where the highest 28 amount of ingested nPS was found. Once nPS enters these plastic particles are transported to 29 mussels tissues, with the gonads and digestive glands being the tissue deposits for these particles. 30 Data on acetylcholinesterase inhibition and those previously obtained on a wide range of cellular 31 biomarkers were elaborated through weighted criteria providing a synthetic assessment of cellular 32 hazard from nPS.

33 34

## 1. Introduction

Concerns about plastic pollution and its toxicity towards animals and people are growing. Global plastic production (GPP) increased to 390.7 Mt in 2021 (PlasticEurope, 2022) and, among the consequences of illegal littering and dumping, waste mismanagement, as well as a low percentage of recycled plastics (8.3%) compared to fossil-based plastics (90.2%), the aquatic environment is the inevitable fate of these materials. In Europe, plastic production was equivalent
to 15% of GPP (57.2 Mt), with applications such as packaging and building and construction
composing 39.1 and 21.3% of Europe's plastic demands, respectively (PlasticEurope, 2022). In
2021, the European Union adopted a directive banning all single-use plastics, except for bottles, as
single-use plastic, and fishing gear which account for 70% of marine litter (Directive (EU)
2019/904).

45 Polystyrene (PS) is a plastic polymer highly produced for packaging purposes, building insulation, medical equipment, and toys as well as single-use plastics (PlasticEurope, 2022). Other 46 47 than being inexpensive and readily available, PS is known for its incredible dimensional stability 48 and water resistance, permitting it to remain consistent in size and shape, along with being certified 49 as safe for use in food and beverages (Block et al., 2017). In day-to-day products, for example, PS 50 can be found in disposable cutlery, food containers, plates, cups, automotive and electrical 51 components, and in household items. However, not all plastic products are recyclable, and PS 52 products are not (The Waste and Resources Action Programme, 2023). With PS products placed in 53 waste bins, waste mismanagement and illegal dumping, with the addition that wastewater treatment 54 plants (WWTP) also lack the correct technique or method for plastic debris removal, the marine 55 environment will be the endpoint for these PS particles. Once in the ocean, plastics undergo five 56 main processes—hydrolysis, mechanical/physical degradation, thermal-oxidative degradation, 57 photo-degradation, and biodegradation (Andrady, 2011) - that break down and fragment plastics 58 into smaller sizes. After much focus on microplastics (>1 and 5 mm), nanoplastics (< 1000 nm; 59 nPS) are gradually gaining priority as nano-sized plastic particles may cross cellular boundaries, 60 increasing their potential toxicity towards marine organisms (Peng et al., 2020). nPS in the marine 61 environment can be either primary or secondary. Primary nPS enter the environment already in the 62 nano-size range, and they are found in daily-use products such as cosmetics, clothing fibres, drug 63 delivery and in 3D ink printers (Bergami et al., 2016; Bessa et al., 2018; Canesi et al., 2015; Dong 64 et al., 2019; Tamminga et al., 2018; Wang et al., 2018). The breakdown and fragmentation of 65 macroplastics (> 5 mm) and microplastics (< 5 mm) generates secondary nPS. Although techniques 66 to detect and quantify nPS in the ocean remain challenging, there is evidence for the presence of 67 plastics in the nano-size range. In the North Atlantic Gyre, a nanoplastic segment was found 68 containing PS in the nano-size range, as well as polyethylene, polyvinyl chloride, and polyethylene 69 terephthalate (Halle et al., 2017); after simulation of coastal activities on a PS-cup and lid, nano-70 sized particles were liberated after 5 minutes (Ekvall et al., 2019).

The toxic effects of nPS in marine organisms have been evaluated (Auguste et al., 2020;
Capolupo et al., 2021; Ferreira et al., 2019; Gonçalves & Bebianno, 2021; Gonçalves et al., 2022;
Kihara et al., 2021), showing that nPS mediated toxicity influences cell growth, larvae
development, embryo malformation, inflammation, and also possible inactivation of photosystems
in algae (Gonçalves & Bebianno, 2021; Kögel et al., 2020). In the marine mussel *Mytilus*

76 galloprovincialis, nPS provoke oxidative stress and damage (10 µg/L; 50 nm; 21-d) (Gonçalves et 77 al., 2022), and lysosomal destabilization (1.5 - 150 ng/L; 50 nm; 21-d) (Capolupo et al., 2021). 78 Although ingestion and translocation of MPs have been evaluated, the assessment of whether nPS 79 toxicity is entirely from their ability to cross cellular boundaries, or if these nanoparticles are 80 ingested, accumulated, and translocated between tissues remains difficult. In this study, a 81 combination of *in vitro* and *in vivo* experiments was carried out to evaluate both acute toxicity and 82 chronic effects of nPS on key model species. An *in vitro* assay on *M. galloprovincialis* haemocytes 83 was performed to evaluate cellular viability after 24 h exposure to 10 µg/L of nPS (50 nm) and 84 acute toxicity was further characterized through the Microtox<sup>®</sup> bioassay (LC<sub>50</sub>). An *in vivo* 28-d 85 exposure of mussels was performed at the same concentration of nPS to evaluate both ingestion of nanoparticles in gills, digestive glands, and gonads, as well as the onset of neurotoxicity in gills. 86 87 These data were further integrated with those previously reported on a wide selection of cellular 88 biomarkers; their overall elaboration through weighted criteria was expected to provide a better 89 evaluation of the toxicological hazard of nPS.

90 91

#### 2. Materials and Methods

92

# 2.1. Polystyrene nanoplastics (nPS)

Fluoresbrite® Plain YG spherical polystyrene nanoplastics of 50 nm in size (CAS 9003-53-6) were purchased from Polysciences, Inc. (Germany). 50 nm particles packed as 2.5% aqueous suspension, with  $3.64 \times 1014$  particles/mL in ultrapure water (7732-18- 5) (CV = 15%, Excitation max. = 441 nm, Emission max. = 486 nm). Gonçalves et al. (2022) provide a full description of the characterization of nPS and demonstrate that the hydrodynamic diameter of nPS rises when distributed in FSW (852 103 nm), which suggests that the high salt content in seawater causes aggregation/agglomeration kinetics. A concentration of 10 µg/L was used for exposure assays.

100 101

#### 2.2. Acute toxicity assay

102 The solid phase Microtox® bioassay (SPT, Azur Environmental, 1998) was used to evaluate 103 polystyrene nanoplastics (nPS; 50 nm) acute toxicity in the bioluminescent marine bacterium 104 Aliivibrio fischeri according to ISO 1134-3: 2007 and ISO 21338: 2010 according to the method 105 described in Gambardella et al., 2019). All reagents and lyophilized A. fischeri bacteria (NRRL B-106 11177) were obtained from Modern Water Ltd (USA). Tests were carried out at 15°C in the 107 Microtox diluent supplied. The initial concentration of nPS used was 3724 µg/L and was 108 sequentially diluted. Acute toxicity of nPS was measured in terms of relative bioluminescence by Microtox<sup>™</sup> 500 luminometer after 5-, 15-, and 30-mins incubation. The bioluminescence inhibition 109 was determined using the Microtox®FX equipment, and data were analysed using the 110

112

111

MicrotoxOmni software. The median lethal concentration ( $LC_{50}$ ) was defined as the concentration that produced a 50% light reduction after 5-, 15-, and 30-mins of contact time for bacteria.

113

## 114 **2.3.** *In vitro* assay

115 Fifteen mussels *M. galloprovincialis* ( $60 \pm 5$  mm) were collected from an offshore aquaculture site in Lagos, Southeast Portugal (A: 37°04'200" N 8°42,800" W, B: 37°04'200" N 8°41'000" W, 116 117 C: 37°03'400" N 8°41'000" W, D: 37°03'400" N 8°42'800" W; Testa & Cunhas Ltd) and 118 transported alive to the laboratory. Mussels (2 mussels/L) were placed into a 10 L tank, with 7.5 L of seawater (S:  $36 \pm 1$ ) and left to acclimatise for 48 hours. Then, mussel haemolymphs were 119 120 extracted from the posterior adductor muscle of mussels with a sterile hypodermic syringe (1 mL) 121 (25 G needle), under aseptic conditions in a vertical laminar airflow cabinet, and kept on ice. Cell 122 extraction and incubation methods were based on the modifications of protocols developed by 123 Gómez-Mendikute & Cajaraville (2003) and Katsumiti et al. (2014). To have sufficient haemolymph for in vitro exposures, five pools of three mussels per pool were used. Firstly, to avoid 124 clumping and aggregation of cells, 10 µL of pooled haemolymph was added to an anti-aggregation 125 solution (0.2M Tris; 171mM NaCl; 24mM EDTA; 0.15% v/v HCl 1 N) (Katsumiti et al., 2014). 126 Trypan blue dye (0.4% in physiological solution; v/v) was added for cell staining in a 1:2 proportion 127 128 (cell suspension: Trypan Blue 0.4%). Using a Neubauer chamber (200 cells per specimen) and with 129 the aid of a haemocytometer and light microscopy (Compound Light Microscopy; 400x), cell 130 viability was determined by the following equation:

131

Concentration (cells 
$$mL^{-1}$$
) =  $\frac{n^{\circ} cells \times 10000}{n^{\circ} squares} \times dilution factor$ 

132 133

134 Following cellular viability, cell suspensions were diluted at a density of 2 x 10<sup>5</sup> haemocyte 135 cells/mL (in anti-aggregation solution), and 100 µL were seeded into 96-well microplates (6 replicated per pool) with cell culture media Dulbecco's Modified Eagle Medium (DMEM, pH 7.4) 136 137 and exposed to 10 µg/L of nPS. Microplates were incubated for 24 h at 18 °C. Afterwards, culture media was discarded, and the neutral red (NR) assay was applied following an adaptation from the 138 protocol described in Katsumiti et al. (2014). The culture medium from the wells was removed and 139 140 the state of the cells was verified under light microscopy (Compound Light Microscopy; 400x). 50 µL of neutral red working solution (0.4%, pH 7.3-7.4) was added to each well, including empty 141 wells for the negative control, and left to incubate for 1 hour in the dark. The microplate was then 142 143 centrifuged at 270 g (4°C, 10 mins) and the supernatant was removed and carefully washed with 144 PBS. The dye was then extracted from the viable cells in acetic acid/ethanol solution (1:100), following samples transference to U-bottom 96-well microplates and centrifuged at 270 g for 10 145

146 147

148

min, at 4 °C. The supernatants were then transferred into flat-bottom microplates and the absorbance was measured at 550 nm (Infinite M200 Pro, TECAN®).

# 149 **2.4.** *In vivo* assay

150 Mussels belonging to the species *M. galloprovincialis*, with a shell size of  $60 \pm 5$  mm were 151 procured from the Ria Formosa Lagoon, Southeast Portugal (37°00'30.6" N 7°59'39.6" W) and 152 brought alive to the laboratory. Following a four-day acclimatisation period, a total of 50 mussels 153 were placed into each 30 L tank, with 25 L of seawater in a triplicate design, to follow the ratio of 154 2 mussels per litre. Mussels were contaminated with 10 µg/L of polystyrene nanoparticles (nPS; 50 155 nm) for 28 days. Seawater was exchanged every two days and nanoparticles were re-dosed. No 156 mortality was encountered in either treatment.

Mussels were collected at 0 days, before exposure, and 3, 7, 14, 21 and 28 days after exposure.
The gills, digestive gland, and gonads of the weighted mussels were dissected and immediately
frozen in liquid nitrogen before being preserved at -80°C for further research on neurotoxicity
(AChE) in the gills, and ingestion of nPS in the three different tissues.

161 162

## 2.5. Quality control and assessment

Each tank was covered with glass lids to limit airborne pollution and glass pipettes were used to administer aeration to avoid plastic contamination. No plastic gloves, tools, or materials were utilised during tissue dissection to avoid further contamination.

166 167

# 2.6. Acetylcholinesterase activity (AChE)

168AChE activity was only assessed in the gills of *M. galloprovincialis* (n=5 per treatment and169time of exposure) following a modification of Ellman's colorimetric method (Ellman et al., 1961).170Firstly, gills were individually homogenized in 5 mL of Tris-HCl buffer (100 mM, pH 8) and 50171 $\mu$ L of Triton – X 100 (0.1%). Following centrifugation (12 000 x g, 4°C, 30 min), the supernatant172was collected, and stored at -80°C until further analysis.

For the determination of AChE activity, 5,5'-dithio-bis(2-nitrobenzoic acid) (DTNB, 0.75 mM) was added to samples and left to incubate at room temperature for 5 mins. Then, acetylthiocholine solution (ATC, 3 mM) was added to trigger the reaction, and the absorbance was read at 412 nm using a Tecan (Infinite 200 Pro) microplate reader, for 5 mins with 30-sec intervals. AChE activity is expressed as nmol ATC min<sup>-1</sup> mg protein<sup>-1</sup>.

178

**2.7. Total protein** 

AChE activity was standardized with the determination of total protein concentrations (mg
 protein g<sup>-1</sup> tissue), calculated using bovine serum albumin (BSA), as a standard, following the
 method described by Bradford (1976), adapted for the microplate reader.

183

### **2.8. Ingestion**

Following the method described by Gagné (2019), ingestion of nPS was evaluated using a 185 186 fluorescence-based methodology using the molecular rotor probe 9-(dicyanovinyl)-julolidine 187 (DCVJ). First and foremost, individual tissues (gills, digestive glands, and gonads) (n=6 per treatment, per time of exposure) of mussels were homogenized at 20% (w/v) in an ice-cold buffer 188 189 solution (50 mM NaCl, 10 mM Hepes - NaOH [pH 7.4], 1 mM EDTA, and 1mM DTT), using a 190 VWR Star-Beater (5 min, 20/s shaking, with grinding balls). Samples were centrifuged for 20 mins (2°C, 15 000 g) to isolate the cytosolic fraction, and the resulting supernatant was immediately 191 192 frozen (-80°C) until further analysis. Samples were then analysed using the spectrofluorometric 193 microplate reader (Berthold Tristar 5) with excitation at 450 nm and emission spectra ranging from 194 400 - 800 nm. The wavelength equalling nPS was 510 nm. Results are expressed as nPS  $\mu$ g/g wet 195 weight relative controls.

196 197

#### 2.9. Statistical Analysis

198 The significance of the differences between treatments and time were determined using the 199 Shapiro-Wilk test for data distribution and variance homogeneity and either parametric tests (ANOVA, followed by Tukey's posthoc test) or non-parametric equivalent tests (Kruskal-Wallis 200 201 and a two-tailed multiple comparisons test). The results were significant if p < 0.05. GraphPad 202 Prism version 9.4.1 (GraphPad Software, Inc. CA) was utilized for the statistical analysis. 203 Additionally, the relationship between treatments (not exposed and exposed to nPS) and the 204 examined tissues (gills, digestive gland, and gonads) was assessed using a Principal Component 205 Analysis (PCA) (Statistica 7.0 software (Statsoft Inc., 2005; USA).

- 206
- 207 208

# 2.10. Weight of Evidence (WOE)

In addition to results on acetylcholinesterase, data on cellular biomarkers (antioxidant enzymes: superoxide dismutase SOD, catalase CAT, glutathione peroxidase GPx, biotransformation enzymes: glutathione S-transferases GST, oxidative damage: lipid peroxidation LPO and genotoxicity: DNA damage) were also obtained from Gonçalves et al. (2022) who exposed mussels to the same exposure to nPS as described in the present work (10  $\mu$ g/L of nPS 50 nm for 21 days). The overall biomarkers results were elaborated through a quantitative Weight Of Evidence model (WOE, Sediqualsoft) that provides a synthetic hazard index based on a specifically developed algorithm and mathematical procedure (Regoli et al., 2019). For each analysed biomarker, the
magnitude of the observed variation is compared to a specific threshold, corrected for the
toxicological relevance of the biological endpoint (weight) and the statistical significance of the
difference concerning controls. In-depth descriptions of complete calculations, intricate flow charts,
the justification for weights, thresholds, and expert opinions have previously been provided by
(Regoli et al., 2019).

222

**3. Results** 

**3.1. nPS Acute Toxicity** 

The LC<sub>50</sub> ( $\mu$ g/L) of *A. fischeri* after 5, 15 and 30 minutes of exposure to nPS was 217, 196, and 180  $\mu$ g/L, respectively (Table 1).

227

225

226

#### **3.2.** *In vitro* assay

229 In *M. galloprovincialis* haemocytes, the 24 h exposure to nPS (10  $\mu$ g/L) caused a significant 230 reduction in cell viability compared to unexposed ones (p < 0.05) (Fig. 1).

231

# **3.3.** AChE activity

Significant differences in unexposed mussels were observed on days 14, 21 and 28 compared to the beginning, while the effects of nPS were particularly evident after 3 and 7 days of the exposure (p < 0.05; Fig 2). AChE activity in the gills of nPS-exposed mussels, significantly decreased after 3 days of exposure (p < 0.05), followed by a further 2-fold decrease measured between 3 and 28 days (p < 0.05).

238

#### **3.4. Ingestion**

Mussel gills presented a significant ingestion of nPS ( $1.7 \pm 0.01 \mu g/w.w.$ ), equivalent to 6.75 x 10<sup>8</sup> particles/mL (Fig. 3A) after 3 days of exposure compared to other exposure times (p < 0.05). In the digestive glands, a significant increase in nPS ingestion occurred after 14 days of exposure (p < 0.05), equalling 5.5 x 10<sup>10</sup> particles/mL (Fig. 3B). Compared to gills and digestive glands, the highest levels of nPS ingestion were found in gonads after 7 and 14 days of exposure, with 2.6 ± 0.5 and 2.0 ± 0.6 µg nPS/g w.w., representing 2.9 x 10<sup>10</sup> and 2.1 x 10<sup>10</sup> particles/mL, respectively (p < 0.05), (Fig. 3C).

247

248

# **3.5. Principal Component Analysis (PCA)**

The ingestion of nPS in the three mussel tissues (gills, digestive gland, and gonads) was described using a PCA, which was applied to all the data collected at each time point throughout the 28-d of exposure (Fig. 4). The two principal components accounted for 74.2% of the overall

- variation (PC1 = 50.37 %, PC2 = 23.82 %; Fig. 3). There is a clear separation between mussel gills 252 253 and the other mussel tissues evaluated, as well as a distinction between unexposed and exposed to 254 polystyrene nanoplastics. The ingestion in the mussel digestive gland and gonads are positively 255 related to the principal component and support the results observed in Fig. 3B and 3C. Moreover, 256 PCA implies that although gills are the first tissue to contact with the nanoplastic, this tissue does 257 not accumulate within its tissues. This suggests that the translocation of the ingested polystyrene 258 nanoplastics in mussels is distributed, once filtered by the gills, to the other tissues, the digestive 259 gland, and gonads the destination of these nano-sized particles. PCA supports that at days 7, 14 and 260 21 of exposure, ingestions of nPS are most critical for mussels, as they are positively related to the 261 1<sup>st</sup> component. Therefore, PCA confirms that ingestion of nPS in *M. galloprovincialis* is tissuedependent and time-dependent. Furthermore, PCA corroborates the different time-point increases 262 263 in nPS ingestion observed in individual tissues.
- 264
- 265

### 266 **3.6. WOE**

The overall WOE elaboration of biomarker results (Table 2) indicated an increase of the cellular hazard after 3 days (Moderate) and even more after 7 and 14 days (Major), while it returned to a Slight level at day 21. The parameters that mostly contributed to the time-course increase of hazard were the antioxidant enzymes (SOD, CAT, GPx and GST) further reflected by severe variations of oxidative damages in terms of DNA and lipid peroxidation.

272 273

### 4. Discussion

274 The haemolymph of mussels reports the functional condition of the organs it perfuses as well 275 as extensive information on the animals' overall physiological status (Digilio et al., 2016). The 276 viability of haemocyte cells in *M. galloprovincialis* haemolymph was affected by the exposure to 277 nPS (10 µg/L; 50 nm; Fig. 1). Functionalized – nPS (PS-COOH and PS-NH<sub>2</sub>), alter immune 278 parameters, such as phagocytosis, lysosome activity, ROS and NO production (Auguste et al., 2020; 279 Canesi et al., 2015). Additionally, an increase in DNA damage in *M. galloprovincialis* haemolymph 280 was caused by exposure to nPS [10  $\mu$ g/L; 50 nm; 14-d (Gonçalves et al., 2022) and 0.05 – 50 mg/L; 281  $106 \pm 10$  nm; 96 h (Brandts et al., 2018)], and a protein corona formation with PS-NH<sub>2</sub> was 282 noticeable in the haemolymph serum of *M. galloprovincialis* [50 nm; 50 µg/mL; 30 mins (Canesi 283 et al., 2015)]. The translocation of nPS from tissues to cells was evaluated, demonstrating that 284 internalization of nPS in haemocytes is size-dependent, being 50 nm nPS mostly internalized 285 compared to 100 nm and 1 µm polystyrene particles (Sendra et al., 2020). Therefore, nPS of 50 nm 286 in size cause cytotoxicity and can have more severe implications on the overall physiological status 287 of *M. galloprovincialis*. Regarding acute toxicity, no effects of microplastics were demonstrated at 288 environmentally relevant concentrations (Booth et al., 2016; Gagné, 2017; Gambardella et al., 289 2019), while exposure of the luminescent bacteria to nPS (50 nm) caused toxic effects after 5, 15 290 and 30 mins of exposure (Table 1). Gambardella et al. (2019) suggested that particles smaller than 291 1µm would be toxic as the pore size of cell walls prevents larger particles from entering (PE-MPs; 292  $1 - 500 \ \mu\text{m}$ ; 25 mg/L), thus confirming the ability of nano-sized particles to cross cellular 293 boundaries; nano-sized plastics also affected the growth of the bacterium Halomonas alkaliphile 294 [nPS; 20 mg/L; 55 nm; (Sun et al., 2018)]. With impacts on cellular viability, growth, and their 295 possible internalization of nPS, nano-sized particles certainly pose a greater risk than micro-sized 296 ones, with still unpredictable consequences on trophic transfer and impacts on the whole ecosystem.

297 In the gills of mussels, neurotoxicity occurred after 3 days of exposure to 10 µg/L of nPS (50 298 nm). Despite the differences observed in controls, a decrease in AChE activity was noticeable 299 throughout the 21-d exposure. A 15 ng/L exposure of nPS (50 nm) for 21 days also caused a 300 decrease in AChE activity in the M. galloprovincialis gills (Capolupo et al., 2021), while no 301 neurotoxicity was observed in the freshwater clam Corbicula fluminea after exposure to nPS (80 302 nm; 0.1 – 5 mg/L; 96 h) (Li et al., 2020). Microfibres of polyethylene terephthalate caused a dose-303 dependent increase of AChE activity in the gills of M. galloprovincialis (PET-MFs; 100 µm; 0.0005 304 - 100 mg/L; Choi et al., 2021). Acetylcholine, a key neurotransmitter in the neurological and 305 sensory systems, is broken down by AChE to convey impulses at cholinergic synapses, essential 306 for neurotransmitter release and synaptic plasticity (Picciotto et al., 2012). Therefore, from all 307 available data, it is deducible that the smaller the plastic particles, the more neurotoxic they are 308 towards mussels, being more effective in seawater compared to freshwater.

309 This is the first data set on the ingestion of polystyrene nanoplastics (50 nm) in three tissues of 310 *M. galloprovincialis* after a 10  $\mu$ g/L exposure for 28 days, that we know of. Considering the 311 pathways of NPs intake for a filter-feeder organism like M. galloprovincialis, due to their small 312 size these particles can directly enter the organism through the gills, or their ability in crossing 313 cellular boundaries. With biological reactivity increasing with decreasing particle size (Peng et al., 314 2020), the relation between NPs ingestion and toxicity towards organisms is a crucial issue. After 315 28 days of exposure to 10 µg/L of nPS (50 nm), M. galloprovincialis gills, digestive gland and gonads presented time-specific ingestion rates. As filter-feeding organisms, the increase in nPS 316 317 ingestion by the gills after 3 days of exposure is expected, as this is the first tissue to encounter 318 contaminants in the surrounding environment. Looking at the same experimental conditions (10 µg/L nPS; 50 nm; 21-d), the gills of *M. galloprovincialis* after 3 days of exposure led to a significant 319 320 decrease of antioxidant enzymes activity until the 14<sup>th</sup> day, leading to oxidative damage (Joanna M. 321 Gonçalves et al., 2022). Results suggested that the internalization of nPS in the gills has a prolonged 322 toxic effect, although an adaptive response was observed after 21 days. In another study, the 323 presence of nPS (50 nm) was also found in *M. galloprovincialis* gills, although larger nPS (100 nm 324 and 1µm) were more abundant after 24h of exposure (Sendra et al., 2020); this was partly expected 325 as larger particles have similar sizes to food sources and are often mistaken as non or low nutritional 326 food (nPS; 30 nm; 0.1 - 0.3 g/L; 8 h) (Wegner et al., 2012).

327 The digestive gland of mussels is known as being responsible for key metabolic functions, such 328 as food intracellular uptake/digestion and as a storage for reserve substances (Faggio et al., 2018). 329 Here, the mussel's digestive gland shows a significant intake of nPS after 14 days of exposure (p < p330 0.05). This suggests that after initial ingestion of the gills at day 3, nPS translocate to the digestive 331 gland, and remains stored until the end of the experiment. Observing toxicity of nPS in the digestive 332 gland after a 21-d exposure to  $10 \,\mu g/L$  (50 nm), the 7<sup>th</sup> day was the most critical, causing inhibition 333 of CAT and GPx activity and consequently leading to oxidative damage (Joanna M. Gonçalves et 334 al., 2022). In this case, results imply that although not significant, the presence of nPS at 7-d is most 335 critical for this tissue and that the increase in ingestion after this time-point truly overwhelm the 336 mussels antioxidant defence system. Moreover, Sendra et al. (2020) found the digestive gland of 337 *M. galloprovincialis* as the common destination for nPS (50 nm; 3 and 24h), despite the difficulties 338 in observing these particles using histological assessments. Also, glitter particles (62.5 particles/L; 339 7-d) ingested by *M. galloprovincialis* accumulated in the digestive tract of mussels, whereby 340 ingested glitter increased with the decrease in particle size (Provenza et al., 2022). Findings confirm 341 that the smaller the size of the nanoplastic, the higher the accumulation of these particles in the 342 digestive system of mussels.

343 Gonads of mussels showed the highest levels of nPS among those measured in the three tissues, 344 particularly after 7 and 14 days of exposure  $(2.6 \pm 0.5 \text{ and } 2.0 \pm 0.6 \text{ } \mu\text{g nPS/g w.w., respectively; } p$ 345 < 0.05). Considering the anatomy of mussels, a hypothesis is that nPS, due to their small size and 346 ability to cross cellular boundaries, may 'diffuse' across tissues besides being transported to 347 different tissues via the haemolymph. A translocation study from cells to tissue showed that 348 internalization of nPS (50 nm, 100 nm and 1 $\mu$ m; 3 – 24 h) into haemocytes was size-dependent, 349 with nPS of 50 nm being the most internalized (Sendra et al., 2020). Therefore, with the same size 350 of particles, the observed ingestion of nPS in gonads may be due to their translocation in the 351 haemocytes, despite their ability to cross cellular boundaries should not be excluded. Looking at 352 the toxic effects of 10 µg/L of nPS pursue in M. galloprovincialis gonads after 21 days of exposure, 353 a similar pattern to the digestive gland is clear (Gonçalves & Bebianno, 2023). The most critical 354 time-point for gonads is at 3 days of exposure, as nPS exposure leads to an 18.6-fold increase in 355 oxidative damage, although the highest ingestion is seen on days 7 and 14. Considering the 356 antioxidant enzyme activities at both 7 and 14 days of exposure, CAT and GPx activity increased, 357 whereas GST activity decreased (Gonçalves & Bebianno, 2023), suggesting that with the increase 358 of ingested nPS in gonads of mussels, the organism's antioxidant defence mechanisms are hindered. 359 The gonads of mussels and their development are crucial for reproductive success and during the 360 gametogenesis cycle, lipid content has been shown to enhance with the ripening of female gonads, 361 as well as been used as an energy source during gametogenesis and embryo-larval development

(Martínez-Pita et al., 2012). More concerning, as nPS of 50 nm internalizes haemocytes, 362 363 spermatozoa and oocytes may also be at risk. In Crassostrea gigas, fertilization rate decreased after 364 gametes were exposed to nPS (50 nm; 1.5 h) (Tallec et al., 2018), and sperm mobility and 365 fertilization success were impaired, as agglomerates of nanoparticles were found attached to spermatozoa and the jelly coating of oocytes (PS-COOH and PS-NH<sub>2</sub>; 0.1 to 100 mg/L; 1, 3 and 5 366 367 h) (González-Fernández et al., 2018). In M. galloprovincialis, malformations of D-larvae were 368 found after embryos were exposed to nPS (50 nm; 10 µg/L; 48 hpf) (Auguste et al., 2021), 369 suggesting the high content of ingested nPS in mussel gonads may pose a serious threat for the 370 reproduction of these organisms.

371 In the present study, the biological significance of the results observed in this and a previous 372 study (Gonçalves et al., 2022) was summarized in specific hazard indices allowing an easier 373 qualitative and quantitative comparison of the effects of nPS at different exposure times. The 374 applied elaboration contemplates the measured endpoints' toxicological importance (weight), as 375 well as the extent and magnitude of shifts, normalised to predetermined thresholds. This model has 376 been validated in several case studies aimed to evaluate biological and environmental risk 377 assessment providing the possibility to integrate large datasets of heterogeneous data and to better 378 interpret asynchronous variations of complex pathways (Mezzelani et al., 2021; Nardi et al., 2022; 379 Pittura et al., 2018; Regoli et al., 2014). The weighted elaboration of biomarker results confirmed 380 the involvement of oxidative stress in modulating the toxicity of nanoparticles with a clear time-381 dependent trend as already outlined by the discussion of individual results (Table 2). The cellular 382 hazard was raised to Moderate after 3 days further increasing on days 7 and 14 when the observed 383 variations were summarized in a Major hazard. After this peak, a counteracting response to 384 oxidative challenge is evidenced by the return to a Slight hazard condition.

385 386

#### 5. Conclusion

387 This dataset provides evidence for the ingestion of polystyrene nanoplastics in the marine 388 mussel *M. galloprovincialis*, as well as neurotoxicity and *in vitro* effects on mussel haemolymph. 389 nPS of 50 nm cause neurotoxicity in mussel gills and reduce viable haemocytes in mussels' 390 haemolymph. Ingestion in mussels is tissue and time specific, whereby gonads present the highest 391 ingested nPS. Biochemical alterations of mussel tissues may be explained by the quantity of 392 ingested nPS, however further investigation is necessary. Moreover, the effects of ingested nPS in 393 the gonads of mussels need more insight into how it can affect gametogenesis, fertilization success, 394 and embryo-larval development. Lastly, the impacts of eating nPS-contaminated shellfish batches 395 can cause adverse effects on human health and should be considered for future analysis.

396 397

#### 6. Acknowledgements

A PhD grant (UI/BD/150758/2020), the JPI Oceans project RESPONSE (FCT JPIOCEANS/0005/2020), which was funded by FCT and the European Regional Development Fund (ERDF) (ALG-01-0145-FEDER-030922) through Portugal 2020 and Portuguese national funds, were all used to support this research. We also value the funding provided by FCT for projects LA/P/0069/2020 awarded to the Associate Laboratory ARNET and UID/00350/2020 awarded to CIMA of the University of Algarve.

- 404
- 405

### 406 **7. References**

- 407 Andrady, A. L. (2011). Microplastics in the marine environment. Marine Pollution Bulletin, 62(8),
  408 1596–1605. https://doi.org/10.1016/j.marpolbul.2011.05.030
- 409 Auguste, M., Balbi, T., Miglioli, A., Alberti, S., Prandi, S., Narizzano, R., Salis, A., Damonte, G., &
- Canesi, L. (2021). Comparison of different commercial nanopolystyrenes: Behavior in exposure
  media, effects on immune function and early larval development in the model bivalve *Mytilus*
- 412 galloprovincialis. Nanomaterials, 11(12). https://doi.org/10.3390/nano11123291
- Auguste, M., Lasa, A., Balbi, T., Pallavicini, A., Vezzulli, L., & Canesi, L. (2020). Impact of
  nanoplastics on hemolymph immune parameters and microbiota composition in *Mytilus*
- 415 *galloprovincialis*. Marine Environmental Research, 159(May), 105017.
- 416 https://doi.org/10.1016/j.marenvres.2020.105017
- 417 Azur Environmental. (1998). *Microtox Acute Toxicity Test*.
- 418 Bergami, E., Bocci, E., Vannuccini, M. L., Monopoli, M., Salvati, A., Dawson, K. A., & Corsi, I.
- 419 (2016). Nano-sized polystyrene affects feeding, behavior and physiology of brine shrimp
   420 Artemia franciscana larvae. Ecotoxicology and Environmental Safety, 123(2016), 18–25.
- 421 https://doi.org/10.1016/j.ecoenv.2015.09.021
- 422 Bessa, F., Barría, P., Neto, J. M., Frias, J. P. G. L., Otero, V., Sobral, P., & Marques, J. C. (2018).
- 423 Occurrence of microplastics in commercial fish from a natural estuarine environment. Marine
- 424Pollution Bulletin, 128(February), 575–584. https://doi.org/10.1016/j.marpolbul.2018.01.044
- 425 Block, C., Brands, B., & Gude, T. (2017). Packaging Materials: Polystyrene for food Packaging
- 426 applications. In ILSI Europe Report Series. http://europe.ilsi.org/publications
- Bradford, M. M. (1976). A rapid and sensitive method for the quantitation of microgram quantities of
  protein utilizing the principle of protein-dye binding. Analytical Biochemistry, 72(1), 248–254.
  https://doi.org/https://doi.org/10.1016/0003-2697(76)90527-3
- 430 Brandts, I., Teles, M., Gonçalves, A. P., Barreto, A., Franco-Martinez, L., Tvarijonaviciute, A.,
- 431 Martins, M. A., Soares, A. M. V. M., Tort, L., & Oliveira, M. (2018). Effects of nanoplastics on
- 432 *Mytilus galloprovincialis* after individual and combined exposure with carbamazepine. Science
- 433 of the Total Environment, 643, 775–784. https://doi.org/10.1016/j.scitotenv.2018.06.257

- 434 Canesi, L., Ciacci, C., Bergami, E., Monopoli, M. P., Dawson, K. A., Papa, S., Canonico, B., & Corsi,
- 435 I. (2015). Evidence for immunomodulation and apoptotic processes induced by cationic
  436 polystyrene nanoparticles in the hemocytes of the marine bivalve *Mytilus*. Marine
- 437 Environmental Research, 111, 34–40. https://doi.org/10.1016/j.marenvres.2015.06.008
- 438 Capolupo, M., Valbonesi, P., & Fabbri, E. (2021). A comparative assessment of the chronic effects of
- 439 micro-and nano-plastics on the physiology of the mediterranean mussel *Mytilus*
- 440 galloprovincialis. Nanomaterials, 11(3), 1–17. https://doi.org/10.3390/nano11030649
- 441 Choi, J. S., Kim, K., Hong, S. H., Park, K. Il, & Park, J. W. (2021). Impact of polyethylene
- terephthalate microfiber length on cellular responses in the Mediterranean mussel *Mytilus*
- 443 galloprovincialis. Marine Environmental Research, 168(March), 105320.
- 444 https://doi.org/10.1016/j.marenvres.2021.105320
- 445 Digilio, G., Sforzini, S., Cassino, C., Robotti, E., Oliveri, C., Marengo, E., Musso, D., Osella, D., &
  446 Viarengo, A. (2016). Haemolymph from *Mytilus galloprovincialis*: Response to copper and
- 447 temperature challenges studied by 1H-NMR metabonomics. Comparative Biochemistry and
- 448 Physiology Part C: Toxicology and Pharmacology, 183–184, 61–71.
- 449 https://doi.org/10.1016/j.cbpc.2016.02.003
- Dong, Z., Zhang, W., Qiu, Y., Yang, Z., Wang, J., & Zhang, Y. (2019). Cotransport of nanoplastics
  (NPs) with fullerene (C60) in saturated sand: Effect of NPs/C60 ratio and seawater salinity.
- 452 Water Research, 148, 469–478. https://doi.org/10.1016/j.watres.2018.10.071
- 453 Ekvall, M. T., Lundqvist, M., Kelpsiene, E., Šileikis, E., Gunnarsson, S. B., & Cedervall, T. (2019).
- 454 Nanoplastics formed during the mechanical breakdown of daily-use polystyrene products.

455 Nanoscale Advances, 1(3), 1055–1061. https://doi.org/10.1039/c8na00210j

- 456 Ellman, G. L., Courtney, K. D., Andres, V., & Featherstone, R. M. (1961). A new and rapid
- 457 colorimetric determination of acetylcholinesterase activity. Biochemical Pharmacology, 7(2),
  458 88–95. https://doi.org/10.1016/0006-2952(61)90145-9
- 459 Faggio, C., Tsarpali, V., & Dailianis, S. (2018). Mussel digestive gland as a model tissue for assessing
  460 xenobiotics: An overview. Science of the Total Environment, 636, 220–229.
- 461 https://doi.org/10.1016/j.scitotenv.2018.04.264
- 462 Ferreira, I., Venâncio, C., Lopes, I., & Oliveira, M. (2019). Nanoplastics and marine organisms: What
  463 has been studied? Environmental Toxicology and Pharmacology, 67(January), 1–7.
- 464 https://doi.org/10.1016/j.etap.2019.01.006
- Gagné, F. (2019). Detection of polystyrene nanoplastics in biological tissues with a fluorescent
  molecular rotor probe. Journal of Xenobiotics, 9, 7–9. https://doi.org/10.4081/xeno.2019.8147
- 467 Gambardella, C., Piazza, V., Albentosa, M., Bebianno, M. J., Cardoso, C., Faimali, M., Garaventa, F.,
- 468 Garrido, S., González, S., Pérez, S., Sendra, M., & Beiras, R. (2019). Microplastics do not affect
- standard ecotoxicological endpoints in marine unicellular organisms. Marine Pollution Bulletin,
- 470 143(February), 140–143. https://doi.org/10.1016/j.marpolbul.2019.04.055

- 471 Gómez-Mendikute, A., & Cajaraville, M. P. (2003). Comparative effects of cadmium, copper,
- 472 paraquat and benzo[a]pyrene on the actin cytoskeleton and production of reactive oxygen
- 473 species (ROS) in mussel haemocytes. Toxicology in Vitro, 17(5–6), 539–546.

474 https://doi.org/10.1016/S0887-2333(03)00093-6

- 475 Gonçalves, J. M., & Bebianno, M. J. (2023). Ecotoxicity of emerging contaminants in the
- 476 reproductive organ of marine mussels *Mytilus galloprovincialis*. Science of The Total
- 477 Environment, 163486. https://doi.org/https://doi.org/10.1016/j.scitotenv.2023.163486
- Gonçalves, J. M., & Bebianno, M. J. (2021). Nanoplastics impact on marine biota: A review.
  Environmental Pollution, 273, 116426. https://doi.org/10.1016/j.envpol.2021.116426
- Gonçalves, J. M., Sousa, V. S., Teixeira, M. R., & Bebianno, M. J. (2022). Chronic toxicity of
  polystyrene nanoparticles in the marine mussel *Mytilus galloprovincialis*. Chemosphere,
- 482 287(January 2022), 132356. https://doi.org/10.1016/j.chemosphere.2021.132356
- 483 González-Fernández, C., Tallec, K., Le Goïc, N., Lambert, C., Soudant, P., Huvet, A., Suquet, M.,
- Berchel, M., & Paul-Pont, I. (2018). Cellular responses of Pacific oyster (*Crassostrea gigas*)
  gametes exposed in vitro to polystyrene nanoparticles. Chemosphere, 208, 764–772.
- 486 https://doi.org/10.1016/j.chemosphere.2018.06.039
- Halle, A. Ter, Jeanneau, L., Martignac, M., Jardé, E., Pedrono, B., Brach, L., & Gigault, J. (2017).
  Nanoplastic in the North Atlantic Subtropical Gyre. Environmental Science and Technology,
  51(23), 13689–13697. https://doi.org/10.1021/acs.est.7b03667
- 490 Katsumiti, A., Gilliland, D., Arostegui, I., & Cajaraville, M. P. (2014). Cytotoxicity and cellular
- 491 mechanisms involved in the toxicity of CdS quantum dots in hemocytes and gill cells of the
- 492 mussel *Mytilus galloprovincialis*. Aquatic Toxicology, 153, 39–52.
- 493 https://doi.org/10.1016/j.aquatox.2014.02.003
- Kihara, S., Köper, I., Mata, J. P., & McGillivray, D. J. (2021). Reviewing nanoplastic toxicology: It's
  an interface problem. Advances in Colloid and Interface Science, 288, 102337.
- 496 https://doi.org/10.1016/j.cis.2020.102337
- Kögel, T., Bjorøy, Ø., Toto, B., Bienfait, A. M., & Sanden, M. (2020). Micro- and nanoplastic
  toxicity on aquatic life: Determining factors. Science of the Total Environment, 709(5817),
  136050. https://doi.org/10.1016/j.scitotenv.2019.136050
- 500 Li, Z., Feng, C., Wu, Y., & Guo, X. (2020). Impacts of nanoplastics on bivalve: Fluorescence tracing
- 501 of organ accumulation, oxidative stress and damage. Journal of Hazardous Materials,
- 502 392(February), 122418. https://doi.org/10.1016/j.jhazmat.2020.122418
- 503 Martínez-Pita, I., Sánchez-Lazo, C., Ruíz-Jarabo, I., Herrera, M., & Mancera, J. M. (2012).
- 504 Biochemical composition, lipid classes, fatty acids and sexual hormones in the mussel *Mytilus*
- 505 *galloprovincialis* from cultivated populations in south Spain. Aquaculture, 358–359, 274–283.
- 506 https://doi.org/10.1016/j.aquaculture.2012.06.003
- 507 Mezzelani, M., Nardi, A., Bernardini, I., Milan, M., Peruzza, L., d'Errico, G., Fattorini, D., Gorbi, S.,

- 508 Patarnello, T., & Regoli, F. (2021). Environmental pharmaceuticals and climate change: The
- 509 case study of carbamazepine in *M. galloprovincialis* under ocean acidification scenario.
- 510 Environment International, 146, 106269. https://doi.org/10.1016/j.envint.2020.106269
- 511 Nardi, A., Mezzelani, M., Costa, S., d'Errico, G., Benedetti, M., Gorbi, S., Freitas, R., & Regoli, F.
- 512 (2022). Marine heatwaves hamper neuro-immune and oxidative tolerance toward carbamazepine
- 513 in *Mytilus galloprovincialis*. Environmental Pollution, 300(February), 118970.
- 514 https://doi.org/10.1016/j.envpol.2022.118970
- Peng, L., Fu, D., Qi, H., Lan, C. Q., Yu, H., & Ge, C. (2020). Micro- and nano-plastics in marine
  environment: Source, distribution and threats A review. Science of the Total Environment,
  698, 134254. https://doi.org/10.1016/j.scitotenv.2019.134254
- 518 Picciotto, M. R., Higley, M. J., & Mineur, Y. S. (2012). Acetylcholine as a Neuromodulator:
- 519 Cholinergic Signaling Shapes Nervous System Function and Behavior. Neuron, 76(1), 116–129.
  520 https://doi.org/10.1016/j.neuron.2012.08.036
- 521 Pittura, L., Avio, C. G., Giuliani, M. E., d'Errico, G., Keiter, S. H., Cormier, B., Gorbi, S., & Regoli,
- F. (2018). Microplastics as vehicles of environmental PAHs to marine organisms: Combined
  chemical and physical hazards to the mediterranean mussels, *Mytilus galloprovincialis*. Frontiers
  in Marine Science, 5(APR). https://doi.org/10.3389/fmars.2018.00103
- 525 PlasticEurope: Plastics The Facts 2022, H. org/knowledge-hub/plastics-the-fact.-2022/. (2022).
  526 *Plastics the Facts 2022. October.*
- 527 Provenza, F., Anselmi, S., Specchiulli, A., Piccardo, M., Barceló, D., Prearo, M., Pastorino, P., &
- 528 Renzi, M. (2022). Sparkling plastic: Effects of exposure to glitter on the Mediterranean mussel
- 529 *Mytilus galloprovincialis*. Environmental Toxicology and Pharmacology, 96(September).
- 530 https://doi.org/10.1016/j.etap.2022.103994
- 531 Regoli, F., d'Errico, G., Nardi, A., Mezzelani, M., Fattorini, D., Benedetti, M., Di Carlo, M.,
- Pellegrini, D., & Gorbi, S. (2019). Application of a weight of evidence approach for monitoring
  complex environmental scenarios: The case-study of off-shore platforms. Frontiers in Marine
  Science, 6(JUL), 1–15. https://doi.org/10.3389/fmars.2019.00377
- 535 Regoli, F., Pellegrini, D., Cicero, A. M., Nigro, M., Benedetti, M., Gorbi, S., Fattorini, D., D'Errico,
- 536 G., Di Carlo, M., Nardi, A., Gaion, A., Scuderi, A., Giuliani, S., Romanelli, G., Berto, D.,
- 537 Trabucco, B., Guidi, P., Bernardeschi, M., Scarcelli, V., & Frenzilli, G. (2014). A
- 538 multidisciplinary weight of evidence approach for environmental risk assessment at the Costa
- 539 Concordia wreck: Integrative indices from Mussel Watch. Marine Environmental Research, 96,
- 540 92–104. https://doi.org/10.1016/j.marenvres.2013.09.016
- 541 Sendra, M., Saco, A., Yeste, M. P., Romero, A., Novoa, B., & Figueras, A. (2020). Nanoplastics:
- 542 From tissue accumulation to cell translocation into *Mytilus galloprovincialis* hemocytes.
- resilience of immune cells exposed to nanoplastics and nanoplastics plus *Vibrio splendidus*
- combination. Journal of Hazardous Materials, 388(November 2019), 121788.

545 https://doi.org/10.1016/j.jhazmat.2019.121788

Sun, X., Chen, B., Li, Q., Liu, N., Xia, B., Zhu, L., & Qu, K. (2018). Toxicities of polystyrene nanoand microplastics toward marine bacterium *Halomonas alkaliphila*. Science of the Total

```
548 Environment, 642, 1378–1385. https://doi.org/10.1016/j.scitotenv.2018.06.141
```

- 549 Tallec, K., Huvet, A., Di Poi, C., González-Fernández, C., Lambert, C., Petton, B., Le Goïc, N.,
- 550 Berchel, M., Soudant, P., & Paul-Pont, I. (2018). Nanoplastics impaired oyster free living stages,
- gametes and embryos. Environmental Pollution, 242, 1226–1235.
- 552 https://doi.org/10.1016/j.envpol.2018.08.020
- Tamminga, M., Hengstmann, E., & Fischer, E. K. (2018). Microplastic analysis in the South Funen
  Archipelago, Baltic Sea, implementing manta trawling and bulk sampling. Marine Pollution
  Bulletin, 128(October 2017), 601–608. https://doi.org/10.1016/j.marpolbul.2018.01.066
- The Waste and Resources Action Programme. (2023). Recycle. https://www.recyclenow.com/recyclean-item/polystyrene
- Wang, W., Yuan, W., Chen, Y., & Wang, J. (2018). Microplastics in surface waters of Dongting Lake
  and Hong Lake, China. Science of the Total Environment, 633(41907352), 539–545.
  https://doi.org/10.1016/j.scitotenv.2018.03.211
- 561 Wegner, A., Besseling, E., Foekema, E. M., Kamermans, P., & Koelmans, A. A. (2012). Effects of
- 562 nanopolystyrene on the feeding behavior of the blue mussel (*Mytilus edulis* L.). Environmental
- 563 Toxicology and Chemistry, 31(11), 2490–2497. https://doi.org/10.1002/etc.1984

564