



UNIVERSITÀ POLITECNICA DELLE MARCHE
Repository ISTITUZIONALE

Historical DNA as a tool to genetically characterize the Mediterranean sand tiger shark (*Carcharias taurus*, Lamniformes: Odontaspidae): A species probably disappeared from this basin

This is the peer reviewed version of the following article:

Original

Historical DNA as a tool to genetically characterize the Mediterranean sand tiger shark (*Carcharias taurus*, Lamniformes: Odontaspidae): A species probably disappeared from this basin / Fioravanti, T.; Bargnesi, F.; Splendiani, A.; Giovannotti, M.; Renzi, F.; Caputo Barucchi, V.. - In: AQUATIC CONSERVATION-MARINE AND FRESHWATER ECOSYSTEMS. - ISSN 1052-7613. - ELETTRONICO. - 30:5(2020), pp. 892-902. [10.1002/aqc.3294]

Availability:

This version is available at: 11566/277654 since: 2024-04-12T16:59:51Z

Publisher:

Published

DOI:10.1002/aqc.3294

Terms of use:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. The use of copyrighted works requires the consent of the rights' holder (author or publisher). Works made available under a Creative Commons license or a Publisher's custom-made license can be used according to the terms and conditions contained therein. See editor's website for further information and terms and conditions.

This item was downloaded from IRIS Università Politecnica delle Marche (<https://iris.univpm.it>). When citing, please refer to the published version.

note finali coverpage

(Article begins on next page)

1 **Historical DNA as a tool to genetically characterize the Mediterranean sand tiger shark**
2 **(*Carcharias taurus*, Lamniformes: Odontaspidae): a species probably disappeared from this**
3 **basin**

4

5 Tatiana Fioravanti¹, Filippo Bargnesi^{1,2}, Andrea Splendiani¹, Massimo Giovannotti¹, Francesca
6 Renzi¹, Vincenzo Caputo Barucchi^{1,*}

7

8 ¹ Dipartimento di Scienze della Vita e dell'Ambiente (DiSVA), Università Politecnica delle
9 Marche, Via Brece Bianche, 60131, Ancona, Italy

10 ² Acquario di Cattolica, Parco Le Navi Soc. Coop., Piazzale delle Nazioni 1, 47841, Cattolica (RN),
11 Italy

12 * Corresponding author: Dipartimento di Scienze della Vita e dell'Ambiente (DiSVA), Università
13 Politecnica delle Marche, Via Brece Bianche, 60131, Ancona, Italy. E-mail: v.caputo@univpm.it

14

15

16

17

18

19

20

21

22

23

24

25 **ABSTRACT**

- 26 1. The sand tiger shark (*Carcharias taurus*) is a coastal species distributed in temperate and sub-
27 tropical waters, classified as “Vulnerable” at global level and “Critically endangered” in Eastern
28 Australia, Southwestern Atlantic Ocean and Mediterranean Sea. Six populations (Northwestern
29 Atlantic, Brazil, South Africa, Japan, Eastern Australia and Western Australia) with low genetic
30 diversity and limited gene flow were identified worldwide, but genetic information for many
31 other geographic areas are still missing. Specifically, this species is listed in several reports as
32 part of the Mediterranean fauna, even if there is a lack of catches and sightings in recent years in
33 this basin. In order to clarify the origin of *C. taurus* individuals caught in the past in the
34 Mediterranean Sea, historical samples were genetically analysed.
- 35 2. Nine samples with a certain Mediterranean origin were collected from different European
36 museums. Genomic DNA was extracted and ~ 600 bp of the mitochondrial DNA control region
37 was amplified using eight overlapping species-specific primer pairs. Sequences obtained were
38 aligned with all the haplotypes globally known so far.
- 39 3. Genetic analysis revealed the misidentification of one museum specimen. Among the remaining
40 Mediterranean historical samples, three different haplotypes were recovered. Two of them
41 previously observed only in South Africa and one described in both South African and Brazilian
42 populations.
- 43 4. Results suggest a genetic relationship between Mediterranean sand tiger sharks and those from
44 the Western Indian Ocean. According to previous studies, we hypothesized that during the
45 Pleistocene the cold Benguela upwelling barrier was temporarily reduced allowing the passage
46 of *C. taurus* individuals from the Indian to Atlantic Ocean. After the restoration of this
47 phylogeographic barrier some individuals were trapped in the Atlantic Ocean and probably
48 migrated northward colonizing the Western African coasts and the Mediterranean Sea.

49

50 **KEYWORDS**

51 ancient DNA, *Carcharias taurus*, endangered species, genetics

52

53

54 1. INTRODUCTION

55 The sand tiger shark (*Carcharias taurus* Rafinesque, 1810) is a lamniform shark characterized by a
56 burly body and protruding teeth. It can be found in coastal temperate and sub-tropical areas, except
57 in the Eastern Pacific Ocean, usually swimming in shallow waters close to sandy or rocky bottoms
58 or submerged reefs (Compagno, 2001). Tracking and tagging studies have demonstrated that,
59 despite the presence of some differences depending on the geographic area examined, this is a
60 phylopatric species and undertakes north-south seasonal migrations (Lucifora *et al.*, 2002; Dicken *et*
61 *al.*, 2007; Bansemer and Bennett, 2011; Kneebone *et al.*, 2014; Teter *et al.*, 2015; Haulsee *et al.*,
62 2018). *C. taurus* reaches sexual maturity at the age of six-seven years in males and nine-ten years in
63 females (Goldman *et al.*, 2006). Gestation lasts between nine-twelve months and, together with
64 intra-uterine cannibalism, leads to the birth of only two newborns every two years (Gilmore, 1993).
65 As for many other sharks, the features of its life cycle (i.e. late sexual maturity, long gestation, low
66 fecundity) make it extremely prone to the risk of extinction (García *et al.*, 2008). This risk is
67 exacerbated by the drastic population decline observed in some areas as a direct consequence of
68 coastal habitat degradation and overexploitation, due to by-catch and intentional fisheries (Pollard
69 *et al.*, 1996; Otway *et al.*, 2004). For these reasons, in 2000, the IUCN classified the sand tiger
70 shark as “Vulnerable” at global level (Pollard and Smith, 2000) and it is currently considered
71 “Critically endangered” in Eastern Australia (Pollard, Gordon, Williams, Flaherty, & McAuley,
72 2003), Southwestern Atlantic Ocean (Chiaramonte *et al.*, 2007) and Mediterranean Sea (Walls and
73 Soldo, 2016).

74 It is well-known that a reduction in size of wild populations leads to a loss of genetic diversity
75 (Frankham, 1996), with a consequent decrease in the ability to adapt to future environmental
76 changes and an increased probability of extinction (Frankham, 2005). In this context, to shed light
77 on the conservation status of threatened sharks, such as *C. taurus*, genetic population analyses are
78 necessary (Dudgeon *et al.*, 2012). Currently, there are a limited number of studies describing levels
79 of genetic variation and connectivity between different populations of this species. The first was
80 performed at regional scale (South Africa, Eastern and Western Australia) by Stow *et al.* (2006)
81 using AFLP loci and the mitochondrial DNA control region (mtDNA CR) as molecular markers.
82 The second one was performed at global level using a longer sequence of the mtDNA CR and six
83 microsatellite loci (Ahonen *et al.*, 2009). Low levels of genetic diversity were demonstrated,
84 probably related to historical processes rather than recent human-mediated bottleneck events (Stow
85 *et al.*, 2006; Ahonen *et al.*, 2009). In addition, a genetic structure with six distinct populations
86 corresponding to different geographic areas (Northwestern Atlantic, Brazil, South Africa, Japan,

87 Eastern Australia and Western Australia) was revealed, with a low gene flow shown only between
88 Southern Africa and Brazilian populations. These results highlighted the necessity to manage the
89 populations of this shark as distinct Evolutionary Significant Units (ESUs; Waples, 1991) for a
90 better conservation of this species (Ahonen *et al.*, 2009).

91 Unfortunately, the genetic characterization of *C. taurus* populations seems to be still incomplete
92 because for some geographic areas the current presence and abundance of this species is unknown,
93 even if their existence has been historically well documented. This is the case of the Mediterranean
94 Sea where the occurrence of this species was known in the past. Since the 1970s, records of *C.*
95 *taurus* have become more and more sporadic until they ceased in the last decade (Fergusson *et al.*,
96 2002). Some of the last catches were made in Sicily (Fergusson *et al.*, 2002), Tunisia (Quignard and
97 Capapé, 1972; Capapé *et al.*, 1976), Croatia (Lipej *et al.*, 2004) and Aegean Sea (Ismen *et al.*,
98 2009). The lack of contemporary records makes the sampling of individuals for genetic studies
99 impossible, however, the analysis of historical samples of *C. taurus* from the Mediterranean area
100 could be very useful to improve the phylogeography of this species.

101 A first attempt to extract good quality DNA from historical shark jaws and teeth, including from *C.*
102 *taurus* specimens, was made by Ahonen & Stow (2008). Two different DNA extraction methods
103 were successfully tested. As expected, a lower amplification success of historical DNA compared to
104 a contemporary one was observed. In fact, the PCR amplification of DNA from ancient samples is
105 usually difficult due to the high degradation and small concentration of DNA extracted and/or by
106 the presence of PCR inhibitors (Pääbo *et al.*, 2004). Subsequently, DNA from historical tissue and
107 jaw cartilage was analysed to confirm the previous hypothesized Indo-Pacific origin of
108 Mediterranean great white sharks (*Carcharodon carcharias* Linnaeus, 1758) (Gubili *et al.*, 2011,
109 2015). In this paper, the mtDNA CR of historical samples of Mediterranean *C. taurus* was
110 amplified and sequenced with the aim to genetically characterize sand tiger sharks observed and
111 caught in the past in the Mediterranean Sea. The Mediterranean haplotypes found were then
112 compared with haplotypes known from the literature in order to assess the presence of haplotypes
113 endemic to the Mediterranean Sea and therefore to understand if the extinction of *C. taurus* in this
114 basin may have affected the global genetic variability of the species.

115

116 **2. METHODS**

117 **2.1 Precautions to work on historical DNA**

118 Genetic analyses on ancient and historical samples are subject to a high risk of contamination by
119 exogenous DNA. In order to avoid this problem, pre- and post- PCR work phases were performed
120 in two separate laboratories located in different buildings (Pääbo *et al.*, 2004; Knapp *et al.*, 2012).
121 In particular, the pre-PCR laboratory was equipped with two hoods provided with UV lamps, the
122 first one dedicated only to DNA extraction and the second one to reagents and PCR preparation
123 (Knapp *et al.*, 2012). The entrance to the pre-PCR area was allowed only to qualified staff equipped
124 with total body coverall, laboratory shoes, safety glasses, face mask and two pairs of gloves (Knapp
125 *et al.*, 2012). All laboratory surfaces were daily cleaned with 10% bleach and wiped with ethanol
126 70%. In addition, they were UV irradiated for 20-30 min before and after every work session.
127 Laboratory equipment (micropipettes, glassware, plasticware, etc...) was exposed to UV light for
128 20-30 min before and after their use. In contrast, the post-PCR area was dedicated only to
129 thermocycling, electrophoretic analysis of amplicons on agarose gel and preparation of samples for
130 sequencing. The thermocycler placed in this area was dedicated only to the amplification of ancient
131 or historical DNA and, after each PCR cycle, was decontaminated with UV light for 30 min.
132 Moreover, each sample was analysed separated from others to avoid cross-contamination and,
133 extraction and PCR controls were always added to detect if contamination occurred during work
134 phases (Pääbo *et al.*, 2004) .

135 **2.2 Sampling and DNA extraction**

136 An overview of the ichthyological collections of the main European museums was done through on-
137 line resources and personal contact with curators in search of *Carcharias taurus* Mediterranean
138 specimens. A total of nine historical samples of *C. taurus* with a certain Mediterranean origin
139 (Table 1) were found and collected. Five samples were powder from jaw cartilage, two were pieces
140 of cartilage and two were teeth (Table 1). All samples were decontaminated prior to DNA
141 extraction to reduce the presence of exogenous DNA and inhibitors from their surface, thus
142 reducing the risk of contamination and the probability of PCR failure (Rohland and Hofreiter,
143 2007). In the case of cartilage powder, the decontamination phase was performed before sampling.
144 Specifically, the sampling area was chosen from an internal portion of the jaws and was previously
145 scratched using sandpaper, washed with bleach and then rinsed with ultrapure sterile water. When
146 the surface was perfectly dry, the cartilage powder was obtained using a drill equipped with a sterile
147 drill bit at very low speed to avoid overheating and additional damage to DNA (Rohland and
148 Hofreiter, 2007). The powder obtained was recovered in a sterile 1.5 ml microcentrifuge tube and
149 the hole produced on the jaws was closed with dental restoration paste to make them invisible for
150 museum visitors. For pieces of cartilage, the decontamination phase was the same as described

151 above for the jaw surface, while teeth were decontaminated using the protocol proposed by Rohland
152 and Hofreiter (2007) with an additional final step consisting in the exposure to UV light for 30 min
153 for each side of the tooth. After decontamination, small pieces of the root were cut using a serrated
154 blade previously washed with DNA AWAY™ Surface Decontaminant (Thermo Scientific) and UV
155 irradiated for 30 min per side. The root was chosen for DNA extraction because it was more
156 accessible than the inner part. In addition, *C. taurus* teeth do not contain a pulp cavity, that usually
157 has a higher quantity of DNA, but both the root and the inside of the tooth are made of osteodentine
158 (Whitenack *et al.*, 2010).

159 Genomic DNA was extracted using the protocol developed for ancient bones by Yang, Cannon, and
160 Saunders (2004) with some modifications. Samples were put in 4 ml of lysis buffer (0.5 M EDTA
161 pH 8.0, 0.5% SDS, 0.5 mg/ml proteinase K) and were incubated overnight at 50°C in a washing
162 bath with gentle orbital oscillation. After incubation, samples were centrifuged to facilitate the
163 deposition of undigested materials, 3 ml of supernatant were recovered and transferred on
164 Amicon Ultra-15 centrifugal filter units (MWCO 30kDa, Merck Millipore) to concentrate samples
165 up to 125 µl. Finally, the recovered volume was purified using QIAquick PCR purification kit
166 (Qiagen) and DNA was eluted in 100 µl of ultrapure sterile water.

167 **2.3 Amplification and Sanger sequencing**

168 A fragment of ~ 600 bp of the mtDNA CR (Ahonen *et al.*, 2009) was analysed in this study. In
169 order to avoid amplification problems related to the low quality and quantity of DNA extracted
170 from historical samples, eight overlapping primer pairs were designed (Table 2, Figure 1) using the
171 software Primer3Plus (Untergasser *et al.*, 2012) and the complete mtDNA genome of *C. taurus*
172 deposited in GenBank (Accession number: KF569943, Chang, Jabado, Lin, & Shao, 2015) as
173 reference sequence.

174 PCRs were performed in a 25 µl reaction volume containing 1X PCR buffer, 1.5 mM MgCl₂, 0.08
175 mM of each dNTP, 0.48 µM of each primer, 4U of Platinum Taq DNA Polymerase (Invitrogen) and
176 3 µl of genomic DNA. All amplifications were performed in a BioRad T100™ Thermal Cycler
177 (BioRad) with an initial denaturation step at 94°C for 7 min, followed by 60 cycles of 20 s at 94°C,
178 30 s at 54°C and 40 s at 72°C, with a final extension at 72°C for 7 min.

179 PCR products were checked on 2% agarose gel stained with GelRed™ (Biotium). All amplicons
180 were sent to BMR Genomics (Padua, Italy) for Sanger sequencing, purified by exoSAP-IT™

181 (Thermo Scientific) and sequenced in both directions using an automated sequencer, ABIPRISM
182 3730XL (Applied Biosystems).

183 **2.4 Alignment and data analysis**

184 For all the samples, sequences obtained using each primer pair were checked by eye and assembled
185 to have the complete sequence of interest. All historical sequences were checked with BLAST
186 (Altschul *et al.*, 1990) and aligned using CLUSTALW (Larkin *et al.*, 2007) with the 11 haplotypes
187 described so far at the global level (Ahonen *et al.*, 2009; Chang *et al.*, 2015; Wynne and Wilding,
188 2018). When necessary, the alignment was manually edited on BioEdit (Hall, 1999). For the sample
189 PA002, the very low amplification success and the lack of a correspondence after the alignment
190 with *C. taurus* sequences suggested a mislabelling of the museum specimen. For this reason, the
191 short and not contiguous sequences obtained from this sample were checked using BLAST
192 (Altschul *et al.*, 1990) and a morphological analysis of the teeth on the jaws was carried out
193 (Compagno, 2001) using pictures taken during the sampling phase.

194 Excluding the PA002 sample, evolutionary relationships between all haplotypes were shown on a
195 Median-Joining Network (Bandelt *et al.*, 1999) using Network 5 (Fluxus Technology Ltd.,
196 www.fluxus-engineering.com), considering also gaps and missing nucleotides. The ϵ parameter was
197 set to zero and information from previous studies about sampled individuals and sampling locations
198 (Ahonen *et al.*, 2009; Chang *et al.*, 2015; Wynne and Wilding, 2018) were added to the analysis.

199

200 **3. RESULTS**

201 DNA was successfully extracted and amplified from all the historical samples of *Carcharias taurus*
202 (Table 1). The complete mtDNA CR sequence of 574 bp in length, previously analysed also by
203 Ahonen *et al.* (2009), was obtained for five samples (FI002, PA001, PA003, PR004, XL002).
204 Amplification failures produced 550 bp for PA004, 507 bp for PR001, 495 bp for XL001 and only
205 198 bp for PA002. Specifically, the primer pairs CtCR2 and CtCR3 failed the amplification of the
206 samples PR001 and XL001, respectively. The sequence produced by the primer pair CtCR6 was not
207 obtained for two samples, PA004 and PR001. For PA002, only CtCR4, CtCR7 and CtCR8 provided
208 a PCR product.

209 Undoubtedly, the primer pair and the sample with the worst amplification success were CtCR6 and
210 PA002, respectively. The comparison of the short not contiguous sequences obtained from PA002

211 with those of *C. taurus* and with all the sequences deposited in data banks did not show any perfect
212 match. The morphological analysis of the jaws showed a probable misidentification of the museum
213 specimen; teeth on the museum jaws have two lateral cusplets on each side of the main cusp, a
214 characteristic of the small-tooth sand tiger shark (*Odontaspis ferox* Risso, 1810) (Compagno, 2001).
215 The lack of the complete mitochondrial genome and/or the mtDNA CR sequence of this species in
216 data banks makes the corroboration of morphological observations impossible and this sample was
217 precautionarily excluded from the following analysis.

218 All sequences obtained from historical samples have been submitted to the GenBank database under
219 accession numbers: MK434273-MK434280. The alignment of all *C. taurus* sequences known so far
220 and those obtained in this study have allowed, on the basis of 18 polymorphic sites, the
221 classification of Mediterranean historical samples into three previously described haplotypes:
222 Haplotypes A, B and I (Table 3). Of the five samples for which the complete sequence of interest
223 was obtained, four belonged to Haplotype A (FI002, PA001, PA003, PR004) and one to Haplotype
224 I (XL002) (Table 3). The affinity to a specific haplotype was also clearly defined for two of the
225 three Mediterranean incomplete sequences. Sample PA004 seems to belong to Haplotype A also in
226 absence of the diagnostic site in 356 and, sample XL001 to Haplotype B also in absence of the
227 diagnostic site 182 (Table 3). The classification of the sample PR001 was more difficult. The
228 amplification failure of the primer pair CtCR2 did not mask any known polymorphic sites (Table 3),
229 while the failure of the primer pair CtCR6 did not allow us to obtain the diagnostic site 356. This
230 latter failure prevented us from understanding if sample PR001 belonged to Haplotype B or
231 Haplotype J (Table 3).

232 The alignment result was also confirmed by the Median Joining network performed to visualize
233 haplotypes relationships (Figure 2). In addition, the inclusion of information about sampling
234 locations from other previous studies (Ahonen *et al.*, 2009; Chang *et al.*, 2015; Wynne and Wilding,
235 2018) was very useful because it showed that Mediterranean historical samples have the same
236 haplotypes as *C. taurus* individuals sampled in South Africa (Western Indian Ocean) and Brazil
237 (Western Atlantic Ocean). Specifically, five Mediterranean samples (FI002, PA001, PA003, PR004,
238 PA004) belonged to Haplotype A and one (XL002) to Haplotype I, previously observed only in
239 individuals sampled in South Africa (Figure 2). The sample XL001 was identified as Haplotype B,
240 which was found in both South Africa and Brazil (Figure 2). The Network 5 software also included
241 the PR001 sample within Haplotype B cluster, on the basis of the maximum parsimony principle
242 (Figure 2).

244 **4. DISCUSSION**

245 The sand tiger shark (*Carcharias taurus*) is considered as “Critically endangered” within the
246 Mediterranean Sea by the IUCN (Walls and Soldo, 2016). However, its presence in this basin is
247 currently uncertain due to the lack of sightings and catches over the last decade, which suggest a
248 probable extinction at regional scale (Fergusson *et al.*, 2002; Walls and Soldo, 2016). The use of
249 DNA extracted from historical samples has allowed us to genetically characterize, for the first time,
250 *C. taurus* individuals that inhabited the Mediterranean waters in the past and to suggest a possible
251 route of colonization of this basin. Only eight specimens of certain Mediterranean origin were
252 sampled and analysed. It was not possible to obtain a larger sample mainly because of the lack of
253 information about the original catch location for most museum specimens and because some
254 institutions do not allow samples to be taken from their collections.

255 MtDNA was successfully extracted from all the nine historical samples using a protocol developed
256 for ancient bones (Yang *et al.*, 2004) and, in contrast to Ahonen and Stow (2008), a higher
257 amplification success was achieved. Ahonen and Stow (2008) tried the DNA extraction and PCR
258 amplification on 34 historical samples (20-40 years old) from different shark species, including of
259 *C. taurus* (cartilage and teeth). The PCR amplification failed for 19 of them highlighting that the
260 use of a single primer pair to amplify a region of ~ 700 bp of the mtDNA CR (Stow *et al.*, 2006) is
261 unsuitable to analyse historical DNA. Instead, the use of overlapping primer pairs delimiting a
262 region of 150-200 bp in length has been able to improve the amplification success of both historical
263 and ancient DNA (Barnett *et al.*, 2014; Splendiani *et al.*, 2016, 2017; Cole *et al.*, 2018) and was
264 successful also in this study. However, for three samples, an incomplete sequence was obtained
265 probably due to the degradation of DNA extracted and/or to the presence of PCR inhibitors (Pääbo
266 *et al.*, 2004). The primer pairs with the lowest amplification success was CtCR6 because it failed
267 the amplification in two *C. taurus* samples. It was designed to amplify a sequence of 213 bp in
268 length, while it is widely known that DNA molecules extracted from ancient samples rarely exceed
269 200 bp (Pääbo *et al.*, 2004). The presence of repeated motifs and a high AT content in the region
270 encompassed by these primers have limited us in primer design. The repetition of a single base or
271 dinucleotide motifs for many times in a DNA sequence can cause the incorrect pairing of the
272 primers on the DNA template. In addition, the presence of AT rich sequences leads to primers with
273 a very low melting temperature (T_m). A low T_m is responsible for pairing of the primers even in

274 regions with several mismatches, thus leading to the amplification of aspecific PCR products
275 (Dieffenbach *et al.*, 1993).

276 Excluding the sample PA002, due to the probable misidentification of the museum specimen, all the
277 other historical jaws and teeth undoubtedly belonged to *C. taurus* individuals. The mtDNA CR
278 sequences obtained here were attributed to two different haplotypes (Haplotype A and I) previously
279 reported only for South Africa and one (Haplotype B) shared by both South Africa and Brazil
280 (Ahonen *et al.*, 2009). The incomplete sequence of PR001 could be attributed to two distinct
281 haplotypes (Haplotype B and J) however, the presence of another Haplotype B among the
282 Mediterranean historical samples (XL001) and the distribution of the Haplotype J only in Abu
283 Dhabi waters (Chang *et al.*, 2015) suggest that the sample PR001 bears Haplotype B, as indicated
284 also by the Median Joining network. The lack of new haplotypes from Mediterranean historical
285 samples was probably due to the limited number of samples analysed or to the low rate of molecular
286 evolution estimated for this species (Stow *et al.*, 2006; Ahonen *et al.*, 2009). Instead, the
287 observation of haplotypes mainly described for South African individuals suggests a genetic
288 relationship between Mediterranean sand tiger sharks and those from the Western Indian Ocean.

289 Ahonen *et al.* (2009) observed the deepest genetic divergence between the Northwest Atlantic
290 population and all the others, while the lowest divergence was identified between South Africa and
291 Brazil, which also share some haplotypes. In the first case, the major divergence was traced back to
292 the formation of the Isthmus of Panama (~ 3 million years ago), which has definitively separated
293 Atlantic and Pacific Oceans (Toonen *et al.*, 2016). On the other hand, the low differentiation
294 between South African and Brazilian populations indicates a relatively recent connection (Ahonen
295 *et al.*, 2009). The belonging of historical samples analysed here to haplotypes already described in
296 the Western Indian Ocean highlights a recent origin also in the case of the Mediterranean sand tiger
297 sharks excluding an ancient origin due to the separation between the Mediterranean Sea and the
298 Indo-Pacific Ocean by the rising of the Isthmus of Suez (11-18 million years ago) (Toonen *et al.*,
299 2016). The Mediterranean Sea was separated many years before the formation of the Isthmus of
300 Panama indicating that if the Mediterranean *C. taurus* are descendant from those trapped after the
301 raising of the Isthmus of Suez, they should have a greater genetic divergence than observed.

302 The connection between the Red and Mediterranean seas was re-established in 1876, after the
303 opening of the Suez Canal, and promoted the entry of Indo-Pacific species into the Mediterranean
304 basin, a phenomenon known as “Lessepsian migration” (Por, 1978). However, this route for
305 colonization by Lessepsian migrants of *C. taurus* is rejected as several evidences indicate that this

306 species was already present in the Mediterranean Sea before the opening of the Suez Canal: i) the
307 species was described for the first time by Rafinesque in 1810, based on an individual caught in
308 Sicilian waters (Compagno, 2001; Fergusson *et al.*, 2002), ii) other catches and sightings were
309 reported in the Mediterranean basin before the 1876 (Fergusson *et al.*, 2002) and iii) our historical
310 samples were mainly from the Western Mediterranean and the collection dates are earlier or close to
311 the date of the opening of the Suez Canal opening. A migration through the Red Sea can also be
312 hypothesized in the opposite direction (anti-Lessepsian migration), from the Mediterranean Sea to
313 the Western Indian Ocean, but anti-Lessepsian migrants are very rare (Por, 1978). In addition, the
314 low genetic diversity observed in the Mediterranean historical samples could be due to a “founder
315 effect” suggesting that the South Africa, characterized by the highest genetic diversity (Ahonen *et*
316 *al.*, 2009), was probably the origin of the Mediterranean population.

317 Thus, the most probable biogeographic way used by the sand tiger sharks to colonize the
318 Mediterranean Sea is along the Western African coasts. Ahonen *et al.* (2009) explained the low rate
319 of genetic differentiation and the gene flow observed between South African and Brazilian
320 populations by the establishment of a recent connection between Indian and Atlantic Ocean. The
321 Southwestern African coast is characterised by the presence of an upwelling zone, caused by the
322 northward flow of the cold Benguela Current, that acts as a phylogeographic barrier (Benguela
323 barrier) (Dudgeon *et al.*, 2012; Toonen *et al.*, 2016). During Pleistocene interglacial periods, the
324 northward cold Benguela current was reduced with a simultaneous expansion of the south-westward
325 warm Agulhas current (Peeters *et al.*, 2004) that seems to have promoted the passage of *C. taurus*
326 individuals from the Western Indian to Atlantic Ocean (Ahonen *et al.*, 2009). A similar pattern of
327 dispersion was also proposed to explain the genetic similarities observed for South Atlantic and
328 Indo-Pacific populations of other shark species such as *Carcharhinus limbatus* (Keeney and Heist,
329 2006), *Carcharhinus longimanus* (Camargo *et al.*, 2016) and *Carcharhinus falciformis* (Domingues
330 *et al.*, 2018).

331 A relatively recent colonization of the Mediterranean Sea by individuals of Indo-Pacific origin was
332 also suggested for the great white shark *Carcharodon carcharias* (Gubili *et al.*, 2011) and
333 confirmed by the analysis of historical samples (Gubili *et al.*, 2015). Contrary to what observed for
334 the Mediterranean sand tiger shark, the great white shark haplotypes from the Mediterranean Sea
335 were more similar to North-Eastern Pacific/Australia/New Zealand haplotypes and not to South
336 African (Western Indian Ocean) ones (Gubili *et al.*, 2011, 2015). This discrepancy is probably
337 related to the life history characteristics of the two species. Both species are characterized by natal
338 philopatry but shows a different migratory behaviour. *C. taurus* is a coastal species that usually

339 accomplish short migration, for example in the South-eastern coast of South Africa a seasonal
340 north-south migration between mating, gestating and parturition areas was observed (Dicken *et al.*,
341 2006). *C. carcharias* instead has a high migratory capacity as documented by the observation of a
342 trans-oceanic migration from South Africa to Western Australia (Bonfil *et al.*, 2005). Gubili *et al.*
343 (2011) estimated that the separation between Mediterranean and Indo-Pacific white shark
344 populations occurred during the Late Pleistocene, a period characterized by climate instability.
345 During a trans-oceanic migration some Indo-Pacific white sharks reached South Africa and,
346 following the expansion of the Agulhas current, were driven to the Eastern Atlantic Ocean. The
347 chase of prey, such as Atlantic bluefin tuna and swordfish, that showed a similar dispersion pattern
348 (Alvarado Bremer *et al.*, 2005) and the propensity to swim eastward to return to natal areas have
349 forced them within the Mediterranean Sea.

350 In the case of *C. taurus*, an immediate colonization of the Mediterranean area seems unlikely
351 because this species usually undertakes short migrations, only in one case a distance travelled of ~
352 2000 km was observed (Dicken *et al.*, 2007). We propose that South African individuals have
353 reached the Atlantic Ocean during the Pleistocene, when the cold Benguela Current was temporarily
354 attenuated and the Agulhas current enhanced. The restoration of the cold Benguela upwelling
355 barrier probably trapped some individuals of sand tiger shark along the Southeast African coasts
356 from which they migrated northward to reach warmer habitats. In fact, *C. taurus* rarely tolerates
357 temperature lower than 15°C (Lucifora *et al.*, 2002; Otway and Ellis, 2011; Smale *et al.*, 2012;
358 Kneebone *et al.*, 2014; Teter *et al.*, 2015). The coastal behaviour of this species together with the
359 propensity to accomplish north-south seasonal migrations probably allowed, following a stepping
360 stone model of dispersion, the colonization of Western African coasts and finally entry into the
361 Mediterranean basin. However, the lack of unique haplotypes among the Mediterranean historical
362 samples and the lack of genetic data for Western Atlantic Ocean do not allow us to understand if
363 Mediterranean sand tiger sharks belonged to a distinct population or if they were visitors from
364 African Atlantic coasts (Fergusson *et al.*, 2002).

365 **4.1 Conclusion**

366 The decline of chondrichthyan species recorded at global scale and in particular in the
367 Mediterranean Sea as a consequence of human activities is alarming (Ferretti *et al.*, 2008; Dulvy *et al.*
368 *et al.*, 2014). In this context, the importance of genetic tools to develop beneficial management and
369 conservation strategies has been largely demonstrated (Dudgeon *et al.*, 2012). However, the
370 difficulty in collecting shark specimens poses a serious limit to conservation genetic studies. This

371 limit can be overcome by the use of historical shark jaws and teeth that represent an alternative
372 source of DNA (Ahonen and Stow, 2008; Gubili *et al.*, 2015; Nielsen *et al.*, 2017). In this study, the
373 genetic analysis of historical samples helped us to genetically characterize Mediterranean sand tiger
374 sharks using historical DNA and to hypothesize a biogeographic scenario for the colonization of the
375 Mediterranean Sea by individuals coming from Western Indian Ocean. However, the limited
376 number of samples and the complete lack of genetic information for some geographic areas (e.g.
377 Eastern Atlantic Ocean) did not allow us to clarify if Mediterranean individuals belonged to a
378 distinct population currently extinct or if they were vagrants from the African Atlantic coast
379 (Fergusson *et al.*, 2002). The identification of previously described haplotypes among historical
380 Mediterranean samples suggests that, if a Mediterranean *C. taurus* population had been lost, there
381 would have not been a loss in terms of global genetic variability. Regarding individuals from
382 African Atlantic coasts, a conservation planning to reduce the threats for this species could allow
383 the recolonization of the Eastern Atlantic coast and probably of the Mediterranean Sea. Shark
384 species of Western Africa have long been subjected to over-exploitation by fishing activities (Diop
385 and Dossa, 2011), this could have led to the reduction of *C. taurus* populations also in this area.
386 Further studies are therefore necessary to clarify the status of the Mediterranean sand tiger shark
387 and to improve the global knowledge on this species. Following the last IUCN assessment for the
388 sand tiger shark (Walls and Soldo, 2016), trends and dynamics in the world populations of this
389 species are still unknown. Data about its distribution range and conservation status are absent or
390 incomplete for several geographic area, as observed for the Mediterranean Sea and Eastern Atlantic
391 Ocean. Fragmentation and isolation are known as factors that may weak subpopulations, and in the
392 case of a species as the sand tiger sharks such vulnerable to coastal human impact (i.e. by-catch,
393 commercial fisheries, habitat degradation), they can strengthen a declining process. Additional
394 information about the distribution range, size of populations, levels of genetic diversity and gene
395 flow between different geographic areas, also by the analysis of historical samples, must be
396 obtained. These data could favour the development of regional and inter-regional conservation
397 policies to prevent the extinction of *C. taurus* at local and global level and, if possible, to encourage
398 the recolonization of areas from which it seems to have disappeared.

399

400 **ACKNOWLEDGMENTS**

401 The authors are very grateful to: Stefano Gridelli (Acquario di Cattolica, Italy) to have promoted the
402 collaboration between the two involved institutions; Sabrina Lo Brutto and Enrico Bellia (Museo di

403 Zoologia “Pietro Doderlein”, Palermo, Italy), Olivier Pauwels and Tom Geerinckx (Royal Belgian
404 Institute of Natural Sciences, Brussels, Belgium), Bernard Seret (Muséum National d’Histoire
405 Naturelle, Paris, France), Stefano Vanni (Museo di Storia Naturale, Sezione di Zoologia “La
406 Specola”, Florence, Italy) for allowing us to collect tissue samples from museum specimens and for
407 providing us with precious information about these specimens; Heidi Ahonen (Norsk Polarinstitut,
408 Tromsø, Norway) for sending us all the sequences from Ahonen et al. (2009), which are not
409 available in GenBank. This research was financed by funds from “Università Politecnica delle
410 Marche” awarded to Vincenzo Caputo Barucchi (Ricerca Scientifica di Ateneo 2018, grant number
411 I36C18004750005 and Progetto Strategico di Ateneo, grant number 040017_R. SCIENT.
412 A_2016_CAPUTO BARUCCHI_V_STRATEGICI).

413

414 REFERENCES

- 415 Ahonen H, Stow AJ. 2008. Shark jaws and teeth: An unexploited resource for population genetic
416 studies. *Journal of Fish Biology* **73**: 450–455.
- 417 Ahonen H, Harcourt RG, Stow AJ. 2009. Nuclear and mitochondrial DNA reveals isolation of
418 imperilled grey nurse shark populations (*Carcharias taurus*). *Molecular Ecology* **18**: 4409–
419 4421.
- 420 Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic Local Alignment Search Tool.
421 *Journal of Molecular Biology* **215**: 403–410.
- 422 Alvarado Bremer JR, Viñas J, Mejuto J, Ely B, Pla C. 2005. Comparative phylogeography of
423 Atlantic bluefin tuna and swordfish: The combined effects of vicariance, secondary contact,
424 introgression, and population expansion on the regional phylogenies of two highly migratory
425 pelagic fishes. *Molecular Phylogenetics and Evolution* **36**: 169–187.
- 426 Bandelt H-J, Forster P, Röhl A. 1999. Median-joining networks for inferring intraspecific
427 phylogenies. *Molecular Biology and Evolution* **16**: 37–48.
- 428 Bansemer CS, Bennett MB. 2011. Sex- and maturity-based differences in movement and migration
429 patterns of grey nurse shark, *Carcharias taurus*, along the eastern coast of Australia. *Marine
430 and Freshwater Research* **62**: 596–606.
- 431 Barnett R, Yamaguchi N, Shapiro B, Ho SYW, Barnes I, Sabin R, Werdelin L, Cuisin J, Larson G.

- 432 2014. Revealing the maternal demographic history of *Panthera leo* using ancient DNA and a
433 spatially explicit genealogical analysis. *BMC evolutionary biology* **14**: 70.
- 434 Bonfil R, Meÿer M, Scholl MC, Johnson R, O'Brien S, Oosthuizen H, Swanson S, Kotze D,
435 Paterson M. 2005. Transoceanic migration, spatial dynamics, and population linkages of white
436 sharks. *Science* **310**: 100–103.
- 437 Camargo SM, Coelho R, Chapman D, Howey-Jordan L, Brooks EJ, Fernando D, Mendes NJ, Hazin
438 FH V., Oliveira C, Santos MN, et al. 2016. Structure and Genetic Variability of the Oceanic
439 Whitetip Shark, *Carcharhinus longimanus*, Determined Using Mitochondrial DNA. *Plos One*
440 **11**: e0155623.
- 441 Capapé C, Chaldi A, Prieto R. 1976. Les Sélaciens dangereux des côtes tunisiennes. *Archives de*
442 *l'Institut Pasteur de Tunis* **53**: 61–108.
- 443 Chang C-H, Jabado RW, Lin Y-S, Shao K-T. 2015. The complete mitochondrial genome of the
444 sand tiger shark, *Carcharias taurus* (Chondrichthyes, Odontaspidae). *Mitochondrial DNA* **26**:
445 728–729.
- 446 Chiaramonte G, Domingo A, Soto J. 2007. *Carcharias taurus* Southwest Atlantic subpopulation. *The*
447 *IUCN Red List of Threatened Species 2007* e.T63163A1.
- 448 Cole TL, Waters JM, Shepherd LD, Rawlence NJ, Joseph L, Wood JR. 2018. Ancient DNA reveals
449 that the 'extinct' Hunter Island penguin (*Tasidyptes hunteri*) is not a distinct taxon. *Zoological*
450 *Journal of the Linnean Society* **182**: 459–464.
- 451 Compagno LJV. 2001. Sharks of the world. An annotated and illustrated catalogue of shark species
452 known to date. Volume 2. Bullhead, mackerel and carpet sharks (Heterodontiformes,
453 Lamniformes and Orectolobiformes). *FAO Species Catalogue for Fishery Purposes* **2**: 269.
- 454 Dicken ML, Smale MJ, Booth AJ. 2006. Spatial and seasonal distribution patterns of the ragged-
455 tooth shark *Carcharias taurus* along the coast of South Africa. *African Journal of Marine*
456 *Science* **28**: 603–616.
- 457 Dicken ML, Booth AJ, Smale MJ, Cliff G. 2007. Spatial and seasonal distribution patterns of
458 juvenile and adult raggedtooth sharks (*Carcharias taurus*) tagged off the east coast of South
459 Africa. *Marine and Freshwater Research* **58**: 127–134.

- 460 Dieffenbach CW, Lowe TMJ, Dveksler GS. 1993. General concepts for PCR primer design.
461 *Genome Research* **3**: S30–S37.
- 462 Diop M, Dossa J. 2011. 30 years of shark fishing in West Africa: Development of fisheries, catch
463 trends, and their conservation status in Sub-Regional Fishing Commission member countries
- 464 Doderlein P. 1879. *Manuale ittiologico del Mediterraneo: ossia sinossi metodica delle varie specie*
465 *di pesci riscontrate nel Mediterraneo ed in particolare nei mari di Sicilia (Vol. 1)*. Tipografia
466 del Giornale di Sicilia: Palermo.
- 467 Domingues RR, Hilsdorf AWS, Shivji MM, Hazin FVH, Gadig OBF. 2018. Effects of the
468 Pleistocene on the mitochondrial population genetic structure and demographic history of the
469 silky shark (*Carcharhinus falciformis*) in the western Atlantic Ocean. *Reviews in Fish Biology*
470 *and Fisheries* **28**: 213–227.
- 471 Dudgeon CL, Blower DC, Broderick D, Giles JL, Holmes BJ, Kashiwagi T, Krück NC, Morgan
472 JAT, Tillett BJ, Ovenden JR. 2012. A review of the application of molecular genetics for
473 fisheries management and conservation of sharks and rays. *Journal of Fish Biology* **80**: 1789–
474 1843.
- 475 Dulvy NK, Fowler SL, Musick JA, Cavanagh RD, Kyne PM, Harrison LR, Carlson JK, Davidson
476 LNK, Fordham S V., Francis MP, et al. 2014. Extinction risk and conservation of the world's
477 sharks and rays. *eLife* **2014**: 1–34.
- 478 Fergusson IK, Vacchi M, Serena F. 2002. Note on the Declining Status of the Sandtiger Shark
479 *Carcharias Taurus* in the Mediterranean Sea. In *Proceedings of the 4th European*
480 *Elasmobranch Association Meeting, Livorno, (Italy) 2000*, Vacchi M, , La Mesa G, , Serena F,
481 , Séret B (eds). 73–76.
- 482 Ferretti F, Myers RA, Serena F, Lotze HK. 2008. Loss of large predatory sharks from the
483 Mediterranean Sea. *Conservation Biology* **22**: 952–964.
- 484 Frankham R. 1996. Relationship of genetic variation to population size in wildlife. *Conservation*
485 *Biology* **10**: 1500–1508.
- 486 Frankham R. 2005. Genetics and extinction. *Biological Conservation* **126**: 131–140.
- 487 García VB, Lucifora LO, Myers RA. 2008. The importance of habitat and life history to extinction

- 488 risk in sharks, skates, rays and chimaeras. *Proceedings of the Royal Society B: Biological*
489 *Sciences* **275**: 83–89.
- 490 Gilmore RG. 1993. Reproductive biology of lamnoid sharks. *Environmental Biology of Fishes* **38**:
491 95–114.
- 492 Goldman KJ, Branstetter S, Musick JA. 2006. A re-examination of the age and growth of sand tiger
493 sharks, *Carcharias taurus*, in the western North Atlantic: The importance of ageing protocols
494 and use of multiple back-calculation techniques. *Environmental Biology of Fishes* **77**: 241–
495 252.
- 496 Gubili C, Bilgin R, Kalkan E, Karhan SU, Jones CS, Sims DW, Kabasakal H, Martin AP, Noble
497 LR. 2011. Antipodean white sharks on a Mediterranean walkabout? Historical dispersal leads
498 to genetic discontinuity and an endangered anomalous population. *Proceedings of the Royal*
499 *Society B: Biological Sciences* **278**: 1679–1686.
- 500 Gubili C, Robinson CEC, Cliff G, Wintner SP, de Sabata E, De Innocentiis S, Canese S, Sims DW,
501 Martin AP, Noble LR, et al. 2015. DNA from historical and trophy samples provides insights
502 into white shark population origins and genetic diversity. *Endangered Species Research* **27**:
503 233–241.
- 504 Guichenot AA. 1850. Exploration Scientifique de l'Algérie: Pendant les Années 1840, 1841, 1842.
505 In *Histoire naturelle des reptiles et des poisson (Vol. 5)* Imprimerie nationale: Paris;
- 506 Hall TA. 1999. BioEdit: a user-friendly biological sequence alignment editor and analysis program
507 for Windows 95/98/NT. *Nucleic Acids Symposium Series* **41**: 95–98.
- 508 Haulsee DE, Breece MW, Brown LM, Wetherbee BM, Fox DA, Oliver MJ. 2018. Spatial ecology
509 of *Carcharias taurus* in the northwestern Mid-Atlantic coastal ocean. *Marine Ecology Progress*
510 *Series* **597**: 191–206.
- 511 Ismen A, Cigdem Yigin C, Altinagac U, Ayaz A. 2009. Length-weight relationships for ten shark
512 species from Saros Bay (North Aegean Sea). *Journal of Applied Ichthyology* **25**: 109–112.
- 513 Keeney DB, Heist EJ. 2006. Worldwide phylogeography of the blacktip shark (*Carcharhinus*
514 *limbatus*) inferred from mitochondrial DNA reveals isolation of western Atlantic populations
515 coupled with recent Pacific dispersal. *Molecular Ecology* **15**: 3669–3679.

- 516 Knapp M, Clarke AC, Horsburgh KA, Matisoo-Smith EA. 2012. Setting the stage - Building and
517 working in an ancient DNA laboratory. *Annals of Anatomy* **194**: 3–6.
- 518 Kneebone J, Chisholm J, Skomal G. 2014. Movement patterns of juvenile sand tigers (*Carcharias*
519 *taurus*) along the east coast of the USA. *Marine Biology* **161**: 1149–1163.
- 520 Larkin MA, Blackshields G, Brown NP, Chenna R, Mcgettigan PA, McWilliam H, Valentin F,
521 Wallace IM, Wilm A, Lopez R, et al. 2007. Clustal W and Clustal X version 2.0.
522 *Bioinformatics* **23**: 2947–2948.
- 523 Lipej L, De Maddalena A, Soldo A. 2004. *Sharks of the Adriatic sea*. Knjiznica Annales Majora.
524 Koper.
- 525 Lucifora LO, Menni RC, Escalante AH. 2002. Reproductive ecology and abundance of the sand
526 tiger shark, *Carcharias taurus*, from the southwestern Atlantic. *ICES Journal of Marine Science*
527 **59**: 553–561.
- 528 Nielsen EE, Morgan JAT, Maher SL, Edson J, Gauthier M, Pepperell J, Holmes BJ, Bennett MB,
529 Ovenden JR. 2017. Extracting DNA from ‘jaws’: high yield and quality from archived tiger
530 shark (*Galeocerdo cuvier*) skeletal material. *Molecular Ecology Resources* **17**: 431–442.
- 531 Otway NM, Ellis MT. 2011. Pop-up archival satellite tagging of *Carcharias taurus*: Movements and
532 depth/temperature-related use of south-eastern Australian waters. *Marine and Freshwater*
533 *Research* **62**: 607–620.
- 534 Otway NM, Bradshaw CJA, Harcourt RG. 2004. Estimating the rate of quasi-extinction of the
535 Australian grey nurse shark (*Carcharias taurus*) population using deterministic age- and stage-
536 classified models. *Biological Conservation* **119**: 341–350.
- 537 Pääbo S, Poinar H, Serre D, Jaenicke-Després V, Hebler J, Rohland N, Kuch M, Krause J, Vigilant
538 L, Hofreiter M. 2004. Genetic Analyses from Ancient DNA. *Annual Review of Genetics* **38**:
539 645–679.
- 540 Peeters FJC, Acheson R, Brummer GJA, De Ruijter WPM, Schneider RR, Ganssen GM, Ufkes E,
541 Kroon D. 2004. Vigorous exchange between the Indian and Atlantic oceans at the end of the
542 past five glacial periods. *Nature* **430**: 661–665.
- 543 Pollard D, Smith A. 2000. *Carcharias taurus*. *The IUCN Red List of Threatened Species 2000*.

- 544 Pollard DA, Lincoln Smith MP, Smith AK. 1996. The biology and conservation status of the grey
545 nurse shark (*Carcharias taurus* Rafinesque 1810) in New South Wales, Australia. *Aquatic*
546 *Conservation: Marine and Freshwater Ecosystems* **6**: 1–20.
- 547 Pollard DA, Gordon I, Williams S, Flaherty A, McAuley R. 2003. *Carcharias taurus* East coast of
548 Australia subpopulation. *The IUCN Red List of Threatened Species 2003* e.T44070A1.
- 549 Por FD. 1978. Lessepsian migration. The influx of Red Sea biota into the Mediterranean by way of
550 the Suez Canal. *Springer Verlag publ., Berlin* **3**: 1–128.
- 551 Quignard J-P, Capapé C. 1972. Complément à la liste commentée des Sélaciens de Tunisie. *Bulletin*
552 *de l'Institut National Scientifique et Technique d'Océanographie et de Pêche de Salammbô* **2**:
553 445–447.
- 554 Rohland N, Hofreiter M. 2007. Ancient DNA extraction from bones and teeth. *Nature Protocols* **2**:
555 1756–1762.
- 556 Smale MJ, Booth AJ, Farquhar MR, Mejer MR, Rochat L. 2012. Migration and habitat use of
557 formerly captive and wild raggedtooth sharks (*Carcharias taurus*) on the southeast coast of
558 South Africa. *Marine Biology Research* **8**: 115–128.
- 559 Splendiani A, Fioravanti T, Giovannotti M, Negri A, Ruggeri P, Olivieri L, Nisi Cerioni P,
560 Lorenzoni M, Caputo Barucchi V. 2016. The effects of paleoclimatic events on mediterranean
561 trout: Preliminary evidences from ancient DNA. *PLoS ONE* **11**: e0157975.
- 562 Splendiani A, Fioravanti T, Giovannotti M, Olivieri L, Ruggeri P, Nisi Cerioni P, Vanni S,
563 Enrichetti F, Caputo Barucchi V. 2017. Museum samples could help to reconstruct the original
564 distribution of *Salmo trutta* complex in Italy. *Journal of Fish Biology* **90**: 2443–2451.
- 565 Stow A, Zenger K, Briscoe D, Gillings M, Peddemors V, Otway N, Harcourt R. 2006. Isolation and
566 genetic diversity of endangered grey nurse shark (*Carcharias taurus*) populations. *Biology*
567 *Letters* **2**: 308–311.
- 568 Teter SM, Wetherbee BM, Fox DA, Lam CH, Kiefer DA, Shivji M. 2015. Migratory patterns and
569 habitat use of the sand tiger shark (*Carcharias taurus*) in the western North Atlantic. *Marine*
570 *and Freshwater Research* **66**: 158–169.
- 571 Toonen RJ, Bowen BW, Iacchei M, Briggs JC. 2016. Biogeography, Marine. In *Encyclopedia of*

- 572 *Evolutionary Biology Vol. 1*, Kliman RM (ed). Academic Press: Oxford; 166–178.
- 573 Untergasser A, Cutcutache I, Koressaar T, Ye J, Faircloth BC, Remm M, Rozen SG. 2012. Primer3-
574 new capabilities and interfaces. *Nucleic Acids Research* **40**: 1–12.
- 575 Vanni S. 1992. Cataloghi del Museo di Storia Naturale dell’Università di Firenze sezione di
576 zoologia «La Specola». XI: Chondrichthyes. In *Atti della Società Toscana di Scienze Naturali*,
577 *Memorie, Serie B* 85–114.
- 578 Walls RHL, Soldo A. 2016. *Carcharias taurus*. *The IUCN Red List of Threatened Species 2016*
579 **e.T3854A16**.
- 580 Waples RS. 1991. Pacific salmon, *Oncorhynchus* spp., and the definition of ‘species’ under the
581 Endangered Species Act. *Marine Fisheries Review* **53**: 11–22.
- 582 Whitenack LB, Simkins DC, Motta PJ, Hirai M, Kumar A. 2010. Young’s modulus and hardness of
583 shark tooth biomaterials. *Archives of Oral Biology* **55**: 203–209.
- 584 Wynne R, Wilding CS. 2018. Mitochondrial DNA haplotype diversity and origin of captive sand
585 tiger sharks (*Carcharias taurus*). *Journal of Zoo and Aquarium Research* **6**: 74–78.
- 586 Yang DY, Cannon A, Saunders SR. 2004. DNA species identification of archaeological salmon
587 bone from the Pacific Northwest Coast of North America. *Journal of Archaeological Science*
588 **31**: 619–631.

589

590 TABLES

591 **TABLE 1** Information about museum specimens of *Carcharias taurus* sampled and analysed in the
592 present study

Genetic code	Institution	Institution code	Description	Sampling location and date	Reference	Sample type
FI002	Museo di Storia Naturale, Sezione di Zoologia “La Specola”, Florence, Italy	INV6136	Taxidermied specimen, male, 170 cm length	Messina, Sicily, 15th November 1879	Vanni, 1992	Cartilage powder
PA001	Museo di Zoologia “Pietro Doderlein”, Palermo, Italy	ID AN 68	Jaws	Sicily, second half of the XIX century	Doderlein, 1879	Cartilage powder
PA002	Museo di Zoologia “Pietro Doderlein”, Palermo, Italy	ID AN 94	Jaws	Sicily, second half of the XIX century	Doderlein, 1879	Cartilage powder

PA003	Museo di Zoologia "Pietro Doderlein", Palermo, Italy	ID AN 60	Jaws	Sicily, second half of the XIX century	Doderlein, 1879	Cartilage powder
PA004	Museo di Zoologia "Pietro Doderlein", Palermo, Italy	ID AN 38	Skeleton	Sicily, second half of the XIX century	Doderlein, 1879	Cartilage powder
PR001	Muséum National d'Histoire Naturelle, Paris, France	A-9685	Taxidermied specimen, female	Algeria, ~ 1840	Guichenot, 1850 B. Seret, pers. comm.	Piece of cartilage
PR004	Muséum National d'Histoire Naturelle, Paris, France	AB-0038	Jaws	Algeria, ~ 1840	Guichenot, 1850 B. Seret, pers. comm.	Piece of cartilage
XL001	Royal Belgian Institute of Natural Sciences, Brussels, Belgium	507β	Jaws	Algeria, end of the XIX century	O. Pauwels, pers. comm.	Tooth
XL002	Royal Belgian Institute of Natural Sciences, Brussels, Belgium	1386β	Teeth collection, erroneously classified as <i>Odontaspis ferox</i>	Tunisia, 1st October 1933	O. Pauwels, pers. comm.	Tooth

593

594 **TABLE 2** Primer pairs designed and used to amplify a portion of the mtDNA CR of *Carcharias*
595 *taurus* historical samples

Primer name	Sequence 5' to 3'	Product length
CtCR1	F CTTCAATCCTTGATCGCGTCA	135 bp
	R CTTCCGGGGAATAGCGATGG	
CtCR2	F TGGCATTTCGTCCCTTGATCG	146 bp
	R TGAGTATGTTAGATAGATGTCGAGGA	
CtCR3	F GGCTGAACTGGGACACTGAG	146 bp
	R TCGAAACTTGCCGACTATGG	
CtCR4	F TGTCAAGTTGACCAAACTGAAA	118 bp
	R CCGGATGGGGGTTAAGAGAG	
CtCR5	F CCATAGTCGGCAAGTTTCGA	148 bp
	R TGCCAGATAAAGTGAAGAATGTGT	
CtCR6	F CTCTCTTAACCCCATCCGG	213 bp
	R GGGTTTTTCGAGGAGTCCGT	
CtCR7	F ACACATTCTTCACTTTATCTGGCA	172 bp

R ATGTCCGGCCCTCGTTTTAG

CtCR8 F ACGGACTCCTCGAAAAACCC

141 bp

R TCATCTTAGCATCTTCAGTGCCA

596

597 **TABLE 3** Polymorphic sites detected after the alignment

	Polymorphic sites																	
	42	131	182	318	330	335	337	339	356	407	408	420	421	427	444	445	562	572
Haplotypes																		
HapA	T	C	A	C	A	G	G	G	T	A	G	-	-	G	G	A	G	A
FI002	-	-
PA001	-	-
PA003	-	-
PR004	-	-
PA004	?	.	.	-	-
HapB	A	-	-
XL001	.	.	?	.	.	A	-	-
PR001	A	.	.	?	.	.	-	-
HapJ	A	.	.	C	.	.	-	-
HapD	G	A	-	-
HapI	A	A	-	-
XL002	A	A	-	-
HapC	A	A	-	-	G
HapH	.	T	G	.	G	A	A	A	.	.	.	-	-	G
HapE	.	.	G	.	G	A	A	A	.	.	.	-	-	G
HapG	.	.	G	.	G	A	A	A	.	G	A	A	T	A	-	-	T	.
HapF	.	.	G	T	G	A	A	A	.	G	A	A	T	A	-	-	T	.
HapK	A	.	G	.	G	A	A	A	.	G	A	A	T	A	-	-	?	?

Haplotypes from previous studies were highlighted in grey (Ahonen *et al.*, 2009; Chang *et al.*, 2015; Wynne and Wilding, 2018). Haplotype A was used as a reference sequence. All identical nucleotides in other sequences are indicated as full stops (.), indels as dashes (-) and missing nucleotides as question marks (?). In the case of historical samples, missing data are due to amplification failures, whereas for Haplotype K (Wynne and Wilding, 2018) they are present because the sequence is shorter than the others (518 bp vs 574 bp).

598

599 **FIGURE LEGENDS**

600 **FIGURE 1** Graphic representation of the eight overlapping primer pairs designed to amplify a
601 portion of the mtDNA CR sequence in Mediterranean historical samples of *Carcharias taurus*. The
602 numeration of the mitochondrial DNA started from the first base of the region studied by Ahonen et
603 al. (2009)

604 **FIGURE 2** Median Joining network showing the relationship between mtDNA CR haplotypes of
605 *Carcharias taurus*. The circle size is related to the number of individuals sampled worldwide for
606 each haplotype. Each colour indicates a different sampling location