



UNIVERSITÀ POLITECNICA DELLE MARCHE
Repository ISTITUZIONALE

Protective effects of Manuka honey on LPS-treated RAW 264.7 macrophages. Part 2: Control of oxidative stress induced damage, increase of antioxidant enzyme activities and attenuation of inflammation

This is a pre print version of the following article:

Original

Protective effects of Manuka honey on LPS-treated RAW 264.7 macrophages. Part 2: Control of oxidative stress induced damage, increase of antioxidant enzyme activities and attenuation of inflammation / Gasparri, Massimiliano; Afrin, Sadia; Forbes-Hernández, Tamara Y.; Cianciosi, Danila; Reboledo-Rodríguez, Patricia; Amici, Adolfo; Battino, Maurizio; Giampieri, Francesca. - In: FOOD AND CHEMICAL TOXICOLOGY. - ISSN 0278-6915. - STAMPA. - 120:(2018), pp. 578-587. [10.1016/j.fct.2018.08.001]

Availability:

This version is available at: 11566/264841 since: 2022-05-25T12:24:22Z

Publisher:

Published

DOI:10.1016/j.fct.2018.08.001

Terms of use:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. The use of copyrighted works requires the consent of the rights' holder (author or publisher). Works made available under a Creative Commons license or a Publisher's custom-made license can be used according to the terms and conditions contained therein. See editor's website for further information and terms and conditions.

This item was downloaded from IRIS Università Politecnica delle Marche (<https://iris.univpm.it>). When citing, please refer to the published version.

(Article begins on next page)

Manuscript Number:

Title: Protective effects of Manuka honey on LPS-treated RAW 264.7 macrophages. Part 2: Control of oxidative stress induced damage, increase of antioxidant enzyme activities and attenuation of inflammation

Article Type: Full Length Article

Keywords: Manuka honey; lipopolysaccharides; reactive oxygen species; anti-oxidative effects; anti-inflammatory effects.

Corresponding Author: Professor Maurizio Battino, PhD, DSc, MS, MD (Hon)

Corresponding Author's Institution: Università Politecnica delle Marche

First Author: Massimiliano Gasparrini

Order of Authors: Massimiliano Gasparrini; Sadia Afrin; Tamara Forbes-Hernandez; Danila Cianciosi; Patricia Reboredo-Rodriguez; Adolfo Amici; Maurizio Battino, PhD, DSc, MS, MD (Hon); Francesca Giampieri

Abstract: The redox-system is altered by oxidative stress that is initiated by oxidative agents such as lipopolysaccharides (LPS) and reactive oxygen species (ROS), which are strongly involved in chronic inflammation. Even if Manuka honey (MH) is a good source of polyphenol rich antioxidants, its antioxidant and anti-inflammatory effects are still elusive. The aim of the present work was to explore the protective effects of MH against E.coli LPS stimulated oxidative stress and inflammatory condition and the underlying mechanisms on murine RAW 264.7 macrophages. Pre-treatment with MH markedly inhibited LPS induced ROS and nitrite accumulation and increased the protection against cellular biomolecules such as lipids, proteins, and DNA. Stimulation by LPS suppressed both antioxidant enzyme activities and expressions which were significantly ($p < 0.05$) increased in the presence of MH. The pro-inflammatory cytokines, such as TNF- α , IL-1 β and IL-6, and other inflammatory mediators (iNOS) were enhanced after LPS treatment, whereas MH suppressed the expression of these inflammatory markers. Moreover, MH also inhibited the expression of TLR4/NF- κ B via I κ B phosphorylation in LPS-stressed RAW 264.7 macrophages. In conclusion, MH acted as a natural agent for preventing oxidative and inflammatory-related diseases.

Suggested Reviewers: Maria Greabu
University Carol Davila, Bucharest
mariagreabu@yahoo.com

Li-Shu Wang
Medical College of Wisconsin
liswang@mcw.edu

Pedro Bullon
University of Seville
pbullon@us.es

Stefano Bompadre
Polytechnic University of Marche
s.bompadre@univpm.it

Jose L Quiles
University of Granada, Spain
jlquiles@ugr.es

Jose M Alvarez-Suarez
Universidad de Las Américas. Ecuador.
jose.alvarez@udla.edu.ec

Lada Živković
University of Belgrade
lada.zivkovic@pharmacy.bg.ac.rs

Andrea Čabarkapa-Pirkovića
University of Belgrade
andrea.cabarkapa@gmail.com

Opposed Reviewers: Luca Tiano
l.tiano@univpm.it
Conflict of interest

Elisabetta Damiani
e.damiani@univpm.it
Conflict of interest



Ancona, July 19th, 2018

Dear Dr Domingo,

as agreed last week, my coworkers and I are submitting the second part of the research concerning **“Protective effects of Manuka honey on LPS-treated RAW 264.7 macrophages”** to be considered for publication in *Food and Chemical Toxicology*, being the full title **“Protective effects of Manuka honey on LPS-treated RAW 264.7 macrophages. Part 2: Control of oxidative stress induced damage, increase of antioxidant enzyme activities and attenuation of inflammation”**.

The manuscript has not been published and is not under consideration for publication elsewhere. All the authors have read the manuscript and have approved this submission. The authors report no conflicts of interest.

Hoping that our paper could be suitable for publication I look forward to hearing from you.

Sincerely yours,

Prof. Maurizio Battino, PhD, DSc, MS, MD



Protective effects of Manuka honey on LPS-treated RAW 264.7 macrophages.

**Part 2: Control of oxidative stress induced damage, increase of antioxidant enzyme activities
and attenuation of inflammation**

Massimiliano Gasparrini^{a,1}, Sadia Afrin^{a,1}, Tamara Y. Forbes-Hernández^a, Danila Cianciosi^a,
Patricia Reboredo-Rodriguez^{a,b}, Adolfo Amici^a, Maurizio Battino^a, Francesca Giampieri^{a,*}

^a*Dipartimento di Scienze Cliniche Specialistiche ed Odontostomatologiche (DISCO)-Sez. Biochimica, Facoltà di Medicina, Università Politecnica delle Marche, 60131, Ancona, Italy*

^b*Departamento de Química Analítica y Alimentaria, Grupo de Nutrición y Bromatología, Universidade de Vigo, Ourense 32004, Spain*

¹These authors contributed equally to this work.

***Correspondence to:** Dr Francesca Giampieri, Dipartimento di Scienze Cliniche Specialistiche ed Odontostomatologiche (DISCO)-Sez. Biochimica, Facoltà di Medicina, Università Politecnica delle Marche, Ancona, Via Ranieri 65, 60131, Italy. E mail: f.giampieri@univpm.it; Tel.: +39-071-220-4136; Fax: +39-071-220-4123.

Abstract

The redox-system is altered by oxidative stress that is initiated by oxidative agents such as lipopolysaccharides (LPS) and reactive oxygen species (ROS), which are strongly involved in chronic inflammation. Even if Manuka honey (MH) is a good source of polyphenol rich antioxidants, its antioxidant and anti-inflammatory effects are still elusive. The aim of the present work was to explore the protective effects of MH against *E.coli* LPS stimulated oxidative stress and inflammatory condition and the underlying mechanisms on murine RAW 264.7 macrophages. Pre-treatment with MH markedly inhibited LPS induced ROS and nitrite accumulation and increased the protection against cellular biomolecules such as lipids, proteins, and DNA. Stimulation by LPS suppressed both antioxidant enzyme activities and expressions which were significantly ($p < 0.05$) increased in the presence of MH. The pro-inflammatory cytokines, such as TNF- α , IL-1 β and IL-6, and other inflammatory mediators (iNOS) were enhanced after LPS treatment, whereas MH suppressed the expression of these inflammatory markers. Moreover, MH also inhibited the expression of TLR4/NF- κ B via I κ B phosphorylation in LPS-stressed RAW 264.7 macrophages. In conclusion, MH acted as a natural agent for preventing oxidative and inflammatory-related diseases.

Keywords: Manuka honey; lipopolysaccharides; reactive oxygen species; anti-oxidative effects; anti-inflammatory effects.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Protective effects of Manuka honey on LPS-treated RAW 264.7 macrophages.

Part 2: Control of oxidative stress induced damage, increase of antioxidant enzyme activities and attenuation of inflammation

Massimiliano Gasparri^{a,1}, Sadia Afrin^{a,1}, Tamara Y. Forbes-Hernández^a, Danila Cianciosi^a,
Patricia Reboredo-Rodriguez^{a,b}, Adolfo Amici^a, Maurizio Battino^a, Francesca Giampieri^{a,*}

^a*Dipartimento di Scienze Cliniche Specialistiche ed Odontostomatologiche (DISCO)-Sez. Biochimica, Facoltà di Medicina, Università Politecnica delle Marche, 60131, Ancona, Italy*

^b*Departamento de Química Analítica y Alimentaria, Grupo de Nutrición y Bromatología, Universidade de Vigo, Ourense 32004, Spain*

¹These authors contributed equally to this work.

***Correspondence to:** Dr Francesca Giampieri, Dipartimento di Scienze Cliniche Specialistiche ed Odontostomatologiche (DISCO)-Sez. Biochimica, Facoltà di Medicina, Università Politecnica delle Marche, Ancona, Via Ranieri 65, 60131, Italy. E mail: f.giampieri@univpm.it; Tel.: +39-071-220-4136; Fax: +39-071-220-4123.

1
2
3
4 **Abstract**
5

6 The redox-system is altered by oxidative stress that is initiated by oxidative agents such as
7 lipopolysaccharides (LPS) and reactive oxygen species (ROS), which are strongly involved in
8 chronic inflammation. Even if Manuka honey (MH) is a good source of polyphenol rich
9 antioxidants, its antioxidant and anti-inflammatory effects are still elusive. The aim of the
10 present work was to explore the protective effects of MH against *E.coli* LPS stimulated oxidative
11 stress and inflammatory condition and the underlying mechanisms on murine RAW 264.7
12 macrophages. Pre-treatment with MH markedly inhibited LPS induced ROS and nitrite
13 accumulation and increased the protection against cellular biomolecules such as lipids, proteins,
14 and DNA. Stimulation by LPS suppressed both antioxidant enzyme activities and expressions
15 which were significantly ($p < 0.05$) increased in the presence of MH. The pro-inflammatory
16 cytokines, such as TNF- α , IL-1 β and IL-6, and other inflammatory mediators (iNOS) were
17 enhanced after LPS treatment, whereas MH suppressed the expression of these inflammatory
18 markers. Moreover, MH also inhibited the expression of TLR4/NF- κ B via I κ B phosphorylation
19 in LPS-stressed RAW 264.7 macrophages. In conclusion, MH acted as a natural agent for
20 preventing oxidative and inflammatory-related diseases.
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

48 **Keywords:** Manuka honey; lipopolysaccharides; reactive oxygen species; anti-oxidative effects;
49 anti-inflammatory effects.
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 **Highlights**
5
6

- 7 • Manuka honey protected RAW 264.7 macrophages against LPS induces oxidative stress.
- 8 • Manuka honey improved antioxidant enzyme activities and expressions.
- 9 • Manuka honey enhanced oxidative defense by activation of Keap1-Nrf2 signaling.
- 10 • Manuka honey suppressed LPS stimulated inflammatory mediators.
- 11 • Manuka honey suppressed LPS induced TLR4/NF-κB signaling.
- 12
- 13
- 14
- 15
- 16
- 17
- 18
- 19
- 20
- 21
- 22
- 23
- 24
- 25
- 26
- 27
- 28
- 29
- 30
- 31
- 32
- 33
- 34
- 35
- 36
- 37
- 38
- 39
- 40
- 41
- 42
- 43
- 44
- 45
- 46
- 47
- 48
- 49
- 50
- 51
- 52
- 53
- 54
- 55
- 56
- 57
- 58
- 59
- 60
- 61
- 62
- 63
- 64
- 65

1. Introduction

Inflammation is the defensive response in the human body to infection caused by microbial pathogens, chemical irritants or damaged cells (Martinon et al., 2009). Several studies have reported that chronic inflammation is the main link for the pathogenesis of various diseases, such as arthritis, age-related diseases, atherosclerosis and cancer (Medzhitov, 2008). In the inflammatory processes, uncontrolled reactive species destruct lipids in cell membranes, affecting their permeability and the structure and function of proteins; they lead to DNA damage and mutations, which have high impact on cellular oxidative state, and decrease antioxidant defense mechanisms (Nogueira and Hay, 2013). Inflammation is also promoted by genetic alterations in the intestinal tract because of the former production of reactive oxygen species (ROS) and reactive nitrogen species (RNS), inflammatory chemokine and cytokine (Munkholm, 2003).

Macrophages are the main immune cells, which are associated with the progression of various inflammatory diseases, and are activated by cytokines, mitogens, radiation or bacterial lipopolysaccharides (LPS); at the same time, activated macrophage release various inflammatory mediators (Laskin et al., 2011). LPS activation of toll like receptor (TLR4) complexes on the outer surface of macrophages triggers intracellular nuclear factor kappa-light-chain-enhancer of activated B cells (NF- κ B) signaling pathways, which lead to transcriptional activation of several genes, including inducible nitric oxide synthase (iNOS), cyclooxygenase 2 (COX-2), prostaglandin E2 (PGE2), tumor necrosis factor alpha (TNF- α), interleukin (IL)-1 β , and IL-6 (Laskin et al., 2011).

Although efficient anti-inflammation drugs are currently accessible, it is still a challenge to discover less toxic and effectual agents for treating the range of acute and chronic inflammatory

1
2
3
4 diseases. From the ancient time, honey has been used as conventional medicine in various
5
6 cultural populations mainly for its wound healing and anti-bacterial properties (Alvarez-Suarez
7
8 et al., 2013; Alvarez-Suarez et al., 2014). Recently, several reports have pointed out multiple
9
10 activities of honey in enhancing anti-inflammatory (Alvarez-Suarez et al., 2010), anti-cancer
11
12 (Afrin et al., 2018a; Afrin et al., 2018b), immune response (Al-Waili and Haq, 2004) and anti-
13
14 cardiovascular (Alvarez-Suarez et al., 2013) effects. Manuka honey (MH), obtained from the
15
16 nectar of honey bees (*Apis mellifera*), comes from New Zealand Manuka tree (*Leptospermum*
17
18 *scoparium*, Family: Myrtaceae) and has been largely studied for its health beneficial effects
19
20 (Alvarez-Suarez et al., 2014). It is a complex mixture of various pharmacological and
21
22 biologically active compounds (Alvarez-Suarez et al., 2016; Afrin et al., 2018a; Afrin et al.,
23
24 2017) and those compounds are thought to be involved in its anti-microbial (Alvarez-Suarez et
25
26 al., 2014), wound healing, anti-oxidative (Alvarez-Suarez et al., 2016) and chemopreventive
27
28 effects (Afrin et al., 2018a; Afrin et al., 2018b), but the anti-inflammatory properties remain to
29
30 be unclear in spite of its high nutritional values.
31
32
33
34
35
36
37

38 The current study is aimed to investigate the antioxidant and anti-inflammatory effects of MH on
39
40 RAW 264.7 macrophages. Therefore, in part 1 of our work we reported the nutritional and
41
42 phytochemical characterization of MH (Afrin et al., 2018c). After that, ROS and nitrite
43
44 accumulation and the activity and the expression of the antioxidant enzymes were estimated.
45
46 Particular attention has been given on the molecular mechanisms by which MH exhibits its anti-
47
48 inflammatory effects that were evaluated in by *E.coli* LPS activated RAW 264.7 macrophages.
49
50
51
52
53
54

55 **2. Materials and methods**

56 *2.1. Chemicals and reagents*

57
58
59
60
61
62
63
64
65

1
2
3
4 MH samples were supplied by EfitSrl (Italy) which and reserved at 4 °C until analysis. All
5
6 chemicals and reagents were used for the experimental analysis were purchased from Sigma-
7
8 Aldrich (Milan, Italy). Media for cell culture were obtained from Carlo Erba Reagents (Milan,
9
10 Italy). The primary antibodies were purchased from Santa Cruz Biotechnology (Dallas, TX,
11
12 USA), while secondary antibody was purchased from Sigma-Aldrich (Milan, Italy).
13
14
15
16
17
18

19 *2.2. RAW 264.7 macrophages cell culture and treatments*

20
21 Murine RAW 264.7 macrophages cells were purchased from the American Type Culture
22
23 Collection (ATCC-TIB71). Cells were cultured in Dulbecco's Modified Eagle Medium (DMEM)
24
25 with 10% heat-inactivated fetal bovine serum, 100 IU/mL penicillin and 100 IU/mL
26
27 streptomycin. Cells were preserved in Hera Cell CO₂ incubator at 37°C with 95% air and 5%
28
29 CO₂. All the experiments were conducted between the passages 4th to 6th. In part 1 of our work,
30
31 we performed the viability assay for choosing the subsequent concentration of MH and LPS
32
33 (Afrin et al., 2018c). Cells were treated for 24 h with different condition: only DMEM (control),
34
35 MH (3 mg/mL and 8 mg/mL), LPS (1 µg/mL), and MH (3 mg/mL)+LPS and MH (8
36
37 mg/mL)+LPS) combination.
38
39
40
41
42
43
44

45 *2.3. Determination of intracellular ROS levels*

46
47 CellROX[®] Oxidative Stress kit (Invitrogen™, Life Techonoliges, Milan, Italy) was used for
48
49 determination of intracellular ROS generation (Giampieri et al., 2018). The cells were seeded in
50
51 a 6 well plate at a density of 1.5 X 10⁵ and treated for 24 h with different treatments as specified
52
53 earlier. At the end of the treatment, the CellROX[®] orange reagents (2 µL/mL) were added and
54
55 incubated for 30 min at 37°C. After the incubation, the cells were centrifuged for removing the
56
57
58
59
60
61
62
63
64
65

1
2
3
4 excess dye and medium, re-suspended the cells with 100 μ L PBS and analyzed by using a Tali®
5
6 Image-Based cytometer (Invitrogen™, Life Technologies, Milan, Italy) and the results were
7
8 expressed as the fold change compared to control.
9

10 11 12 13 14 *2.4. Determination of nitrite accumulation*

15
16 Nitrite production was demonstrated by the Griess method (Pekarova et al., 2009). The cells
17
18 were seeded in a T75 flask at a density of 1×10^6 and treated for 24 h with different treatments
19
20 as specified earlier. At the end of the treatment, cells supernatants were mixed with equal volume
21
22 of Griess reagent (1% sulfanilamide and 0.1% naphthyl-ethylenediamine in 5% phosphoric acid)
23
24 and incubated 10 min in the dark at room temperature. The absorbance was measured on a
25
26 microplate reader (ThermoScientific Multiskan EX) at 540 nm. The nitrite concentration was
27
28 determined by using a sodium nitrite standard curve (0.1 to 6.25 μ M). Each experiment was
29
30 carried out in three replicates and the results were expressed as nmol nitrite /mg protein, while
31
32 the final results were expressed as the fold change compared to control.
33
34
35
36
37
38

39 40 *2.5. Determination of biomarkers of oxidative stress and antioxidant enzyme activities*

41
42 RAW 264.7 macrophages were treated for 24 h as previously described. The cellular lysates
43
44 were prepared by using the RIPA buffer (Sigma-Aldrich, Milan, Italy) for determination of
45
46 glutathione (GSH), lipid peroxidation (thiobarbituric acid-reactive substance, TBARS), protein
47
48 carbonyl content and antioxidant enzyme activities (glutathione peroxidase (GPx), glutathione
49
50 reductase (GR), glutathione transferase (GST), superoxide dismutase (SOD) and catalase (CAT))
51
52 following to the earlier described method (Giampieri et al., 2016; Gasparini et al., 2017). The
53
54 results of GSH, TBARS, and carbonyl content were expressed as nmol /mg protein, the results of
55
56 GPx and GR were expressed as nmole NADPH oxidized/mg protein/min, the results of GST
57
58
59
60
61
62
63
64
65

1
2
3
4 were expressed as nmole 1-chloro-2,4-dinitrobenzene (CDNB)-reduced GSH conjugate/mg
5
6 protein/min and the results of SOD and CAT were expressed as U/mg protein/min. The final
7
8 results were expressed as the fold change compared to control.
9

10 11 12 13 14 *2.6. Isolation of RNA and RT-PCR analysis*

15
16 RAW 264.7 macrophages were treated for 24 h as previously described. By using PureLink®
17
18 RNA Mini Kit (Invitrogen, Carlsbad, CA, USA), the total RNA content of the cells was
19
20 determined. Microplate spectrophotometer system (BioTek Synergy HT, Winooski, USA) were
21
22 used for checking the RNA concentration and purity. 5× All-In-One RT MasterMix kit (Applied
23
24 Biological Materials Inc. Canada) were used for the synthesized cDNA from 75 ng RNA
25
26 according to reverse transcription. Real-time PCR (RT-PCR) amplification (Corbett Life
27
28 Science, Rotor-Gene 3000, Mortlake, Australia) was performed by using EvaGreen 2X qPCR
29
30 MasterMix (EvaGreen 2X qPCR MasterMix kit, Applied Biological Materials Inc. Canada) of
31
32 forward and reverse primers (Table 1). Glyceraldehyde-3-phosphate dehydrogenase (GADPH)
33
34 was used for normalizing quantitative data and the results were calculated by the $2^{-\Delta\Delta ct}$ method,
35
36 while the final results were expressed as the fold change compared to control.
37
38
39
40
41
42
43
44

45 46 *2.7. Immunoblotting analysis*

47
48 RAW 264.7 macrophages were treated for 24 h as previously described. Cell lysis buffer (120
49
50 mmol/L NaCl, 40 mmol/L Tris [pH 8], 0.1% NP40 with protease inhibitor cocktails (Sigma))
51
52 were used for the lysis of protein from cellular pellets. Proteins from cellular supernatants were
53
54 separated on 8 to 10% polyacrylamide gel and transferred into a nitrocellulose membrane by
55
56 using the trans-blot SD semidry electrophoretic transfer cell (Bio-Rad, Hercules, CA, USA). Tris
57
58
59
60
61
62
63
64
65

1
2
3
4 HCl buffered saline with Tween 20 (TBST) with 5% non-fat-milk were used for blocking the
5
6 membrane. The primary antibodies 8-Oxoguanine glycosylase (OGG1), nuclear factor E2-related
7
8 factor 2 (Nrf2), Kelch-like ECH-associated protein 1 (Keap1), SOD, CAT, heme oxygenase 1
9
10 (HO-1), TLR4, NF- κ B, phosphorylated nuclear factor of kappa light polypeptide gene enhancer
11
12 in B-cells inhibitor alpha (p-I κ B α), iNOS, TNF- α , IL-1 β , IL-6, IL-10 and GADPH (1:500
13
14 dilutions) were used after all night incubation (4 °C). TBST were used for washing the
15
16 membrane and incubated with secondary antibodies (1:80 000) for another 1 h. Immunolabeled
17
18 proteins were recognized by using a chemiluminescence method (C-DiGit Blot Scanner, LICOR,
19
20 Bad Homburg, Germany) and bands were quantified by using studio digits software 3.1 (C-DiGit
21
22 Blot Scanner, LICOR, Bad Homburg, Germany).

2.8. Statistical analysis

23
24
25
26
27
28
29
30
31
32
33 The statistical analysis was performed by using STATISTICA software (Statsoft Inc., Tulsa, OK,
34
35 USA) and the results are expressed as the mean values with the standard deviation of three
36
37 independent repetitions. One-way analysis of variance (ANOVA) followed by Tukey's honest
38
39 significant difference (HSD) post hoc test ($p < 0.05$) were used for the significant differences and
40
41 symbolized by letters.

3. Results and discussion

3.1. MH treatment decreases LPS induced intracellular ROS and nitrite accumulation

42
43
44
45
46
47
48
49
50
51
52
53 ROS play a vital role in modulating multiple molecular targets which are related to acute and
54
55 chronic inflammation, growth differentiation, proliferation and apoptosis (Nogueira and Hay,
56
57 2013). In the present study, we first investigated whether MH treatment could induce any effects
58
59
60
61
62
63
64
65

1
2
3
4 on ROS accumulation in RAW 264.7 macrophages. The cells were exposed to 3 mg/mL and 8
5 mg/mL of MH for 24 h. As shown in Fig. 1a no significant differences ($p < 0.05$) were observed
6
7 after the treatment of MH compared to control. Furthermore, ROS production increased nearly
8
9 2.5 fold after LPS treatment, while exposure of MH significantly (p -value < 0.05) decreased LPS
10
11 induced ROS generation up to 1.8 to 1.5 fold (Fig. 1a). In agreement with our results, MH
12
13 treatment significantly reduced the ROS levels in 2,2'-Azobis(2-amidinopropane)
14
15 dihydrochloride (AAPH) stressed human dermal fibroblast (HDF) cells (Alvarez-Suarez et al.,
16
17 2016). Decreased ROS generation was also observed in RAW 264.7 and HDF cells by
18
19 polyphenol-rich food which improved endotoxin induced inflammatory stress (Choi et al., 2007;
20
21 Gasparrini et al., 2017; Gasparrini et al., 2018).

22
23
24
25
26
27
28 Nitric oxide is a vital bioregulatory molecule in the immune systems and it modulates various
29
30 pathophysiological conditions inside the cells. Acute and chronic inflammation is induced by
31
32 overproduction of nitric oxide (Blaise et al., 2005), and therefore more awareness is being paid to
33
34 the development of natural potent inhibitor of its production. In RAW 264.7 macrophages, no
35
36 significant differences were found in nitrite generation after the treatment with MH compared to
37
38 control (Fig. 1b). After incubation with LPS, nitrite production increased nearly 1.9 fold
39
40 compared to control, while MH treatments significantly ($p < 0.05$) decreased LPS induced nitrite
41
42 accumulation up to 1.5 to 1.4 fold (Fig. 1b). Inhibitory effects were also observed in LPS
43
44 stimulated RAW 264.7 macrophages after the treatment with other phenolic compounds (Wang
45
46 and Mazza, 2002). The authors reported that all the flavonoids decreased nitric oxide production
47
48 and that the decrease was strongly correlated to the concentration of total polyphenols.
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 *3.2. MH improves GSH levels against oxidative damage*
5

6
7 GSH is a prototype antioxidant, which is able to protect the cellular components from the
8
9 excessive oxidative stress (free radicals, peroxides, and heavy metals), acting directly or as a
10
11 cofactor (Pompella et al., 2003). Increased GSH levels were observed after MH treatment in
12
13 RAW 264.7 macrophages (Fig. 2a). LPS treated cells exhibited higher oxidative stress by
14
15 significantly ($p < 0.05$) lowering GSH values (0.48 fold) compared to control (1.00 fold) and also
16
17 compared to MH (1.34 fold) (Fig. 2a). Interestingly, pre-treatment with MH protected the LPS-
18
19 induced stressed cells by restoring the GSH values up to 0.69 to 0.91 fold in the dose-dependent
20
21 manner (Fig. 2a). These results were consistent with previous reports that highlighted increased
22
23 GSH levels exerted by other food matrix or isolated compounds after LPS treatment in murine
24
25 RAW 264.7 macrophages (Gasparrini et al., 2017; Li et al., 2018).
26
27
28
29
30
31
32

33 *3.3. MH treatment decreases LPS stimulated lipid, protein and DNA damage*
34

35
36 Disproportion in the redox system increases ROS or RNS production, damages immune system
37
38 and may influence injuries to proteins, lipids, and DNA (Nogueira and Hay, 2013). ROS
39
40 commences lipid peroxidation by a series of reaction that generates free radicals and additional
41
42 substances (hydroperoxides, malondialdehydes, lipoperoxides and toxic aldehydes), which affect
43
44 membrane permeability and augment inflammation (Mena et al., 2009).
45
46
47

48 To determine whether the oxidative stress was related to the damage to main biomolecules, we
49
50 evaluated the markers of lipid (TBARS), protein (carbonyl content) and DNA damage (OGG1
51
52 expression) after the treatment of MH alone and in the presence of the oxidative agent (LPS)
53
54 (Fig. 2b-2d). We observed that the highest concentration of MH (8 mg/mL) significantly ($p <$
55
56 0.05) decreased TBARS levels (0.41 fold) and carbonyl content (0.44 fold) in RAW 264.7
57
58
59
60
61
62
63
64
65

1
2
3
4 macrophages compared to untreated cells (1.00 fold) (Fig. 2b and 2c). After the incubation of
5
6 LPS, the levels of lipid (1.85 fold) and protein (1.48 fold) damage were significantly higher
7
8 compared to control group (Fig. 2b and 2c), while MH treatment significantly ($p < 0.05$)
9
10 protected lipid and protein damage in LPS stressed cells by decreasing TBARS and protein
11
12 carbonyl content up to 1.21 fold and 1.12 fold (Fig. 2b and 2c). In agreement with our results,
13
14 MH has also shown a significant reduction of TBARS levels and protein carbonyl content
15
16 triggered by AAPH in HDF cells (Alvarez-Suarez et al., 2016). Moreover, polyphenol-
17
18 containing food demonstrated similar effects by reducing LPS-induced overproduction of
19
20 TBARS and protein carbonyl content in RAW 264.7 macrophages (Gasparrini et al., 2017).
21
22

23
24
25
26 OGG1, a bifunctional glycosylase, represents a DNA damage marker that is capable to break the
27
28 DNA backbone strand and split the glycosidic bond in the mutagenic area (Jacobs and Schär,
29
30 2012). Several studies exposed that the polyphenol-containing food have the ability to suppress
31
32 the OGG1 expression in a stressed condition both *in vitro* and *in vivo* models (Giampieri et al.,
33
34 2016; Gasparrini et al., 2017). In the present study, no significant differences were observed in
35
36 the expression of OGG1 after the treatment of MH compared to control, whereas after LPS
37
38 treatment the OGG1 expression was significantly increased compared to the control (Fig. 2d).
39
40 Pre-incubation with MH lead to significantly reduced OGG1 expression against LPS generated
41
42 damaged cells.
43
44
45
46
47
48
49

50 51 *3.4. MH treatment improve antioxidant enzyme activity and expression against oxidative damage*

52
53 The antioxidant enzyme GPS, GR, GST, SOD and CAT activities are showed in Fig. 3. On the
54
55 one hand, the antioxidant system upholds the redox system by scavenging cellular ROS and
56
57 decreasing the ratio of oxidized glutathione (GSH/GSSG) or NADPH/NADP⁺. On the other
58
59
60
61
62
63
64
65

1
2
3
4 hand, the protective effects of the antioxidant enzyme are hampered due to excess ROS or other
5
6 oxidants (Nogueira and Hay, 2013). Several studies reported that the natural compounds have the
7
8 ability to protect the cells from oxidative damage condition by increasing the endogenous
9
10 antioxidant enzyme activity and expression for restoring the normal condition of the cells (Lobo
11
12 et al., 2010).
13

14
15
16 Therefore, we evaluated the endogenous antioxidant enzyme activity and expression after the
17
18 treatment with MH alone or in the presence of LPS. The activity of GPx, GR and GST were
19
20 dose-dependently increased after the treatment of MH compared to control group in RAW 264.7
21
22 cells (Fig. 3a). Furthermore, when the cells were incubated with LPS a significant ($p < 0.05$)
23
24 reduction of above antioxidant enzyme activities were observed. MH was able to protect cells
25
26 from this damage condition in a dose-dependent manner by improving the activity of GPx and
27
28 GR, while GST activity was improved only for the highest concentration (8 mg/mL of MH) (Fig.
29
30 3a). No significant differences were found in SOD activity compared to control after both
31
32 concentration of MH, while the CAT activity significantly ($p < 0.05$) increased in the presence of
33
34 both doses of MH (Fig. 3b). Treatment of LPS decreased the activity of SOD and CAT, while
35
36 pre-treatment of MH significantly increased the SOD (3 and 8 mg/mL of MH) and CAT activity
37
38 (8 mg/mL of MH) (Fig. 3b). In a recent report by Alvarez-Suarez *et al.* MH showed the ability of
39
40 MH to protect AAPH stimulated HDF cells by increasing antioxidant enzyme SOD and CAT
41
42 activity (Alvarez-Suarez et al., 2016). Concomitantly, Gelam honey protected human diploid
43
44 fibroblast against gamma radiation by improving the activity of SOD, CAT and GPx (Ahmad et
45
46 al., 2013).
47
48
49
50
51
52
53

54
55 The transcription factors Nrf2 plays a vital role in cellular defense mechanism by stimulating
56
57 various antioxidant and detoxification enzymes at the transcriptional level through the
58
59
60
61
62
63
64
65

1
2
3
4 antioxidant response element (ARE). Under quiescent conditions, Nrf2 is sequestered by Keap1
5
6 in the cytoplasm, while during oxidative stress, Nrf2 is separated from Keap1 and translocates to
7
8 the nucleus, thus stimulating genes transcription having an ARE (Battino et al., 2018).
9

10
11 The protein expression of Nrf2-Keap1 signaling and other antioxidant enzyme SOD, CAT and
12
13 HO-1 were determined after the treatment of MH alone or in the presence of LPS in RAW 264.7
14
15 macrophages. We found that MH treatment increased the expression of Nrf2 and similar trend
16
17 was observed after LPS incubation, while co-treatment with MH significantly ($p < 0.05$)
18
19 increased the activities Nrf2 compared to control (Fig. 4). The protein levels of Keap1 were
20
21 increased after the exposure to MH or LPS (Fig. 4), consequently, the MH+LPS treatment
22
23 induced a significant upregulation of Keap1 in the RAW 264.7 cells (Fig. 4) compared to control
24
25 group. The expression of other antioxidant enzyme SOD, CAT and HO-1 were upregulated after
26
27 the MH treatment in a dose-dependent manner. In addition, in case of LPS, the expression of
28
29 SOD and HO-1 increased more compared to MH treatment while for CAT it induced similar
30
31 effects like MH (Fig. 4). At the same time, MH+LPS combination led to a significant (p -value <
32
33 0.05) higher increase in the expression of these genes compared to control and also other
34
35 treatments (Fig. 4). Similar trends were observed in AAPH stressed HDF cells, where MH
36
37 treatment increased the expression of Nrf2, SOD, and CAT for protecting the oxidative damage
38
39 condition (Alvarez-Suarez et al., 2016). Other natural compounds such as epigallocatechin-3-
40
41 gallate protected TNF- α induced NF- κ B activation by increasing Nrf2-Keap1 pathway in
42
43 macrophage foam cells (Jiang et al., 2012), whereas fisetin induced anti-inflammatory effects
44
45 demonstrated by HO-1 activation in LPS stimulated RAW 264.7 macrophages (Li et al., 2018).
46
47 Furthermore, polyphenol-rich foodstuff acted against oxidative damage through increasing both
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 SOD and CAT activity and expression in RAW macrophages and HDF cells (Gasparrini et al.,
5
6
7 2017; Gasparrini et al., 2018).
8
9

10 11 *3.5. MH treatment suppresses key inflammatory biomarkers induced by LPS*

12
13
14 Since chronic inflammation has been associated with the development of cancer, atherosclerosis,
15
16 and aging-related diseases, we evaluated the anti-inflammatory activity of MH (Fig. 5 and Fig. 6)
17
18 (Medzhitov, 2008). The transcription factor NF κ B acts as a key role in inflammation by
19
20 modulating numerous genes responsible for the generation of inflammatory mediators. In
21
22 presence of multiple pro-inflammatory stimuli, inhibitor of kappa B (I κ B) is rapidly degraded
23
24 and NF κ B subunits translocated into the nucleus to stimulate the transcription of various genes
25
26 encoding chemokines, cytokines, anti-apoptotic factors and growth factors (Lawrence, 2009). In
27
28 the colonic epithelial barrier, TLR4 activation leads to increasing intracellular NF- κ B signaling
29
30 and other pro-inflammatory cytokine (IL-1 β , IL-6, iNOS, COX-2, PGE2 etc.) which lead to
31
32 progression of chronic inflammatory diseases (Foran et al., 2010).
33
34
35
36
37

38
39 In our work, we quantified the mRNA and protein expression of several inflammation markers to
40
41 highlight if MH has a protective effect against the inflammation induced by LPS in RAW 264.7
42
43 macrophages. No significant differences were observed in the mRNA expression of NF- κ B,
44
45 iNOS, TNF- α , and IL-1 β after the treatment of MH, while all the mRNA expression were
46
47 significantly increased after the LPS incubation (Fig. 5). Interestingly, pre-treatment with MH at
48
49 both concentrations were able to decrease LPS induced over activation of NF- κ B, iNOS, TNF- α ,
50
51 and IL-1 β and at the highest concentration of MH values similar to the control were found (Fig.
52
53
54
55
56 5). Our results are very similar to those obtained in RAW264.7 mouse macrophages in which
57
58
59
60
61
62
63
64
65

1
2
3
4 plant polyphenol treatment suppressed LPS stimulated iNOS, TNF- α , COX-2, and IL-6 mRNA
5
6 expression (Kanno et al., 2006).
7

8
9 The anti-inflammatory effects were also confirmed by evaluating the protein expression of
10
11 TLR4, NF- κ B, p-I κ B α , iNOS, TNF- α , IL-1 β , IL-6 and IL-10 by Western Blot (Fig. 6). MH
12
13 treatment significantly (p -value < 0.05) decreased the expression of the above inflammatory
14
15 markers after 24 h treatment in RAW 264.7 cells (Fig. 6). Incubation with LPS significantly (p <
16
17 0.05) increased the protein levels of all the inflammatory markers, while in the presence of MH
18
19 suppressed expressions were observed (Fig. 6). Earlier reports have suggested that stimulation by
20
21 LPS induced the production of several inflammatory biomarkers, such as NF- κ B, p-I κ α , TNF- α ,
22
23 iNOS, COX-2, IL-6 and IL-1 β , whereas co-treatment with nutritionally rich foodstuff (Kim et
24
25 al., 2016; Gasparrini et al., 2017; Limtrakul et al., 2015) or biologically active compounds from
26
27 plants sources (Hou et al., 2015; Terra et al., 2007; Kanno et al., 2006) protected RAW 264.7
28
29 macrophages by suppressing the pro-inflammatory production. A different expression was
30
31 observed of IL-10, which was significantly (p < 0.05) increased after the LPS treatments even in
32
33 the presence of MH alone or in the combinations (Fig. 6). This result is quite similar to those
34
35 obtained in RAW macrophages cells which found a greater amount of inflammatory cytokines
36
37 stimulated by LPS, improved by the co-treatment of other plant polyphenols (Zong et al., 2012;
38
39 Lee et al., 2013; Mo et al., 2014).
40
41
42
43
44
45
46
47
48
49

50 **4. Conclusions**

51
52 This study explored the anti-oxidative and anti-inflammatory effects of MH against oxidative
53
54 damage in inflammation mediatory immune cells, macrophages. Anti-oxidative effects of MH
55
56 were related to the suppression of ROS and nitrite production as well as the protection of lipid,
57
58
59
60
61
62
63
64
65

1
2
3
4 protein and DNA damage. This mechanism was accompanied by increased antioxidant enzyme
5
6 activities, consequently to the enhanced Nrf2 expression and its synchronized enzymes SOD,
7
8 CAT and HO-1 expression. Subsequent analysis confirmed that MH effectively suppressed the
9
10 expression of inflammation mediators such as TNF- α , IL-1 β , IL-6 and iNOS. Finally, we also
11
12 found that MH effectively inhibited the expression of TLR4 and NF κ B pathway via suppression
13
14 of p-I κ B α . Taken together, these promising findings may increase the potential efficacy of MH
15
16 as a useful natural compound for inhibiting the inflammation via enhancing antioxidant activity
17
18 and consequently suppressing key inflammation mediators. Therefore, future research is required
19
20 to validate the anti-oxidative and anti-inflammatory approach *in vivo* models.
21
22
23
24
25
26
27

28 **Conflicts of interest**

29 The authors declare no conflicts of interest.
30
31
32
33
34
35

36 **Acknowledgment**

37 Patricia Reboredo-Rodriguez acknowledges Xunta de Galicia for her post-doctoral contract.
38
39
40
41
42

43 **Abbreviations**

44 AAPH, 2,2'-Azobis(2-amidinopropane) dihydrochloride; ARE, antioxidant responsive element;
45
46 COX-2, cyclooxygenase 2; CAT, catalase; DMEM, dulbecco's modified eagle medium;
47
48 GADPH, glyceraldehyde-3-phosphate dehydrogenase; GSH, glutathione; GST, glutathione
49
50 transferase; GPx, glutathione peroxidase; GR, glutathione reductase; HDF, human dermal
51
52 fibroblast, HO-1, heme oxygenase 1; IL-1 β , interleukin 1-beta; IL-6, interleukin 6; IL-10,
53
54 interleukin 10; iNOS, inducible nitric oxide synthase; Keap1, kelch-like ECH-associated protein
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 1; LPS, lipopolysaccharide; MH, Manuka honey; NF-kB, Nuclear factor kappa-light-chain-
5
6 enhancer of activated B cells; Nrf2, nuclear factor E2-related factor 2; OGG1, 8-Oxoguanine
7
8 glycosylase; PGE2, prostaglandin E2; p-IkBa, phosphorylated nuclear factor of kappa light
9
10 polypeptide gene enhancer in B-cells inhibitor alpha; RNS, reactive nitrogen species; ROS,
11
12 reactive oxygen species; SD, standard deviation; SOD, superoxide dismutase; TBARS,
13
14 thiobarbituric acid-reactive substances; TBST, Tris HCl buffered saline with Tween 20; TLR4,
15
16 toll like receptor 4; TNF-a, tumor necrosis factor alpha.
17
18
19
20
21
22

23 **References**

- 24
25
26 Afrin, S., Forbes-Hernandez, T.Y., Gasparrini, M., Bompadre, S., Quiles, J.L., Sanna, G., Spano,
27
28 N., Giampieri, F., Battino, M., 2017. Strawberry-tree honey induces growth inhibition of
29
30 human colon cancer cells and increases ROS generation: a comparison with Manuka honey. *Int.*
31
32 *J. Mol. Sci.* 18, 613.
33
34
35
36 Afrin, S., Giampieri, F., Gasparrini, M., Forbes-Hernández, T.Y., Cianciosi, D., Reboredo-
37
38 Rodriguez, P., Amici, A., Quiles, J.L., Battino, M., 2018a. The inhibitory effect of Manuka
39
40 honey on human colon cancer HCT-116 and LoVo cell growth. Part 1: the suppression of cell
41
42 proliferation, promotion of apoptosis and arrest of the cell cycle. *Food Funct.* 9, 2145-2157.
43
44
45 Afrin, S., Giampieri, F., Gasparrini, M., Forbes-Hernández, T.Y., Cianciosi, D., Reboredo-
46
47 Rodriguez, P., Manna, P.P., Zhang, J., Quiles, J.L., Battino, M., 2018b. The inhibitory effect of
48
49 Manuka honey on human colon cancer HCT-116 and LoVo cell growth. Part 2: Induction of
50
51 oxidative stress, alteration of mitochondrial respiration and glycolysis, and suppression of
52
53 metastatic ability. *Food Funct.* 9, 2158-2170.
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 Afrin, S., Gasparini, M., Forbes-Hernandez, T.Y., Cianciosi, D., Reboredo-Rodriguez, P.,
5
6 Manna, P.P., Battino M., Giampieri, F., 2018c. Protective effects of Manuka honey on LPS-
7
8 treated RAW 264.7 macrophages. Part 1: Enhancement of cellular viability, regulation of
9
10 cellular apoptosis and improvement of mitochondrial functionality. Food Chem. Toxicol.
11
12 Submitted.
13
14

15
16 Ahmad, T.A.F.T., Jubri, Z., Rajab, N.F., Rahim, K.A., Yusof, Y.A.M., Makpol, S., 2013. Gelam
17
18 honey protects against gamma-irradiation damage to antioxidant enzymes in human diploid
19
20 fibroblasts. Molecules 18, 2200-2211.
21
22

23
24 Al-Waili, N.S., Haq, A., 2004. Effect of honey on antibody production against thymus-dependent
25
26 and thymus-independent antigens in primary and secondary immune responses. J. Med. Food 7,
27
28 491-494.
29
30

31
32 Alvarez-Suarez, J.M., Gasparini, M., Forbes-Hernández, T.Y., Mazzoni, L., Giampieri, F.,
33
34 2014. The composition and biological activity of honey: a focus on Manuka honey. Foods 3,
35
36 420-432.
37

38
39 Alvarez-Suarez, J.M., Giampieri, F., Battino, M., 2013. Honey as a source of dietary
40
41 antioxidants: structures, bioavailability and evidence of protective effects against human
42
43 chronic diseases. Curr. Med. Chem. 20, 621-638.
44

45
46 Alvarez-Suarez, J.M., Giampieri, F., Cordero, M., Gasparini, M., Forbes-Hernández, T.Y.,
47
48 Mazzoni, L., Afrin, S., Beltrán-Ayala, P., González-Paramás, A.M., Santos-Buelga, C., 2016.
49
50 Activation of AMPK/Nrf2 signalling by Manuka honey protects human dermal fibroblasts
51
52 against oxidative damage by improving antioxidant response and mitochondrial function
53
54 promoting wound healing. J Funct. Foods 25, 38-49.
55
56
57
58
59
60
61
62
63
64
65

- 1
2
3
4 Alvarez-Suarez, J.M., Tulipani, S., Díaz, D., Estevez, Y., Romandini, S., Giampieri, F., Damiani,
5
6 E., Astolfi, P., Bompadre, S., Battino, M., 2010. Antioxidant and antimicrobial capacity of
7
8 several monofloral Cuban honeys and their correlation with color, polyphenol content and other
9
10 chemical compounds. *Food Chem. Toxicol.* 48, 2490-2499.
11
12
13
14 Battino, M., Giampieri, F., Pistollato, F., Sureda, A., de Oliveira, M.R., Pittalà, V., Mehterov, N.,
15
16 Sarafian, V., Fallarino, F., Nabavi, S.F., 2018. Nrf2 as regulator of innate immunity: A
17
18 molecular Swiss army knife! *Biotechnol. Adv.* 36, 358-370.
19
20
21 Blaise, G.A., Gauvin, D., Gangal, M., Authier, S., 2005. Nitric oxide, cell signaling and cell
22
23 death. *Toxicology* 208, 177-192.
24
25
26 Choi, S.-Y., Hwang, J.-H., Ko, H.-C., Park, J.-G., Kim, S.-J., 2007. Nobiletin from citrus fruit
27
28 peel inhibits the DNA-binding activity of NF- κ B and ROS production in LPS-activated RAW
29
30 264.7 cells. *J. Ethnopharmacol.* 113, 149-155.
31
32
33
34 Foran, E., Garrity-Park, M.M., Mureau, C., Newell, J., Smyrk, T.C., Limburg, P.J., Egan, L.J.,
35
36 2010. Upregulation of DNA methyltransferase-mediated gene silencing, anchorage-
37
38 independent growth, and migration of colon cancer cells by interleukin-6. *Mol. Cancer Res.* 8,
39
40 471-481.
41
42
43 Gasparrini, M., Forbes-Hernandez, T.Y., Giampieri, F., Afrin, S., Alvarez-Suarez, J.M.,
44
45 Mazzoni, L., Mezzetti, B., Quiles, J.L., Battino, M., 2017. Anti-inflammatory effect of
46
47 strawberry extract against LPS-induced stress in RAW 264.7 macrophages. *Food Chem.*
48
49 *Toxicol.* 102, 1-10.
50
51
52
53 Gasparrini, M., Giampieri, F., Forbes-Hernandez, T.Y., Afrin, S., Cianciosi, D., Reboredo-
54
55 Rodriguez, P., Varela-Lopez, A., Zhang, J., Quiles, J.L., Mezzetti, B., Bompadre, S., Battino,
56
57 M., 2018. Strawberry extracts efficiently counteract inflammatory stress induced by the
58
59
60
61
62
63
64
65

1
2
3
4 endotoxin lipopolysaccharide in Human Dermal Fibroblast. *Food Chem. Toxicol.* 114, 128-
5
6
7 140.

8
9 Giampieri, F., Alvarez-Suarez, J.M., Gasparri, M., Forbes-Hernandez, T.Y., Afrin, S.,
10
11 Bompadre, S., Rubini, C., Zizzi, A., Astolfi, P., Santos-Buelga, C., 2016. Strawberry
12
13 consumption alleviates doxorubicin-induced toxicity by suppressing oxidative stress. *Food*
14
15
16
17 *Chem. Toxicol.* 94, 128-137.

18
19 Giampieri, F., Quiles, J.L., Orantes-Bermejo, F.J., Gasparri, M., Forbes-Hernandez, T.Y.,
20
21 Sánchez-González, C., Llopis, J., Rivas-García, L., Afrin, S., Varela-López, A., Cianciosi, D.,
22
23 Reboledo-Rodríguez, P., Fernández-Piñar, C.T., Iglesias, R.C., Ruiz, R., Aparicio, S., Crespo,
24
25
26
27 J., Lopez, L.D., Xiao, J., Battino, M., 2018. Are by-products from beeswax recycling process a
28
29 new promising source of bioactive compounds with biomedical properties? *Food Chem.*
30
31
32 *Toxicol.* 112, 126-133.

33
34 Hou, X.L., Tong, Q., Wang, W.Q., Shi, C.Y., Xiong, W., Chen, J., Liu, X., Fang, J.G., 2015.
35
36 Suppression of inflammatory responses by dihydromyricetin, a flavonoid from *ampelopsis*
37
38 *grossedentata*, via inhibiting the activation of NF- κ B and MAPK signaling pathways. *J.Nat.*
39
40
41
42 *Prod.* 78, 1689-1696.

43
44 Jacobs, A.L., Schär, P., 2012. DNA glycosylases: in DNA repair and beyond. *Chromosoma* 121,
45
46
47 1-20.

48
49 Jiang, J., Mo, Z.-C., Yin, K., Zhao, G.-J., Lv, Y.-C., Ouyang, X.-P., Jiang, Z.-S., Fu, Y., Tang,
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65 C.-K., 2012. Epigallocatechin-3-gallate prevents TNF- α -induced NF- κ B activation thereby
upregulating ABCA1 via the Nrf2/Keap1 pathway in macrophage foam cells. *Int. J. Mol. Med.*
29, 946-956.

- 1
2
3
4 Kanno, S.-i., Shouji, A., Tomizawa, A., Hiura, T., Osanai, Y., Ujibe, M., Obara, Y., Nakahata,
5
6 N., Ishikawa, M., 2006. Inhibitory effect of naringin on lipopolysaccharide (LPS)-induced
7
8 endotoxin shock in mice and nitric oxide production in RAW 264.7 macrophages. *Life Sci.* 78,
9
10 673-681.
11
12
13
14 Kim, Y.-S., Ahn, C.-B., Je, J.-Y., 2016. Anti-inflammatory action of high molecular weight
15
16 *Mytilus edulis* hydrolysates fraction in LPS-induced RAW264. 7 macrophage via NF- κ B and
17
18 MAPK pathways. *Food Chem.* 202, 9-14.
19
20
21 Laskin, D.L., Sunil, V.R., Gardner, C.R., Laskin, J.D., 2011. Macrophages and tissue injury:
22
23 agents of defense or destruction? *Annu. Rev. Pharmacol. Toxicol.* 51, 267-288.
24
25
26 Lawrence, T., 2009. The Nuclear Factor NF-kB Pathway in Inflammation. Cold Spring Harb.
27
28 *Perspect. Biol.* 1, a001651.doi:001610.001101/cshperspect.a001651.
29
30
31 Lee, J., Kim, S., Namgung, H., Jo, Y.-H., Bao, C., Choi, H.-K., Auh, J.-H., Lee, H.J., 2013.
32
33 Ellagic acid identified through metabolomic analysis is an active metabolite in strawberry
34
35 ('Seolhyang') regulating lipopolysaccharide-induced inflammation. *J. Agric. Food Chem.* 62,
36
37 3954-3962.
38
39
40
41 Li, K.K., Shen, S.S., Deng, X., Shiu, H.T., Siu, W.S., Leung, P.C., Ko, C.H., Cheng, B.H., 2018.
42
43 dihydrofisetin exerts its anti-inflammatory effects associated with suppressing Erk/p38 Mapk
44
45 and Heme Oxygenase-1 activation in lipopolysaccharide-stimulated Raw 264.7 macrophages
46
47 and carrageenan-induced mice paw edema. *Int. Immunopharmacol.* 54, 366-374.
48
49
50
51 Limtrakul, P., Yodkeeree, S., Pitchakarn, P., Punfa, W., 2015. Suppression of inflammatory
52
53 responses by black rice extract in RAW 264.7 macrophage cells via downregulation of NF-kB
54
55 and AP-1 signaling pathways. *Asian Pac. J. Cancer Prev.* 16, 4277-4283.
56
57
58
59
60
61
62
63
64
65

- 1
2
3
4 Lobo, V., Patil, A., Phatak, A., Chandra, N., 2010. Free radicals, antioxidants and functional
5
6 foods: Impact on human health. *Pharmacogn. Rev.* 4, 118-126.
7
8
9 Martinon, F., Mayor, A., Tschopp, J.r., 2009. The inflammasomes: guardians of the body. *Annu*
10
11 *Rev. Immunol.* 27, 229-265.
12
13
14 Medzhitov, R., 2008. Origin and physiological roles of inflammation. *Nature* 454, 428.
15
16 Mena, S., Ortega, A., Estrela, J.M., 2009. Oxidative stress in environmental-induced
17
18 carcinogenesis. *Mutat. Res. Genet. Toxicol. Environ. Mutagen.* 674, 36-44.
19
20
21 Mo, C., Wang, L., Zhang, J., Numazawa, S., Tang, H., Tang, X., Han, X., Li, J., Yang, M.,
22
23 Wang, Z., 2014. The crosstalk between Nrf2 and AMPK signal pathways is important for the
24
25 anti-inflammatory effect of berberine in LPS-stimulated macrophages and endotoxin-shocked
26
27 mice. *Antioxid. Redox Signal.* 20, 574-588.
28
29
30
31 Munkholm, P., 2003. The incidence and prevalence of colorectal cancer in inflammatory bowel
32
33 disease. *Aliment. Pharmacol. Ther.* 18, 1-5.
34
35
36 Nogueira, V., Hay, N., 2013. Molecular pathways: reactive oxygen species homeostasis in
37
38 cancer cells and implications for cancer therapy. *Clin. Cancer Res.* 19, 4309-4314.
39
40
41 Pekarova, M., Kralova, J., Kubala, L., Ciz, M., Papezikova, I., Macickova, T., Pecivova, J.,
42
43 Nosal, R., Lojek, A., 2009. Carvedilol and adrenergic agonists suppress the lipopolysaccharide-
44
45 induced NO production in RAW 264.7 macrophages via the adrenergic receptors. *Acta Physiol.*
46
47 *Polonica* 60, 143.
48
49
50
51 Pompella, A., Visvikis, A., Paolicchi, A., De Tata, V., Casini, A.F., 2003. The changing faces of
52
53 glutathione, a cellular protagonist. *Biochem. Pharmacol.* 66, 1499-1503.
54
55
56 Terra, X., Valls, J., Vitrac, X., Mérrillon, J.-M., Arola, L., Ardèvol, A., Bladé, C., Fernández-
57
58 Larrea, J., Pujadas, G., Salvadó, J., Blay, M., 2007. Grape-seed procyanidins act as
59
60
61
62
63
64
65

1
2
3
4 antiinflammatory agents in endotoxin-stimulated RAW 264.7 macrophages by inhibiting NFkB
5
6 signaling pathway. *J. Agric. Food Chem.* 55, 4357-4365.
7
8

9 Wang, J., Mazza, G., 2002. Inhibitory effects of anthocyanins and other phenolic compounds on
10 nitric oxide production in LPS/IFN- γ -activated RAW 264.7 macrophages. *J. Agric. Food*
11
12 *Chem.* 50, 850-857.
13
14
15

16 Zong, Y., Sun, L., Liu, B., Deng, Y.-S., Zhan, D., Chen, Y.-L., He, Y., Liu, J., Zhang, Z.-J., Sun,
17
18 J., 2012. Resveratrol inhibits LPS-induced MAPKs activation via activation of the
19
20 phosphatidylinositol 3-kinase pathway in murine RAW 264.7 macrophage cells. *PloS One* 7,
21
22 e44107.
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Tables

Table 1. Sequences of real-time PCR oligonucleotide primers.

| Gene (symbol) | Direction | Sequence (5'-3') |
|--------------------|-----------|---------------------------------|
| NF- κ B p65 | Forward | AGGCTTCTGGGCCTTATGTG |
| | Reverse | TGCTTCTCTCGCCAGGAATAC |
| iNOS | Forward | CCCTTCCGAAGTTTCTGGCAGC |
| | Reverse | GGCTGTCAGAGCCTCGTGGCTT |
| TNF- α | Forward | ATGAGCACAGAA AG CATGATC |
| | Reverse | TACAGGCTTGTCACTCGA ATT |
| IL-1 β | Forward | TGCAGAGTTCCCCAACTGGTACATC |
| | Reverse | GTGCTGCCTAATGTCCCCTTGAATC |
| GADPH | Forward | CAC TCA CGG CAA ATT CAA CGG CAC |
| | Reverse | GAC TCC ACG ACA TAC TCA GCA |

1
2
3
4 **Figure captions**
5

6 **Fig. 1. Intracellular ROS and nitrite levels.** RAW 264.7 macrophages treated with MH and
7 LPS alone or in combination for 24 h. (a) Intracellular ROS levels were determined by using the
8 Tali™ Image-based Cytometer assay. Representative fluorescence image of RAW 264.7
9 macrophages cells: non-fluorescent upon no ROS and red fluorescence upon oxidation by ROS.
10
11 (b) Nitrite levels were determined by using the Griess method. Results are expressed as a fold
12 change in comparison with control (no treatment). All data are indicated as the mean ± standard
13 deviation (SD) (n=3). Different letters indicate significant differences (p -value < 0.05) between
14 each treatment.
15
16
17
18
19
20
21
22
23
24
25
26
27
28

29 **Fig. 2. Levels of glutathione (GSH) and biomarkers of oxidative damage (lipid, protein, and**
30 **DNA).** (a) GSH levels, (b) lipid damage levels (TBARS), (c) protein carbonyl levels and (d)
31 protein expression of OGG1 (DNA damage) were determined in RAW 264.7 macrophages
32 treated with MH and LPS alone or in combination for 24 h. Results are expressed as a fold
33 change in comparison with control (no treatment). All data are indicated as the mean ± SD (n=3).
34 Different letters indicate significant differences (p -value < 0.05) between each treatment.
35
36
37
38
39
40
41
42
43
44

45 **Fig. 3. Antioxidant enzyme activities.** The activity of antioxidant enzymes (a) GPx, GR, GST
46 and (b) SOD, CAT were determined in RAW 264.7 macrophages treated with MH and LPS
47 alone or in combination for 24 h. Results are expressed as a fold change in comparison with
48 control (no treatment). All data are indicated as the mean ± SD (n=3). Different letters indicate
49 significant differences (p -value < 0.05) between each treatment.
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 **Fig. 4. Protein expression levels of Nrf2-Keap1 and other antioxidant enzymes.** RAW 264.7
5
6 macrophages treated with MH and LPS alone or in combination for 24 h. The protein expression
7
8 of Keap1, Nrf2, SOD, CAT, and HO-1 were analyzed by western blotting. GADPH was utilized
9
10 as a loading control. Results are expressed as a fold change in comparison with control (no
11
12 treatment). All data are indicated as the mean \pm SD (n=3). Different letters indicate significant
13
14 differences (p -value $<$ 0.05) between each treatment.
15
16
17
18
19
20

21 **Fig. 5. mRNA levels of inflammation markers.** RAW 264.7 macrophages treated with MH and
22
23 LPS alone or in combination for 24 h. The mRNA expression of NF- κ B, iNOS, TNF- α , and IL-
24
25 1 β were analyzed by RT-PCR. GADPH was utilized as a loading control. Results are expressed
26
27 as a fold change in comparison with control (no treatment). All data are indicated as the mean \pm
28
29 SD (n=3). Different letters indicate significant differences (p -value $<$ 0.05) between each
30
31 treatment.
32
33
34
35
36
37

38 **Fig. 6. Protein expression levels of inflammation markers.** RAW 264.7 macrophages treated
39
40 with MH and LPS alone or in combination for 24 h. The protein expression of (a) TLR4, NF- κ B,
41
42 p-I κ B α , iNOS and (b) TNF- α , IL-1 β , IL-6, IL-10 were analyzed by western blotting. GADPH
43
44 was utilized as a loading control. Results are expressed as a fold change in comparison with
45
46 control (no treatment). All data are indicated as the mean \pm SD (n=3). Different letters indicate
47
48 significant differences (p -value $<$ 0.05) between each treatment.
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 **Graphical Abstract**
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65




Figure 1
[Click here to download high resolution image](#)










Figure 2
[Click here to download high resolution image](#)



| | | | | | | |
|---------------|---|---|---|---|---|---|
| Control | - | - | - | - | - | - |
| MH (3 mg/mL) | - | + | - | - | + | - |
| MH (8 mg/mL) | - | - | + | - | - | + |
| LPS (1 µg/mL) | - | - | - | + | + | + |

Figure 3
[Click here to download high resolution image](#)




Figure 4

[Click here to download high resolution image](#)





Figure 5
[Click here to download high resolution image](#)

Levels of inflammation markers









Figure 6
[Click here to download high resolution image](#)



Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting http://www.adobe.com/go/reader_download.

For more assistance with Adobe Reader visit <http://www.adobe.com/go/acrreader>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.

Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting http://www.adobe.com/go/reader_download.

For more assistance with Adobe Reader visit <http://www.adobe.com/go/acrreader>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.

Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting http://www.adobe.com/go/reader_download.

For more assistance with Adobe Reader visit <http://www.adobe.com/go/acrreader>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.

Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting <http://www.adobe.com/products/acrobat/readstep2.html>.

For more assistance with Adobe Reader visit <http://www.adobe.com/support/products/acrreader.html>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.

Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting http://www.adobe.com/go/reader_download.

For more assistance with Adobe Reader visit <http://www.adobe.com/go/acrreader>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.

Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting http://www.adobe.com/go/reader_download.

For more assistance with Adobe Reader visit <http://www.adobe.com/go/acrreader>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.

Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting http://www.adobe.com/go/reader_download.

For more assistance with Adobe Reader visit <http://www.adobe.com/go/acrreader>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.

Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting http://www.adobe.com/go/reader_download.

For more assistance with Adobe Reader visit <http://www.adobe.com/go/acrreader>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.