

UNIVERSITÀ POLITECNICA DELLE MARCHE Repository ISTITUZIONALE

Invariant metrics for the quaternionic Hardy space

This is the peer reviewd version of the followng article:

Original

Invariant metrics for the quaternionic Hardy space / Arcozzi, Nicola; Sarfatti, Giulia. - In: THE JOURNAL OF GEOMETRIC ANALYSIS. - ISSN 1050-6926. - STAMPA. - 25:(2015), pp. 2028-2059. [10.1007/s12220-014-9503-4]

Availability:

This version is available at: 11566/272342 since: 2022-05-27T11:58:13Z

Publisher:

Published DOI:10.1007/s12220-014-9503-4

Terms of use:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. The use of copyrighted works requires the consent of the rights' holder (author or publisher). Works made available under a Creative Commons license or a Publisher's custom-made license can be used according to the terms and conditions contained therein. See editor's website for further information and terms and conditions. This item was downloaded from IRIS Università Politecnica delle Marche (https://iris.univpm.it). When citing, please refer to the published version.

note finali coverpage

(Article begins on next page)

Invariant metrics for the quaternionic Hardy space

Nicola Arcozzi, Giulia Sarfatti

This is a post-print version, the reference for the published version is: J Geom Anal (2015) 25:2028–2059 DOI 10.1007/s12220-014-9503-4

Licence: Publisher's Bespoke License

Articles, books and chapters published within the Springer Nature group of companies which are made available through academic repositories remain subject to copyright. Any further use is subject to permission from Springer Nature. The following restrictions on use of such articles apply:

Academic research only

• 1. Archived content may only be used for academic research. Any content downloaded for text based experiments should be destroyed when the experiment is complete.

Use must not be for Commercial Purposes

• 2. Archived content may not be used for purposes that are intended for or directed towards commercial advantage or monetary compensation by means of sale, resale, licence, loan, transfer or any other form of commercial exploitation ("Commercial Purposes").

Wholesale re-publishing is prohibited

- 3. Archived content may not be published verbatim in whole or in part, whether or not this is done for Commercial Purposes, either in print or online.
- 4. This restriction does not apply to reproducing normal quotations with an appropriate citation. In the case of text-mining, individual words, concepts and quotes up to 100 words per matching sentence may be used, whereas longer paragraphs of text and images cannot (without specific permission from Springer Nature).

Moral rights

- 5. All use must be fully attributed. Attribution must take the form of a link using the article DOI to the published article on the journal's website.
- 6. All use must ensure that the authors' moral right to the integrity of their work is not compromised.

Third party content

• 7. Where content in the document is identified as belonging to a third party, it is the obligation of the user to ensure that any use complies with copyright policies of the owner.

Use at own risk

• 8. Any use of Springer Nature content is at your own risk and Springer Nature accepts no liability arising from such use.

Invariant metrics for the quaternionic Hardy space

Nicola Arcozzi, Giulia Sarfatti[†]

Abstract

We find Riemannian metrics on the unit ball of the quaternions, which are naturally associated with reproducing kernel Hilbert spaces. We study the metric arising from the Hardy space in detail. We show that, in contrast to the one-complex variable case, no Riemannian metric is invariant under all regular self-maps of the quaternionic ball.

KEY WORDS AND PHRASES: Hardy space on the quaternionic ball; functions of a quaternionic variable; invariant Riemannian metric.

MATHEMATICS SUBJECT CLASSIFICATION: 30G35, 46E22, 58B20.

Notation. The symbol \mathbb{H} denotes the set of the quaternions $q = x_0 + x_1i + x_2j + x_3k = \operatorname{Re}(q) + \operatorname{Im}(q)$, with $\operatorname{Re}(q) = x_0$ and $\operatorname{Im}(q) = x_1i + x_2j + x_3k$; where the x_j 's are real numbers and the imaginary units i, j, k are subject to the rules ij = k, jk = i, ki = j and $i^2 = j^2 = k^2 = -1$. We identify the quaternions q whose imaginary part vanishes, $\operatorname{Im}(q) = 0$, with real numbers, $\operatorname{Re}(q) \in \mathbb{R}$; and, similarly, we let $\mathbb{I} = \mathbb{R}i + \mathbb{R}j + \mathbb{R}k$ be the set of the imaginary quaternions. The norm $|q| \ge 0$ of q is $|q| = \sqrt{\sum_{l=0}^{3} x_l^2} = (q\bar{q})^{1/2}$, where $\bar{q} = x_0 - x_1i - x_2j - x_3k$ is the conjugate of q. The open unit ball \mathbb{B} in \mathbb{H} contains the quaternions q such that |q| < 1. The boundary of \mathbb{B} in \mathbb{H} is denoted by $\partial\mathbb{B}$. By the symbol \mathbb{S} we denote the unit sphere of the imaginary quaternions having the form q = x + yI, with x, y in \mathbb{R} . If f is a real differentiable function on a domain $\Omega \subseteq \mathbb{H}$, we denote its real differential at a point $w \in \Omega$ by the symbol $f_*[w]$.

1 Introduction

Let \mathbb{H} be the skew-field of the quaternions. The quaternionic Hardy space $H^2(\mathbb{B})$ consists of the formal power series of the quaternionic variable q,

$$f(q) = \sum_{n=0}^{\infty} q^n a_n$$

such that the sequence of quaternions $\{a_n\}$ satisfies

$$||f||_{H^2(\mathbb{B})} := \left(\sum_{n=0}^{\infty} |a_n|^2\right)^{1/2} < \infty.$$
(1)

^{*}Partially supported by the PRIN project Real and Complex Manifolds of the Italian MIUR and by INDAM-GNAMPA †Partially supported by INDAM-GNSAGA and by the PRIN project Real and Complex Manifolds of the Italian MIUR

It is easily verified that the series converges to a function $f : \mathbb{B} = \{q \in \mathbb{H} : |q| < 1\} \to \mathbb{H}$. The function f is *slice regular* [14] in the sense of Gentili and Struppa, who developed a version of complex function theory which holds in the quaternionic setting. See the monograph [13] for a detailed account of the theory. The norm (1) can be polarized to obtain an inner product with values in the quaternions,

$$\left\langle \sum q^n a_n, \sum q^n b_n \right\rangle_{H^2(\mathbb{B})} := \sum_{n=0}^{\infty} \overline{b_n} a_n.$$

The space $H^2(\mathbb{B})$ is a reproducing kernel Hilbert space, in the quaternionic sense: for w in \mathbb{B} and f in $H^2(\mathbb{B})$ we have

$$f(w) = \langle f, k_w \rangle_{H^2(\mathbb{B})}$$
, where $k_w(q) = k(w, q) = \sum_{n=0}^{\infty} q^n \overline{w}^n$

There is a rich interplay between reproducing kernel Hilbert spaces and distance functions. See [3] for an overview and several examples from one-variable holomorphic function space theory. In [10] the connection between metric theory and operator theory is analyzed at a very deep level, and the case of the Hardy space is a model example of that. The seminal article [4] by Aronszajn is still an excellent introduction to the theory of reproducing kernel Hilbert spaces.

In this article we are mainly interested in studying metrics on \mathbb{B} which are associated with the function space $H^2(\mathbb{B})$. We also provide evidence that the metric properties of the space reflect the behavior of functions in $H^2(\mathbb{B})$. The first metric we consider measures the distance between projections of kernel functions in the unit sphere of the Hilbert space $H^2(\mathbb{B})$:

$$\delta(p,q) := \sqrt{1 - \left| \left\langle \frac{k_q}{\|k_q\|_{H^2(\mathbb{B})}}, \frac{k_p}{\|k_p\|_{H^2(\mathbb{B})}} \right\rangle_{H^2(\mathbb{B})} \right|^2}.$$
(2)

In the holomorphic case of $H^2(\Delta)$ one obtains this way the pseudo-hyperbolic metric $\delta'(z, w) = \left| \frac{z-w}{1-\overline{w}z} \right|$. A calculation, see Proposition 4.2 below, gives a formally similar result in the quaternionic case:

$$\delta(z,w) = |(1-q\overline{w})^{-*} * (q-w)|_{|_{q=z}},$$

for z, w in \mathbb{B} . Here, the product f(q) * g(q) and the multiplicative inverse $f(q)^{-*}$ are not pointwise product and pointwise inverse: they are *-product and *-inverse, which are defined so that the usual convolution rule for coefficients of power series' products holds. See [13], and Section 2 where we summarize some background material on slice regular functions.

The infinitesimal version of the pseudo-hyperbolic metric in the complex disc, is the hyperbolic metric in the Riemann-Beltrami-Poincaré disc model: $ds^2 = \frac{|dz|^2}{(1-|z|^2)^2}$. By infinitesimal version of a distance δ_0 , we mean the length metric associated with δ_0 (see e.g. [16]). The infinitesimal metric associated with δ is a Riemannian metric g on \mathbb{B} .

Theorem 1.1. (I) The length metric associated with (2) is the Riemannian metric g defined below.

For any $w \in \mathbb{B}$, let us identify the tangent space $T_w\mathbb{B}$ with \mathbb{H} . For any vector $d \in T_w\mathbb{B}$, where $w = x + yI_w$ lies in L_{I_w} , we decompose $d = d_1 + d_2$ with d_1 in L_{I_w} and d_2 in $L_{I_w}^{\perp}$, the orthogonal

complement of L_{I_w} with respect to the Euclidean metric in \mathbb{H} . The length of d with respect to g is:

$$|d|_{g(w)}^{2} = \frac{1}{(1-|w|^{2})^{2}}|d_{1}|^{2} + \frac{1}{|1-w^{2}|^{2}}|d_{2}|^{2}.$$
(3)

(II) The isometry group of g is the one generated by the following classes of self-maps of \mathbb{B} :

(a) regular Möbius transformations of the form

$$q \mapsto M_{\lambda}(q) = (1 - q\lambda)^{-*} * (q - \lambda) = \frac{q - \lambda}{1 - \lambda q},$$

with λ in (-1, 1);

(b) isometries of the sphere of the imaginary units,

$$q = x + yI \mapsto T_A(q) = x + yA(I),$$

where $A : \mathbb{S} \to \mathbb{S}$ is an isometry of \mathbb{S} ;

(c) the reflection in the imaginary hyperplane,

$$q \mapsto R(q) = -\overline{q}.$$

In the metric (3), the first, "large" summand is the hyperbolic metric on a slice, while the second "small" summand is peculiarly quaternionic: it measures the variation of a quaternionic Hardy function in the direction orthogonal to the slices. Its small size reflects in quantitative, geometric terms the fact that regular functions are affine in the S variable, see [13].

The special rôle of the real axis in the theory of slice regular functions is here reflected in the fact that all isometries of the metric g fix $\mathbb{R} \cap \mathbb{B}$. In particular, contrary to the case of the complex disc, the action is not transitive. Other, more precise, properties of the metric will be stated and proved on route to the proof of Theorem 1.1. We will study geodesics, geodesically complete submanifolds and other geometric properties of the metric. For instance, we will prove that the radius of injectivity is infinite at points of the real diameter of \mathbb{B} , and finite elsewhere. The metric has neither positive, nor negative sectional curvature.

The proof of Theorem 1.1 splits into two steps. First, we compute the Riemannian metric associated with rather general reproducing kernel quaternionic Hilbert spaces \mathcal{H} ; that is the "infinitesimal" length metric $g_{\mathcal{H}}$ associated with the distance function

$$\delta_{\mathcal{H}}(w,z) = \sqrt{1 - \left| \left\langle \frac{k_w}{\|k_w\|_{\mathcal{H}}}, \frac{k_z}{\|k_z\|_{\mathcal{H}}} \right\rangle_{\mathcal{H}} \right|^2}.$$

We prove

Theorem 1.2. Let Ω be a symmetric slice domain and let \mathcal{H} be a reproducing kernel Hilbert space of regular functions on Ω . Suppose that k is slice preserving: for any $w \in \Omega$ the kernel function k_w preserves the slice L_{I_w} identified by w. Then the length of a tangent vector $d = d_1 + d_2 \in L_{I_w} + L_{I_w}^{\perp} \cong$ $T_w \Omega$ with respect to the Riemannian metric $g_{\mathcal{H}}$ associated with \mathcal{H} is given by

$$|d|_{g_{\mathcal{H}}(w)}^{2} = \frac{\left(\|k_{w}\|_{\mathcal{H}}^{2} \left\|\overline{\partial_{c}k_{q}(w)}\right\|_{\mathcal{H}}^{2} - \left|\overline{\partial_{c}k_{w}(w)}\right|^{2}\right)}{\|k_{w}\|_{\mathcal{H}}^{4}} |d_{1}|^{2} + \frac{\left(\|k_{w}\|_{\mathcal{H}}^{2} \left\|\overline{\partial_{s}k_{q}(w)}\right\|_{\mathcal{H}}^{2} - \left|\overline{\partial_{s}k_{w}(w)}\right|^{2}\right)}{\|k_{w}\|_{\mathcal{H}}^{4}} |d_{2}|^{2} + \frac{\left(\|k_{w}\|_{\mathcal{H}}^{2} \left\|\overline{\partial_{s}k_{q}(w)}\right\|_{\mathcal{H}}^{2} - \left(\|k_{w}\|_{\mathcal{H}}^{2} \left\|\overline{\partial_{s}k_{q}(w)}\right\|_{\mathcal{H}}^{2} - \left(\|k_{w}\|_{\mathcal{H}}^{2} \left\|\overline{\partial_{s}k_{q}(w)}\right\|_{\mathcal{H}}^{2} - \left(\|k_{w}\|_{\mathcal{H}}^{2} \left\|\overline{\partial_{s}k_{q}(w)}\right\|_{\mathcal{H}}^{2} - \left(\|k_{w}\|_{\mathcal{H}}^{2} \left\|k_{w}\|_{\mathcal{H}}^{2} - \left(\|k_{w}\|_{\mathcal{H}}^{2} \left\|\overline{\partial_{s}k_{q}(w)}\right\|_{\mathcal{H}}^{2} - \left(\|k_{w}\|_{\mathcal{H}}^{2} \left\|k_{w}\|_{\mathcal{H}}^{2} - \left(\|k_{w}\|_{\mathcal{H}}^{2} \left\|k_{w}\|_{\mathcal{H}}^{2} - \left(\|k_{w}\|_{\mathcal{H}}^{2} \left\|k_{w}\|_{\mathcal{H}}^{2} - \left(\|k_{w}\|_{\mathcal{H}}^{2} \right\|k_{w}\|_{\mathcal{H}}^{2} - \left(\|k_{w}\|_{\mathcal{H}}^{2} - \left(\|k_$$

Symmetric slice domains are domains in \mathbb{H} endowed with a natural symmetry, see Section 2. Examples of Hilbert spaces satisfying the assumptions of Theorem 1.2 are the Hardy space on \mathbb{B} and on the right half-space $\mathbb{H}^+ = \{q \in \mathbb{H} \mid \text{Re}(q) > 0\}$, and the Bergman space on \mathbb{B} . The formula in Theorem 1.2 should be compared with that in Theorem 1.17 in [10]. See also formula (0.3) in [17]. Part I of Theorem 1.1 is a special case of Theorem 1.2.

The classification of the isometries in part II of Theorem 1.1 is carried out in Section 4.

The Riemannian metric g has a rather small group of isometries, compared with the state of things in the unit disc of the complex plane, or even in the unit ball in several complex variables, with the Bergman-Kobayashi metric. The latter metrics have a transitive group of isometries and, more, the space is isotropic; whereas all isometries of the former have to fix the real line. One might expect that something better is possible. Unfortunately, there is no Riemannian metric on \mathbb{B} which is invariant under any *regular* Möbius function, unless the Möbius function is already an isometry for the metric defined in Theorem 1.1. If a geometric invariant for slice regular functions on the quaternionic ball exists, it has to be other than a Riemannian metric.

Theorem 1.3. Let a be a point of $\mathbb{B} \setminus \mathbb{R}$. There is no Riemannian metric m on \mathbb{B} having as isometry the regular Möbius transformation $q \mapsto (1 - q\overline{a})^{-*} * (a - q)$.

The proof will be given in Subsection 4.3. We mention here that Bisi and Gentili proved in [5] that the usual Poincaré metric on \mathbb{B} is invariant under classical (non-regular) Möbius maps.

A first relationship between the space $H^2(\mathbb{B})$ and the metric g concerns the H^2 norm itself. Let $r\mathbb{S}^3$ be the sphere of radius 0 < r < 1 in \mathbb{B} , with respect to the usual Euclidean metric; containing quaternions $q = re^{tI}$, with I in \mathbb{S} and t in $[0, \pi]$. The restriction of g to $r\mathbb{S}^3$ induces a volume form $dVol_{r\mathbb{S}^3}$. Let f be in $H^2(\mathbb{B})$. Then,

$$\|f\|_{H^2(\mathbb{B})}^2 = \lim_{r \to 1} (1 - r^2) \frac{1}{Vol_{r\mathbb{S}^3}(r\mathbb{S}^3)} \int_{r\mathbb{S}^3} |f|^2 dVol_{r\mathbb{S}^3},$$

a relation similar to the definition of the Hardy norm in the unit disc by means of the Poincaré metric.

In Section 5 we use the Caley map $C : q \mapsto (1-q)^{-1}(1+q)$ to write down the metric g in coordinates living in right half-space $\mathbb{H}^+ := C(\mathbb{B}) = \{q \in \mathbb{H} \mid \operatorname{Re}(q) > 0\}$. This makes it easier to prove a bilateral estimate for the distance function associated with g, Theorem 5.4. As an application, we further investigate the "rigidity" of the metric g, by showing that the only inner functions which are Lipschitz continuous with respect to g have to be slice preserving. A regular function $f : \mathbb{B} \to \mathbb{H}$ is *inner* if (i) it maps \mathbb{B} into \mathbb{B} ; (ii) the limit as $r \to 1$ of f along the radius $r \mapsto ru$ exists for a.e. u in $\partial \mathbb{B}$ and it has unitary norm. A function defined from $\Omega \subseteq \mathbb{H}$ to \mathbb{H} is *slice preserving* if it maps $L_I \cap \Omega$ to L_I for all I in \mathbb{S} .

Theorem 1.4. Let $f : \mathbb{B} \to \mathbb{B}$ be an inner function. Then, f is Lipschitz with respect to the metric g if and only if it is slice preserving. In this case, it is a contraction.

In particular, Lipschitz inner functions have to fix the real diameter of \mathbb{B} .

We also consider in Subsection 5.1 four equivalent definitions of the Hardy space $H^2(\mathbb{H}^+)$ on \mathbb{H}^+ . First, functions f in $H^2(\mathbb{H}^+)$ might be characterized, pretty much as in the one-dimensional complex case, as inverse Fourier transforms -in the quaternionic sense- of functions $F : [0, \infty) \to \mathbb{H}$ with finite L^2 -norm

$$||F||_{L^2} = \left(\int_0^\infty |F(\zeta)|^2 d\zeta\right)^{1/2}.$$

In Proposition 5.1 we show that $H^2(\mathbb{H}^+)$ is the Hilbert space having reproducing kernel $k_w(q) = (q + \overline{w})^{-*}$. This second viewpoint has the advantage of relating $H^2(\mathbb{H}^+)$ and $H^2(\mathbb{B})$. We show in fact in Proposition 5.2 that the reproducing kernel for $H^2(\mathbb{H}^+)$ is a *rescaling* of the reproducing kernel for $H^2(\mathbb{B})$:

$$k_{H^2(\mathbb{B})}(C^{-1}(w), C^{-1}(z)) = \frac{1}{2}(1+z)k_{H^2(\mathbb{H}^+)}(w, z)(1+\overline{w}).$$

Here we use the symbols $k_{H^2(\mathbb{B})}$ and $k_{H^2(\mathbb{H}^+)}$ for the reproducing kernels on \mathbb{B} and \mathbb{H}^+ , respectively. Hence, third, the functions in $H^2(\mathbb{H}^+)$ might be defined as the rescaled versions of functions in $H^2(\mathbb{B})$. In Proposition 5.3 we finally show that, fourth, the norm of f in $H^2(\mathbb{H}^+)$ can be computed as the limit of the integrals of $|f|^2$ on "horocycles" in $H^2(\mathbb{H}^+)$, when these are endowed with the natural volume form induced by the metric g. The space $H^2(\mathbb{H}^+)$ was defined in [2], using a fifth definition, which is shown to give rise to the same reproducing kernel. Our contribution here is mainly relating $H^2(\mathbb{H}^+)$ and the geometry of \mathbb{H}^+ .

There are different extensions of the notion of Hardy space to the quaternionic setting, or to the more general context of Clifford algebras. Let us spend a few words to motivate the one we have here adopted. We might interpret the sequence $\{a_n\}$ as a quaternion valued signal in positive, discrete time $n \ge 0$, and its ℓ^2 norm as a notion of energy, exactly as in the classical case of complex valued signals. The function $f(q) = \sum_{n=0}^{\infty} q^n a_n$ is then the natural notion of generating function (the "q-transform") for the signal. The *-product arises in connection with convolution (or, to put it in the language of signal processing, of linear and time invariant filtering); in general, slice regularity is the analytic constraint satisfied by power series of a quaternionic variable. The geometry studied in this paper is related to the time shift $f(q) \mapsto qf(q)$, the most natural operator arising from the viewpoint of signal theory. In a forthcoming paper, we will return to the interpretation of $H^2(\mathbb{B})$ in terms of signals in more detail.

A different and very interesting extension of the Hardy theory has been developed by Marius Mitrea and other researchers over the past twenty years, relying on a different notion of regular functions. The literature on the topic is vast. We direct the interested reader to the monographs [7] and [18] for an introduction to, respectively, function theory on Clifford algebras and the corresponding harmonic analysis; and to [13], especially the introduction, and the recent article [20], for a comparison between the "Clifford" and the slice viewpoints.

Constructive criticism from an anonimous referee encouraged us to improve Theorem 1.3, which is now stronger than in the previous version of the article, and proved by a different argument.

2 Preliminaries

We recall the definition of slice regularity, together with some basic results that hold for this class of functions. We refer to the book [13] for all details and proofs. Let \mathbb{H} denote the four-dimensional (non-commutative) real algebra of quaternions and let \mathbb{S} denote the two-dimensional sphere of imaginary units of \mathbb{H} , $\mathbb{S} = \{q \in \mathbb{H} \mid q^2 = -1\}$. One can "slice" the space \mathbb{H} in copies of the complex plane that intersect along the real axis,

$$\mathbb{H} = \bigcup_{I \in \mathbb{S}} (\mathbb{R} + \mathbb{R}I), \qquad \mathbb{R} = \bigcap_{I \in \mathbb{S}} (\mathbb{R} + \mathbb{R}I),$$

where $L_I := \mathbb{R} + \mathbb{R}I \cong \mathbb{C}$, for any $I \in \mathbb{S}$. Then, each element $q \in \mathbb{H}$ can be expressed as $q = x + yI_q$, where x, y are real (if $q \in \mathbb{R}$, then y = 0) and I_q is an imaginary unit. Let $\Omega \subseteq \mathbb{H}$ be a subset of \mathbb{H} . For any $I \in S$, we will denote by Ω_I the intersection $\Omega \cap L_I$. We can now recall the definition of slice regular functions, in the sequel simply called *regular* functions.

Definition 2.1. Let Ω be a domain (open connected subset) in \mathbb{H} . A function $f : \Omega \to \mathbb{H}$ is said to be (slice) regular if for any $I \in \mathbb{S}$ the restriction f_I of f to Ω_I has continuous partial derivatives and it is such that

$$\overline{\partial}_I f_I(x+yI) = \frac{1}{2} \left(\frac{\partial}{\partial x} + I \frac{\partial}{\partial y} \right) f_I(x+yI) = 0$$

for all $x + yI \in \Omega_I$.

A wide class of examples of regular functions is given by power series with quaternionic coefficients of the form $\sum_{n=0}^{\infty} q^n a_n$ which converge in open balls centered at the origin.

Theorem 2.2. A function f is regular on $B(0, R) = \{q \in \mathbb{H} \mid |q| < R\}$ if and only if f has a power series expansion $f(q) = \sum_{n=0}^{\infty} q^n a_n$ converging in B(0, R).

For regular functions, it is possible to define an appropriate notion of derivative:

Definition 2.3. Let f be a regular function on a domain $\Omega \subseteq \mathbb{H}$. The slice (or Cullen) derivative of f is the regular function defined as

$$\partial_c f(x+yI) = \frac{1}{2} \left(\frac{\partial}{\partial x} - I \frac{\partial}{\partial y} \right) f_I(x+yI).$$

We will consider domains in certain restricted classes.

Definition 2.4. Let $\Omega \subseteq \mathbb{H}$ be a domain.

- 1. Ω is called a slice domain if it intersects the real axis and if, for any $I \in S$, Ω_I is a domain in L_I .
- 2. Ω is called a symmetric domain if for any point $x + yI \in \Omega$, with $x, y \in \mathbb{R}$ and $I \in S$, the entire *two-sphere* x + yS *is contained in* Ω .

The ball \mathbb{B} and the right half-space $\mathbb{H}^+ = \{q = x + Iy : I \in \mathbb{S}, x > 0, y \in \mathbb{R}\}$ are symmetric slice domains.

Slice regular functions defined on symmetric slice domains have a peculiar property.

Theorem 2.5 (Representation Formula). Let f be a regular function on a symmetric slice domain Ω and let $x + y \mathbb{S} \subset \Omega$. Then, for any $I, J \in \mathbb{S}$,

$$f(x+yJ) = \frac{1}{2}[f(x+yI) + f(x-yI)] + J\frac{I}{2}[f(x-yI) - f(x+yI)].$$

In particular, there exist $b, c \in \mathbb{H}$ such that f(x + yJ) = b + Jc for any $J \in \mathbb{S}$.

When restricted to a sphere of the form x + yS, a regular function is actually affine in the variable q. This nice geometric property leads to the following definition

Definition 2.6. Let f be a regular function on a symmetric slice domain Ω . The spherical derivative of f is defined as

$$\partial_s f(q) = (q - \overline{q})^{-1} \left(f(q) - f(\overline{q}) \right).$$

A basic result that establishes a relation between regular functions and holomorphic functions of one complex variable is the following.

Lemma 2.7 (Splitting Lemma). Let f be a regular function on a slice domain $\Omega \subseteq \mathbb{H}$. Then for any $I \in \mathbb{S}$ and for any $J \in \mathbb{S}$, $J \perp I$ there exist two holomorphic functions $F, G : \Omega_I \to L_I$ such that

$$f(x+yI) = F(x+yI) + G(x+yI)J$$

for any $x + yI \in \Omega_I$.

In general, the pointwise product of functions does not preserve slice regularity. It is possible to introduce a new multiplication operation, which, in the special case of power series, can be defined as follows.

Definition 2.8. Let $f(q) = \sum_{n=0}^{\infty} q^n a_n$, and $g(q) = \sum_{n=0}^{\infty} q^n b_n$ be regular functions on B(0, R). Their regular product (or *-product) is

$$f * g(q) = \sum_{n \ge 0} q^n \sum_{k=0}^n a_k b_{n-k},$$

regular on B(0, R) as well.

The *-product is related to the standard pointwise product by the following formula.

Proposition 2.9. Let f, g be regular functions on a symmetric slice domain Ω . Then

$$f * g(q) = \begin{cases} 0 & \text{if } f(q) = 0\\ f(q)g(f(q)^{-1}qf(q)) & \text{if } f(q) \neq 0 \end{cases}$$

The reciprocal f^{-*} of a regular function f with respect to the *-product can be defined.

Definition 2.10. Let $f(q) = \sum_{n=0}^{\infty} q^n a_n$ be a regular function on B(0, R), $f \neq 0$. Its regular reciprocal *is*

$$f^{-*}(q) = \frac{1}{f * f^c(q)} f^c(q),$$

where $f^c(q) = \sum_{n=0}^{\infty} q^n \overline{a}_n$. The function f^{-*} is regular on $B(0,R) \setminus \{q \in B(0,R) \mid f * f^c(q) = 0\}$ and $f * f^{-*} = 1$ there.

For example, in the case of the *reproducing kernel* for the quaternionic Hardy space $H^2(\mathbb{B})$, we have **Remark 2.11.** The *reproducing kernel* for $H^2(\mathbb{B})$ is

$$k_w(q) = \sum_{n=0}^{\infty} q^n \overline{w}^n = (1 - q\overline{w})^{-*}$$

Then we have a natural definition of regular quotients of regular functions, which satisfy

Proposition 2.12. Let f and g be regular functions on a symmetric slice domain Ω and denote by $Z = \{q \in \Omega \mid f * f^c(q) = 0\}$. If $T_f : \Omega \setminus Z \to \Omega \setminus$ is defined as

$$T_f(q) = f^c(q)^{-1}qf^c(q),$$

then

$$f^{-*} * g(q) = f(T_f(q))^{-1}g(T_f(q))$$
 for every $q \in \Omega \setminus Z_{f^s}$.

Important examples of regular quotients that will appear in the sequel are the *regular Möbius transformations*, of the form

$$M_a(q) = (1 - q\overline{a})^{-*} * (q - a),$$

where $a \in \mathbb{B}$, which are regular self-maps of the quaternionic unit ball \mathbb{B} . After multiplication on the right by unit-norm quaternions, they are the only self-maps of \mathbb{B} which are regular, with regular inverse. They were introduced by Stoppato in [21]. See also [13].

3 Metrics associated with quaternionic reproducing kernel Hilbert spaces

Let $\Omega \subseteq \mathbb{H}$ be a symmetric slice domain and let \mathcal{H} be a reproducing kernel Hilbert space of regular functions on Ω . For the definition and all basic results concerning quaternionic Hilbert spaces see, e.g., [15] and references therein. For the properties we are interested in, the same results hold in quaternion valued Hilbert spaces and complex valued Hilbert spaces, and the proofs are very similar. It is possible to define a metric $\delta_{\mathcal{H}}$ on Ω in terms of the distance between projections of kernel functions in the unit sphere of the Hilbert space \mathcal{H} . Namely, if $k(w, q) = k_w(q)$ denotes the reproducing kernel of \mathcal{H} , then $\delta_{\mathcal{H}} : \Omega \times \Omega \to \mathbb{R}^+$ can be defined as

$$\delta_{\mathcal{H}}(w,z) = \sqrt{1 - \left| \left\langle \frac{k_w}{\|k_w\|_{\mathcal{H}}}, \frac{k_z}{\|k_z\|_{\mathcal{H}}} \right\rangle_{\mathcal{H}} \right|^2}.$$
(4)

Proposition 3.1. Let Ω be a symmetric slice domain and let \mathcal{H} be a reproducing kernel Hilbert space of regular functions on Ω . Let $w \in \Omega \cap L_{I_w}$ and let $d \in \mathbb{H}$ be such that $w + d \in \Omega$. Consider the decomposition $d = d_1 + d_2$, where $d_1 \in L_{I_w}$ and $d_2 \in L_{I_w}^{\perp}$. Then

$$\delta_{\mathcal{H}}^2(w,w+d) = \frac{\|k_w\|_{\mathcal{H}}^2 \left\|\overline{d_1\partial_c k_q(w) + d_2\partial_s k_q(w)}\right\|_{\mathcal{H}}^2 - \left|\left\langle k_w, \overline{d_1\partial_c k_q(w) + d_2\partial_s k_q(w)}\right\rangle_{\mathcal{H}}\right|^2}{\|k_w\|_{\mathcal{H}}^4} + O(|d|^2).$$

Proof. Recalling the definition (4) of $\delta_{\mathcal{H}}$, we get

$$\delta_{\mathcal{H}}^{2}(w, w+d) = \frac{\|k_{w}\|_{\mathcal{H}}^{2} \|k_{w+d}\|_{\mathcal{H}}^{2} - |\langle k_{w}, k_{w+d} \rangle_{\mathcal{H}}|^{2}}{\|k_{w}\|_{\mathcal{H}}^{2} \|k_{w+d}\|_{\mathcal{H}}^{2}}.$$
(5)

We want to have a better description of the numerator of (5). Using the properties of the kernel functions and the fact that regular functions are real analytic functions of 4 real variables, we can write

$$k_{w+d}(q) - k_w(q) = \overline{k_q(w+d) - k_q(w)} = \overline{(k_q)_*[w](d)} + O(|d|^2)$$

where $(k_q)_*[w](d)$ denotes the real differential of k_q at the point w, applied to the vector d. We identify here the tangent space $T_w\Omega$ with \mathbb{H} . Thanks to the decomposition properties of the real differential of regular functions in terms of slice and spherical derivatives, see Remark 8.15 in [13], we have

$$k_{w+d}(q) - k_w(q) = \overline{d_1 \partial_c k_q(w) + d_2 \partial_s k_q(w)} + O(|d|^2),$$

hence,

$$\|k_{w+d}\|_{\mathcal{H}}^2 = \|k_w\|_{\mathcal{H}}^2 + \left\|\overline{d_1\partial_c k_q(w) + d_2\partial_s k_q(w)}\right\|_{\mathcal{H}}^2 + 2\operatorname{Re}\left\langle k_w, \overline{d_1\partial_c k_q(w) + d_2\partial_s k_q(w)}\right\rangle_{\mathcal{H}} + O(|d|^2)$$
and

$$\begin{aligned} |\langle k_w, k_{w+d} \rangle_{\mathcal{H}}|^2 &= \left| \|k_w\|_{\mathcal{H}}^2 + \left\langle k_w, \overline{d_1 \partial_c k_q(w) + d_2 \partial_s k_q(w)} \right\rangle_{\mathcal{H}} \right|^2 + O(|d|^2) \\ &= \|k_w\|_{\mathcal{H}}^4 + \left| \left\langle k_w, \overline{d_1 \partial_c k_q(w) + d_2 \partial_s k_q(w)} \right\rangle_{\mathcal{H}} \right|^2 \\ &+ 2\|k_w\|_{\mathcal{H}}^2 \operatorname{Re} \left\langle k_w, \overline{d_1 \partial_c k_q(w) + d_2 \partial_s k_q(w)} \right\rangle_{\mathcal{H}} + O(|d|^2). \end{aligned}$$

Therefore

$$\delta_{\mathcal{H}}^2(w,w+d) = \frac{\|k_w\|_{\mathcal{H}}^2 \left\|\overline{d_1\partial_c k_q(w) + d_2\partial_s k_q(w)}\right\|_{\mathcal{H}}^2 - \left|\left\langle k_w, \overline{d_1\partial_c k_q(w) + d_2\partial_s k_q(w)}\right\rangle_{\mathcal{H}}\right|^2}{\|k_w\|_{\mathcal{H}}^4} + O(|d|^2).$$

Proposition 3.1 mirrors the complex case situation, see Theorem 1.17 in [10], or formula (0.3) in [17].

In fact the functions $\overline{\partial_c k_q(w)}$ and $\overline{\partial_s k_q(w)}$ are regular with respect to the variable q, and they reproduce respectively the slice and the spherical derivative of any regular function $f : \Omega \to \mathbb{H}$. In fact, for any $w \in \Omega_{I_w}$, if $h \in L_{I_w}$, we can write

$$\partial_c f(w) = \lim_{h \to 0, h \in L_{I_w}} h^{-1}(f(w+h) - f(w)) = \lim_{h \to 0, h \in L_{I_w}} h^{-1}(\langle f, k_{w+h} \rangle_{\mathcal{H}} - \langle f, k_w \rangle_{\mathcal{H}})$$
$$= \lim_{h \to 0, h \in L_{I_w}} h^{-1} \left\langle f, \overline{k_q(w+h) - k_q(w)} \right\rangle_{\mathcal{H}} = \lim_{h \to 0, h \in L_{I_w}} h^{-1} \left\langle f, \overline{h\partial_c k_q(w)} \right\rangle_{\mathcal{H}} = \left\langle f, \overline{\partial_c k_q(w)} \right\rangle_{\mathcal{H}}$$

and

$$\partial_s f(w) = (w - \overline{w})^{-1} (f(w) - f(\overline{w})) = (w - \overline{w})^{-1} \left(\langle f, k_w \rangle_{\mathcal{H}} - \langle f, k_{\overline{w}} \rangle_{\mathcal{H}} \right)$$
$$= (w - \overline{w})^{-1} \left\langle f, k_w - k_{\overline{w}} \rangle_{\mathcal{H}} = (w - \overline{w})^{-1} \left\langle f, \overline{(w - \overline{w})} \partial_s k_q(w) \right\rangle_{\mathcal{H}} = \left\langle f, \overline{\partial_s k_q(w)} \right\rangle_{\mathcal{H}}.$$

Proposition 3.1 allows us to define a Riemannian metric $g_{\mathcal{H}}$ on the symmetric slice domain Ω . For each point $w \in \Omega$, let us identify the tangent space $T_w \Omega$ with $\mathbb{H} = L_{I_w} + L_{I_w}^{\perp}$. Then the length of a tangent vector $d = d_1 + d_2 \in L_{I_w} + L_{I_w}^{\perp}$ is

$$|d|_{g_{\mathcal{H}}(w)}^{2} = \frac{\|k_{w}\|_{\mathcal{H}}^{2} \left\|\overline{d_{1}\partial_{c}k_{q}(w) + d_{2}\partial_{s}k_{q}(w)}\right\|_{\mathcal{H}}^{2} - \left|\left\langle k_{w}, \overline{d_{1}\partial_{c}k_{q}(w) + d_{2}\partial_{s}k_{q}(w)}\right\rangle_{\mathcal{H}}\right|^{2}}{\|k_{w}\|_{\mathcal{H}}^{4}}.$$
 (6)

Theorem 1.2 now follows from specializing Proposition 3.1 to the case of a slice preserving kernel, where formula (6) simplifies.

Proof of Theorem 1.2. We begin by working out the numerator in equation (6). We have:

$$\left\| \overline{d_1 \partial_c k_q(w) + d_2 \partial_s k_q(w)} \right\|_{\mathcal{H}}^2 = \left\langle \overline{d_1 \partial_c k_q(w) + d_2 \partial_s k_q(w)}, \overline{d_1 \partial_c k_q(w) + d_2 \partial_s k_q(w)} \right\rangle_{\mathcal{H}}$$
$$= \left\| \overline{\partial_c k_q(w)} \right\|_{\mathcal{H}}^2 |d_1|^2 + \left\| \overline{\partial_s k_q(w)} \right\|_{\mathcal{H}}^2 |d_2|^2 + 2 \operatorname{Re} \left\langle \overline{d_1 \partial_c k_q(w)}, \overline{d_2 \partial_s k_q(w)} \right\rangle_{\mathcal{H}}$$

and

$$\left|\left\langle k_w, \overline{d_1\partial_c k_q(w) + d_2\partial_s k_q(w)}\right\rangle_{\mathcal{H}}\right|^2 = \left|\overline{d_1\partial_c k_w(w) + d_2\partial_s k_w(w)}\right|^2$$
$$= \left|\overline{\partial_c k_w(w)}\right|^2 |d_1|^2 + \left|\overline{\partial_s k_w(w)}\right|^2 |d_2|^2 + 2\operatorname{Re}\left(\overline{d_1\partial_c k_w(w)}d_2\partial_s k_w(w)\right)$$

Hence we are left to prove that both

$$\operatorname{Re}\left\langle \overline{d_1 \partial_c k_q(w)}, \overline{d_2 \partial_s k_q(w)} \right\rangle_{\mathcal{H}} = \operatorname{Re}\left(d_2 \left\langle \overline{\partial_c k_q(w)}, \overline{\partial_s k_q(w)} \right\rangle_{\mathcal{H}} \overline{d_1} \right)$$

and

$$\operatorname{Re}\left(\overline{d_1\partial_c k_w(w)}d_2\partial_s k_w(w)\right) = \operatorname{Re}\left(d_2\partial_s k_w(w)\overline{\partial_c k_w(w)}\,\overline{d_1}\right)$$

equal zero. Now notice that if k_w maps Ω_{I_w} to L_{I_w} , the same holds true for both $\partial_c k_w$ and $\partial_s k_w$. The fact that $d_1 \in L_{I_w}$ and $d_2 \in L_{I_w}^{\perp}$ leads us to conclude.

The hypothesis about kernel functions required in Theorem 1.2 is satisfied by the quaternionic analogues of Hardy and Bergman spaces; see [1, 9].

4 Invariant metrics associated with the Hardy space $H^2(\mathbb{B})$

In this section, we turn our attention to the special example of the Hardy space $H^2(\mathbb{B})$. We will study the corresponding Riemannian metric $g := g_{H^2(\mathbb{B})}$. Recalling that for any w the kernel function $k_w(q) = \sum_{n=0}^{\infty} q^n \overline{w}^n$ preserves the slice L_{I_w} , we can directly apply Theorem 1.2 to find the expression of g, thus proving the first part of Theorem 1.1.

Proposition 4.1. For any $w \in \mathbb{B}$, let us identify the tangent space $T_w\mathbb{B}$ with \mathbb{H} . For any vector $d \in T_w\mathbb{B}$, if w lies in L_{I_w} and we decompose $d = d_1 + d_2$ with d_1 in L_{I_w} and d_2 in $L_{I_w}^{\perp}$, then the length of d with respect to g is given by

$$|d|_{g(w)}^{2} = \frac{1}{(1-|w|^{2})^{2}}|d_{1}|^{2} + \frac{1}{|1-w^{2}|^{2}}|d_{2}|^{2}.$$
(7)

Proof. The following equalities can, by their nature, be reduced to simple calculations in the complex plane:

$$\|k_w\|_{H^2(\mathbb{B})}^2 = \frac{1}{1 - |w|^2}, \quad \left|\overline{\partial_c k_w(w)}\right|^2 = \frac{|w|^2}{(1 - |w|^2)^4}, \quad \left|\overline{\partial_s k_w(w)}\right|^2 = \frac{|w|^2}{(1 - |w|^2)^2 |1 - w^2|^2},$$
$$\left\|\overline{\partial_c k_q(w)}\right\|_{H^2(\mathbb{B})}^2 = \left\|\sum_{n \ge 0} n^2 q^n \overline{w}^{n-1}\right\|_{H^2(\mathbb{B})}^2 = \frac{1 + |w|^2}{(1 - |w|^2)^3},$$

$$\left\|\overline{\partial_s k_q(w)}\right\|_{H^2(\mathbb{B})}^2 = \frac{1}{|w - \overline{w}|^2} \left(\frac{2}{1 - |w|^2} - \frac{1}{1 - w^2} - \frac{1}{1 - \overline{w}^2}\right).$$

A direct application of Theorem 1.2, then, yields that, with respect to coordinates $(d_1, d_2) \in (L_{I_w}, L_{I_w}^{\perp})$,

$$|d|_{g(w)}^{2} = \frac{1}{(1-|w|^{2})^{2}}|d_{1}|^{2} + \frac{1}{|1-w^{2}|^{2}}|d_{2}|^{2}.$$

The volume form $dVol_g$ associated with the metric g at any point $w = x_0 + x_1i + x_2j + x_3k \in \mathbb{B}$ is then

$$dVol_g(w) = \frac{dVol_{Euc}(w)}{(1 - |w|^2)^2 |1 - w^2|^2},$$

where $dVol_{Euc}(w) = dx_0 dx_1 dx_2 dx_3$ is the usual Euclidean volume element.

Proposition 4.2. Let $\delta := \delta_{H^2(\mathbb{B})}$ be defined as in (4). For any $w, z \in \mathbb{B}$, $\delta(z, w)$ coincides both with the value at z of the regular Möbius transformation M_w associated with w and with the value at w of the regular Möbius transformation M_z associated with z, namely

$$\delta(w,z) = \left| (1 - q\overline{z})^{-*} * (q - z) \right|_{|q=w} = \left| (1 - q\overline{w})^{-*} * (q - w) \right|_{|q=z}.$$

Proof. Let w, z be two points in \mathbb{B} . By Proposition 2.12,

$$|\langle k_w, k_z \rangle_{H^2(\mathbb{B})}| = |k_w(z)| = |(1 - q\overline{w})^{-*}|_{|q=z} = |1 - \hat{z}\overline{w}|^{-1}$$

where $\hat{z} = (1 - zw)^{-1}z(1 - zw)$, which implies

$$\left| \left\langle \frac{k_w}{\|k_w\|_{H^2(\mathbb{B})}}, \frac{k_z}{\|k_z\|_{H^2(\mathbb{B})}} \right\rangle_{H^2(\mathbb{B})} \right|^2 = |1 - \hat{z}\overline{w}|^{-2} \left(1 - |w|^2\right) \left(1 - |z|^2\right).$$

Thus, since $|\hat{z}| = |z|$, we get

$$\begin{split} \delta^{2}(w,z) &= 1 - \left| \left\langle \frac{k_{w}}{\|k_{w}\|_{H^{2}(\mathbb{B})}}, \frac{k_{z}}{\|k_{z}\|_{H^{2}(\mathbb{B})}} \right\rangle_{H^{2}(\mathbb{B})} \right|^{2} \\ &= \left| 1 - \hat{z}\overline{w} \right|^{-2} \left(\left| 1 - \hat{z}\overline{w} \right|^{2} - \left(1 - |w|^{2} \right) \left(1 - |z|^{2} \right) \right) \\ &= \left| 1 - \hat{z}\overline{w} \right|^{-2} \left(\left(1 - \hat{z}\overline{w} \right) \left(1 - w\overline{\hat{z}} \right) - \left(1 - w\overline{w} \right) \left(1 - \hat{z}\overline{\hat{z}} \right) \right) = \left| 1 - \hat{z}\overline{w} \right|^{-2} \left(\hat{z} - w \right) \left(\overline{\hat{z}} - \overline{w} \right) \\ &= \left| 1 - \hat{z}\overline{w} \right|^{-2} \left| \hat{z} - w \right|^{2} = \left| \left(1 - \hat{z}\overline{w} \right)^{-1} \left(\hat{z} - w \right) \right|^{2} = \left| \left(1 - q\overline{w} \right)^{-*} * \left(q - w \right) \right|_{|q=z}^{2} \end{split}$$

where the last equality follows from Proposition 2.12.

	-	-	-	-	
L					
L					
L					

The previous relation between the metric δ (which is the finite version of the metric g) and regular Möbius transformations, is not unexpected. In fact, as studied in [6], the real differential $(M_w)_*$ of the regular Möbius map M_w associated with a point $w \in \mathbb{B}_{I_w}$ acts on L_{I_w} by right multiplication by $(1 - |w|^2)^{-1}$ and on $L_{I_w}^{\perp}$ by right multiplication by $(1 - \overline{w}^2)^{-1}$. Looking at equation (7), we see that the coefficients of the metric g at the point w with respect to coordinates $(L_{I_w}, L_{I_w}^{\perp})$ coincide in modulus with the components of $(M_w)_*$. Moreover, the fact that g(w) measures vectors in L_{I_w} by multiplying their Euclidean length by $\frac{1}{1-|w|^2}$ means that the restriction of g to a slice L_I is the classical Poincaré metric in the unit disc \mathbb{B}_I .

Using spherical coordinates, $\mathbb{B} = \{re^{tI} \mid r \in [0, 1), t \in [0, \pi], I \in \mathbb{S}\}$, if $q = re^{tI}$ and we decompose the lenght element $dq = dq_1 + dq_2 \in L_{I_w} + L_{I_w}^{\perp}$, then, since dI is orthogonal to I (because I is unitary) we have $|d_1|^2 = dr^2 + r^2 dt^2$ and $|d_2|^2 = r^2 \sin^2 t |dI|^2$ where |dI| denotes the usual twodimensional sphere round metric on $\mathbb{S} \cong \mathbb{S}^2$. Therefore we get the expression of the metric tensor ds_g^2 associated with g in spherical coordinates:

$$ds_g^2 = \frac{dr^2 + r^2 dt^2}{(1 - r^2)^2} + \frac{r^2 \sin^2 t |dI|^2}{(1 - r^2)^2 + 4r^2 \sin^2 t}.$$
(8)

That is, g is a *warped product* of the hyperbolic metric g_{hyp} on the complex unit disc with the standard round metric $g_{\mathbb{S}}$ on the two-dimensional sphere [19].

4.1 Isometries and geodesics of (\mathbb{B}, g)

From the expression (7) of g, it is clear that three families of functions act isometrically on (\mathbb{B}, g) :

(a) regular Möbius transformation of the form

$$q \mapsto M_{\lambda}(q) = (1 - q\lambda)^{-*} * (q - \lambda) = \frac{q - \lambda}{1 - q\lambda},$$

with λ in (-1, 1);

(b) isometries of the sphere of imaginary units, which in polar coordinates $r \ge 0, t \in [0, \pi], I \in \mathbb{S}$ read as

$$q = re^{tI} \mapsto T_A(q) = re^{tA(I)},$$

where $A : \mathbb{S} \to \mathbb{S}$ is an isometry of \mathbb{S} ;

(c) the reflection in the imaginary hyperplane,

$$q \mapsto R(q) = -\overline{q}.$$

Our goal is to prove the following classification result, thus proving the second part of Theorem 1.1.

Theorem 4.3. The group Γ of isometries of (\mathbb{B}, g) is generated by maps of type (a), (b) and (c).

The proof requires a few steps. To begin with, we identify three classes of totally geodesic submanifolds of \mathbb{B} , each one related to a class of isometries.

The first family is the one related to isometries of type (a).

Lemma 4.4. For any $I \in \mathbb{S}$, the two-dimensional submanifold of \mathbb{B}

$$\mathbb{B}_{I} = \mathbb{B} \cap L_{I} = \{ re^{tI} \in \mathbb{B} \mid r \in [0, 1), t \in [0, 2\pi] \}$$

is totally geodesic. In particular, for any $I \in S$, \mathbb{B}_I is an hyperbolic disc.

Proof. Fix $I \in \mathbb{S}$ and let g_{hyp}^{I} be the restriction of the metric g to \mathbb{B}_{I} , which is just the classical hyperbolic metric in the unit disc. We will show that each geodesic of $(\mathbb{B}_{I}, g_{hyp}^{I})$ is still a geodesic of (\mathbb{B}, g) . Pick two points w, z in \mathbb{B}_{I} and let γ be the (hyperbolic) geodesic in \mathbb{B}_{I} joining w with z, and $\alpha(\tau) = r(\tau) (\cos(t(\tau)) + \sin(t(\tau))I(\tau))$ be a parametrized curve which joins $w = \alpha(\tau_{0})$ with $z = \alpha(\tau_{1})$. If $\pi_{I}(\alpha)$ denotes the piecewise regular curve obtained by projecting α on \mathbb{B}_{I} ,

$$\pi_I(\alpha)(\tau) = r(\tau) \left(\cos(t(\tau)) + \sin(t(\tau))I \right),$$

since |dI| is orthogonal to \mathbb{B}_I , we conclude

$$\operatorname{length}(\alpha) \geq \operatorname{length}(\pi_I(\alpha)) \geq \operatorname{length}(\gamma).$$

		1

The second family of totally geodesic submanifolds is related to isometries of type (b). For any $I \in \mathbb{S}$, we denote by $\mathcal{C}(I)$ the great circle obtained intersecting \mathbb{S} with the plane L_I^{\perp} .

Lemma 4.5. For any $J \in \mathbb{S}$, the three-dimensional submanifold of \mathbb{B}

$$\mathbb{B}(\mathcal{C}(J)) = \{ re^{tI} \in \mathbb{B} \mid r \in [0,1), t \in [0,\pi], I \in \mathcal{C}(J) \}$$

is totally geodesic.

Proof. We will prove the statement by showing that the imaginary units identified by all points lying on a same geodesic always belong to the same great circle of S. More precisely, let $\gamma(\tau) = r(\tau) \left(\cos(t(\tau)) + \sin(t(\tau))I(\tau)\right)$ be a parametrized geodesic of (\mathbb{B}, g) such that

$$\begin{cases} \gamma(\tau_0) = x_0 + y_0 I_0 \\ \gamma'(\tau_0) = v_0 + w_0 J_0 \end{cases}$$

We want to show that, for any τ , the imaginary unit $I(\tau)$ of $\gamma(\tau)$ belongs to the great circle of S identified by I_0 and J_0 , namely that, for any τ , $I(\tau) \in \mathcal{C} := \mathcal{C}(I_0 \times J_0)$. Let $\psi : \mathbb{S} \to \mathbb{S}$ be the reflection of S with respect to \mathcal{C} . Then the curve $\tilde{\gamma}(\tau) = r(\tau)(\cos(t(\tau)) + \sin(t(\tau))\psi(I(\tau)))$ is a geodesic of (\mathbb{B}, g) such that

$$\begin{bmatrix} \tilde{\gamma}(\tau_0) = \gamma(\tau_0) \\ \tilde{\gamma}'(\tau_0) = \gamma'(\tau_0) \end{bmatrix}$$

since ψ fixes I_0 and J_0 . By the uniqueness of geodesics with assigned initial conditions, we get that $\tilde{\gamma}(\tau) = \gamma(\tau)$ and hence that ψ fixes $I(\tau)$ for any τ . Therefore we conclude that $I(\tau) \in C$ for any $\tau \in I$.

The third totally geodesic submanifold is the one related to the last class of isometries, type (c).

Lemma 4.6. *The three-dimensional submanifold of* \mathbb{B}

$$\mathbb{B}(\pi/2) = \{ re^{tI} \in \mathbb{B} \mid r \in [0,1), t = \pi/2, I \in \mathbb{S} \} = \{ rI \mid r \in [0,1), I \in \mathbb{S} \}$$

is totally geodesic.

Proof. The statement can be proven following the line of the proof of Lemma 4.5. The ingredients are the fact that the map $R : \mathbb{B} \to \mathbb{B}$, $q \mapsto -\overline{q}$ is an isometry which fixes (punctually) $\mathbb{B}(\pi/2)$, and the uniqueess of geodesics with assigned initial conditions.

Considering the intersection of totally geodesic submanifolds of type $\mathbb{B}(C(J))$ with $\mathbb{B}(\pi/2)$ allows us to identify another family of totally geodesic submanifolds of \mathbb{B} .

Corollary 4.7. Let C(J) be a great circle in S. Then the two-dimensional submanifold $D(\pi/2, C(J)) \subset \mathbb{B}(\pi/2)$, defined as

$$D(\pi/2, \mathcal{C}(J)) = \{ re^{tI} \in \mathbb{B} \mid r \in [0, 1), t = \pi/2, I \in \mathcal{C}(J) \} = \{ rI \in \mathbb{B} \mid r \in [0, 1), I \in \mathcal{C}(J) \},\$$

is totally geodesic.

Remark 4.8. Notice that for the two-dimensional submanifold $D(\pi/2, C(J))$ the following orthogonality relation holds:

$$D(\pi/2, \mathcal{C}(J)) \cap \mathbb{B}_J = \{0\} \text{ and } T_0 D(\pi/2, \mathcal{C}(J)) = T_0 \mathbb{B}_J^{\perp}$$

Moreover, applying Möbius maps of the form M_{λ} to $D(\pi/2, C(J))$, we can extend the orthogonality relation from the origin to all points in $\mathbb{B} \cap \mathbb{R}$. In this way we obtain a family of totally geodesic submanifolds

$$D(t, \mathcal{C}(J)) = M_{\lambda(t)} \left(D(\pi/2, \mathcal{C}(J)) \right)$$

that, for $t \in [0, \pi]$ and $J \in \mathbb{S}/\{\pm 1\}$, defines a foliation of the manifold \mathbb{B} .

In order to have some understanding of the (global) behavior of the metric g, let us investigate some metric properties of the discs of the type $D\left(\frac{\pi}{2}, \mathcal{C}(J)\right)$. Since the imaginary units taken into account belong to $\mathcal{C}(J) \cong \mathbb{S}^1$, we can change coordinates, setting $I = e^{i\theta}$ and $|dI| = d\theta$, so that the metric g, on $D\left(\frac{\pi}{2}, \mathcal{C}(J)\right)$, reduces to

$$ds_D^2 = \frac{dr^2}{(1-r^2)^2} + \frac{r^2 d\theta^2}{(1+r^2)^2}.$$

It is actually convenient to parametrize $D\left(\frac{\pi}{2}, \mathcal{C}(J)\right) \subset \mathbb{I} \cong \mathbb{R}^3$ as a surface of revolution of the form $(\Phi(\rho), \Psi(\rho) \cos \theta, \Psi(\rho) \sin \theta)$, where ρ is the arc length of the generating curve. Setting

$$\rho = \rho(r) = \frac{1}{2} \log \frac{1+r}{1-r},$$

we get

$$\frac{dr^2}{(1-r^2)^2} = d\rho^2$$
 and $\frac{r^2}{(1+r^2)^2} = \frac{1}{4} \tanh^2(2\rho)$

and hence, in coordinates (ρ, θ) , we get that the metric is expressed as

$$ds_D^2 = d\rho^2 + \frac{1}{4} \tanh^2(2\rho) d\theta^2 = d\rho^2 + \Psi^2(\rho) d\theta^2.$$
(9)

Remark 4.9. The Gaussian curvature K of the two-dimensional submanifold $D\left(\frac{\pi}{2}, C(J)\right)$ is positive. In fact, see e.g. [12], with respect to coordinates (ρ, θ) it can be computed as

$$K = \frac{-\Psi''(\rho)}{\Psi(\rho)}$$

which is a non-negative quantity since $\Psi(\rho) = \frac{1}{2} \tanh(2\rho) \ge 0$ and $\Psi''(\rho) \le 0$. This in particular implies that the sectional curvature of (\mathbb{B}, g) is positive on all sections $D\left(\frac{\pi}{2}, \mathcal{C}(J)\right)$, while it is negative on all slices \mathbb{B}_I .

It is possible to study geodesics of $D\left(\frac{\pi}{2}, \mathcal{C}(J)\right)$ by means of the Euler-Lagrange equations

$$\begin{cases} \frac{\partial}{\partial \theta} L = \frac{d}{dt} \frac{\partial}{\partial \dot{\theta}} L\\ \frac{\partial}{\partial \rho} L = \frac{d}{dt} \frac{\partial}{\partial \dot{\rho}} L\end{cases}$$

associated with the Lagrangian

$$L(\rho,\theta,\dot{\rho},\dot{\theta},\tau) = \frac{1}{2} \left(\dot{\rho}^2 + \frac{\tanh^2(2\rho)}{4} \dot{\theta}^2 \right),$$

namely

$$\begin{cases} 0 = \frac{d}{dt} \left(\frac{\tanh^2(2\rho)}{4} \dot{\theta} \right) \\ \tanh(2\rho) \frac{1-\tanh^2(2\rho)}{4} \dot{\theta}^2 = \ddot{\rho}. \end{cases}$$
(10)

The first equation in (10) yields

$$\frac{\tanh^2(2\rho)}{4}\dot{\theta} = A$$

for some constant A. If A = 0, we get $\dot{\theta} = 0$ and hence the second equation in (10) implies that $\ddot{\rho} = 0$. If otherwise $A \neq 0$ we get $\dot{\theta} = \frac{4A}{\tanh^2(2\rho)}$ which implies $|\dot{\theta}| > 4|A|$. Therefore all generating curves, with $\dot{\theta} = 0$, are geodesics of D. Which is not surprising since they correspond to radii $\gamma(r) = re^{\frac{\pi}{2}I}$ for $I \in C(J)$. The other important fact that arises is that for any point $q \in D(\frac{\pi}{2}, C(J)) \setminus \{0\}$ any geodesic corresponding to $A \neq 0$ intersects in finite time the "radial" geodesic through q. This leads to the following result.

Lemma 4.10. Let $J \in \mathbb{S}$. For any $q \in D\left(\frac{\pi}{2}, \mathcal{C}(J)\right)$ such that $q \neq 0$, the injectivity radius at q is finite. On the other hand, the injectivity radius at q = 0 is infinite.

This important metric property of the point q = 0 is useful to classify the isometries of (\mathbb{B}, g) . First of all it tells us that isometries map the real diameter of \mathbb{B} to itself.

Lemma 4.11. Let Γ be the group of isometries of (\mathbb{B}, g) . Then, for any $\phi \in \Gamma$, $\phi(\mathbb{B} \cap \mathbb{R}) = \mathbb{B} \cap \mathbb{R}$.

Proof. Consider first q = 0. Since the injectivity radius at q = 0 is infinite, then, for any $\phi \in \Gamma$, the injectivity radius at $\phi(0)$ is infinite as well. By post-composing ϕ with a regular Möbius transformation of type $(a) \ M_{\lambda}$ we can map 0 to $D\left(\frac{\pi}{2}, C(J)\right)$ (for some $J \in \mathbb{S}$) and hence Lemma 4.10 yields that $M_{\lambda}(\phi(0)) = 0$. Since M_{λ} preserves the real diameter of \mathbb{B} , we get that $\phi(0) \in \mathbb{R}$. To conclude, notice that we can map each point of $\mathbb{B} \cap \mathbb{R}$ to 0 by means of a regular Möbius map of type (a).

We can finally prove the Classification Theorem for isometries of (\mathbb{B}, g) .

Proof of Theorem 4.3. Let $\Phi \in \Gamma$ be an isometry of (\mathbb{B}, g) . Up to composition with a regular Möbius transformation of type (a) and with the map $R : q \mapsto -\overline{q}$, we can suppose that $\Phi(0) = 0$ and that, by Lemma 4.11, $\Phi(\mathbb{B} \cap \mathbb{R}^+) = \mathbb{B} \cap \mathbb{R}^+$.

We now show that Φ fixes $\mathbb{B}(\pi/2)$. Set $\tilde{B}(\pi/2) = \Phi(\mathbb{B}(\pi/2))$. Since Φ is an isometry, Lemma 4.6 implies that $\tilde{B}(\pi/2)$ is a totally geodesic submanifold of \mathbb{B} . Moreover, since $\Phi(0) = 0$, since the geodesics starting at 0 lie on slices, and since, by Lemma 4.4, the slices carry the usual hyperbolic-Poincaré metric: we have that Φ maps radii $\gamma_I(r) = re^{\frac{\pi}{2}I}$ to radii of the form $\Phi(\gamma_I(r)) = re^{\theta(I)\psi(I)}$ with $\theta(I) \in [0, \pi]$, and $\psi(I) \in \mathbb{S}$. Let us show that θ is actually constant on \mathbb{S} .

If d_g denotes the distance function on \mathbb{B} associated with g, recalling equation (8), on the one hand we have

$$d_{g}\left(\Phi(\gamma_{I_{1}}(r)), \Phi(\gamma_{I_{2}}(r))\right) = d_{g}\left(\gamma_{I_{1}}(r), \gamma_{I_{2}}(r)\right) = d_{g}\left(re^{\frac{\pi}{2}I_{1}}, re^{\frac{\pi}{2}I_{2}}\right)$$

$$\leq \frac{r}{1+r^{2}}d_{\mathbb{S}}(I_{1}, I_{2}) \xrightarrow{r \to 1} \frac{1}{2}d_{\mathbb{S}}(I_{1}, I_{2}), \qquad (11)$$

where $d_{\mathbb{S}}$ denotes the usual spherical distance on the unit sphere \mathbb{S} . In particular we deduce that

 $d_g\left(\Phi(\gamma_{I_1}(r)), \Phi(\gamma_{I_2}(r))\right)$

is bounded as a function of r. On the other hand, if $\alpha(\tau) = r(\tau)e^{t(\tau)I(\tau)}$ is a parametrized geodesic joining $\alpha(\tau_1) = \Phi(\gamma_{I_1}(r))$ and $\alpha(\tau_2) = \Phi(\gamma_{I_2}(r))$, we have

$$d_{g}\left(\Phi(\gamma_{I_{1}}(r)), \Phi(\gamma_{I_{2}}(r))\right) = d_{g}\left(re^{\theta(I_{1})\psi(I_{1})}, re^{\theta(I_{2})\psi(I_{2})}\right) = \text{length}\left(\alpha([\tau_{1}, \tau_{2}])\right)$$
$$= \int_{\tau_{1}}^{\tau_{2}} \sqrt{\frac{r'(\tau)^{2} + r^{2}t'(\tau)^{2}}{1 - r(\tau)^{2}}} + \frac{r(\tau)^{2}\sin^{2}(t(\tau))I'(\tau)^{2}}{(1 - r(\tau)^{2})^{2} + 4r(\tau)^{2}\sin^{2}(t(\tau))}} d\tau$$
$$\geq \int_{\tau_{1}}^{\tau_{2}} \sqrt{\frac{r'(\tau)^{2} + r(\tau)^{2}t'(\tau)^{2}}{1 - r(\tau)^{2}}} d\tau \geq d_{hyp}(re^{\theta(I_{1})I}, re^{\theta(I_{2})I})$$

where *I* is any fixed imaginary unit and d_{hyp} the hyperbolic distance associated to the restriction g_{hyp}^{I} of the metric *g* to $\mathbb{B} \cap L_{I}$. If, by contradiction, $\theta(I_{1}) \neq \theta(I_{2})$, then the distance $d_{hyp}(re^{\theta(I_{1})I}, re^{\theta(I_{2})I})$ tends to infinity as *r* goes to 1, contradicting (11). Then, $\theta(I_{1}) = t(\tau_{1}) = t(\tau_{2}) = \theta(I_{2})$: θ is constant on \mathbb{S} .

Therefore we have that $\tilde{B}(\pi/2)$ is ruled by radii of the form $\tilde{\gamma}(r) = re^{t_0 I}$ for some constant t_0 . If $t_0 = \pi/2$, then we are done. Suppose then $t_0 \neq \pi/2$. Since $\tilde{B}(\pi/2)$ and $\mathbb{B}(\pi/2)$ intersect at 0 and they are three-dimensional submanifolds in \mathbb{H} , the intersection V of their respective tangent spaces at 0 must have dimension 2 or 3. Let v be a vector in V and let $r \mapsto re^{Jt}$ be the reparametrized geodesic with initial velocity v. The geodesic lies on both $\tilde{B}(\pi/2)$ and $\mathbb{B}(\pi/2)$, hence $t_0 = t = \pi/2$. (A different proof consists in showing that, if $t_0 \neq \pi/2$, then $\tilde{B}(\pi/2)$ is not smooth at the origin).

The next step is to show that the restriction of Φ to $\mathbb{B}(\pi/2)$ is an isometry T_A of type (b) for some isometry A of the sphere S. We have that

$$d_g\left(\Phi(\gamma_{I_1}(r)), \Phi(\gamma_{I_2}(r))\right) = d_g\left(re^{\frac{\pi}{2}\psi(I_1)}, re^{\frac{\pi}{2}\psi(I_2)}\right).$$
(12)

We now prove an improvement of (11).

Lemma 4.12.

$$\lim_{r \to 1} d_g \left(r e^{\frac{\pi}{2}I_1}, r e^{\frac{\pi}{2}I_2} \right) = \frac{1}{2} d_{\mathbb{S}}(I_1, I_2).$$

Proof of the lemma. Only the case $I_1 \neq I_2$ is interesting. Let $D(\pi/2, \mathcal{C}(J))$ be the two-dimensional manifold introduced in Lemma 4.7 which contains the reparametrized geodesics $r \mapsto re^{\frac{\pi}{2}I_j}$, j = 1, 2. The metric g restricted to the totally geodesic surface $D(\pi/2, \mathcal{C}(J))$ was discussed earlier in this subsection, where we gave it the expression (9). Since $\Psi'(\rho) = 1/\cosh(2\rho) \leq 1$, the surface can be isometrically imbedded as a surface S in \mathbb{R}^3 , with parametric equations $(u(s, \theta), v(s, \theta), z(s, \theta)) = \chi(s, \theta)$, where:

$$\begin{cases} u = p(s)\cos(\theta); \\ v = p(s)\sin(\theta); \\ z = s. \end{cases}$$

Here $s \ge 0$, $\theta \in [-\pi, \pi]$ and $p: [0, +\infty) \to [0, 1/2)$ is a smooth, increasing function such that p(0) = 0and $\lim_{s\to\infty} p(s) = 1/2$. The relationship between p and Ψ is the following: if $\int_0^s \sqrt{p'(\sigma)^2 + 1} d\sigma = \rho$, then $p(s) = \psi(\rho)$. Now, $r = constant \to 1$ corresponds to $s = constant \to \infty$, and the choice of I_1 and I_2 corresponds to a choice of θ_1 and θ_2 . Let k be the metric on the surface. It is elementary that

$$\lim_{s \to \infty} d_k(\chi(s, \theta_1), \chi(s, \theta_2)) = \frac{1}{2} d_{\mathbb{S}^1}(\theta_1, \theta_2)$$

is one-half the distance between θ_1 and θ_2 on the unit circle, which is the same as one-half the distance between I_1 and I_2 in \mathbb{S} .

Equations (11) and (12) together with Lemma 4.12 imply that that $\psi : \mathbb{S} \to \mathbb{S}$ is an isometry of the sphere \mathbb{S} , i.e. $\Phi|_{\mathbb{B}(\pi/2)} = T_{\psi}|_{\mathbb{B}(\pi/2)}$. In conclusion, $T_{\psi}^{-1} \circ \Phi$ is an isometry that fixes $\mathbb{R} \cap \mathbb{B}$ and $\mathbb{B}(\pi/2)$ and hence its (real) differential at the origin $(T_{\psi}^{-1} \circ \Phi)_*[0] : T_0\mathbb{B} \to T_0\mathbb{B}$ is the identity map. Therefore $T_{\psi}^{-1} \circ \Phi$ is the identity map as well and the theorem is proved.

4.2 Volume form on the boundary of \mathbb{B}

If we restrict the metric g to a three-dimensional sphere rS^3 of radius r, in spherical coordinates we get

$$ds_{r\mathbb{S}^3}^2 = \frac{r^2}{(1-r^2)^2} dt^2 + \frac{r^2 \sin^2 t}{(1-r^2)^2 + 4r^2 \sin^2(t)} |dI|^2$$

whose corresponding volume form is

$$dVol_{rS^{3}}(re^{tI}) = \frac{r^{3}\sin^{2}t}{(1-r^{2})((1-r^{2})^{2}+4r^{2}\sin^{2}(t))}dtdA_{S}(I)$$

where $dA_{\mathbb{S}}$ denotes the area element of the two-dimensional sphere \mathbb{S} . This volume form (after a normalization) induces a volume form on the boundary \mathbb{S}^3 of the unit ball: if $u = e^{sJ} \in \mathbb{S}^3$, we have

$$dVol_{\mathbb{S}^3}(u) := \lim_{r \to 1^-} (1 - r^2) dVol_{r\mathbb{S}^3}(ru) = \lim_{r \to 1^-} \frac{(1 - r^2)r^3 \sin^2 s}{(1 - r^2)((1 - r^2)^2 + 4r^2 \sin^2(s))} dt dA_{\mathbb{S}}(I)$$
$$= \frac{1}{4} dt dA_{\mathbb{S}}(I).$$

Notice that $dVol_{\mathbb{S}^3}$ is the product of the usual spherical metric on the two-dimensional sphere \mathbb{S} with the metric dt on circles \mathbb{S}^3_I which appears in the definition of Hardy spaces given in [11]. Moreover in [11] it is proven that any $f \in H^2(\mathbb{B})$ has radial limit along almost any radius and hence, denoting (with a slight abuse of notation) the radial limit by f itself, we have

Proposition 4.13. *If* $f \in H^2(\mathbb{B})$ *, then*

$$\int_{\mathbb{S}^3} |f(u)|^2 dVol_{\mathbb{S}^3}(u) = \frac{1}{8} \int_{\mathbb{S}} \int_0^{2\pi} |f(e^{tI})|^2 dt dA_{\mathbb{S}}(I) = \frac{\pi}{2} ||f||^2_{H^2(\mathbb{B})}.$$

An analogous relation holds for the Hardy space on the right half space. See Section 5.

4.3 **Proof of Theorem 1.3**

Let $a \in \mathbb{B}$, $a \notin \mathbb{R}$, and consider the map $\phi(q) = (1 - q\bar{a})^{-*} * (a - q)$. Then, we can write

$$\phi(q) = \left(\sum_{n=0}^{\infty} q^n \overline{a}^n\right) * (a-q) = \sum_{n=1}^{\infty} q^n \overline{a}^{n-1}(|a|^2 - 1) + a.$$

Taking into account that q and \bar{q} commute and manipulating geometric series, we find expressions for the spherical derivative of ϕ :

$$\partial_s \phi(q) = (q - \bar{q})^{-1} \sum_{n=1}^{\infty} (q^n - \bar{q}^n) \bar{a}^{n-1} (|a|^2 - 1)$$

= $\sum_{n=1}^{\infty} q^{n-1} \sum_{k=0}^{n-1} (q^{-1}\bar{q})^k \bar{a}^{n-1} (|a|^2 - 1) = \sum_{k=0}^{\infty} \bar{q}^k \sum_{m=0}^{\infty} q^m \bar{a}^m \bar{a}^k (|a|^2 - 1),$

and for the Cullen derivative of ϕ :

$$\partial_c \phi(q) = \sum_{n=1}^{\infty} nq^{n-1} \bar{a}^{n-1} (|a|^2 - 1).$$

Thus,

$$\partial_c \phi(0) = \partial_s \phi(0) = |a|^2 - 1$$

and (see also Remark 4 in [6])

$$\partial_c \phi(a) = (|a|^2 - 1)^{-1}$$
 and $\partial_s \phi(a) = (\bar{a}^2 - 1)^{-1}$.

Consider now a tangent vector $d \in T_0 \mathbb{B}$ (identified with the space of quaternions) such that $d \in L_{I_a}^{\perp}$ (where L_{I_a} is the unique slice containing the point *a*). Recalling the decomposition of the real differential of a regular function in terms of spherical and slice derivatives, we have

$$\phi_*[0](d) = d\partial_s \phi(0) = d(|a|^2 - 1) \in T_a \mathbb{B} \cap L_{I_a}^{\perp},$$

and, if we apply to this new tangent vector the real differential of ϕ at $a, \phi_*[a] : T_a \mathbb{B} \to T_0 \mathbb{B}$, we get

$$\phi_*[a](\phi_*[0](d)) = d \frac{1 - |a|^2}{1 - \bar{a}^2}.$$

Since $d \in T_0 \mathbb{B} \cap L_{I_a}^{\perp}$ and $(\bar{a}^2 - 1) \in L_{I_a}$, the vector $\phi_*[a](\phi_*[0](d))$ lies in $T_0 \mathbb{B} \cap L_{I_a}^{\perp}$. Now, $\left|\frac{1-|a|^2}{1-\bar{a}^2}\right| < 1$ if $a \in \mathbb{B} \setminus \mathbb{R}$. Let B_e be the unit disc in $T_0 \mathbb{B} \cap L_{I_a}^{\perp}$, with respect to the Euclidean metric induced by the identification of $T_0 \mathbb{B}$ and \mathbb{H} . The linear map $\phi_*[a] \circ \phi_*[0]$ maps B_e into a proper subset of itself, hence it can not be isometric.

5 Metric in the right half space \mathbb{H}^+

Consider the right half space $\mathbb{H}^+ = \{q \in \mathbb{H} \mid \operatorname{Re}(q) > 0\}$. The Cayley map $C : \mathbb{B} \to \mathbb{H}^+$,

$$C(q) = (1-q)^{-1}(1+q),$$

is a regular bijection from the quaternionic unit ball onto the quaternionic right half space with regular inverse the function $C^{-1}: \mathbb{H}^+ \to \mathbb{B}$,

$$C^{-1}(q) = (1+q)^{-1}(q-1).$$

The aim of this section is to study the image (\mathbb{H}^+, h) of (\mathbb{B}, g) under the map C, where h is the pullback of the metric g by the map C^{-1} . In the introduction we labeled g and h by the same letter, since C is by definition an isometry from (\mathbb{B}, g) to (\mathbb{H}^+, h) . Let $u \in \mathbb{H}^+$ and let $v = v_1 + v_2$ be a tangent vector in $T_u \mathbb{H}^+ \cong L_{I_u} + L_{I_u}^{\perp}$. The length of v with respect to h is

$$|v|_{h(u)} = \left| (C^{-1})_*[u](v) \right|_{g(C^{-1}(u))}$$

where $(C^{-1})_*[u]$ is the real differential of C^{-1} at the point $u \in \mathbb{H}^+$. Recalling the decomposition of the real differential of a regular function in terms of its slice and spherical derivatives, if $v = v_1 + v_2 \in L_{I_u} + L_{I_u}^{\perp}$ we can write

$$(C^{-1})_*[u](v) = v_1 \partial_c C^{-1}(u) + v_2 \partial_s C^{-1}(u) = v_1 \frac{2}{(1+u)^2} + v_2 \frac{2}{|1+u|^2}$$

Hence $(C^{-1})_*[u]$ preserves the decomposition $T_u \mathbb{H}^+ = L_{I_u} + L_{I_u}^{\perp}$ and we get

$$\begin{aligned} |v|_{h(u)}^{2} &= \frac{1}{(1 - |C^{-1}(u)|^{2})^{2}} \frac{4}{|1 + u|^{4}} |v_{1}|^{2} + \frac{1}{|1 - C^{-1}(u)^{2}|^{2}} \frac{4}{|1 + u|^{4}} |v_{2}|^{2} \\ &= \frac{1}{4 \operatorname{Re}(u)^{2}} |v_{1}|^{2} + \frac{1}{4|u|^{2}} |v_{2}|^{2}. \end{aligned}$$

If $v \in L_{I_u}$ then its length is, not surprisingly, the hyperbolic length in the hyperbolic half plane $\mathbb{H}_{I_u}^+ = \{x + yI_u | x > 0, y \in \mathbb{R}\}$. Notice that C maps \mathbb{B}_I to \mathbb{H}_I^+ for any $I \in \mathbb{S}$ and it maps the totally geodesic submanifold $\mathbb{B}(\pi/2)$ to $\mathbb{H}^+(\pi/2) := C(\mathbb{B}(\pi/2)) = \{q \in \mathbb{H}^+ | |q| = 1\}$ i.e. the right half of the three-dimensional unit sphere \mathbb{S}^3 . Then it is not difficult to verify that the isometry group of (\mathbb{H}^+, h) is generated by the images under C of isometry of (\mathbb{B}, g) of type (a), (b) and (c),

(a') linear maps preserving the positive real half-axis,

$$q \mapsto q\lambda$$
,

with $\lambda > 0$;

(b') isometries of the sphere of the imaginary units, which in polar coordinates $r \ge 0, t \in [0, \pi/2), I \in \mathbb{S}$ read as

$$q = re^{tI} \mapsto T_A(q) = re^{tA(I)},$$

where $A : \mathbb{S} \to \mathbb{S}$ is an isometry of \mathbb{S} ;

(c') the inversion in the three-dimensional unit (half) sphere

$$q \mapsto \frac{1}{\overline{q}}$$

Acting on $\mathbb{H}^+(\pi/2)$ by means of isometries of type (a') we obtain totally geodesic regions of the form $\{q \in \mathbb{H}^+ | |q| = R\}$ for R > 0, that can be sliced in totally geodesic two-dimensional submanifolds, corresponding to submanifolds of type D(t, C(J)) in the ball case.

In this setting it is possible to introduce *horocycles*, i.e. hyperplanes of points with constant real part, $H_c = \{q \in \mathbb{H}^+ | \operatorname{Re}(q) = c\}$ for some constant c > 0. They deserve the name of horocycles because their intersection with each slice L_I is a proper horocycle in the hyperbolic half plane \mathbb{H}_I^+ . Isometries of type (a') map horocycles one into another.

If we restrict the metric h to horocycles we obtain that the length of a vector $v = v_1 + v_2 \in T_u H_c \cong \mathbb{R}I_u + L_{I_u}^{\perp}$ tangent to the horocycle H_c at the point $u \in H_c$, can be written as

$$|v|_{H_c}^2 = \frac{1}{4c^2}|v_1|^2 + \frac{1}{4(c^2 + |\operatorname{Im}(u)|^2)}|v_2|^2$$

and the corresponding volume form at $u = c + x_1 i + x_2 j + x_3 k$ is

$$dVol_{H_c}(u) = \frac{dVol_{Euc}(u)}{8c(c^2 + |\operatorname{Im}(u)|^2)},$$

(since the component in L_{I_u} is one-dimensional) where $dVol_{Euc}(u) = dx_1 dx_2 dx_3$ is the standard Euclidean volume element. If we want to define a (non-degenerate) volume form $dVol_{\partial\mathbb{H}^+}$ on the boundary $\partial\mathbb{H}^+$ of the quaternionic right half space we can not directly take the limit of $dVol_{H_c}$ as c approaches 0, we need indeed first to normalize it. For any $u \in \partial\mathbb{H}^+$, we define

$$dVol_{\partial \mathbb{H}^+}(u) := \lim_{c \to 0^+} c \left(dVol_{H_c}(u+c) \right) = \lim_{c \to 0^+} \frac{dVol_{Euc}(u+c)}{8(c^2 + |\operatorname{Im}(u+c)|^2)} = \frac{dVol_{Euc}(u)}{8|\operatorname{Im}(u)|^2} = \frac{dVol_{Euc}(u)}{8|u|^2}.$$
(13)

5.1 Hardy space on \mathbb{H}^+

We will show that, as in the case of the metric g on \mathbb{B} , the invariant metric h on \mathbb{H}^+ introduced in the previous section and in particular the corresponding volume form (13), is related with the quaternionic Hardy space on the right half space \mathbb{H}^+ . It is possible to define the Hardy space $H^2(\mathbb{H}^+)$ on \mathbb{H}^+ as the space of regular functions $f: \mathbb{H}^+ \to \mathbb{H}$ of the form,

$$f(q) = \int_0^{+\infty} e^{-\zeta q} F(\zeta) d\zeta, \qquad (14)$$

with $F : \mathbb{R}^+ \to \mathbb{H}$, such that

$$||f||_{H^2(\mathbb{H}^+)}^2 := \int_0^{+\infty} |F(\zeta)|^2 d\zeta < +\infty.$$
(15)

Proposition 5.1. The quaternionic Hilbert space defined by (15) has reproducing kernel

$$k(q,w) = (q + \overline{w})^{-*}.$$

Proof. Let w be a point in \mathbb{H}^+ . The reproducing function of $H^2(\mathbb{H}^+)$ at w, if it exists, has to be a function

$$k(q,w) = k_w(q) = \int_0^{+\infty} e^{-\zeta q} G(\zeta) d\zeta$$

where $G: \mathbb{R}^+ \to \mathbb{H}$ belongs to $L^2(\mathbb{R}^+)$ and it satisfies

$$f(w) = \langle f, k_w \rangle_{H^2(\mathbb{H}^+)} = \int_0^{+\infty} \overline{G(\zeta)} F(\zeta) d\zeta = \int_0^{+\infty} e^{-\zeta w} F(\zeta) d\zeta.$$

Hence $\overline{G(\zeta)} = e^{-\zeta w}$, which implies $G(\zeta) = e^{-\zeta \bar{w}}$. That is, the kernel function is

$$k_w(q) = \int_0^{+\infty} e^{-\zeta q} e^{-\zeta \bar{w}} d\zeta.$$

To obtain a closed expression of $k_w(q)$, let first q be a (positive) real number. In this case q commutes with all points in \mathbb{H}^+ and we can write

$$k_w(q) = \int_0^{+\infty} e^{-\zeta(q+\overline{w})} d\zeta = \frac{1}{q+\overline{w}}$$

Consider now the function $q \mapsto (q + \overline{w})^{-*}$ (here the regular reciprocal is defined with a slight generalization of Definition 2.10, see [13]). This function is regular and it coincides with $q \mapsto (q + \overline{w})^{-1}$ on real numbers. Thanks to the Identity Principle for regular functions, Theorem 1.12 in [13], we obtain that the reproducing kernel is $k_w(q) = (w + \overline{q})^{-*} = \int_0^{+\infty} e^{-\zeta w} e^{-\zeta \overline{q}} d\zeta$.

Another way to obtain the reproducing kernel on $H^2(\mathbb{H}^+)$ is the following.

Proposition 5.2. Denote by $k_{H^2(\mathbb{B})}$ and by $k_{H^2(\mathbb{H}^+)}$ the reproducing kernels of the Hardy space on the unit ball $H^2(\mathbb{B})$ and on the right half-space $H^2(\mathbb{H}^+)$ respectively. Let $C : \mathbb{B} \to \mathbb{H}^+$ be the Cayley map, $C(q) = (1-q)^{-1}(1+q)$. For any $z, w \in \mathbb{H}^+$, the function $k_{H^2(\mathbb{B})}(C^{-1}(w), C^{-1}(z))$ is a rescaling of the reproducing kernel of $H^2(\mathbb{H}^+)$:

$$k_{H^2(\mathbb{B})}(C^{-1}(w), C^{-1}(z)) = \frac{1}{2}(1+z)k_{H^2(\mathbb{H}^+)}(w, z)(1+\overline{w}).$$

Proof. The map C^{-1} , having real coefficients, is slice preserving. Hence, we can compose $k_{H^2(\mathbb{B})}$ with C^{-1} preserving (left) regularity in the first variable and (right) "anti-regularity" in the second one. We

have, then,

$$\begin{aligned} k_{H^{2}(\mathbb{B})}(C^{-1}(z),C^{-1}(w)) &= (1-q\overline{C^{-1}(w)})_{|_{q=C^{-1}(z)}}^{-*} \\ &= (1-2C^{-1}(z)\operatorname{Re}(C^{-1}(w)) + C^{-1}(z)^{2}|C^{-1}(w)|^{2})^{-1} \left(1-C^{-1}(z)C^{-1}(w)\right) \\ &= (1+z)^{2}|1+w|^{2} \left(|1+w|^{2}(1+z)^{2}-2(1-z^{2})(1-|w|^{2}) + (1-z)^{2}|1-w|^{2}\right)^{-1} \\ &\quad \cdot (1+z)^{-1} \left((1+z)(1+w) - (1-z)(1-w)\right) (1+w)^{-1} \\ &= (1+z)\frac{1}{4} \left(z^{2}+2z\operatorname{Re}(w) + |w|^{2}\right)^{-1} 2 \left(z+w\right) \left(1+\overline{w}\right) \\ &= \frac{1}{2}(1+z) \left(z+\overline{w}\right)^{-*} \left(1+\overline{w}\right) = \frac{1}{2}(1+z)k_{H^{2}(\mathbb{H}^{+})}(w,z)(1+\overline{w}). \end{aligned}$$

Now we want to show that the volume form on $\partial \mathbb{H}^+$ obtained in (13) is the natural volume form for the Hardy space $H^2(\mathbb{H}^+)$. In fact, let $f(q) = \int_0^{+\infty} e^{-\zeta q} F(\zeta) d\zeta \in H^2(\mathbb{H}^+)$. For any $I \in \mathbb{S}$, we can decompose the function F as $F(\zeta) = F_1(\zeta) + F_2(\zeta)J$ where J is an imaginary unit orthogonal to I and $F_1, F_2 : \mathbb{R} \to L_I$. It is possible to prove (see [2]) that functions in $H^2(\mathbb{H}^+)$ have limit at the boundary for almost any point $yI \in \partial \mathbb{H}^+ = \{vJ \mid v > 0, J \in \mathbb{S}\}$. If we denote by $dA_{\mathbb{S}}$ the usual surface element of the unit two-dimensional sphere \mathbb{S} , thanks to equation (13) and to the orthogonality of I and J, we can write

$$\begin{split} &\int_{\partial\mathbb{H}^{+}} |f(yI)|^{2} dVol_{\partial\mathbb{H}^{+}}(yI) = \int_{\partial\mathbb{H}^{+}} |f(yI)|^{2} \frac{dVol_{Euc}(yI)}{8y^{2}} \\ &= \int_{0}^{+\infty} \left(\int_{\mathbb{S}} \left| \int_{0}^{+\infty} e^{-\zeta yI} F(\zeta) d\zeta \right|^{2} \frac{y^{2} dA_{\mathbb{S}}(I)}{8y^{2}} \right) dy \\ &= \frac{1}{8} \int_{\mathbb{S}} \left(\int_{0}^{+\infty} \left| \int_{0}^{+\infty} e^{-\zeta yI} (F_{1}(\zeta) + F_{2}(\zeta)J) d\zeta \right|^{2} dy \right) dA_{\mathbb{S}}(I) \\ &= \frac{1}{8} \int_{\mathbb{S}} \left(\int_{0}^{+\infty} \left| \int_{0}^{+\infty} e^{-\zeta yI} F_{1}(\zeta) d\zeta + \int_{0}^{+\infty} e^{-\zeta yI} F_{2}(\zeta)J d\zeta \right|^{2} dy \right) dA_{\mathbb{S}}(I) \\ &= \frac{1}{8} \int_{\mathbb{S}} \left(\int_{0}^{+\infty} \left| \int_{0}^{+\infty} e^{-\zeta yI} F_{1}(\zeta) d\zeta \right|^{2} dy + \int_{0}^{+\infty} \left| \int_{0}^{+\infty} e^{-\zeta yI} F_{2}(\zeta) d\zeta \right|^{2} dy \right) dA_{\mathbb{S}}(I) \\ &= \frac{2\pi}{8} \int_{\mathbb{S}} \left(\int_{0}^{+\infty} |F_{1}(\zeta)|^{2} d\zeta + \int_{0}^{+\infty} |F_{2}(\zeta)| |^{2} d\zeta \right) dA_{\mathbb{S}}(I) \end{split}$$

where the last equality is due to the classical Plancherel Theorem. Therefore, thanks again to the orthogonality of I and J,

Proposition 5.3. If f belongs to $H^2(\mathbb{H}^+)$ and F is as in (14), we have the equalities:

$$\int_{\partial \mathbb{H}^+} |f(yI)|^2 dV ol_{\partial \mathbb{H}^+}(yI) = \frac{\pi}{4} \int_{\mathbb{S}} \left(\int_0^{+\infty} |F(\zeta)|^2 d\zeta \right) dA_{\mathbb{S}}(I) = \pi^2 ||f||^2_{H^2(\mathbb{H}^+)}.$$
 (16)

Propositions 5.1, 5.2, and 5.3 provide four isometric definitions of $H^2(\mathbb{H}^+)$.

5.2 A bilateral estimate for the distance and the proof of Theorem 1.4

In the right half space model it is easier to prove a bilateral estimate for the distance associated with the invariant metric h. Fix a imaginary unit I_0 and define the projection

$$\pi: x + yI \mapsto x + yI_0,\tag{17}$$

with x real and $y \ge 0$. Let d_{hyp} be hyperbolic distance in $\mathbb{H}_{I_0}^+ = \{x + yI_0 : x > 0, y \in \mathbb{R}\}$: d_{hyp} is the distance associated with the Riemannian metric tensor $ds_{hyp}^2 = (dx^2 + dy^2)/(4x^2)$. Let now $d_{\mathbb{S}}$ be the usual spherical distance on the unit two-dimensional sphere \mathbb{S} , associated with the metric tensor $ds_{\mathbb{S}}^2$. Then, the metric tensor associated with h can be decomposed as

$$ds_h^2 = ds_{hyp}^2 + \frac{y^2}{4(x^2 + y^2)} ds_{\mathbb{S}}^2$$

Theorem 5.4. Let $q_j = x_j + y_j I_j$, j = 1, 2, be points in \mathbb{H}^+ : $x_j > 0$, $y_j \ge 0$. The following estimate for the distance function d_h associated with the metric h holds:

$$d_h(q_1, q_2) \approx d_{hyp}(\pi(q_1), \pi(q_2)) + \min\left\{\frac{y_j}{|q_j|}: j = 1, 2\right\} d_{\mathbb{S}}(I_1, I_2),$$

where \approx means that we have a lower and an upper estimate for the right hand side in terms of the left hand side, with multiplicative constants C_1, C_2 independent of q_1, q_2 .

Proof. We may suppose that $y_1/|q_1| \le y_2/|q_2|$.

The upper estimate is elementary. Let γ be a curve going from q_1 to $x_1 + y_1 I_2 \in \mathbb{H}_{I_2}^+$ leaving $x = x_1$ and $y = y_1$ fixed, and varying the imaginary unit I only. Suppose, more, that I varies along a geodesic on \mathbb{S} , which joins I_1 and I_2 . Then,

$$\operatorname{length}(\gamma) = \int_{\gamma} \frac{y_1}{2|q_1|} ds_{\mathbb{S}} = \frac{y_1}{2|q_1|} d_{\mathbb{S}}(I_1, I_2).$$

Let now δ be a hyperbolic geodesic in L_{I_2} , joining $x_1 + y_1I_2$ and q_2 : length(δ) = $d_{hyp}(\pi(q_1), \pi(q_2))$, which proves the estimate.

The lower estimate is more delicate. Let γ be a curve in \mathbb{H}^+ joining q_1 and q_2 . Then,

$$\operatorname{length}(\gamma) = \int_{\gamma} ds_h \ge \int_{\pi(\gamma)} ds_{hyp} \ge d_{hyp}(\pi(q_1), \pi(q_2)).$$
(18)

We have then to show that

$$\int_{\gamma} ds_h \gtrsim \frac{y_1}{|q_1|} d_{\mathbb{S}}(I_1, I_2). \tag{19}$$

Since the right hand side of (19) is bounded, and we have already proved (18), it suffices to show that (19) holds when $d_{hyp}(\pi(q_1), \pi(q_2)) \leq 1$. By elementary hyperbolic geometry, see the "sixth model" in [8], and using the fact that dilations $p \mapsto \lambda p$ are isometric for $\lambda > 0$, we can assume that $\pi(q_1)$ and $\pi(q_2)$ both lie in the square $Q_n = \{x + yI_0 : 1 \leq x \leq 2, n \leq y \leq n+1\} \subset L_{I_0}$, for some integer $n \geq 0$. Consider now $q_3 = x_3 + y_3I_3$, $y_3 \geq 0$ the point along γ which minimizes $y_3/|q_3|$. We can assume that $\pi(\gamma)$ (hence,

 $\pi(q_3)$) is contained in $\tilde{Q}_n = \{z = x + yI_0 : x > 0, y \ge 0, 1/2 \le x \le 2, n - 1/2 \le y \le n + 3/2\}$, otherwise length $(\gamma) \ge 1$ (which would imply the estimate (19) we are proving). Let $t \ge 0$ be the angle between the positive real half axis \mathbb{R}^+ and the half line originating at 0 and passing through $\pi(q_3)$. For j = 1, 2, 3:

$$t_j \approx \sin(t_j) = y_j / |q_j|.$$

We have two cases. Either $y_3/|q_3| \ge 1/2 \cdot y_1/|q_1|$, but then we are done because

$$\int_{\gamma} \frac{y}{2|q|} ds_{\mathbb{S}}^2 \ge \frac{y_1}{2|q_1|} d_{\mathbb{S}}(I_1, I_2).$$

Or $y_3/|q_3| \le 1/2 \cdot y_1/|q_1|$. Then n = 0, and

$$\begin{split} \int_{\gamma} ds_h &\geq \quad \text{length}(\pi(\gamma)) \gtrsim \max(|\pi(z_1) - \pi(z_3)|, |\pi(z_2) - \pi(z_3)|) \geq |\pi(z_1) - \pi(z_3)| \\ &\gtrsim \quad y_1 \gtrsim \frac{y_1}{|z_1|} d_{\mathbb{S}}(I_1, I_2). \end{split}$$

Overall, $\int_{\gamma} ds_h \gtrsim \frac{y_1}{|z_1|} d_{\mathbb{S}}(I_1, I_2)$, as wished.

Changing coordinates from the right half plane to the ball, we have the same bilateral estimate in the ball model.

Corollary 5.5. Let d_g be the invariant distance associated with the metric g in the ball model and let q_1, q_2 be points of \mathbb{B} . If π is defined as in (17), then:

$$d_g(q_1, q_2) \approx d_{hyp}(\pi(q_1), \pi(q_2)) + \min\left\{\frac{y_j}{|1 - q_j^2|} : j = 1, 2\right\} d_{\mathbb{S}}(I_1, I_2).$$

We collect some facts which will be used in the proof of Theorem 1.4.

It is well known (see [11]) that regular, bounded functions have radial limits along almost all radii,

$$f(e^{tI}) := \lim_{r \to 1} f(re^{tI})$$

exists for *a.e.* $(t, I) \in [0, \pi] \times S$. We start with a Lemma which might have independent interest; for instance, it provides a different route to prove the classification of the isometries for the metric *g*.

Lemma 5.6. If $\varphi : \mathbb{B} \to \mathbb{B}$ is Lipschitz with respect to the metric g and

$$\lim_{r \to 1} \varphi(re^{tI_1}) = e^{sJ_1} \in \partial \mathbb{B}$$
⁽²⁰⁾

exists, with $s \in [0, \pi]$ and $J_1 \in \mathbb{S}$; then for each I_2 in \mathbb{S} , if the limit $\lim_{r \to 1} \varphi(re^{I_2 t})$ exists, then

$$\lim_{r \to 1} \varphi(re^{tI_2}) = e^{sJ_2} \tag{21}$$

for some J_2 in S. The values of t and s in (21) are the same as in (20).

Proof. Let $u_j = e^{tI_j}$, with the same $t \in [0, \pi]$. By Lipschitz continuity,

$$d(\varphi(ru_1),\varphi(ru_2)) \lesssim d(ru_1,ru_2) \\\approx \frac{rt|I_1-I_2|}{(1-r)+rt} \lesssim |I_1-I_2| \\\leq 1.$$
(22)

By the lower estimate in Corollary 5.5 and (22),

$$d_{hyp}\left(\pi\left(\varphi(ru_1)\right),\pi\left(\varphi(ru_2)\right)\right) \lesssim 1$$

But this and elementary hyperbolic geometry imply that, if $\lim_{r\to 1} \varphi(ru_1) = e^{sJ_1}$, then the limit $\lim_{r\to 1} \pi(\varphi(ru_2)) = L$ exists and $L = e^{sI_0}$ (recall that $\pi : \mathbb{B} \to L_{I_0}$). Since $\lim_{r\to 1} \varphi(re^{tI_2}) = L$ exists by hypothesis and π is continuous, it must be $\pi(L) = e^{sI_0}$, then $L = e^{sJ_2}$ for some J_2 in \mathbb{S} .

The statement of Lemma 5.6 can be sharpened in several ways. For instance, the Lipschitz assumption might be weakened to a sub-exponential growth assumption. We apply it to show that inner Lipschitz functions are scarce.

Proof of Theorem 1.4. Being inner, f has boundary limits along radii $r \mapsto re^{tI}$ for *a.e.* I in \mathbb{S} and t in $[0, \pi]$. We write for such couples of (t, I): $f(e^{tI}) := \lim_{r \to 1} f(re^{tI})$. We can assume without loss of generality that the limit exists for two antipodal imaginary units L and -L, and hence, in view of the Representation Formula 2.5, for any $L \in \mathbb{S}$. If f is regular and Lipschitz with respect to the distance d_g , thanks on the one hand to the Representation Formula 2.5, on the other hand to Lemma 5.6, we have that for any $I \in \mathbb{S}$

$$b(t) + Ic(t) = f(e^{tI}) = e^{s(t)J(s,I)}$$

where $b(t), c(t) \in \mathbb{H}$, and $s(t) \in [0, 2\pi]$ and $J(t, I) \in \mathbb{S}$. Then $\operatorname{Re}(f(e^{tI})) = \operatorname{Re}(f(e^{tL}))$ for any $L \in \mathbb{S}$ and in particular for L = -I, which gives

$$\operatorname{Re}(b(t)) - \langle I, c(t) \rangle = \operatorname{Re}(b(t) + Ic(t)) = \operatorname{Re}(b(t) - Ic(t)) = \operatorname{Re}(b(t)) + \langle I, c(t) \rangle$$

(where $\langle \cdot, \cdot \rangle$ denotes the standard scalar product in \mathbb{R}^4). Since *I* is any imaginary unit, we necessarily have that $c(t) \in \mathbb{R}$.

Also, comparing imaginary parts, for any $L_1, L_2 \in \mathbb{S}$ we have $|\operatorname{Im}(f(e^{tL_1}))| = |\operatorname{Im}(f(e^{tL_2}))|$. Then, if $b = b_0 + b_1 K$ with $b_0, b_1 \in \mathbb{R}$, $K \in \mathbb{S}$ (omitting the dependence on t), when $L_1 = K$ and $L_2 = -K$ we get

$$|b_1 + c| = |\operatorname{Im}(b_0 + b_1 K + cK)| = |\operatorname{Im}(f(e^{tK}))| = |\operatorname{Im}(f(e^{-tK}))| = |\operatorname{Im}(b_0 + b_1 K - cK)| = |b_1 - c|.$$

Therefore almost every $t \in [0, \pi]$ belongs to $D \cup E$:

$$D = \{t : c(t) = 0\}, E = \{t : b_1(t) = 0\}.$$

Consider first the case when $t \in D$ holds *a.e.*. Then $f(e^{tI}) = b(t)$ for almost every t. Since boundary values uniquely identify f (see [11]) and by invariance under rotations of \mathbb{S} , we deduce that

$$f(re^{tI}) = \Phi(r, t),$$

for some function Φ . In particular, f can not be open $(dim(f(\mathbb{B})) \leq 2)$, hence (see Theorem 7.4 in [13]) it must be constant; thus it is not inner. Then E has positive measure. For t in E,

$$b(t) + Ic(t) = f(e^{tI}) = e^{s(t)J(t,I)}$$

with b and c real valued, hence J = I:

$$f(e^{tI}) = e^{s(t)I} \tag{23}$$

for t in F. By the Splitting Lemma 2.7, if $J \perp I$ is fixed in S, then there are holomorphic functions F, G on \mathbb{B}_I such that

$$f(re^{\tau I}) = F(re^{\tau I}) + G(re^{\tau I})J_{\tau}$$

By (23), $G(e^{tI}) = 0$ for t in E. Since E has positive measure, this implies that G vanishes identically and hence $f(re^{\tau I}) = F(re^{\tau I})$ for all $0 \le r < 1$ and $0 \le \tau \le \pi$. That is, f is slice preserving.

We have to verify that f is a contraction with respect to the metric g, and this can be verified at the infinitesimal level. Let q be a point in a fixed slice $\mathbb{B} \cap L_I$. (i) Since f is slice preserving, its restriction to $\mathbb{B} \cap L_I$ is an inner function in the one dimensional sense, hence it is a contraction of the Poincaré-hyperbolic metric on $\mathbb{B} \cap L_I$. (ii) On the other hand, preserving the slices, f acts isometrically in the \mathbb{S} variables, with respect to the spherical metric on \mathbb{S} . (iii) Now, the space tangent to $\mathbb{B} \cap L_I$ at q and the space tangent to $\mathbb{R} e q + \mathbb{S}$ at q form an orthogonal decomposition, with respect to the metric g, of the space tangent to \mathbb{B} . From the expression for g given in (8) and facts (i)-(iii) one easily deduces that g is a contraction.

References

- [1] D. Alpay, F. Colombo, I. Sabadini, *Schur functions and their realizations in the slice hyperholomorphic setting*, Integral Equations Operator Theory **72** (2012), 25–289.
- [2] D. Alpay, F. Colombo, I. Lewkowicz, I. Sabadini, *Realizations of slice hyperholomorphic generalized contractive and positive functions*, Preprint arXiv:1310.1035v1 [math.CV] (2013).
- [3] N. Arcozzi, R. Rochberg, E. Sawyer, B. D. Wick, *Distance functions for reproducing kernel Hilbert spaces*, Function spaces in modern analysis, 25–53, Contemp. Math., 547, Amer. Math. Soc., Providence, RI, 2011.
- [4] N. Aronszajn, Theory of reproducing kernels, Trans. Amer. Math. Soc. 68, (1950) 337–404.
- [5] C. Bisi, G. Gentili, *Möbius transformations and the Poincaré distance in the quaternionic setting*, Indiana Univ. Math. J., **58** (2009), 2729–2764.
- [6] C. Bisi, C. Stoppato, *Regular vs. classical Möbius transformations of the quaternionic unit ball*, Advances in hypercomplex analysis, Springer INdAM Ser. 1, Springer, Milan, 2013, 1–13.
- [7] F. Brackx, R. Delanghe and F. Sommen, *Clifford Analysis*, Pitman Publishing, Boston-London-Melbourne, (1982).
- [8] J. W. Cannon, W. J. Floyd, R. Kenyon, W. R. Parry, *Hyperbolic geometry*, in Flavors of geometry, Math. Sci. Res. Inst. Publ., 31, Cambridge Univ. Press, Cambridge (1997), 59–115,.

- [9] F. Colombo, J. O. Gonzalez-Cervantes, M. E. Luna-Elizarraras, I. Sabadini, M. V. Shapiro, *On two approaches to the Bergman theory for slice regular functions*, Advances in hypercomplex analysis, Springer INdAM Ser. 1, Springer, Milan, 2013, 39–54.
- [10] M. J. Cowen, R. G. Douglas, *Complex geometry and operator theory*, Acta Math. 141 (1978), 187–261.
- [11] C. de Fabritiis, G. Gentili, G. Sarfatti, *Quaternionic Hardy spaces*, Preprint, www.math.unifi.it/users/sarfatti/Hardy.pdf, (2013).
- [12] M. P. do Carmo, Differential geometry of curves and surfaces, Prentice-Hall, Inc., Englewood Cliffs, N.J., 1976.
- [13] G. Gentili, C. Stoppato, D. C. Struppa, *Regular functions of a quaternionic variable*, Springer Monographs in Mathematics, Springer, Berlin-Heidelberg, 2013.
- [14] G. Gentili, D. C. Struppa, *A new theory of regular functions of a quaternionic variable*, Adv. Math., 216 (2007), 279–301.
- [15] R. Ghiloni, V. Moretti, A. Perotti, *Continuous slice functional calculus in quaternionic Hilbert spaces*, Rev. Math. Phys. 25 (2013).
- [16] M. Gromov, *Metric structures for Riemannian and non-Riemannian spaces*, Based on the 1981 French original, translated from the French by Sean Michael Bates. Progress in Mathematics, 152, Birkhäuser Boston, Inc., Boston, MA, 1999.
- [17] J. E. McCarthy, Boundary values and Cowen-Douglas curvature, J. Funct. Anal. 137 (1996), 1–18.
- [18] M. Mitrea, *Clifford wavelets, singular integrals, and Hardy spaces*, Lecture Notes in Mathematics, 1575, Springer Verlag, Berlin, (1994).
- [19] B. O'Neill, Semi-Riemannian geometry, with applications to relativity, Pure and Applied Mathematics, 103. Academic Press, Inc., New York, 1983.
- [20] A. Perotti, *Fueter regularity and slice regularity: meeting points for two function theories*, Advances in hypercomplex analysis, Springer INdAM Ser. 1, Springer, Milan, 2013, 93–117.
- [21] C. Stoppato, Regular Moebius transformations of the space of quaternions, Ann. Global Anal. Geom. 39 (2011), 387–401.

Nicola Arcozzi

Dipartimento di Matematica, Università di Bologna Piazza di Porta San Donato 5, 40126 Bologna, Italy, nicola.arcozzi@unibo.it

Giulia Sarfatti

Dipartimento di Matematica, Università di Bologna

Piazza di Porta San Donato 5, 40126 Bologna, Italy, giulia.sarfatti@unibo.it