



UNIVERSITÀ POLITECNICA DELLE MARCHE
Repository ISTITUZIONALE

Slice-polynomial functions and twistor geometry of ruled surfaces in CP 3

This is the peer reviewed version of the following article:

Original

Slice-polynomial functions and twistor geometry of ruled surfaces in CP 3 / Altavilla, A.; Sarfatti, G.. - In: MATHEMATISCHE ZEITSCHRIFT. - ISSN 0025-5874. - 291:3-4(2019), pp. 1059-1092. [10.1007/s00209-018-2225-8]

Availability:

This version is available at: 11566/272056 since: 2022-05-27T11:54:35Z

Publisher:

Published

DOI:10.1007/s00209-018-2225-8

Terms of use:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. The use of copyrighted works requires the consent of the rights' holder (author or publisher). Works made available under a Creative Commons license or a Publisher's custom-made license can be used according to the terms and conditions contained therein. See editor's website for further information and terms and conditions.

This item was downloaded from IRIS Università Politecnica delle Marche (<https://iris.univpm.it>). When citing, please refer to the published version.

note finali coverpage

(Article begins on next page)

Slice-polynomial functions and twistor geometry of ruled surfaces in CP^3

A. Altavilla, G. Sarfatti

This is a post-print version, the reference for the published version is:

Mathematische Zeitschrift (2019) 291:1059–1092

<https://doi.org/10.1007/s00209-018-2225-8>

Licence: Publisher's Bespoke License

Articles, books and chapters published within the Springer Nature group of companies which are made available through academic repositories remain subject to copyright. Any further use is subject to permission from Springer Nature. The following restrictions on use of such articles apply:

Academic research only

- 1. Archived content may only be used for academic research. Any content downloaded for text based experiments should be destroyed when the experiment is complete.

Use must not be for Commercial Purposes

- 2. Archived content may not be used for purposes that are intended for or directed towards commercial advantage or monetary compensation by means of sale, resale, licence, loan, transfer or any other form of commercial exploitation ("Commercial Purposes").

Wholesale re-publishing is prohibited

- 3. Archived content may not be published verbatim in whole or in part, whether or not this is done for Commercial Purposes, either in print or online.
- 4. This restriction does not apply to reproducing normal quotations with an appropriate citation. In the case of text-mining, individual words, concepts and quotes up to 100 words per matching sentence may be used, whereas longer paragraphs of text and images cannot (without specific permission from Springer Nature).

Moral rights

- 5. All use must be fully attributed. Attribution must take the form of a link - using the article DOI - to the published article on the journal's website.
- 6. All use must ensure that the authors' moral right to the integrity of their work is not compromised.

Third party content

- 7. Where content in the document is identified as belonging to a third party, it is the obligation of the user to ensure that any use complies with copyright policies of the owner.

Use at own risk

- 8. Any use of Springer Nature content is at your own risk and Springer Nature accepts no liability arising from such use.

SLICE-POLYNOMIAL FUNCTIONS AND TWISTOR GEOMETRY OF RULED SURFACES IN \mathbb{CP}^3

A. ALTAVILLA^{†,‡} AND G. SARFATTI^{†,*}

ABSTRACT. In the present paper we introduce the class of slice-polynomial functions: slice regular functions defined over the quaternions, outside the real axis, whose restriction to any complex half-plane is a polynomial. These functions naturally emerge in the twistor interpretation of slice regularity introduced in [15] and developed in [2]. To any slice-polynomial function P we associate its *companion* P^\vee and its *extension* to the real axis $P_{\mathbb{R}}$, that are quaternionic functions naturally related to P . Then, using the theory of twistor spaces, we are able to show that for any quaternion q the cardinality of simultaneous pre-images of q via P , P^\vee and $P_{\mathbb{R}}$ is generically constant, giving a notion of degree. With the brand new tool of slice-polynomial functions, we compute the twistor discriminant locus of a cubic scroll \mathcal{C} in \mathbb{CP}^3 and we conclude by giving some qualitative results on the complex structures induced by \mathcal{C} via the twistor projection.

CONTENTS

1.	Introduction and motivation	2
1.1.	Orthogonal complex structures and twistor theory of \mathbb{S}^4	2
1.2.	Relation between twistor theory and slice regularity	3
2.	Slice-polynomial functions	5
2.1.	Extension to the real line	10
2.2.	Companion of a slice regular function	11
3.	Twistor fibration and blow up of the real line	12
3.1.	Blow-up of the real line	15
4.	Pre-images of a slice-polynomial function	16
5.	Discriminant locus of a cubic scroll	20
5.1.	A first description of the OCS's induced by \mathcal{C}	26
	References	29

Date: November 15, 2018.

2010 Mathematics Subject Classification. Primary 53C28, 30G35; secondary 32A30, 14J26.

Key words and phrases. Twistor spaces, Slice regular functions, Functions of hypercomplex variables, Rational and ruled surfaces, Slice-polynomial functions.

[†]Partially supported by GNSAGA of INdAM, FIRB 2012 *Geometria differenziale e teoria geometrica delle funzioni* and by SIR 2014 *AnHyC - Analytic aspects in complex and hypercomplex geometry* n. RBSI14DYEB. [‡]Partially supported by SIR grant *NEWHOLITE - New methods in holomorphic iteration* n. RBSI14CFME. * Partially supported by Finanziamento Premiale FOE 2014 *SUNRISE - Splines for accurate numerics: adaptive models for simulation environments*. The second author wishes to thank the Institut Montpelliérain Alexander Grothendieck where part of this project was carried out.

1. INTRODUCTION AND MOTIVATION

A fascinating aspect of the recent theory of slice regular functions over the quaternions is that it proved to have a fruitful interaction with the theory of twistor geometry. As first stated in [15] and then developed in [2], one can describe the theory of slice regularity using the language of complex geometry and thus use this description to associate with any injective slice regular function an orthogonal complex structure (OCS in the sequel) defined on its image. The relevance of this construction relies on the fact that the classification of orthogonal complex structures on general open domains in \mathbb{R}^4 is an open problem, [25]. In the present work we use this twistor interpretation to analyse a special family of slice regular functions defined outside the real axis, that we named *slice-polynomial functions*. The interest in this class of functions is motivated by the fact that they furnish a new tool to describe the twistor geometry of a wide family of ruled surfaces in \mathbb{CP}^3 . Let us begin by introducing the general contexts where we will work in.

1.1. Orthogonal complex structures and twistor theory of \mathbb{S}^4 . An OCS on a Riemannian manifold (M, g) is an (integrable) complex structure which is Hermitian, i.e., compatible with the metric g . For our purposes we will restrict our attention to orientation preserving OCS's. Two global OCS's J_1, J_2 are said to be *independent* if there exists some $x \in M$ such that $J_1(x) \neq \pm J_2(x)$. Moreover they are said to be *strongly independent* if $J_1(x) \neq \pm J_2(x)$ for any $x \in M$ (see [5, 14, 23, 24]). It is therefore a relevant question to understand if, given a Riemannian manifold (M, g) , it admits any OCS and, if this is the case, to describe the set of all OCS's defined on it. The appropriate tool to deal with this problem is the so called *twistor space* Z of M , which is defined as the total space of the bundle containing all the OCS's defined on M . This space contains informations useful not only to characterise the existence of such structures (see, for instance, [10, 25]), but also to study other interesting geometric problems as the existence of minimal surfaces (see e.g. [9]).

Notice that, given any OCS J on a Riemannian manifold (M, g) , if we replace the metric g by a conformally equivalent one $e^f g$, the orthogonality of J is not affected. Thus, the problem of finding an OCS on (M, g) is more related to the conformal class of g than to g itself. From this perspective we are interested in conformal transformations of Z i.e. automorphisms of Z induced by conformal transformations of the base space M .

Example 1.1. For $n > 1$ the sphere \mathbb{S}^{2n} has no OCS J relative to its standard metric, [22]. However, let $x \in \mathbb{S}^{2n}$, then, the induced metric on $\mathbb{S}^{2n} \setminus \{x\}$ is conformally equivalent to the flat metric on \mathbb{R}^{2n} and the latter admits infinitely many OCS's including the constant ones parameterised by the homogeneous space $O(2n)/U(n)$. In particular, $\mathbb{S}^4 \setminus \{\infty\}$ admits only constant OCS's, parameterised by $O(4)/U(2)$.

For the remainder of the paper, we restrict our attention to the case of the 4-sphere \mathbb{S}^4 which can be identified with the quaternionic projective line \mathbb{HP}^1 . The twistor space in this case is \mathbb{CP}^3 and the associated bundle structure is given by $\pi : \mathbb{CP}^3 \rightarrow \mathbb{HP}^1$ with fibre \mathbb{CP}^1 (see Section 3 for further details). It is known (see, e.g., [25, Section 2.6]), that any complex surface in \mathbb{CP}^3 transverse to the fibres of π induces an OCS on subdomains of \mathbb{R}^4 whenever such a surface is a single valued graph (with respect to the twistor projection). Vice versa, any OCS on a domain $\Omega \subset \mathbb{S}^4$ corresponds to a holomorphic surface in \mathbb{CP}^3 . Since it is not possible to define any OCS on the whole \mathbb{S}^4 , then no such surface in \mathbb{CP}^3 can intersect every fibre of the twistor fibration in exactly one point.

In this context, conformal transformations of \mathbb{CP}^3 are described in, e.g., [6, Section 2]. In a series of papers [4, 6, 7, 8, 15, 25], the authors have studied algebraic surfaces in \mathbb{CP}^3 from this point of view. In particular, in [25] non-singular quadric surfaces are fully classified under conformal transformations, while in [6, 7, 8] qualitative and quantitative results on the twistor geometry of non-singular cubics are given. In the paper [15] the authors study a (singular) quartic scroll by means of slice regularity. All

the results mentioned in the previous papers have inferred the existence (and in some case the explicit construction), of OCS's over explicit open dense domains of \mathbb{S}^4 .

1.2. Relation between twistor theory and slice regularity. The definition of slice regular functions, due to G. Gentili and D. C. Struppa, is the most recent attempt to generalise the notion of holomorphicity to the quaternionic setting in order to include polynomials of the form $\sum_{n=0}^N q^n a_n$, with quaternionic coefficients, see [17]. The main idea is the following: if \mathbb{S} is the two dimensional sphere of quaternionic imaginary units $\mathbb{S} := \{q \in \mathbb{H} \mid q^2 = -1\}$, then any $q \in \mathbb{H}$ can be written as $q = \alpha + I\beta$ for $\alpha, \beta \in \mathbb{R}$ and $I \in \mathbb{S}$. In this way we obtain the *slice* decomposition $\mathbb{H} = \bigcup_{I \in \mathbb{S}} \mathbb{C}_I$, where $\mathbb{C}_I := \text{Span}_{\mathbb{R}}(1, I)$. Sets of the form \mathbb{C}_I are called *slices* while the half-planes \mathbb{C}_I^+ , identified by positive complex imaginary part, are called *semi-slices*. For any quaternion $q = \alpha + I\beta$, its real part is the real number $\text{Re}(q) = \alpha$, its imaginary part is the purely imaginary number $\text{Im}(q) = I\beta$ and its conjugate is $\bar{q} = \alpha - I\beta$, so that its modulus can be computed as $|q|^2 = q\bar{q}$.

Definition 1.2. Let $\Omega \subset \mathbb{H}$ be a domain. A differentiable function $f : \Omega \rightarrow \mathbb{H}$ is said to be *slice regular* if for all $I \in \mathbb{S}$ the restriction $f|_{\mathbb{C}_I} : \mathbb{C}_I \rightarrow \mathbb{H}$ is a holomorphic function with respect to the complex structure defined by quaternionic left multiplication by I .

Several results hold for this class of functions, in analogy with the complex case: just as some examples we can mention a Cauchy integral representation formula, a Maximum and Minimum Modulus Principle, an Open Mapping Theorem. On the contrary other aspects are very different, as the nature of zeros and poles that can be either isolated points or isolated 2-dimensional spheres. See [16] and references therein for an extensive account to the theory and to [1, 3] for some generalizations.

Many steps forward in the theory are due to the work of R. Ghiloni and A. Perotti and their idea of using *stem functions*, [19]. The formalism of stem functions allows, in particular, to enlarge the class of domains on which the functions can be defined (including domains that do not intersect the real axis) and to extend the theory to the more general setting of alternative algebras.

Let us now describe the interplay between slice regular functions and OCS's. Consider the manifold $X = \mathbb{H} \setminus \mathbb{R}$, endowed with the Euclidean metric. Using the language of quaternions, it is possible to define a non-constant OCS on X : consider a point $p = \alpha + I_p\beta \in X$, with $\beta > 0$, and identify $T_p X \simeq \mathbb{H}$. Then we can define the OCS \mathbb{J} over X as

$$\mathbb{J}_p : T_p X \rightarrow T_p X, \quad v \mapsto \frac{\text{Im}(p)}{|\text{Im}(p)|} v = I_p v,$$

where v is a tangent vector to X in p and $I_p v$ denotes the quaternionic multiplication between I_p and v . It is proven in [25] that $\pm \mathbb{J}$ are the only non-constant OCS's, up to conformal transformation of \mathbb{S}^4 , that can be defined on $\mathbb{H} \setminus \mathbb{R}$.

Slice regular functions enter in the picture as follows: first of all $(\mathbb{H} \setminus \mathbb{R}, \mathbb{J})$ is biholomorphic to a suitable open subset \mathcal{Q}^+ of the *Segre Quadric* $\mathcal{Q} \subset \mathbb{C}\mathbb{P}^3$. Let now $f : \Omega \rightarrow \mathbb{H}$ be a slice regular function on a circular domain (i.e. a domain which is symmetric with respect to the real axis). Then f admits a *twistor lift* to $\pi^{-1}(\Omega \setminus \mathbb{R}) \cap \mathcal{Q}^+$, i.e.: there exists a holomorphic function $\tilde{f} : \pi^{-1}(\Omega \setminus \mathbb{R}) \cap \mathcal{Q}^+ \rightarrow \mathbb{C}\mathbb{P}^3$, such that $\pi \circ \tilde{f} = f \circ \pi$ (see Section 3 for the details). Suppose now that the function f is also injective and let $p = \alpha + I_p\beta \in \Omega \setminus \mathbb{R}$. Then it is possible to define an OCS \mathbb{J}^f on the image of f as

$$\mathbb{J}_{f(p)}^f v = \frac{\text{Im}(p)}{|\text{Im}(p)|} v = I_p v$$

where $\mathbb{J}^f := (df)\mathbb{J}(df)^{-1}$ denotes the push-forward of \mathbb{J} via f . See [2, 15]. In [2] it is given a first account on the family of algebraic surfaces in $\mathbb{C}\mathbb{P}^3$ that can be parameterised by the twistor lift of a slice regular function. This family is composed by surfaces *ruled by lines* called also *scrolls*. In particular, this family

contains all hyperplanes and all non-singular quadrics. Moreover, up to projective transformations, for any quadric surface and cubic scroll $\mathcal{S} \subset \mathbb{C}\mathbb{P}^3$ there exists a slice regular function f such that its twistor lift \tilde{f} has image in \mathcal{S} .

An issue of this construction consists in the impossibility, in general, of extending the twistor lift of a slice regular function to the whole quadric \mathcal{Q} . See for instance [2, Remark 17] for an explicit simple example. This issue is solved in the present paper in which we start the study of what we call *slice-polynomial functions*.

Definition 1.3. A slice regular function $P : \mathbb{H} \setminus \mathbb{R} \rightarrow \mathbb{H}$ is a slice-polynomial function if, for any $I \in \mathbb{S}$, the restriction of P to the semi-slice \mathbb{C}_I^+ is a polynomial.

(See Definition 2.9 and Proposition 2.11 for the details). For this class of functions we ask ourselves and answer a natural basic question: *does a Fundamental Theorem of Algebra hold for slice-polynomial functions?* The answer is positive in the class of slice regular polynomials of a quaternionic variable (see [18] and [19] for a generalization). In the case of slice-polynomial functions the answer is positive as well but it requires the introduction of new techniques. In fact, we need to consider, together with a slice-polynomial function P , its *companion* P^\vee , defined as a slice-polynomial function which is a sort of dual of P (see Definition 2.17), and its *extension* to the real axis $P_{\mathbb{R}}$ (see Subsection 2.1). Given any quaternion $q \in \mathbb{H}$, using twistor theory, we are able to study the simultaneous pre-images of q via a slice-polynomial function, its companion and its extension to the reals, see Corollary 4.2. In particular we get the following result.

Theorem 1.4. *Let P be a slice-polynomial function, not slice-constant and let q be any quaternion. Then, generically,*

$$\#\{P^{-1}(q)\} + \#\{P^\vee^{-1}(q)\} + \#\{P_{\mathbb{R}}^{-1}(q)\} = d$$

where d is the twistor degree of P .

In the previous statement *generically* means outside a real Zariski closed subset of \mathbb{H} . The twistor degree of a slice-polynomial function is a suitable notion of degree in this context (see Definition 4.3) and furthermore it is strictly related to the degree of the algebraic surface where the twistor lift of the slice-polynomial function lies (see Corollary 4.8). As a consequence of Corollary 4.2, we prove the following version of the Fundamental Theorem of Algebra.

Theorem 1.5. *Let P be a slice-polynomial function, not slice-constant. Then*

$$P(\mathbb{H} \setminus \mathbb{R}) \cup P^\vee(\mathbb{H} \setminus \mathbb{R}) \cup P_{\mathbb{R}}(\mathbb{R}) = \mathbb{H}.$$

The introduction of slice-polynomial functions provides us with a useful tool to study the twistor geometry of a certain class of surfaces in $\mathbb{C}\mathbb{P}^3$. Given an algebraic surface $\mathcal{S} \subset \mathbb{C}\mathbb{P}^3$ of degree d its *twistor discriminant locus* $Disc(\mathcal{S})$ is defined as the following subset of the 4-sphere

$$Disc(\mathcal{S}) := \{p \in \mathbb{S}^4 \mid \#(\pi^{-1}(p) \cap \mathcal{S}) \neq d\}.$$

The importance of the discriminant locus lies in the fact that its topology is a conformal invariant of the surface \mathcal{S} (see e.g. [7, 25]). In particular, the number of twistor fibres contained in \mathcal{S} is a conformal invariant. Twistor fibres can be effectively computed by means of slice regular functions: in fact, if f is a slice regular function which is constant on a 2-sphere of the form $\alpha + \mathbb{S}\beta$, then $\pi^{-1}(f(\mathbb{S}_{\alpha+I\beta}))$ is a twistor fibre.

In this paper, as an application, we study the particular case of a cubic scroll \mathcal{C} in $\mathbb{C}\mathbb{P}^3$, showing some of its beautiful topological properties. With the newly introduced theory of slice-polynomial functions we are able to compute the discriminant locus of \mathcal{C} in a couple of pages giving therefore a new method to analyse something that is in general hard to compute (see [8, Section 2]).

Theorem 1.6. *The discriminant locus of the cubic scroll*

$$\mathcal{C} = \{[X_0, X_1, X_2, X_3] \in \mathbb{C}\mathbb{P}^3 : X_0X_3^2 + X_1^2X_2 = 0\}$$

consists in a 2-sphere with 6 handles, pinched at one pole and at a third root of -1 lying on an equator:

$$\text{Disc}(\mathcal{C}) = \Sigma = \hat{\mathbb{C}}_i \bigcup_{\substack{P=0,\infty \\ z_m^3=-1, z_m \in \mathbb{C}_i}} \Sigma_{P,z_m},$$

where Σ_{P,z_m} denotes a handle pinched to $\hat{\mathbb{C}}_i$ at P and z_m .

As we mentioned at the beginning of this introduction, there is not a complete classification of OCS's in subdomains on \mathbb{S}^4 . Another interesting consequence of our study, it is the fact that we are able to show the existence of 3 different OCS's defined outside the discriminant locus of \mathcal{C} .

Corollary 1.7. *The manifold $\mathbb{S}^4 \setminus \Sigma$ admits 3 non-constant strongly independent OCS's.*

The paper is structured as follows. In Section 2 we introduce slice-polynomial functions with their various representations. We describe how to extend a slice-polynomial function to the real line and we introduce the notion of companion. In Section 3 we recall the main tools in twistor geometry of the 4-sphere and we generalise and adapt some construction of [2, 15]. Both Sections 2 and 3 are quite technical but introduces several tools that are used in what follows. We point out that most of the results contained in these sections are in fact stated for a more general class of slice regular functions defined outside the real axis. In Section 4 we prove Theorems 1.4 and 1.5 using the technology of twistor spaces, In Section 5 we use the introduced tools to determine the discriminant locus of a cubic scroll, thus proving Theorems 1.6 and 1.7.

2. SLICE-POLYNOMIAL FUNCTIONS

Let us begin by recalling some preliminary facts about the approach to slice regularity using stem functions. See [19] for the details. Let $D \subseteq \mathbb{C}$ be an open domain of the complex plane, such that $D = \bar{D}$. In the quaternionic setting, a *stem function* is a function of the form $F = F_1 + \iota F_2 : D \subseteq \mathbb{C} \rightarrow \mathbb{H} \otimes \mathbb{C}$, such that, for any $z \in D$, $F(\bar{z}) = \overline{F(z)}$. Any stem function F induces a *slice function* $f = \mathcal{I}(F) : \Omega_D \rightarrow \mathbb{H}$, defined over the quaternionic domain $\Omega_D := \{\alpha + I\beta \mid \alpha + i\beta \in D, I \in \mathbb{S}\}$, as $f(\alpha + I\beta) = F_1(\alpha + i\beta) + IF_2(\alpha + i\beta)$. Sets of the form Ω_D are called *circular domains* (in the literature they are also called *slice symmetric domains* or *product domains*, depending whether they intersect the real axis). In this paper we will always consider circular domains as domains of definition of our slice regular functions. For this reason, we will sometimes drop out the subscript D to simplify the notation. If the stem function F is holomorphic, then the induced $f = \mathcal{I}(F)$ is a *slice regular function*.

We recall that, given a slice regular function $f = \mathcal{I}(F_1 + \iota F_2) : \Omega_D \subset \mathbb{H} \rightarrow \mathbb{H}$, it is possible to define its *regular conjugate* as the slice regular function $f^c : \Omega_D \rightarrow \mathbb{H}$ defined by $f^c := \mathcal{I}(\overline{F_1} + \iota \overline{F_2})$ (see, e.g., [16, 19]).

A general key result in slice regularity is the *Representation Formula* which allows us to restore a slice regular function from its values on a complex slice.

Theorem 2.1. *Let $\Omega \subseteq \mathbb{H}$ be a circular domain and let $f : \Omega \rightarrow \mathbb{H}$ be a slice regular function. Then, for any $J \in \mathbb{S}$, f is uniquely determined by its values over \mathbb{C}_J by the following formula:*

$$f(\alpha + I\beta) = \frac{1}{2} [f(\alpha + J\beta) + f(\alpha - J\beta) - IJ(f(\alpha + J\beta) - f(\alpha - J\beta))].$$

For any $I \in \mathbb{S}$, let $D \subset \mathbb{C}_I$ be a domain such that $\bar{D} = D$. Then any holomorphic function $f : D \rightarrow \mathbb{H}$ can be uniquely extended to a slice regular function $\text{ext}(f) : \Omega_D \rightarrow \mathbb{H}$, called *regular extension*, by means of the Representation Formula.

Given now any slice regular function on a circular domain $f : \Omega \rightarrow \mathbb{H}$ it is possible to define its *spherical derivative* $\partial_s f : \Omega \setminus \mathbb{R} \rightarrow \mathbb{H}$ as

$$\partial_s f(q) := \frac{1}{2} \text{Im}(q)^{-1} (f(q) - f(\bar{q})).$$

Since Ω is circular, given any $q = \alpha + I\beta \in \Omega \setminus \mathbb{R}$, the 2-sphere centred at α with radius β is fully contained in Ω :

$$\mathbb{S}_q := \{\alpha + J\beta \mid J \in \mathbb{S}\} \subset \Omega.$$

If $q \in \Omega \setminus \mathbb{R}$ is such that $\partial_s f(q) = 0$, then $f|_{\mathbb{S}_q} \equiv f(q)$, and \mathbb{S}_q is said to be a *degenerate sphere* for f . Notice that if $f = \mathcal{I}(F_1 + \iota F_2)$, then $\partial_s f(\alpha + I\beta) = 0$ if and only if $F_2(\alpha + i\beta) = 0$. The set of degenerate spheres of a non-constant slice regular function f is closed, with empty interior and denoted by \mathcal{D}_f .

As shown in [3], a slice regular function f defined on a domain without real points can be constant also on another type of 2-dimensional subsets called *wings*¹. A wing for a slice regular function f is a surface $\Sigma \subset \Omega_D$ biholomorphic to $D \cap \mathbb{C}^+$ such that $f|_{\Sigma}$ is constant. As proved in [3], the set of wings of a slice regular function f is also closed and with empty interior (or the full domain if f is slice-constant, see Definition 2.2). We will denote it by \mathcal{W}_f . A very important topological result for slice regular function is the following.

Theorem 2.2 (Open Mapping Theorem). *Let $\Omega \subseteq \mathbb{H}$ be a circular domain and let $f : \Omega \rightarrow \mathbb{H}$ be any slice regular function. Then the restriction $f|_{\Omega \setminus (\mathcal{D}_f \cup \mathcal{W}_f)}$ is open.*

This statement of the Open Mapping Theorem generalises the one in [16], where it is assumed that $\Omega \cap \mathbb{R} \neq \emptyset$ and the one in [1] where \mathcal{W}_f is supposed to be only composed by semi-slices of the form $\Omega \cap \mathbb{C}_I^+$ for suitable $I \in \mathbb{S}$. The proof in this case can be performed suitably adapting that of [1, Theorem 5.1]. Another version of the Open Mapping Theorem was recently given in [20]. Another fundamental result is the following *Splitting Lemma* which allows to split any slice regular function into a suitable sum of holomorphic functions.

Lemma 2.1 (Splitting Lemma). *Let $\Omega \subseteq \mathbb{H}$ be a circular domain and let $f : \Omega \rightarrow \mathbb{H}$ be a slice regular function. Then, for any $I, J \in \mathbb{S}$ such that $I \perp J$, there exist two holomorphic functions $G, H : \Omega \cap \mathbb{C}_I \rightarrow \mathbb{C}_I$, such that*

$$f|_{\Omega \cap \mathbb{C}_I} = G + HJ$$

The point-wise product between two slice regular functions is not, in general, slice regular. In the theory of slice regularity is more natural to use the $*$ -product, which does preserve regularity, defined as follows. Let $f = \mathcal{I}(F)$ and $g = \mathcal{I}(G)$ be two slice regular functions. Then their $*$ -product is defined to be the function induced by the point-wise product of their stem functions, i.e.: $f * g = \mathcal{I}(FG)$. A slice regular function $f : \Omega \rightarrow \mathbb{H}$ is said to be \mathbb{C}_I -preserving if, for some $I \in \mathbb{S}$, it holds $f(\Omega \cap \mathbb{C}_I) \subseteq \mathbb{C}_I$, while it is called *slice preserving* if the previous inclusion holds for any $I \in \mathbb{S}$. If f and g are both \mathbb{C}_I -preserving slice regular functions then $f * g = g * f$, while if f is slice preserving and g is any slice regular function then $f * g = g * f = fg$.

A special class of functions, introduced and studied in [1, 2, 3], which is peculiarly quaternionic, is the class of the so called slice-constant functions.

Definition 2.2. A slice function $f = \mathcal{I}(F) : \Omega \rightarrow \mathbb{H}$ is said to be *slice-constant* if its stem function F is locally constant.

¹The name ‘‘wing’’ is due to R. Ghiloni and A. Perotti.

If a slice-constant function is defined on a connected domain intersecting the real axis then it is constant. As proven in [1, Theorem 3.4], if f is a slice-constant function, then it is regular and its *Cullen derivative* is identically zero (see [16, Definition 1.7]).

A fundamental example of slice-constant functions is the following.

Definition 2.3. Fix any imaginary unit $J \in \mathbb{S}$. Let $L_+^J, L_-^J : \mathbb{C} \setminus \mathbb{R} \rightarrow \mathbb{H}_{\mathbb{C}}$ be the stem functions defined as

$$L_+^J(z) := \begin{cases} \frac{1-iJ}{2}, & \text{if } z \in \mathbb{C}^+ \\ \frac{1+iJ}{2}, & \text{if } z \in \mathbb{C}^- \end{cases}, \quad L_-^J(z) := \begin{cases} \frac{1+iJ}{2}, & \text{if } z \in \mathbb{C}^+ \\ \frac{1-iJ}{2}, & \text{if } z \in \mathbb{C}^- \end{cases}.$$

The previous stem functions induce the slice-constant functions,

$$\ell_+^J(\alpha + I\beta) = \mathcal{I}(L_+^J)(\alpha + I\beta) = \frac{1 - IJ}{2}, \quad \ell_-^J(\alpha + I\beta) = \mathcal{I}(L_-^J)(\alpha + I\beta) = \frac{1 + IJ}{2}.$$

For $J = i$ we will use the simpler notation

$$\ell_+ := \ell_+^i, \quad \ell_- := \ell_-^i.$$

Remark 2.4. The family of functions just defined is of particular interest since any of its element ℓ_+^J is idempotent with respect to the $*$ -product. More precisely, from the definition and straightforward computations, we have the following equalities:

$$(\ell_+^J)^c = \ell_-^J, \quad \ell_+^J * \ell_+^J = \ell_+^J, \quad \ell_+^J * \ell_-^J \equiv 0, \quad \ell_+^J + \ell_-^J \equiv 1.$$

Moreover for any $J \in \mathbb{S}$, the two functions ℓ_+^J and ℓ_-^J are \mathbb{C}_J -preserving. Notice that $\ell_+^J * \ell_-^J$ equals the so-called *symmetrization* of ℓ_+^J , see, e.g., [16].

Remark 2.5. Given any quaternion $q \in \mathbb{H}$, for any fixed $J \in \mathbb{S}$, we denote by q^\top and q^\perp the orthogonal projections of q on \mathbb{C}_J and \mathbb{C}_J^\perp , respectively. Therefore, since J commutes with q^\top and anti-commutes with q^\perp , for any couple $a, b \in \mathbb{H}$ we have, by direct computation, that,

$$(a^\top + a^\perp) * \ell_+^J + (b^\top + b^\perp) * \ell_-^J = \ell_+^J(a^\top + b^\perp) + \ell_-^J(a^\perp + b^\top).$$

This family of idempotent functions generate the space of slice-constant functions in the following sense: a slice function g is slice-constant if and only if for any $J \in \mathbb{S}$, g is a linear combination of the two functions ℓ_+^J and ℓ_-^J , i.e. there exist $a, b \in \mathbb{H}$ such that $g = a * \ell_+^J + b * \ell_-^J = \ell_+^J(a^\top + b^\perp) + \ell_-^J(a^\perp + b^\top)$. See [2, Proposition 15]. In particular, since ℓ_+^J and ℓ_-^J are slice-constant, for any other $K \in \mathbb{S}$, we have (in accordance with the Representation Formula)

$$(2.1) \quad \ell_+^J = \ell_+^K \left(\frac{1 - KJ}{2} \right) + \ell_-^K \left(\frac{1 + KJ}{2} \right) \quad \ell_-^J = \ell_+^K \left(\frac{1 + KJ}{2} \right) + \ell_-^K \left(\frac{1 - KJ}{2} \right).$$

Let $f : \Omega \rightarrow \mathbb{H}$, with $\Omega \cap \mathbb{R} = \emptyset$, be any slice regular function. Thanks to the Splitting Lemma, for any $J \in \mathbb{S}$, the restriction of f to $\Omega \cap \mathbb{C}_J$ can be written as $f_J(v) = G(v) + H(v)K$, where $K \in \mathbb{S}, K \perp J$ and $G, H : \Omega \cap \mathbb{C}_J \rightarrow \mathbb{C}_J$ are holomorphic functions of the form

$$(2.2) \quad G(v) := \begin{cases} g(v) & v \in \Omega \cap \mathbb{C}_J^+ \\ \widehat{g}(\overline{v}) & v \in \Omega \cap \mathbb{C}_J^- \end{cases}, \quad H(v) := \begin{cases} h(v) & v \in \Omega \cap \mathbb{C}_J^+ \\ \widehat{h}(\overline{v}) & v \in \Omega \cap \mathbb{C}_J^- \end{cases},$$

where $g, \widehat{g}, h, \widehat{h}$ are holomorphic functions defined on $\Omega \cap \mathbb{C}_J^+$. The expression of G and H in Formula (2.2) is chosen to be convenient for what follows. Note that, since \widehat{g} and \widehat{h} are holomorphic functions, then the two functions $v \mapsto \widehat{g}(\overline{v}), v \mapsto \widehat{h}(\overline{v})$ are holomorphic as well and if f_J is the restriction of a slice regular function defined over the whole algebra \mathbb{H} then $\widehat{g}(\overline{v}) = g(v)$ and $\widehat{h}(\overline{v}) = h(v)$.

Viceversa, given G, H holomorphic functions on $\Omega \cap \mathbb{C}_J$ defined as above, we can define a slice regular function $f : \Omega \rightarrow \mathbb{H}$ which splits over \mathbb{C}_J as

$$(2.3) \quad f(v) = G(v) + H(v)K = \begin{cases} g(v) + h(v)K & v \in \Omega \cap \mathbb{C}_J^+ \\ \widehat{g}(\bar{v}) + \widehat{h}(\bar{v})K & v \in \Omega \cap \mathbb{C}_J^- \end{cases},$$

by means of the Representation Formula: for any $\alpha + I\beta \in \Omega$, with $\beta > 0$,

$$(2.4) \quad \begin{aligned} f(\alpha + I\beta) &= \frac{1 - IJ}{2} f(\alpha + J\beta) + \frac{1 + IJ}{2} f(\alpha - J\beta) \\ &= \frac{1 - IJ}{2} (g(\alpha + J\beta) + h(\alpha + J\beta)K) + \frac{1 + IJ}{2} (\widehat{g}(\alpha + J\beta) + \widehat{h}(\alpha + J\beta)K). \end{aligned}$$

After having identified g, h, \widehat{g} and \widehat{h} with their extensions on $\mathbb{C}_J \setminus \mathbb{R}$ by Schwarz reflection (i.e. $g(\bar{v}) = \overline{g(v)}$), we can consider their regular extensions $\text{ext}(g), \text{ext}(h), \text{ext}(\widehat{g}), \text{ext}(\widehat{h})$. Then we have that

$$\begin{aligned} &(\text{ext}(g) + (\text{ext}(\widehat{h}))^c K) * \ell_+^J + ((\text{ext}(\widehat{g}))^c + \text{ext}(h)K) * \ell_-^J \\ &= \ell_+^J * (\text{ext}(g) + \text{ext}(h)K) + \ell_-^J * ((\text{ext}(\widehat{g}))^c + (\text{ext}(\widehat{h}))^c K). \end{aligned}$$

The previous expression restricted to $\Omega \cap \mathbb{C}_J^+$ coincides with $g + hK$, while restricted to $\Omega \cap \mathbb{C}_J^-$ coincides with $\widehat{g} + \widehat{h}K$, therefore, thanks to Identity Principle [1, Theorem 3.6], we get that

$$f = f_+ * \ell_+^J + f_- * \ell_-^J,$$

where

$$(2.5) \quad f_+ = \text{ext}(g) + (\text{ext}(\widehat{h}))^c K \quad \text{and} \quad f_- = (\text{ext}(\widehat{g}))^c + \text{ext}(h)K.$$

Summarising we have proved the following result.

Theorem 2.6. *Let $f : \Omega \rightarrow \mathbb{H}$ be a slice regular function on a circular domain Ω such that $\Omega \cap \mathbb{R} = \emptyset$. Then, for any $J \in \mathbb{S}$, there exist and are unique $f_+, f_- : \Omega \rightarrow \mathbb{H}$ slice regular functions, such that $f = f_+ * \ell_+^J + f_- * \ell_-^J$.*

Remark 2.7. The previous theorem is the restatement, in the context of slice regularity, of the well-known *Peirce decomposition* (see [21, Chapter 7, §21]).

Remark 2.8. If $\Omega \cap \mathbb{R} \neq \emptyset$, then, given any slice regular function $f : \Omega \rightarrow \mathbb{R}$, its restriction $f|_{\Omega \setminus \mathbb{R}}$ can be written as

$$f|_{\Omega \setminus \mathbb{R}} = f * \ell_+^J + f * \ell_-^J = f * (\ell_+^J + \ell_-^J),$$

for any $J \in \mathbb{S}$.

The class of functions that, as we will see, furnishes a new tool in twistor geometry is the following.

Definition 2.9 (slice-polynomial functions). A slice regular function $P : \mathbb{H} \setminus \mathbb{R} \rightarrow \mathbb{H}$ is said to be a *slice-polynomial function* if there exist $J \in \mathbb{S}$ and two quaternionic polynomials $P_+, P_- : \mathbb{H} \rightarrow \mathbb{H}$ such that

$$P = P_+ * \ell_+^J + P_- * \ell_-^J.$$

The next proposition shows that Definition 2.9 is well posed.

Proposition 2.10. *Let $P : \mathbb{H} \setminus \mathbb{R} \rightarrow \mathbb{H}$ be a slice-polynomial function, then for any $K \in \mathbb{S}$, there exist two quaternionic polynomials $Q_+, Q_- : \mathbb{H} \rightarrow \mathbb{H}$ such that*

$$P = Q_+ * \ell_+^K + Q_- * \ell_-^K.$$

Proof. Thanks to Remark 2.5, for any $K \in \mathbb{S}$, there exist $a, b, c, d \in \mathbb{H}$ such that

$$\ell_+^J = a * \ell_+^K + b * \ell_-^K, \quad \ell_-^J = c * \ell_+^K + d * \ell_-^K.$$

Thanks to Equation (2.1), by standard computations, $a = d$ and $b = c$, therefore

$$\begin{aligned} P &= P_+ * \ell_+^J + P_- * \ell_-^J \\ &= P_+ * (a * \ell_+^K + b * \ell_-^K) + P_- * (b * \ell_+^K + a * \ell_-^K) \\ &= (P_+ a + P_- b) * \ell_+^K + (P_+ b + P_- a) * \ell_-^K. \end{aligned}$$

□

As a consequence, we get that the restriction of a slice-polynomial function to any semi-slice is a polynomial and moreover we have the following characterization.

Proposition 2.11. *Let $P : \mathbb{H} \setminus \mathbb{R} \rightarrow \mathbb{H}$ be a slice regular function. P is a slice polynomial if and only if for any $K \in \mathbb{S}$ the restriction $P|_{\mathbb{C}_K^+}$ is a polynomial in the variable $\alpha + K\beta \in \mathbb{C}_K^+$.*

Proof. First of all notice that, if $q = \alpha + I\beta \in \mathbb{H}$ and $v = \alpha + i\beta \in \mathbb{C}_i$, then, by direct computation

$$q * \ell_+ = \ell_+ v, \quad q * \ell_- = \ell_- \bar{v}.$$

Therefore, for any $k, h \in \mathbb{N}$ and any $a, b \in \mathbb{H}$ we have that

$$\begin{aligned} (q^k a) * \ell_+ + (q^h b) * \ell_- &= (q^k a^\top + q^k a^\perp) * \ell_+ + (q^h b^\top + q^h b^\perp) * \ell_- \\ &= \ell_+(v^k a^\top + v^h b^\perp) + \ell_-(\bar{v}^k a^\perp + \bar{v}^h b^\top). \end{aligned}$$

Using the last computation, if P is a slice-polynomial function, we obtain the following equality

$$\begin{aligned} P(q) &= \left(\sum_{k=0}^N q^k a_k \right) * \ell_+ + \left(\sum_{h=0}^M q^h b_h \right) * \ell_- \\ &= \left(\sum_{k=0}^N q^k a_k^\top + q^k a_k^\perp \right) * \ell_+ + \left(\sum_{h=0}^M q^h b_h^\top + q^h b_h^\perp \right) * \ell_- \\ &= \ell_+ \left(\sum_{k=0}^N v^k a_k^\top + \sum_{h=0}^M v^h b_h^\perp \right) + \ell_- \left(\sum_{k=0}^N \bar{v}^k a_k^\perp + \sum_{h=0}^M \bar{v}^h b_h^\top \right). \end{aligned}$$

Therefore, thanks to Proposition 2.10 and the fact that, for any $K \in \mathbb{S}$, $\ell_+^K \equiv 1$ and $\ell_-^K \equiv 0$ on \mathbb{C}_K^+ , we immediately have the thesis. The opposite implication follows by reading the previous chain of equalities in the opposite direction. □

Thanks to Remark 2.8, any quaternionic polynomial $P : \mathbb{H} \rightarrow \mathbb{H}$ is a slice-polynomial function if restricted to $\mathbb{H} \setminus \mathbb{R}$. In fact, it holds

$$P|_{\mathbb{H} \setminus \mathbb{R}} = P * \ell_+ + P * \ell_-.$$

Example 2.12. It is easy to generate examples of slice-polynomial functions. However we recall here some slice-polynomial functions naively introduced in past researches by the first author [1, 3, 2]. In [1] the function $P(q) = q * \ell_+$ was introduced and analysed for its topological properties, showing in particular that it admits the whole semi-slice \mathbb{C}_i^- as zero set. Then in [3] slice regular functions defined over circular domains non intersecting the real axis were studied from a differential point of view and it was shown that the function $Q(q) = (q + j) * \ell_+$ admits two non-compact surfaces, both biholomorphic to the half-plane \mathbb{C}^+ , where it is constant. Finally in [2] the twistor geometry induced by $P(q)$ was studied and the

function $R(q) = -q^2 * \ell_+ + q * \ell_-$ was proposed as an example of slice regular function whose twistor lift parameterises a cubic scroll. All the functions listed in this example are slice-polynomial functions.

Thanks to Proposition 2.10, from now on, if not differently specified, we will always write slice-polynomial functions with respect to the imaginary unit i :

$$(2.6) \quad P = P_+ * \ell_+ + P_- * \ell_-.$$

We want now to express slice-polynomial functions in terms of the Splitting Lemma.

If P is a slice-polynomial function which splits on \mathbb{C}_i as

$$(2.7) \quad P(v) := \begin{cases} g(v) + h(v)j & v \in \mathbb{C}_i^+ \\ \widehat{g}(\bar{v}) + \widehat{h}(\bar{v})j & v \in \mathbb{C}_i^-, \end{cases}$$

we have that g, \hat{g}, h, \hat{h} are polynomials with coefficients in \mathbb{C}_i . Moreover, from the computations in the proof of Proposition 2.11 and Equation (2.4) we have that

$$\begin{aligned} P(\alpha + I\beta) &= \frac{1 - Ii}{2}(g(\alpha + i\beta) + h(\alpha + i\beta)j) + \frac{1 + Ii}{2}(\widehat{g}(\overline{(\alpha + i\beta)}) + \widehat{h}(\overline{(\alpha + i\beta)})j) \\ &= ((g + \hat{h}^c j) * \ell_+ + (\hat{g}^c + hj) * \ell_-)(\alpha + I\beta) \end{aligned}$$

where in the last equality, with a slight abuse of notation, we are identifying g, \hat{g}, h, \hat{h} with their regular extensions. Comparing the last representation with Formula (2.6), we obtain that $P_+ = g + \hat{h}^c j$ and $P_- = \hat{g}^c + hj$.

Example 2.13. Let us consider the polynomial $P(q) = q^2 + qi$, then $h = \hat{h} \equiv 0$ and $g(v) = \widehat{g}(\bar{v}) = v^2 + vi$. Therefore $\widehat{g}(\bar{v}) = \bar{v}^2 + \bar{v}i$

$$P(q) = q^2 + qi = \ell_+(v^2 + vi) + \ell_-(\bar{v}^2 + \bar{v}i).$$

The three slice-polynomial functions introduced in Example 2.12 can be written as follows

$$P(q) = \ell_+ v, \quad Q(q) = \ell_+ v + \ell_- j, \quad R(q) = \ell_+(-v^2) + \ell_- \bar{v}.$$

2.1. Extension to the real line. Given a slice-polynomial function $P = \mathcal{I}(F_1 + \iota F_2)$ we want to analyse its behaviour approaching the real line. Of course, since for any $I \in \mathbb{S}$ the function $P|_{\mathbb{C}_I^+}$ is a polynomial, then it can be extended to \mathbb{R} but the values that it reaches strongly depend on I . The right set to consider is therefore $\mathbb{S} \times \mathbb{R}$ and not merely \mathbb{R} . An element $(I, \alpha) \in \mathbb{S} \times \mathbb{R}$ will also be denoted by α_I . We define then the following function

$$P_{\mathbb{R}} : \mathbb{S} \times \mathbb{R} \rightarrow \mathbb{H}$$

$$(2.8) \quad P_{\mathbb{R}}(\alpha_I) = P_{\mathbb{R}}(I, \alpha) := \lim_{\beta \rightarrow 0^+} P(\alpha + I\beta) = \lim_{\beta \rightarrow 0^+} F_1(\alpha + i\beta) + IF_2(\alpha + i\beta) = F_1(\alpha) + IF_2(\alpha)$$

where the limit always exists, since $P|_{\mathbb{C}_I^+}$ is a polynomial on \mathbb{C}_I^+ . We define, then, the following sets

$$P_{\mathbb{R}}(\alpha) := \bigcup_{I \in \mathbb{S}} P_{\mathbb{R}}(\alpha_I), \quad P_{\mathbb{R}}(\mathbb{R}) := P_{\mathbb{R}}(\mathbb{S} \times \mathbb{R})$$

Notice that, if $P = \mathcal{I}(F_1 + \iota F_2)$ is a slice-polynomial function and there exists $\alpha \in \mathbb{R}$ such that $P(\alpha) = \{q\}$ for some $q \in \mathbb{H}$, then, for any $I, J \in \mathbb{S}$,

$$F_1(\alpha) + IF_2(\alpha) = q = F_1(\alpha) + JF_2(\alpha),$$

which entails that $F_2(\alpha) = 0$. For this reason the slice-polynomial function P can be extended to $\alpha \in \mathbb{R}$ as $P(\alpha) = q$.

If P is a quaternionic polynomial, then, thanks to the odd character of F_2 with respect to β , for any $\alpha \in \mathbb{R}$, $F_2(\alpha) = 0$ and so $P_{\mathbb{R}}(\alpha) = \{P(\alpha)\}$ and $P_{\mathbb{R}}(\mathbb{R}) = P(\mathbb{R})$. If P is not a quaternionic polynomial, then, $P_{\mathbb{R}}(\alpha)$ can be of dimension 2.

Example 2.14. Consider the slice-polynomial function $P(q) = q * \ell_+$, or, explicitly, $P(\alpha + I\beta) = (\alpha + I\beta) * \ell_+ = (\alpha + I\beta) \frac{1-Ii}{2}$. Therefore $P(\alpha + I\beta) = \frac{1}{2}(\alpha + I\beta - \alpha Ii + \beta i)$ and for any $I \in \mathbb{S}$ and any $\alpha \in \mathbb{R}$

$$P_{\mathbb{R}}(\alpha_I) = \lim_{\beta \rightarrow 0^+} P(\alpha + I\beta) = \alpha \frac{(1 - Ii)}{2}.$$

If $I = Ai + Bj + Ck$, then $P|_{\mathbb{R}}(\alpha_I)$ belongs to the real hyperplane $\{q_0 + q_1i + q_2j + q_3k \in \mathbb{H} \mid q_1 = 0\}$. Hence, for any $I \in \mathbb{S} \setminus \{-i\}$, $P|_{\mathbb{C}_I^+}(\mathbb{R})$ is a real line and for any $\alpha \neq 0$, $P_{\mathbb{R}}(\alpha)$ is a 2-sphere. It is not difficult to see that $P_{\mathbb{R}}(\mathbb{R})$ covers the whole hyperplane $\{q_0 + q_1i + q_2j + q_3k \in \mathbb{H} \mid q_1 = 0\}$ (see [2, Theorem 34]).

With similar computations, if $Q(q) = (q + j) * \ell_+$, $R(q) = -q^2 * \ell_+ + q * \ell_-$ and $q = \alpha + I\beta$, then

$$Q|_{\mathbb{R}}(\alpha_I) = \lim_{\beta \rightarrow 0^+} Q(\alpha + I\beta) = \alpha \frac{(1 - Ii)}{2} + \frac{(1 + Ii)}{2} j.$$

and

$$R|_{\mathbb{R}}(\alpha_I) = \lim_{\beta \rightarrow 0^+} R(\alpha + I\beta) = -\alpha^2 \frac{(1 - Ii)}{2} + \alpha \frac{(1 + Ii)}{2}.$$

For any fixed $I \in \mathbb{S} \setminus \{-i\}$ the set $Q_{\mathbb{R}}(\alpha_I)$ parameterises an affine line in \mathbb{H} passing through the point $\frac{(1+Ii)}{2} j$, therefore the resulting set $Q_{\mathbb{R}}(\mathbb{R})$ is a ruled 3-dimensional manifold. Finally, for any $I \in \mathbb{S}$, $R_{\mathbb{R}}(\alpha_I)$ parameterises a conic in \mathbb{H} . In particular $R_{\mathbb{R}}(\alpha_i)$ is a parabola while $R_{\mathbb{R}}(\alpha_{-i})$ is a straight line, both passing through the origin.

Remark 2.15. Given a slice-polynomial function $P = \mathcal{I}(F_1 + \iota F_2)$, the sphere \mathbb{S}_v is degenerate for P if and only if $\partial_s P(v) = 0$. Writing the quantity $P(v) - P(\bar{v})$ in terms of the splitting (2.7) we get,

$$\overline{\hat{g}(v)} + \hat{h}(v)j - g(v) - h(v)j = 0,$$

that is equivalent to the system

$$(2.9) \quad \begin{cases} \overline{\hat{g}(v)} - g(v) = 0 \\ \hat{h}(v) - h(v) = 0. \end{cases}$$

Since g, h, \hat{g} and \hat{h} are polynomials, the last notion can be extended to \mathbb{R} (and so to $P_{\mathbb{R}}$), saying that a real point α is degenerate for P if and only if $P_{\mathbb{R}}(\alpha) = \{q\}$ i.e. P can be defined at α as $P(\alpha) = q$. In particular, with this notion, any quaternionic polynomial P defined on \mathbb{H} is such that the real line is degenerate.

Remark 2.16. We point out that all the results proved in this subsection can be restored for any slice regular function suitable defined, i.e. for any slice regular function $f = \mathcal{I}(F_1 + \iota F_2)$ defined on a circular domain $\Omega \subseteq \mathbb{H} \setminus \mathbb{R}$ not intersecting the real axis, such that F_1 and F_2 extend continuously to \mathbb{R} .

2.2. Companion of a slice regular function. Let us introduce the notion of *companion* of a slice regular function. Given a slice regular function f , its companion is in some sense dual to f and, besides its algebraic definition, we will see that it naturally arises in the geometric construction described in Section 3.

Definition 2.17. Let $f : \Omega \rightarrow \mathbb{H}$ be a slice regular function and suppose that $f|_{\Omega \setminus \mathbb{R}} = f_+ * \ell_+ + f_- * \ell_-$. We define the *companion* of f as the slice regular function $f^\vee : \Omega \setminus \mathbb{R} \rightarrow \mathbb{H}$, defined by $f^\vee = f_- * \ell_+ + f_+ * \ell_-$.

Remark 2.18. The previous definition is well posed. Suppose in fact that $f = f_+ * \ell_+ + f_- * \ell_- = g_+ * \ell_+^J + g_- * \ell_-^J$, for some $J \in \mathbb{S}$, $J \neq i$. Then, if $\ell_+ = a * \ell_+^J + b * \ell_-^J$ (and so $\ell_- = b * \ell_+^J + a * \ell_-^J$), for suitable $a, b \in \mathbb{H}$, we have that

$$\begin{aligned} f &= f_+ * \ell_+ + f_- * \ell_- = f_+ * (a * \ell_+^J + b * \ell_-^J) + f_- * (b * \ell_+^J + a * \ell_-^J) \\ &= [f_- b + f_+ a] * \ell_+^J + [f_- a + f_+ b] * \ell_-^J = g_+ * \ell_+^J + g_- * \ell_-^J \end{aligned}$$

and hence that

$$f^\vee = [f_- a + f_+ b] * \ell_+^J + [f_- b + f_+ a] * \ell_-^J = g_- * \ell_+^J + g_+ * \ell_-^J.$$

Thanks to Remark 2.8, if $\Omega \cap \mathbb{R} \neq \emptyset$, then the companion f^\vee of $f : \Omega \rightarrow \mathbb{H}$ is f itself, i.e.: $f^\vee = f$.

Remark 2.19. Let $f : \Omega \rightarrow \mathbb{H}$ be any slice regular function that splits on \mathbb{C}_J as $f|_{\mathbb{C}_J} = G(v) + H(v)K$, as in Formula (2.3). Then, thanks to Equations (2.4) and (2.5), we get that its companion $f^\vee : \Omega \rightarrow \mathbb{H}$ splits on \mathbb{C}_J as

$$f^\vee(v) := \begin{cases} \overline{\hat{g}(\bar{v})} + \overline{\hat{h}(\bar{v})}K & v \in \mathbb{C}_J^+ \\ g(v) + h(v)K & v \in \mathbb{C}_J^- \end{cases}$$

In particular if P is a slice-polynomial function, then its companion P^\vee is a slice-polynomial function, and if P can be written at $\alpha + I\beta$ as $P(\alpha + I\beta) = \ell_+^J p_+(\alpha + J\beta) + \ell_-^J p_-(\alpha - J\beta)$, then $P^\vee = \ell_+^J p_-(\alpha + J\beta) + \ell_-^J p_+(\alpha - J\beta)$.

Remark 2.20. The degenerate sets of f and f^\vee do not necessarily coincide. Consider for instance the slice-polynomial function $P(q) = (q - q_0) * \ell_+ + (q - \bar{q}_0) * \ell_-$, with $q_0 \in \mathbb{C}_i^+$. The degenerate set of P consists of the sphere \mathbb{S}_{q_0} , where it is identically equal to zero. Its companion is $P^\vee(q) = (q - \bar{q}_0) * \ell_+ + (q - q_0) * \ell_-$, and it is not identically vanishing, neither constant on the same sphere. In fact $P^\vee(q_0) = q_0 - \bar{q}_0$ and $P^\vee(\bar{q}_0) = \bar{q}_0 - q_0$.

Proposition 2.21. *For any slice-polynomial function P we have that $P_{\mathbb{R}}(\mathbb{R}) = P_{\mathbb{R}}^\vee(\mathbb{R})$.*

Proof. Fix any $J \in \mathbb{S}$, then, thanks to Remark 2.19, we have that

$$P_{\mathbb{R}}(\alpha_J) = \lim_{\beta \rightarrow 0^+} P_+(\alpha + J\beta) = P_{\mathbb{R}}^\vee(\alpha_{-J}).$$

□

Remark 2.22. As before, the last proposition can be restored for any slice regular function suitable defined, i.e. for any $f = \mathcal{I}(F_1 + \iota F_2) : \Omega \subseteq \mathbb{H} \setminus \mathbb{R} \rightarrow \mathbb{H}$ such that F_1 and F_2 extend continuously to \mathbb{R} .

3. TWISTOR FIBRATION AND BLOW UP OF THE REAL LINE

In this section we recall some fact concerning the twistor interpretation of slice regular functions and we extend some known results contained in [2, 15].

Consider the *left* quaternionic projective space $\mathbb{H}\mathbb{P}^1$ as the set of equivalence classes $[q_1, q_2] \sim [pq_1, pq_2]$, for any $p \in \mathbb{H} \setminus \{0\}$. The twistor fibration: $\pi : \mathbb{C}\mathbb{P}^3 \rightarrow \mathbb{H}\mathbb{P}^1$ is defined as $\pi[X_0, X_1, X_2, X_3] = [X_0 + X_1j, X_2 + X_3j]$. We embed \mathbb{H} in $\mathbb{H}\mathbb{P}^1$ as $q \mapsto [1, q]$. So that if $q = q_1 + q_2j \in \mathbb{H}$ and $q_1, q_2 \in \mathbb{C}_i$, then

$$\pi[X_0, X_1, X_2, X_3] = [1, q] \Leftrightarrow [1, (X_0 + X_1j)^{-1}(X_2 + X_3j)] = [1, q_1 + q_2j],$$

and the last equality is equivalent to the following system which, therefore, describes the fibers of π

$$(3.1) \quad \begin{cases} X_2 = X_0q_1 - X_1\bar{q}_2 \\ X_3 = X_0q_2 + X_1\bar{q}_1. \end{cases}$$

Let $\mathcal{Q} \simeq (\mathbb{C}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^1) \subset \mathbb{C}\mathbb{P}^3$ be the non-singular *Segre quadric* defined by the equation $X_0X_3 = X_1X_2$, and define the *slice complex structure* over $\mathbb{H} \setminus \mathbb{R}$ as $\mathbb{J}_{\alpha+I\beta}(v) = Iv$, for $\beta > 0$ and for any $v \in T_{\alpha+I\beta}(\mathbb{H} \setminus \mathbb{R}) \simeq \mathbb{H}$. Then, as proven in [15], the quadric \mathcal{Q} contains two open subset \mathcal{Q}^+ and \mathcal{Q}^- both biholomorphic, via the map π , to $(\mathbb{H} \setminus \mathbb{R}, \mathbb{J})$. The set $\mathcal{Q}^+ \simeq (\mathbb{C}\mathbb{P}^1 \times \mathbb{C}^+)$ consists of points for which at least one of the following holds

- $X_0 \neq 0$ and $X_2/X_0 \in \mathbb{C}^+$;
- $X_1 \neq 0$ and $X_3/X_1 \in \mathbb{C}^+$.

Analogously $\mathcal{Q}^- \simeq (\mathbb{C}\mathbb{P}^1 \times \mathbb{C}^-) \subset \mathcal{Q}$ is defined as the set of points for which at least one of the following holds

- $X_0 \neq 0$ and $X_2/X_0 \in \mathbb{C}^-$;
- $X_1 \neq 0$ and $X_3/X_1 \in \mathbb{C}^-$.

To find coordinates for the fiber $\pi^{-1}(\alpha + I\beta)$ intersected with \mathcal{Q} we will suppose $X_0 \neq 0$. Denote then $I = Ai + Bj + Ck$ and $v = \alpha + i\beta$, with $\beta > 0$. Let $q = \alpha + I\beta = q_1 + q_2j$, with $q_1 = \alpha + \beta Ai$ and $q_2 = \beta B + \beta Ci$. Then the system defining the intersection,

$$\begin{cases} X_0X_3 = X_1X_2 \\ X_2 = X_0q_1 - X_1\bar{q}_2 \\ X_3 = X_0q_2 + X_1\bar{q}_1 \end{cases}$$

becomes

$$X_0^2q_2 + X_0X_1[\bar{q}_1 - q_1] + X_1^2\bar{q}_2 = 0$$

and imposing $X_0 = 1$, we get

$$q_2 + X_1[\bar{q}_1 - q_1] + X_1^2\bar{q}_2 = 0.$$

Solving the last equation we find $X_1 \in \left\{ u = -i\frac{B+iC}{1+A}, w = i\frac{B+iC}{1-A} \right\}$, and hence

$$\pi^{-1}(\alpha + I\beta) \cap \mathcal{Q}^+ = [1, u, v, uv], \quad \pi^{-1}(\alpha + I\beta) \cap \mathcal{Q}^- = [1, w, \bar{v}, w\bar{v}].$$

Analogously we get

$$\pi^{-1}(\alpha - I\beta) \cap \mathcal{Q}^+ = [1, w, v, wv], \quad \pi^{-1}(\alpha - I\beta) \cap \mathcal{Q}^- = [1, u, \bar{v}, u\bar{v}].$$

Notice that $w = -\bar{u}^{-1}$.

Consider now a slice regular function $f : \mathbb{H} \setminus \mathbb{R} \rightarrow \mathbb{H}$ which splits on \mathbb{C}_i as in Formula (2.3). As proven in [2, 15], f can be lifted on \mathcal{Q}^+ with the following non-homogeneous parameterisation:

$$(3.2) \quad [1, f(q)] = \pi[\tilde{f}[\pi^{-1}(q)]] = \pi[\tilde{f}[1, u, v, uv]] = \pi[1, u, g(v) - u\hat{h}(v), h(v) + u\hat{g}(v)].$$

With similar computations as in [2, 15] it is possible to prove that any slice regular function $f : \Omega \rightarrow \mathbb{H}$ defined on a circular domain $\Omega \subseteq \mathbb{H} \setminus \mathbb{R}$, can be lifted to a holomorphic map $\tilde{f}_- : \mathcal{Q}^- \cap \pi^{-1}(\Omega) \rightarrow \mathbb{C}\mathbb{P}^3$. The aim of the following lemma is to exhibit an explicit parameterisation for \tilde{f}_- .

Lemma 3.1. *Let $\Omega \subseteq \mathbb{H} \setminus \mathbb{R}$ and let $f : \Omega \rightarrow \mathbb{H}$ be a slice regular function that splits on \mathbb{C}_i as in Formula 2.3. Then, the parameterisation of the twistor lift $\tilde{f}_- : (\mathcal{Q}^- \cap \pi^{-1}(\Omega)) \rightarrow \mathbb{C}\mathbb{P}^3$ has coordinates*

$$\tilde{f}_-[1, w, v, wv] = [1, w, \overline{\hat{g}(v)} - w\overline{\hat{h}(v)}, \overline{\hat{h}(v)} + w\overline{\hat{g}(v)}].$$

Proof. Let $F_1 + \iota F_2$ be the stem function inducing f , and let $q = \alpha + I\beta \in \Omega$, $v = \alpha + i\beta \in \Omega \cap \mathbb{C}_i$ and $Q_w = 1 + wj$ be so that $\alpha + I\beta = Q_w^{-1}\bar{v}Q_w$. In the same spirit of [15, Theorem 5.3] and of [2, Theorem

24], the thesis is a consequence of the following sequence of equalities.

$$\begin{aligned}
[1, f(q)] &= [1, f(Q_w^{-1}(\alpha - i\beta)Q_w)] \\
&= [1, f(\alpha - Q_w^{-1}(i)Q_w\beta)] \\
&= [1, F_1(\alpha - i\beta) + Q_w^{-1}(i)Q_wF_2(\alpha - i\beta)] \\
&= [Q_w, Q_wF_1(\alpha - i\beta) + iQ_wF_2(\alpha - i\beta)] \\
&= [1 + wj, (1 + wj)F_1(\alpha - i\beta) + i(1 + wj)F_2(\alpha - i\beta)] \\
&= [1 + wj, f(\alpha - i\beta) + wj f(\alpha + i\beta)] \\
&= [1 + wj, f(\bar{v}) + wj f(v)] \\
&= [1 + wj, \overline{\hat{g}(v)} + \hat{h}(v)j + wj(g(v) + h(v)j)] \\
&= \pi[1, w, \overline{\hat{g}(v)} - w\overline{h(v)}, \overline{\hat{h}(v)} + w\overline{g(v)}].
\end{aligned}$$

□

Both Theorems 5.3 in [15] (or its extension [2, Theorem 24]), and Lemma 3.1 give a result that in the case of functions defined outside the real axis is only half-satisfactory, in the following sense. Consider a slice regular function $f : \mathbb{H} \setminus \mathbb{R} \rightarrow \mathbb{H}$ that splits as in Formula (2.3) such that $\overline{\hat{g}(v)} \neq g(v)$ or $\overline{\hat{h}(v)} \neq h(v)$. Then the commutation of the following diagrams given in [2, 15] does not extends to \mathcal{Q} (see [2, Section 6]).

$$\begin{array}{ccc}
\mathcal{Q}^+ & \xrightarrow{\tilde{f}_+} & \tilde{f}_+(\mathcal{Q}^+) \\
\pi \downarrow & & \downarrow \pi \\
\mathbb{H} \setminus \mathbb{R} & \xrightarrow{f} & f(\mathbb{H} \setminus \mathbb{R})
\end{array}
\qquad
\begin{array}{ccc}
\mathcal{Q}^- & \xrightarrow{\tilde{f}_-} & \tilde{f}_-(\mathcal{Q}^-) \\
\pi \downarrow & & \downarrow \pi \\
\mathbb{H} \setminus \mathbb{R} & \xrightarrow{f} & f(\mathbb{H} \setminus \mathbb{R})
\end{array}$$

On the contrary, if one considers a slice regular function defined on the whole \mathbb{H} , there is no such an issue (see [15, Section 7]). The aim of what follows is to overcome this problem and to unify the theory.

Proposition 3.2. *Let $f : \mathbb{H} \setminus \mathbb{R} \rightarrow \mathbb{H}$ be a slice regular function that splits on \mathbb{C}_i as in Formula (2.3). Suppose that the lift $\tilde{f}_+ : \mathcal{Q}^+ \rightarrow \mathbb{C}\mathbb{P}^3$ given by the parameterisation in (3.2) can be extended to \mathcal{Q} as $\mathcal{F} : \mathcal{Q} \rightarrow \mathbb{C}\mathbb{P}^3$ by varying the variable $v \in \mathbb{C}\mathbb{P}^1$. Then $\mathcal{F}|_{\mathcal{Q}^-} = \tilde{f}_-^\vee$, where $f^\vee : \mathbb{H} \setminus \mathbb{R} \rightarrow \mathbb{H}$ is the companion of f .*

Proof. Let $f : \mathbb{H} \setminus \mathbb{R} \rightarrow \mathbb{H}$ be a slice regular function as in the hypothesis and define $\mathcal{F} : \mathcal{Q} \rightarrow \mathbb{C}\mathbb{P}^3$ to be the extension of its lift on \mathcal{Q}^+ parameterised as

$$\mathcal{F}[1, u, v, uv] = [1, u, g(v) - u\hat{h}(v), h(v) + u\hat{g}(v)].$$

The possibility to extend \tilde{f}_+ , in particular implies that g, \hat{g}, h and \hat{h} can be extended to \mathbb{C} . Consider now a slice regular function φ that splits on \mathbb{C}_i as

$$\varphi := \begin{cases} l(v) + m(v)j & v \in \mathbb{C}_i^+ \\ \overline{\hat{l}(v)} + \overline{\hat{m}(v)}j & v \in \mathbb{C}_i^- \end{cases}$$

Then, according to Lemma 3.1, the lift of φ on \mathcal{Q}_- is given by

$$\tilde{\varphi}_-[1, u, v, uv] = [1, u, \overline{\hat{l}(v)} - u\overline{\hat{m}(v)}, \overline{\hat{m}(v)} + u\overline{\hat{l}(v)}].$$

For $\tilde{\varphi}_-$ to agree with $\mathcal{F}|_{\mathcal{Q}^-}$ we then have that, for any $v \in \mathbb{C}^-$

$$\overline{\hat{l}(v)} = g(v), \quad \overline{\hat{m}(v)} = \hat{h}(v), \quad \overline{\hat{m}(v)} = h(v), \quad \overline{\hat{l}(v)} = \hat{g}(v),$$

therefore, the splitting of φ is given by

$$\varphi := \begin{cases} \overline{\hat{g}(\bar{v})} + \overline{\hat{h}(\bar{v})}j & v \in \mathbb{C}_i^+ \\ g(v) + h(v)j & v \in \mathbb{C}_i^-, \end{cases}$$

that is $\varphi = f^\vee$, and we have proved the proposition. \square

3.1. Blow-up of the real line. We now want to give a better description of the role of the real line in this framework. Since $\mathbb{H} \setminus \mathbb{R} \simeq \mathbb{S} \times \mathbb{C}^+$, we consider the following embedding

$$\begin{aligned} \mathbb{H} \setminus \mathbb{R} &\hookrightarrow \mathbb{S} \times \overline{\mathbb{C}^+}, \\ \alpha + I\beta &\mapsto (I, \alpha + i\beta). \end{aligned}$$

As recalled at the beginning of this section, the complex manifold $(\mathbb{H} \setminus \mathbb{R}, \mathbb{J})$ is biholomorphic to the two open subsets \mathcal{Q}^+ and \mathcal{Q}^- endowed with the standard complex structure. Computing the pre-images of the real line by means of the projection π , one has that $\mathcal{Q}_{\mathbb{R}} := \pi^{-1}(\mathbb{R}) \cap \mathcal{Q} \simeq \mathbb{C}\mathbb{P}^1 \times \mathbb{S}^1$. Clearly $\overline{\mathcal{Q}^+} = \mathcal{Q}^+ \cup \mathcal{Q}_{\mathbb{R}}$ and $\overline{\mathcal{Q}^-} = \mathcal{Q}^- \cup \mathcal{Q}_{\mathbb{R}}$. As explained in [25, Section 7.3], $\mathcal{Q}_{\mathbb{R}}$ is a 3-real-dimensional submanifold of \mathcal{Q} , which disconnects \mathcal{Q} into two components, namely \mathcal{Q}^+ and \mathcal{Q}^- . Taking care of these observations, we now extend the twistor projection $\pi|_{\mathcal{Q}}$ to $\mathbb{S} \times \overline{\mathbb{C}^+}$ as follows. If $\alpha \in \mathbb{R}$ and $[s, u] \in \mathbb{C}\mathbb{P}^1$ corresponds to the element $I \in \mathbb{S}$, then $\pi[s, u, s\alpha, u\alpha] = (I, \alpha) = \alpha_I$, where we use the same symbol π since there will not be any confusion.

$$\begin{aligned} \pi : \quad \overline{\mathcal{Q}^+} &\rightarrow \mathbb{S} \times \overline{\mathbb{C}^+} \\ [s, u, sv, uv] &\mapsto \begin{cases} \alpha + I\beta & \text{if } v \in \mathbb{C}^+ \\ \alpha_I & \text{if } v \in \mathbb{R} \end{cases} \end{aligned}$$

The map π just defined, obviously extends the twistor projection and is bijective everywhere. For this reason we may exploit it for our purposes.

Given a slice regular function defined away from the reals, such that the functions defining its splitting can be extended to \mathbb{R} , we want to give an explicit parameterisation of its lift to $\pi^{-1}(\mathbb{R}) \cap \mathcal{Q}$, by means of the just defined extension of π to $\mathbb{S} \times \overline{\mathbb{C}^+}$.

Lemma 3.3. *Let $f : \mathbb{H} \setminus \mathbb{R} \rightarrow \mathbb{H}$ be a slice regular function. Assume that f splits on \mathbb{C}_i as in Formula 2.3 and that g, \hat{g}, h, \hat{h} can be extended to \mathbb{R} . Then defining $f_{\mathbb{R}} : \mathbb{S} \times \mathbb{R} \rightarrow \mathbb{H}$ as in Equation 2.8, there exists a differentiable function $\tilde{f}_{\mathbb{R}} : \mathcal{Q}_{\mathbb{R}} \rightarrow \mathbb{C}\mathbb{P}^3$ such that $f_{\mathbb{R}} = \pi \circ \tilde{f}_{\mathbb{R}} \circ \pi^{-1}$. Moreover, for any $\alpha \in \mathbb{R}$ and for any $I \in \mathbb{S}$ (corresponding to $[1, u] \in \mathbb{C}\mathbb{P}^1$), the parameterisation of $\tilde{f}_{\mathbb{R}}$ is given by*

$$\tilde{f}_{\mathbb{R}}[1, u, \alpha, u\alpha] = [1, u, g(\alpha) - u\hat{h}(\alpha), h(\alpha) + u\hat{g}(\alpha)].$$

Proof. As in the proof of Lemma 3.1, let $F_1 + \iota F_2$ be the stem function inducing f , and let $q = \alpha + I\beta \in \Omega$, $v = \alpha + i\beta \in \Omega \cap \mathbb{C}_i$ and $Q_u = 1 + uj$ be such that $\alpha + I\beta = Q_u^{-1}vQ_u$. Then, taking into account the definitions of $f_{\mathbb{R}}$ and of the extension of π to $\overline{\mathcal{Q}^+}$ and the fact that π is bijective, the thesis is a consequence of the following sequence of equalities.

$$\begin{aligned} [1, f(\alpha_I)] &= [1, \lim_{\beta \rightarrow 0^+} f(Q_u^{-1}(\alpha + i\beta)Q_u)] \\ &= [1, F_1(\alpha) + Q_u^{-1}(i)Q_u F_2(\alpha)] \\ &= [Q_u, Q_u F_1(\alpha) + iQ_u F_2(\alpha)] \\ &= [1 + uj, (1 + uj)F_1(\alpha) + i(1 + uj)F_2(\alpha)] \\ &= [1 + uj, F_1(\alpha) + iF_2(\alpha) + uj(F_1(\alpha) - iF_2(\alpha))] \\ &= [1 + uj, f(\alpha_i) + uj(f(\alpha_{-i}))] \\ &= [1 + uj, g(\alpha) + h(\alpha)j + uj(\overline{\hat{g}(\alpha)} + \overline{\hat{h}(\alpha)}j)] \\ &= \pi[1, u, g(\alpha) - u\hat{h}(\alpha), h(\alpha) + u\hat{g}(\alpha)]. \end{aligned}$$

□

Remark 3.4. What we have obtained in the previous lemma can be synthesised by saying that, given a slice regular function $f : \mathbb{H} \setminus \mathbb{R} \rightarrow \mathbb{H}$ that splits as in Formula 2.3 and such that the functions g, \hat{g}, h and \hat{h} can be extended to \mathbb{R} , then the following diagram commutes

$$\begin{array}{ccc} \overline{\mathcal{Q}^+} & \xrightarrow{\tilde{f}_+} & \tilde{f}_+(\overline{\mathcal{Q}^+}) \\ \pi \downarrow & & \downarrow \pi \\ \mathbb{S} \times \overline{\mathbb{C}^+} & \xrightarrow{f} & f(\mathbb{S} \times \overline{\mathbb{C}^+}) \end{array} \quad \begin{array}{ccc} [1, u, v, uv] & \xrightarrow{\tilde{f}_+} & [1, u, g(v) - u\hat{h}(v), h(v) + u\hat{g}(v)] \\ \pi \downarrow & & \downarrow \pi \\ \alpha + I\beta & \xrightarrow{f} & f(\alpha + I\beta) \end{array}$$

Analogous considerations hold for \tilde{f}_- .

4. PRE-IMAGES OF A SLICE-POLYNOMIAL FUNCTION

Let P be a slice-polynomial function which splits on \mathbb{C}_i as in Formula (2.7). We denote by $\tilde{P}_+ : \mathcal{Q}^+ \rightarrow \mathbb{C}\mathbb{P}^3$ the lift of P on \mathcal{Q}^+ , by $\tilde{P}_-^\vee : \mathcal{Q}^- \rightarrow \mathbb{C}\mathbb{P}^3$ the lift of P^\vee on \mathcal{Q}^- and by $\tilde{P}_\mathbb{R} : \mathcal{Q}_\mathbb{R} \rightarrow \mathbb{C}\mathbb{P}^3$ the lift of $P_\mathbb{R}$ on $\mathcal{Q}_\mathbb{R}$.

Definition 4.1. We define the *extended twistor lift* of P as the function $\mathcal{P} : \mathcal{Q} \rightarrow \mathbb{C}\mathbb{P}^3$ given by

$$\mathcal{P}(X) := \begin{cases} \tilde{P}_+(X), & \text{if } X \in \mathcal{Q}^+, \\ \tilde{P}_\mathbb{R}(X), & \text{if } X \in \mathcal{Q}_\mathbb{R}, \\ \tilde{P}_-^\vee(X), & \text{if } X \in \mathcal{Q}^-. \end{cases}$$

Thanks to [15, Theorem 5.3] and its extensions (the already cited [2, Theorem 24] and Lemmas 3.1 and 3.3), we are able to give a parameterisation of \mathcal{P} by means of the splitting of P :

$$\mathcal{P}[1, u, v, uv] = [1, u, g(v) - u\hat{h}(v), h(v) + u\hat{g}(v)],$$

where $u, v \in \mathbb{C}_i$. By its definition u parameterises all the imaginary units of \mathbb{S} except for $-i$. For this reason, recalling the *Segre embedding* $(\mathbb{C}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^1) \simeq \mathcal{Q}$ given by $([s, u], [t, v]) \mapsto [st, ut, sv, uv]$, it is useful to consider the parameterisation of \mathcal{P} homogenised with respect to the variable u

$$\mathcal{P}[s, u, sv, uv] = [s, u, sg(v) - u\hat{h}(v), sh(v) + u\hat{g}(v)].$$

At this stage, the idea is to study simultaneously the pre-images of a point $q \in \mathbb{H}$ via P, P^\vee and $P_\mathbb{R}$ by means of their extended twistor lift \mathcal{P} . This is possible having in mind the commutative properties of the following three diagrams

$$\begin{array}{ccc} \mathcal{Q}^+ & \xrightarrow{\tilde{P}_+} & \tilde{P}_+(\mathcal{Q}^+) & \mathcal{Q}^- & \xrightarrow{\tilde{P}_-^\vee} & \tilde{P}_-^\vee(\mathcal{Q}^-) & \mathcal{Q}_\mathbb{R} & \xrightarrow{\tilde{P}_\mathbb{R}} & \tilde{P}_\mathbb{R}(\mathcal{Q}_\mathbb{R}) \\ \pi \downarrow & & \downarrow \pi & \pi \downarrow & & \downarrow \pi & \pi \downarrow & & \downarrow \pi \\ \mathbb{H} \setminus \mathbb{R} & \xrightarrow{P} & P(\mathbb{H} \setminus \mathbb{R}) & \mathbb{H} \setminus \mathbb{R} & \xrightarrow{P^\vee} & P^\vee(\mathbb{H} \setminus \mathbb{R}) & \mathbb{S} \times \mathbb{R} & \xrightarrow{P_\mathbb{R}} & P_\mathbb{R}(\mathbb{S} \times \mathbb{R}) \end{array}$$

and the fact that $\mathcal{Q}_\mathbb{R}$ disconnects \mathcal{Q} into \mathcal{Q}^+ and \mathcal{Q}^- . Let us denote by $\mathcal{S} = \mathcal{P}(\mathcal{Q}) \subset \mathbb{C}\mathbb{P}^3$ the surface parameterised by the extended lift of P . If a point $q \in \mathbb{H}$ lifted to $\mathcal{S} = \tilde{P}_+(\mathcal{Q}^+) \cup \tilde{P}_-^\vee(\mathcal{Q}^-) \cup \tilde{P}_\mathbb{R}(\mathcal{Q}_\mathbb{R})$ has a pre-image via \mathcal{P} in $\mathcal{Q}^+, \mathcal{Q}^-$ or $\mathcal{Q}_\mathbb{R}$, then it will have a pre-image in $\mathbb{S} \times \overline{\mathbb{C}^+}$ with respect to P, P^\vee or $P_\mathbb{R}$, respectively.

Since the fibres of π are parameterised by System (3.1), then, for any $q = q_1 + q_2j \in \mathbb{H}$, we have that the intersection $\pi^{-1}(q) \cap \mathcal{S}$ is parameterised by

$$\begin{cases} sg(v) - u\hat{h}(v) = sq_1 - u\bar{q}_2 \\ sh(v) + u\hat{g}(v) = sq_2 + u\bar{q}_1 \end{cases}$$

and, therefore, we are left to find solutions $([s, u], v) \in \mathbb{C}\mathbb{P}^1 \times \mathbb{C}$ of the system

$$(4.1) \quad \begin{cases} s(g(v) - q_1) - u(\hat{h}(v) - \bar{q}_2) = 0 \\ s(h(v) - q_2) - u(\bar{q}_1 - \hat{g}(v)) = 0. \end{cases}$$

Let us denote by $A_q(v)$ the two by two complex matrix associate to the previous system,

$$A_q(v) = \begin{pmatrix} g(v) - q_1 & -\hat{h}(v) + \bar{q}_2 \\ h(v) - q_2 & -\bar{q}_1 + \hat{g}(v) \end{pmatrix},$$

by $D_q(v)$ the determinant of $A_q(v)$,

$$(4.2) \quad D_q(v) = -(g(v) - q_1)(\bar{q}_1 - \hat{g}(v)) + (h(v) - q_2)(\hat{h}(v) - \bar{q}_2),$$

and by $d = \max_{q \in \mathbb{H}} \{\deg D_q(v)\}$ its degree. This complex polynomial depends on q but its degree is generically constant and equal to d . The following theorem describes the set of solutions of System (4.1) for a fixed q .

Theorem 4.1. *Let P be a slice-polynomial function, not slice-constant, which splits on \mathbb{C}_i^+ as*

$$P(v) := \begin{cases} g(v) + h(v)j & v \in \mathbb{C}_i^+ \\ \hat{g}(\bar{v}) + \hat{h}(\bar{v})j & v \in \mathbb{C}_i^-, \end{cases}$$

let $q = q_1 + q_2j \in \mathbb{H}$ be any quaternion and let $A_q(v)$, $D_q(v)$ and d be defined as above. Then

- if there exists $v_0 \in \mathbb{C}_i$ such that $A_q(v_0) = 0$, then $([s, u], v_0)$ is a solution of System (4.1) for any $[s, u] \in \mathbb{C}\mathbb{P}^1$;
- if $D_q(v) \equiv 0$ then, for any v such that $A_q(v) \neq 0$, there exists and is unique $[s(v), u(v)] \in \mathbb{C}\mathbb{P}^1$ such that $([s(v), u(v)], v)$ is a solution of System (4.1);
- if $D_q(v) \not\equiv 0$ and v_1, \dots, v_d are the (possibly repeated) roots of $D_q(v)$, then, if $A_q(v_k) \neq 0$, there exists and is unique $[s_k, u_k] \in \mathbb{C}\mathbb{P}^1$ such that $([s_k, u_k], v_k)$ is a solution of System (4.1).

Proof. Note that the fact that P is not slice-constant implies that $A_q(v)$ is not a constant matrix. In order to get solutions $[s, u] \in \mathbb{C}\mathbb{P}^1$ of System (4.1), the determinant $D_q(v)$ must vanish. If there exists v_0 such that $A_q(v_0) = 0$, then it is easy to see that for any $[s, u] \in \mathbb{C}\mathbb{P}^1$ the pair $([s, u], v_0)$ is a solution of System (4.1) in \mathcal{Q} .

Suppose now that $D_q(v)$ vanishes identically. This is equivalent to say that the rows $(A_q)_1(v) = (g(v) - q_1, -\hat{h}(v) + \bar{q}_2)$ and $(A_q)_2(v) = (h(v) - q_2, -\bar{q}_1 + \hat{g}(v))$ of $A_q(v)$ are identically linearly dependent. Suppose then that, without loss of generality, $(A_q)_1(v) \neq (0, 0)$. Then the unique solution of System (4.1) in $\mathbb{C}\mathbb{P}^1 \times \mathbb{C}$ is given by $([s(v), u(v)], v) = ([\hat{h}(v) - \bar{q}_2, g(v) - q_1], v)$.

If otherwise $D_q(v)$ is not identically zero, then it has exactly d roots v_1, \dots, v_d (possibly repeated according to their multiplicity). If $A_q(v_k) \neq 0$ (supposing without loss of generality that $(A_q)_1(v_k) \neq (0, 0)$) the pair $([\hat{h}(v_k) - \bar{q}_2, g(v_k) - q_1], v_k)$ is a solution of System (4.1) in \mathcal{Q} . \square

The previous result can be interpreted in terms of pre-images of slice-polynomial functions. In fact, in the following corollary, we give a complete description of the simultaneous pre-image of a point $q \in \mathbb{H}$ via a slice-polynomial function, its companion and its extension to the reals.

Corollary 4.2. *Let P be a slice-polynomial function, not slice-constant, which splits on \mathbb{C}_i^+ as in Formula (2.7), let q be any quaternion and let $A_q(v)$, $D_q(v)$ and d be defined as above. Then*

- *if there exists $v_0 \in \mathbb{C}_i$ such that $A_q(v_0) = 0$, then if $v_0 \notin \mathbb{R}$, then $P(\mathbb{S}_{v_0}) \equiv q$ or $P^\vee(\mathbb{S}_{v_0}) \equiv q$, otherwise, if $v_0 \in \mathbb{R}$, then $P_{\mathbb{R}}(v_0) = q$ and P, P^\vee can be extended continuously to v_0 as $P(v_0) = P^\vee(v_0) = q$;*
- *if $D_q(v) \equiv 0$ then there exist two real surfaces $\Sigma, \Sigma^\vee \subset \mathbb{H} \setminus \mathbb{R}$, both biholomorphic to \mathbb{C}^+ , such that $P(\Sigma) \equiv P^\vee(\Sigma^\vee) \equiv q$;*
- *if $D_q(v) \not\equiv 0$, let v_1, \dots, v_d be the roots of $D_q(v)$ repeated according to their multiplicity. If $A_q(v_k) \neq 0$ for all k , then*

$$\#\{P^{-1}(q)\} + \#\{P^{\vee-1}(q)\} + \#\{P_{\mathbb{R}}^{-1}(q)\} = d.$$

Proof. Suppose first that there exists $q \in \mathbb{H}$ and $v_0 \in \mathbb{C}$ such that $A_q(v_0) = 0$. As shown in Theorem 4.1, for any $[s, u] \in \mathbb{C}\mathbb{P}^1$, $([s, u], v_0)$ is a solution of the System (4.1). If $v_0 \in \mathbb{C} \setminus \mathbb{R}$, since $\pi : \mathcal{Q}^+ \rightarrow \mathbb{H} \setminus \mathbb{R}$ and $\pi : \mathcal{Q}^- \rightarrow \mathbb{H} \setminus \mathbb{R}$ are both biholomorphisms, we have that $\mathbb{S}_{v_0} = P^{-1}(q)$ or $\mathbb{S}_{v_0} = P^{\vee-1}(q)$. If $v_0 \in \mathbb{R}$, then, thanks to Remark 2.15 and to Lemma 3.3, $\mathbb{S} \times \{v_0\} = P_{\mathbb{R}}^{-1}(q)$, that is $P_{\mathbb{R}}(v_0) = q$.

If now $D_q(v) \equiv 0$, then, for any $v \in \mathbb{C}$ such that $A_q(v) \neq 0$, there exists $[s(v), u(v)] \in \mathbb{C}\mathbb{P}^1$, such that $([s(v), u(v)], v)$ is a solution of System (4.1). Suppose for simplicity that the first line $(A_q)_1(v)$ is not identically zero, then, $([\tilde{h}(v) - \bar{q}_2, g(v) - q_1], v) \subset \mathbb{C}\mathbb{P}^1 \times \mathbb{C}$ defines a holomorphic surface \mathcal{S} . If for some v_0 $A_q(v_0) = 0$, then thanks Theorem 4.1, for all $[s, u] \in \mathbb{C}\mathbb{P}^1$, $([s, u], v_0)$ is a solution of the System (4.1), and so, in particular the surface $([\tilde{h}(v) - \bar{q}_2, g(v) - q_1], v) \subset \mathbb{C}\mathbb{P}^1 \times \mathbb{C}$ can be extended to v_0 holomorphically by taking $([\tilde{h}(v_0) - \bar{q}_2, g(v_0) - q_1], v_0)$. Now, since, again π is a biholomorphism if restricted to \mathcal{Q}^+ or to \mathcal{Q}^- , then $\Sigma := \pi(\mathcal{S} \cap \mathcal{Q}^+) \subseteq P^{-1}(q)$ and $\Sigma^\vee := \pi(\mathcal{S} \cap \mathcal{Q}^-) \subseteq P^{\vee-1}(q)$ are two holomorphic surfaces.

In the remaining case, let v_1, \dots, v_d be the roots of $D_q(v)$ repeated according to their multiplicity. Suppose that $A_q(v_k) \neq 0$ for all k , then by Theorem 4.1, there exists and is unique $[s_k, u_k] \in \mathbb{C}\mathbb{P}^1$ such that $\pi(\mathcal{P}[s_k, u_k, s_k v_k, u_k v_k]) = q$. But then, depending whether v_k belongs to $\mathbb{C}^+, \mathbb{C}^-$ or to \mathbb{R} , it will provide a pre-image for P, P^\vee or $P_{\mathbb{R}}$, respectively and these three options are obviously mutually exclusive. \square

Thanks to the previous corollary we are able to give the following definition.

Definition 4.3. Let P be a slice-polynomial function. We define the *twistor degree* of P as $\widetilde{\deg}(P) = d = \deg D_q$, where D_q is defined as in Formula (4.2).

The two surfaces Σ and Σ^\vee in the statement of Corollary 4.2 glue together at the real line, in the following sense: the surface \mathcal{S} defined in the proof of the Corollary parameterises a holomorphic surface in $\mathbb{C}\mathbb{P}^1 \times \mathbb{C} \subset \mathcal{Q}$. But $\pi : \mathcal{Q} \rightarrow \mathbb{H}$ is a class \mathcal{C}^∞ map whose restrictions to \mathcal{Q}^+ and \mathcal{Q}^- have Σ and Σ^\vee as images, respectively and these, of course, glue together at their boundary, given by $\pi(\mathcal{S} \cap \mathcal{Q}_{\mathbb{R}})$.

Remark 4.4. Given a quaternionic polynomial Q of degree n we have that $\overline{\hat{g}(\bar{v})} = g(v)$ and $\overline{\hat{h}(\bar{v})} = h(v)$, hence $\widetilde{\deg}(Q) = 2n$. Moreover, thanks to these symmetries, to any solution $v \in \mathbb{C}_i^+$ of $D_q(v) = 0$ corresponds another solution in \mathbb{C}_i^- , coherently with the fact that $Q^\vee = Q$. In particular, for this reason, it was not possible to see all this peculiar aspects of the theory in the context of slice regular functions defined over the reals as in the case studied in [15].

Remark 4.5. If P is a slice-constant function, then g, \hat{g}, h, \hat{h} are constant functions and then System (4.1) reduces to

$$\begin{cases} sA - uB = 0 \\ sC - uD = 0. \end{cases}$$

fore some constants $A, B, C, D \in \mathbb{C}_i$. Hence the system has admissible solutions if and only if the corresponding determinant $D_q(v) = -AD + BC = 0$. In this case for any $v \in \mathbb{C}_i$ the unique solution is given by $[s, u] = [B, -A] \in \mathbb{CP}^1$.

Example 4.6. Let us exhibit some examples of analysis of the System in 4.1.

- Consider the slice-constant function ℓ_+ . For this function the splitting is given by $g \equiv 1$ and $\hat{g} \equiv h \equiv \hat{h} \equiv 0$. Therefore, for any $v \in \mathbb{C}_i$, the matrix A_q is given by

$$A_q(v) = \begin{pmatrix} 1 - q_1 & -\bar{q}_2 \\ -q_2 & \bar{q}_1 \end{pmatrix},$$

and $D_q(v) = \bar{q}_1 - |q_1|^2 - |q_2|^2$. This is a constant function and is equal to zero only when $\bar{q}_1 \in \mathbb{R}$ and $q_1 = |q|^2$. If $q = \alpha + I\beta$, then, translating the last condition, the image of ℓ_+ can be expressed as follows

$$\ell_+(\mathbb{H} \setminus \mathbb{R}) := \{\alpha + I\beta \in \mathbb{H} \mid I \perp i, \alpha = \alpha^2 + \beta^2\},$$

representing, for any $I \in \mathbb{S}$ a circle of radius $1/2$ passing through 0 and 1 with center in $1/2$. As expressed in Remark 4.5, for any $q \in \ell_+(\mathbb{H} \setminus \mathbb{R})$ there is an element $[s, u] \in \mathbb{CP}^1$ which solves System (4.1) for any $v \in \mathbb{C}$. The last implies that any $q \in \ell_+(\mathbb{H} \setminus \mathbb{R})$ has a semi-slice of pre-images.

- We pass now to the function $q\ell_+$. This function splits as $g(v) = v$ and $\hat{g} \equiv h \equiv \hat{h} \equiv 0$. The matrix A_q is given by

$$A_q(v) = \begin{pmatrix} v - q_1 & -\bar{q}_2 \\ -q_2 & \bar{q}_1 \end{pmatrix},$$

and $D_q(v) = v\bar{q}_1 - |q_1|^2 - |q_2|^2$. This is identically zero for $q = 0$ and for this value, the system in Formula (4.1) as a solution for $s = 0$, showing that $q\ell_+|_{\mathbb{C}_i^+} \equiv 0$. For any $q \neq 0$ there exists and is unique the solution $([s, u], v)$ of System (4.1). Notice moreover that, for $q = 0$ it holds $A_0(0) \equiv 0$, therefore $q\ell_+$ can be defined in 0 and 0 is degenerate for the function.

- We study now the slice-polynomial function $(q + j) * \ell_+ = \ell_+v + \ell_-j$. The matrix $A_q(v)$ for this function is the following

$$A_q(v) = \begin{pmatrix} v - q_1 & 1 - \bar{q}_2 \\ -q_2 & \bar{q}_1 \end{pmatrix},$$

and $D_q(v) = v\bar{q}_1 + q_2 - |q_1|^2 - |q_2|^2$. This polynomial is again identically zero for $q = 0$ but also for $q = j$. For these two values for any $v \in \mathbb{C}$ there exists $[s, u] \in \mathbb{CP}^1$ that solve the system in (4.1). In particular, for any v , the point $q = 0$ is obtained for $[1, v] \in \mathbb{CP}^1$, while $q = j$, for $[0, 1]$. For this function $A_q(v) \neq 0$ for any $q \in \mathbb{H}$.

Corollary 4.7. *Let P be a slice-polynomial function, not slice-constant. Then*

$$P(\mathbb{H} \setminus \mathbb{R}) \cup P^\vee(\mathbb{H} \setminus \mathbb{R}) \cup P(\mathbb{R}) = \mathbb{H}.$$

Proof. It is a direct consequence of Theorem 4.1 and Corollary 4.2, keeping in mind that, if P is not slice-constant, then at least one among g, \hat{g}, h and \hat{h} is not a constant polynomial. Therefore, for any $q \in \mathbb{H}$ $D_q(v) = 0$ has at least a root and the System (4.1) can be solved. \square

Since the lift \mathcal{P} of a slice-polynomial function is parameterised by polynomial functions, then its image lies on an algebraic surface \mathcal{K} . The twistor projection has \mathbb{CP}^1 as fibres and therefore, they generically intersect \mathcal{K} in $\deg \mathcal{K}$ points. Hence we have the following corollary.

Corollary 4.8. *Let P be a slice-polynomial function, which splits on \mathbb{C}_i as in Formula (2.7) and let $d = \widetilde{\deg}(Q)$ be its twistor degree. Assume that the extended twistor lift \mathcal{P} of P is generically $k : 1$ on its image, then the image of \mathcal{P} lies in an algebraic surface of degree d/k .*

Proof. The thesis is trivial and we just add the following synthesising graph.

$$\begin{array}{ccc}
 Q & \xrightarrow[\mathcal{P}]{k:1} & \mathcal{K} \\
 & \searrow^{d:1} & \downarrow \pi \\
 & & \mathbb{H}
 \end{array}$$

□

Remark 4.9. The last corollary clarifies some issues of the twistor interpretation of slice regularity. First of all, thanks to Remark 4.4, given any quaternionic polynomial Q of degree n , one has that $d = \widetilde{\deg}(Q) = 2n$. Therefore, if the extended lift \mathcal{Q} of Q is generically $1:1$ and it has image on an algebraic surface \mathcal{K} , then $\deg \mathcal{K} = d = 2n$. Hence, given an algebraic surface $\mathcal{K} \subset \mathbb{C}\mathbb{P}^3$ of odd degree, it is not possible to parameterise it by means of the twistor lift of a slice regular function in a generically injective way (see e.g. [15, Sections 6 and 7] where a quaternionic polynomial of degree 2 allows to study a rational quartic scroll). Moreover, the content of the present section completely solves the problem exposed in [2, Remark 17] where it was pointed out that it is not sufficient to consider the twistor lift of a slice regular function to parameterise even simple algebraic surfaces. In particular, the *ghost function* proposed in [2, Remark 17] is what we call *companion*.

Remark 4.10. Most of the techniques introduced in this section can be applied to a generic slice regular function $f: \mathbb{H} \setminus \mathbb{R} \rightarrow \mathbb{H}$ for the study of its pre-images obtaining general results. For instance the Open Mapping Theorem does not imply that the image of a slice regular function defined on a domain without real points is an open set (see [1, Remark 5.2]). However Corollary 4.7 shows that the union of the three images $P(\mathbb{H} \setminus \mathbb{R}) \cup P^\vee(\mathbb{H} \setminus \mathbb{R}) \cup P(\mathbb{R})$ is an open set. This property is likely to be generalised to the case of slice regular functions. This study goes beyond the aims of the present work but it would be really interesting to explore it in the future.

5. DISCRIMINANT LOCUS OF A CUBIC SCROLL

The aim of this section is to describe the discriminant locus of a cubic scroll in $\mathbb{C}\mathbb{P}^3$ by means of the introduced techniques on slice-polynomial functions. In [2] it is proved that the family of surfaces that can be parameterised by the twistor lift of a slice regular function consists only of scrolls (i.e.: surfaces ruled by lines). Let us study, as a meaningful example, the cubic scroll $\mathcal{C} \subset \mathbb{C}\mathbb{P}^3$ defined by the set of points $[X_0, X_1, X_2, X_3] \in \mathbb{C}\mathbb{P}^3$ satisfying the equation

$$(5.1) \quad X_0 X_3^2 + X_1^2 X_2 = 0.$$

In the Cayley classification [11, 12, 13], the cubic \mathcal{C} is denoted by $S(1,1)$ (or by $S(1,1,3)$). Using the notation $m_{ij} := \{[X_0, X_1, X_2, X_3] \mid X_i = 0 = X_j\}$, we have that \mathcal{C} is a non-normal surface, singular along the line m_{13} and it has two cuspidal points at $p_0 = [1, 0, 0, 0]$ and $p_2 = [0, 0, 1, 0]$. It has two directrix lines, namely m_{13} and m_{02} .

In [2] the first author constructs a slice regular function whose twistor lift parameterises \mathcal{C} . The mentioned function is one of the slice-polynomial functions introduced in Example 2.12:

$$R(q) = -q^2 \ell_+ + q \ell_- = \ell_+(-v^2) + \ell_-(\bar{v}).$$

By the second expression of R we immediately see that its splitting of R with respect to \mathbb{C}_i is given by the functions $g(v) = -v^2$, $\hat{g}(v) = v$ and $h = \hat{h} \equiv 0$. Therefore its twistor degree is $\widetilde{\deg}(R) = 3$ and the extended twistor lift $\mathcal{R}: \mathcal{Q} \rightarrow \mathbb{C}\mathbb{P}^3$ of R is given by

$$\mathcal{R}[s, u, sv, uv] = [s, u, -sv^2, uv].$$

Considering the homogenisation with respect to the variable v , the map \mathcal{R} is given by:

$$\mathcal{R}([s, u], [t, v]) = \mathcal{R}[st, ut, sv, uv] = [st^2, ut^2, -sv^2, uv].$$

In the same spirit of [15, Theorem 7.1] we state the following.

Theorem 5.1. *The mapping \mathcal{R} is a birational equivalence between \mathcal{Q} and \mathcal{C} . In fact, \mathcal{R} maps \mathcal{Q} onto $(\mathcal{C} \setminus m_{01}) \cup \{p_2\}$ and is injective on the pre-image of the complement of $m_{13} \setminus \{p_0\}$ where it has 2 pre-images except for the point p_2 , where $\mathcal{R}^{-1}(p_2) \simeq \mathbb{C}\mathbb{P}^1$.*

Proof. First of all, if $X_0 \neq 0$, $X_1 \neq 0$, $X_2 \neq 0$ and $X_3 \neq 0$, we can define

$$\frac{s}{u} = \frac{X_0}{X_1}, \quad \text{and} \quad \frac{v}{t} = \frac{X_1 X_2}{X_0 X_3}$$

and $[X_0, X_1, X_2, X_3]$ has a unique inverse image. The other cases are the following:

- if $X_0 = 0$, then $X_1^2 X_2 = 0$, obtaining the lines m_{01} and m_{02} .
- If $X_1 = 0$, then $X_0 X_3^2 = 0$, obtaining the lines m_{01} and m_{13} .
- If $X_2 = 0$, then $X_0 X_3^2 = 0$, obtaining the lines m_{02} and m_{23} .
- If $X_3 = 0$, then $X_1^2 X_2 = 0$, obtaining the lines m_{13} and m_{23} .

The four lines $m_{01}, m_{02}, m_{13}, m_{23}$ intersect at $p_0, p_2, p_3 = [0, 0, 0, 1]$ and $p_1 = [0, 1, 0, 0]$ (see Figure 1). The line $m_{02} \setminus \{p_3, p_1\}$ can be obtained for $s = 0$ and, in this case the map \mathcal{R} can be inverted by setting

$$\frac{v}{t} = \frac{X_1}{X_3}.$$

The point p_1 is obtained for $s = 0 = v$ and so it has just a pre-image, while the point p_3 has no pre-images since $X_0 = X_1 = 0$ forces $t = 0$. The line $m_{13} \setminus \{p_0, p_2\}$ can be obtained for $u = 0$ and the parameterisation gives two pre-images by

$$\left(\frac{v}{t}\right)^2 = \frac{X_2}{X_0}.$$

The point p_0 is obtained for $u = 0 = v$ and so it has only a pre-image, while the point p_2 can be obtained for $t = 0$ and since there are no condition on $[s, u]$, then $\mathcal{R}^{-1}(p_2) \simeq \mathbb{C}\mathbb{P}^1$. The line $m_{23} \setminus \{p_0, p_1\}$ can be obtained for $v = 0$ and each point is given by

$$\frac{u}{s} = \frac{X_1}{X_0}.$$

Lastly, the line m_{01} would be obtained for $t = 0$, but this forces $X_3 = 0$. Therefore m_{01} is not in the image of \mathcal{R} except for the point p_2 . □

Given an algebraic surface \mathcal{K} in $\mathbb{C}\mathbb{P}^3$ of degree d , its intersection with a twistor fiber consists in exactly d points except at points belonging to the so called discriminant locus, where it can be either of lesser cardinality or it can be an entire fiber $\mathbb{C}\mathbb{P}^1$. More precisely, we recall the following definition.

Definition 5.1. Let \mathcal{K} be a surface of degree d in $\mathbb{C}\mathbb{P}^3$. We define the *discriminant locus* of \mathcal{K} as the following set

$$Disc(\mathcal{K}) := \{q \in \mathbb{S}^4 \simeq \mathbb{H}\mathbb{P}^1 \mid \#(\pi^{-1}(q) \cap \mathcal{K}) \neq d\}.$$

We denote by G_0 the subset of $Disc(\mathcal{K})$ such that for any $q \in G_0$, the twistor fibre $\pi^{-1}(q) \subset \mathcal{K}$ and, for any $n < d$ we define the following sets

$$G_n := \{q \in \mathbb{S}^4 \simeq \mathbb{H}\mathbb{P}^1 \mid \#(\pi^{-1}(q) \cap \mathcal{K}) = n\}.$$

Hence $Disc(\mathcal{K}) = G_0 \cup G_1 \cup \dots \cup G_{d-1}$. In the particular case in which $d = 3$, we have $Disc(\mathcal{K}) = G_0 \cup G_1 \cup G_2$ and the set G_1 is called the set of *triple points* and G_2 the set of *double points*.




FIGURE 1. In black the set $(m_{02} \cup m_{23}) \setminus \{p_3\}$ of points with one pre-image; in blue the set $m_{13} \setminus \{p_0, p_2\}$ of points with 2 pre-images; in red the set $m_{01} \setminus \{p_2\}$ of points with no pre-image and in green the point p_2 with a $\mathbb{C}\mathbb{P}^1$ of pre-images.

Now, in view of the interest related to the discriminant locus of an algebraic surface in the twistor space (see [2, 6, 7, 8, 15, 25]), we are going to describe it for the cubic scroll \mathcal{C} . To do that we will exploit the new techniques on slice regularity concerning slice-polynomial functions and their pre-images. Thanks to Theorem 5.1, we know that the parameterisation \mathcal{R} is 1 : 1 outside a Zariski-closed subset of \mathcal{C} , namely $m_{01} \cup m_{13}$. Therefore, outside this set, we can describe the discriminant locus of \mathcal{C} by means of Theorem 4.1. The set $m_{01} \cup m_{13}$ will be studied by its own in the following remark.

Remark 5.2. Recalling the definition of π , it is immediate to see that the line m_{01} projects to ∞ , which entails that m_{01} is a twistor fiber and that ∞ belongs to G_0 .

Consider now the intersection between m_{13} and the twistor fibers in (3.1). We have that $X_2 = X_0 q_1$ and $0 = X_0 q_2$ implying $q_2 = 0$. Hence m_{13} projects on $\hat{\mathbb{C}}_i$. Since $m_{13} \subset \pi^{-1}(\hat{\mathbb{C}}_i) \cap \mathcal{C}$ and \mathcal{R}^{-1} is not defined on m_{13} we study here separately this set of fibers.

Consider the intersection between the twistor fibre (3.1) above $[1, q_1 + q_2 j] \in \mathbb{H}\mathbb{P}$ and the cubic \mathcal{C} ,

$$(5.2) \quad X_0^3 q_2^2 + 2X_0^2 X_1 \bar{q}_1 q_2 + X_0 X_1^2 (\bar{q}_1^2 + q_1) - X_1^3 \bar{q}_2 = 0.$$

For any $q \in \mathbb{H}$ such that $q_2 = 0$, Equation (5.2) becomes $X_0 X_1^2 (\bar{q}_1^2 + q_1) = 0$. The set of points q_1 such that $\bar{q}_1^2 + q_1 = 0$ clearly consists of elements in G_0 , $q_1 \in \{0, -1, 1/2 + \sqrt{3}/2, 1/2 - \sqrt{3}/2\} \subset G_0$. Therefore, for any q_1 such that $(\bar{q}_1^2 + q_1) \neq 0$, there are two solutions, namely $X_0 = 0$ and $X_1 = 0$ with multiplicity two and so $(\hat{\mathbb{C}}_i \setminus G_0) \subset G_2$.

Having analysed the two lines m_{01} and m_{13} , we now pass to look at the discriminant locus of \mathcal{C} by means of the parameterisation \mathcal{R} .

First we recall some fundamental notion on cubic polynomials. Given any complex polynomial of degree 3, $p(x) = ax^3 + bx^2 + cx + d$, its discriminant is given by the following quantity

$$\Delta_p = 18abcd - 4b^3d + b^2c^2 - 4ac^3 - 27a^2d^2.$$

The polynomial $p(x)$ has a multiple root if and only if $\Delta_p = 0$. Moreover, if we denote by Δ'_p the quantity

$$\Delta'_p = b^2 - 3ac,$$

we have that p has a triple root if and only if $\Delta_p = \Delta'_p = 0$.

Let us begin our analysis of $Disc(\mathcal{C})$ by the set of triple points.

Proposition 5.3. *Let $\mathcal{C} \subset \mathbb{C}\mathbb{P}^3$ be the cubic of equation (5.1). Then the set of triple points G_1 of \mathcal{C} is empty.*

Proof. Having the explicit parameterisation \mathcal{R} , for any $q \in \mathbb{H}$, we compute, according to Formula (4.2), the associated determinant

$$D_q(v) = v^3 - v^2\bar{q}_1 + vq_1 - |q_1|^2 - |q_2|^2.$$

For this polynomial the two quantities Δ_{D_q} and Δ'_{D_q} are respectively

$$-\Delta_{D_q} = 4q_1^3 + 8|q_1|^4 + 4\bar{q}_1^3|q_1|^2 + 36|q_1|^2|q_2|^2 + 4\bar{q}_1^3|q_2|^2 + 27|q_2|^4, \text{ and } \Delta'_{D_q} = \bar{q}_1^2 - 3q_1.$$

Decomposing the two equations $\Delta_{D_q} = 0$ and $\Delta'_{D_q} = 0$ in their real and imaginary parts and setting $z := |q_2|$ and $q_1 = x + iy$, we obtain the following two systems:

$$(5.3) \quad \begin{cases} \Delta_{D_q} = 0 & \text{if and only if} \\ \begin{cases} 27z^4 + z^2(36(x^2 + y^2) + 4x(x^2 - 3y^2)) + 4x(x^2 - 3y^2)(x^2 + y^2 + 1) + 8(x^2 + y^2)^2 = 0 \\ 4y(3x^2 - y^2)(1 - (x^2 + y^2) - z^2) = 0. \end{cases} \end{cases}$$

and

$$\Delta'_{D_q} = 0 \quad \text{if and only if} \quad \begin{cases} x^2 - y^2 - 3x = 0 \\ 2xy + 3y = 0. \end{cases}$$

The last system represents the intersection of a hyperbola with the product of two lines (see Figure 2). The solutions of the equation $\Delta'_{D_p} = 0$ are then $q = q_2, 3 + q_2, -3/2 \pm 3\sqrt{3}/2i + q_2$ for any $q_2 \in \mathbb{C}$.




FIGURE 2. The set $\Delta'_{D_p} = 0$ is given by the intersection of the blue hyperbola $x^2 - y^2 - 3x = 0$ with the two yellow straight lines $2xy + 3y = 0$.

We study the set of triple points by substituting these four solutions in System (5.3).

- For $q_1 = 0$, the discriminant Δ_{D_q} vanishes only for $q_2 = 0$, but the point $q = 0$ belongs to G_0 .
- For $q_1 = 3$, the equation $\Delta_{D_q} = 0$ becomes

$$27z^4 + 432z^2 + 1728 = 0,$$

that has no real solution.

- For $q_1 = -3/2 \pm 3\sqrt{3}/2i$, it holds $|q_1| = 3$ and $(-3/2 \pm 3\sqrt{3}/2i)^3 = 27$, therefore, in both cases, the equation $\Delta_{D_q} = 0$ becomes, as before,

$$27z^4 + 432z^2 + 1728 = 0,$$

which, again, has no real solution.

Therefore, we have proved that $G_1 = \emptyset$. □

We now pass to the study of twistor fibers.

Proposition 5.4. *Let $\mathcal{C} \subset \mathbb{C}\mathbb{P}^3$ be the cubic of equation (5.1). Then the set of twistor fibres G_0 of \mathcal{C} is equal to*

$$G_0 = \left\{ 0, -1, \frac{1}{2} + i\frac{\sqrt{3}}{2}, \frac{1}{2} - i\frac{\sqrt{3}}{2}, \infty \right\} \subset \hat{\mathbb{C}}_i.$$

Proof. In Remark 5.2 we have already seen that $\{0, -1, \frac{1}{2} + i\frac{\sqrt{3}}{2}, \frac{1}{2} - i\frac{\sqrt{3}}{2}, \infty\} \subset G_0$. We now show that there are no other elements in G_0 using the parameterisation given by \mathcal{R} . For any $q \in \mathbb{H}$, the matrix A_q associated with \mathcal{R} , is given by

$$A_q(v) = \begin{pmatrix} -v^2 - q_1 & -\bar{q}_2 \\ -q_2 & \bar{q}_1 - v \end{pmatrix},$$

We have seen in Theorem 4.1, that, on the affine set $[1, q] \subset \mathbb{H}\mathbb{P}^1$, the vanishing of the matrix A_q at some point v_0 implies the existence of a $\mathbb{C}\mathbb{P}^1$ of solution of the System (4.1) and so of a twistor line contained in \mathcal{C} . After simple computations it is clear that there exist $v_0 \in \mathbb{C}$ such that $A_q(v_0) \equiv 0$ if and only if $q \in \{0, -1, \frac{1}{2} + i\frac{\sqrt{3}}{2}, \frac{1}{2} - i\frac{\sqrt{3}}{2}\}$. Then the only other element that possibly lie in G_0 is ∞ which was already found in Remark 5.2. □

We conclude the study of $Disc(\mathcal{C})$ by describing the set of double points.

Proposition 5.5. *Let $\mathcal{C} \subset \mathbb{C}\mathbb{P}^3$ be the cubic of equation (5.1). Then the set of double points G_2 of \mathcal{C} is homeomorphic to a 2-sphere with six handles pinched at G_0 without the attaching points, i.e.*

$$G_2 = \left(\hat{\mathbb{C}}_i \bigcup_{\substack{P=0,\infty \\ z_m^3 = -1, z_m \in \mathbb{C}_i}} \Sigma_{P, z_m} \right) \setminus G_0,$$

where Σ_{P, z_m} denotes a handle pinched to $\hat{\mathbb{C}}_i$ at P and z_m .

Proof. First of all, thanks to Remark 5.2, the set $(\mathbb{C}_i \setminus G_0) \subset G_2$. Then, since the set of triple points is empty, it is sufficient to study the zero set of the discriminant of D_q , i.e. the set of solutions of System (5.3). Since the second equation of the system factorises into three simpler parts, let us study them separately.

First case: $y = 0$. We get the following equation

$$(5.4) \quad p_1(x, z) = 27z^4 + z^2[4x^2(4+x)] + 4x^3(x+1)^2 = 0.$$

This equation defines a curve which intersects the x -axis only at $(x, z) = (0, 0), (-1, 0)$ (and at ∞). By Equation (5.4), it is not difficult to see that the curve is contained in the half-plane $x \leq 0$. Now for $x \leq 0$ we can solve Equation (5.4) for z^2 , obtaining

$$z^2 = \frac{2}{27}(-x^2(4+x) \pm \sqrt{x^4(4+x)^2 - 27x^3(x+1)^2}).$$

Since for $x \leq 0$, $-x^2(4+x) \leq \sqrt{x^4(4+x)^2 - 27x^3(x+1)^2}$ and equality holds for $x = 0, -1$, then, the only admissible solution is

$$z^2 = \frac{2}{27}(-x^2(4+x) + \sqrt{x^4(4+x)^2 - 27x^3(x+1)^2}) = f(x),$$

For $x \leq 0$, the function $f(x)$ appearing on the right hand side of the previous equation is always non-negative, differentiable for any $x \neq 0, -1$ where it is equal to 0 and it diverges at $+\infty$ for x that goes to $-\infty$. The curve $p_1(x, z) = 0$ is then given by the union of the graphs of $\pm\sqrt{f(x)}$, for $x \leq 0$ (see Figure 3). Recall now that the variable z equals $|q_2|$. Hence, the curve $p_1(x, z) = 0$ represents in fact the surface




FIGURE 3. $\{p_1(x, z) = 0\}$.

$p_1(x, |q_2|) = 0$ in \mathbb{S}^4 , intersecting $\hat{\mathbb{C}}_i$ only at $0, -1$ and at ∞ . Removing these three points its connected components (two handles) are both homeomorphic to a cylinder, thus the surface is homeomorphic to a sphere with an equator collapsed to a point.

Second case: $y^2 = 3x^2$. The set of points satisfying $y^2 - 3x^2 = 0$ is the union of the two lines $y = \pm\sqrt{3}x$ in the xy -plane. These are the images of the line $y = 0$ under the rotations around the z -axis of $2\pi/3$ and of $4\pi/3$. Notice that the first equation in System (5.3) is invariant under these two rotations. Therefore the set of solutions of $\Delta_p = 0$ for $y^2 - 3x^2 = 0$ consists exactly of two copies of the curve $p_1(x, z) = 0$ obtained by rotations of angles $2\pi/3$ and $4\pi/3$ around the z -axis. In particular these curves intersect the xy -plane at the origin and at $(1/2, -\sqrt{3}/2)$ and $(1/2, \sqrt{3}/2)$, respectively. As in the first case, since $z = |q_2|$, the two curves represent the other four handles, obtained rotating $p_1(x, |q_2|) = 0$ of angles $2\pi/3$ and $4\pi/3$ around the z -axis.

Third case: $y^2 = 1 - x^2 - z^2$. In this case, System (5.3) reduces to

$$z^4 - 4(5 + 6x)z^2 - 8(4x^3 - 3x + 1) = 0.$$

Solving for z^2 we get

$$z^2 = 2(5 + 6x \pm \sqrt{(3 + 2x)^3}).$$

For any $|x| < 1$, $5 + 6x - \sqrt{(3 + 2x)^3} \leq 0$ and equals zero only for $x = 1/2$, giving again the points $(1/2, \pm\sqrt{3}/2)$ in the xy -plane. On the other hand $0 \leq 5 + 6x + \sqrt{(3 + 2x)^3} \leq 1 - x^2$ if and only if $x = -1$, corresponding to the already found point $(-1, 0)$ in the xy -plane. Thus this case does not give any new contribution to the set of double points. \square

Collecting the results contained in Remark 5.2 and in Propositions 5.3, 5.4 and 5.5, we can state the main theorem of this section.

Theorem 5.2. *The discriminant locus of the cubic scroll $\mathcal{C} \subset \mathbb{C}\mathbb{P}^3$ of equation (5.1) is given by the singular surface*

$$\text{Disc}(\mathcal{C}) = \Sigma = \hat{\mathbb{C}}_i \cup_{\substack{P=0,\infty \\ z_m^3=-1, z_m \in \mathbb{C}_i}} \Sigma_{P,z_m},$$

where Σ_{P,z_m} denotes an handle pinched to $\hat{\mathbb{C}}_i$ at P and z_m . In particular we have that

$$G_0 = \left\{ 0, -1, \frac{1}{2} + i\frac{\sqrt{3}}{2}, \frac{1}{2} - i\frac{\sqrt{3}}{2}, \infty \right\} \subset \mathbb{C}_i, \quad G_1 = \emptyset, \quad G_2 = \Sigma \setminus G_0.$$




FIGURE 4. The set $\Sigma \subset \mathbb{S}^4$. The big black round sphere represents $\hat{\mathbb{C}}_i$. The two red handles on the left represent $\Sigma_{0,-1}$ and $\Sigma_{\infty,-1}$. The two green handles on the back represent $\Sigma_{0,1/2-\sqrt{3}/2}$ and $\Sigma_{\infty,1/2-\sqrt{3}/2}$. The two blue handles on the right represent $\Sigma_{0,1/2+\sqrt{3}/2}$ and $\Sigma_{\infty,1/2+\sqrt{3}/2}$.

5.1. A first description of the OCS's induced by \mathcal{C} . We will now give some qualitative results on the nature of the 3 OCS's induced by the cubic scroll \mathcal{C} .

A slightly different approach in twistor theory relies on the definition of twistor fibers by means of a map j which swaps any OCS J into $-J$. In our case j is the transformation defined on $\mathbb{C}\mathbb{P}^3$ induced, via π , by the quaternionic multiplication by j on $\mathbb{H}\mathbb{P}^1$. To be more precise $j : \mathbb{C}\mathbb{P}^3 \rightarrow \mathbb{C}\mathbb{P}^3$ is the map defined as

$$j : [X_0, X_1, X_2, X_3] \mapsto [-\bar{X}_1, \bar{X}_0, -\bar{X}_3, \bar{X}_2].$$

This map j is an antiholomorphic involution and it has no fixed points. Starting with such a map j , one can recover the twistor fibres in the following way: consider a point $X \in \mathbb{C}\mathbb{P}^3$ and the unique projective line l connecting X and $j(X)$. All the lines constructed in this way form the set of twistor fibres. In particular, a line l in $\mathbb{C}\mathbb{P}^3$ is a fibre for π if and only if $l = j(l)$.

Thanks to the explicit description of the map j , we are able to give a first qualitative result on the three OCS's induced by the cubic \mathcal{C} .

Proposition 5.6. *Let $\mathcal{C} \subset \mathbb{C}\mathbb{P}^3$ be the cubic scroll of equation (5.1). Then the intersection of \mathcal{C} with $j(\mathcal{C})$ consists of the following set*

$$j(\mathcal{C}) \cap \mathcal{C} = m_{02} \cup m_{13} \cup \pi^{-1}(G_0).$$

where, in particular $j(m_{02}) = m_{13}$.

Proof. By the definition of the map j , we have that $[X_0, X_1, X_2, X_3] \in j(\mathcal{C}) \cap \mathcal{C}$ if and only if

$$\begin{cases} X_0 X_3^2 + X_1^2 X_2 = 0 \\ X_0^2 X_3 + X_1 X_2^2 = 0. \end{cases}$$

It is immediate to verify that the lines m_{01}, m_{02}, m_{13} and m_{23} belong to the intersection $j(\mathcal{C}) \cap \mathcal{C}$. The fact that $j(m_{02}) = m_{13}$, $\pi(j(m_{02})) = \pi(m_{13}) = \hat{\mathbb{C}}_i$, $\pi(m_{01}) = \infty \in \mathbb{S}^4$ and $\pi(m_{23}) = 0$ is trivial. Suppose now that $X_n \neq 0$ for all $n = 0, 1, 2, 3$, then from the first equation we have that

$$(5.5) \quad X_0 = -\frac{X_1^2 X_2}{X_3^2}.$$

Hence, from the second equation we obtain

$$X_1 X_2^2 [X_3^3 + X_1^3] = 0 \quad \Leftrightarrow \quad X_3^3 = -X_1^3.$$

Therefore we obtain the three planes defined by the following equations

$$X_3 = -X_1, \quad X_3 = \left(\frac{1}{2} - \frac{\sqrt{3}}{2}i\right) X_1, \quad X_3 = \left(\frac{1}{2} + \frac{\sqrt{3}}{2}i\right) X_1.$$

Substituting the last three equations in Formula (5.5), we get the three lines projecting on $-1, \frac{1}{2} \pm \frac{\sqrt{3}}{2}i \in \mathbb{S}^4$ via π :

$$m_{-1} : \begin{cases} X_3 = -X_1 \\ X_2 = -X_0 \end{cases}, \quad m_{\frac{1}{2} - \frac{\sqrt{3}}{2}i} : \begin{cases} X_3 = \left(\frac{1}{2} - \frac{\sqrt{3}}{2}i\right) X_1 \\ X_2 = -\left(\frac{1}{2} - \frac{\sqrt{3}}{2}i\right)^2 X_0 \end{cases}, \quad m_{\frac{1}{2} + \frac{\sqrt{3}}{2}i} : \begin{cases} X_3 = \left(\frac{1}{2} + \frac{\sqrt{3}}{2}i\right) X_1 \\ X_2 = -\left(\frac{1}{2} + \frac{\sqrt{3}}{2}i\right)^2 X_0. \end{cases}$$

□

A picture of the set $j(\mathcal{C}) \cap \mathcal{C}$ is given in Figure 5, where it is also highlighted its twistor projection.

In the language of independent OCS's, the last proposition has as main implication the following result.

Corollary 5.7. *The manifold $\mathbb{S}^4 \setminus \Sigma$ admits 3 non-constant strongly independent OCS's.*


Proof. Thanks to Theorem 5.2, the cubic \mathcal{C} induces three OCS's on \mathbb{S}^4 outside Σ . Moreover, thanks to Proposition 5.6, outside of $m_{02} \cup m_{13} = \pi^{-1}(\hat{\mathbb{C}}_i)$, the intersection between \mathcal{C} and $j(\mathcal{C})$ is empty, meaning that if J is an OCS induced by \mathcal{C} , then $-J$ (that would be obtained by $j(\mathcal{C})$), is none of the other 2 possible OCS's induced by the cubic. □

Thanks to [2, Theorem 22], if $q = q_1 + q_2 j \in \mathbb{S}^4 \setminus \Sigma$ and if

$$\{x_n = \alpha_n + I_n \beta_n, n = 1, 2, 3\} = \pi(\mathcal{R}^{-1}(\pi^{-1}(q))),$$

then the three OCS's are defined to be

$$\mathbb{J}_{n,q}(v) = I_n v,$$

FIGURE 5. The set $j(\mathcal{C}) \cap \mathcal{C}$ and its twistor projection.

where $v \in T_q(\mathbb{S} \setminus \Sigma) \simeq \mathbb{H}$ and $I_n v$ denote the quaternionic multiplication. It is in general difficult to write an explicit expression of the OCS's \mathbb{J}_n , but at a fixed point q the matrix representing $\mathbb{J}_{n,q}$ can be computed as in [2, Remark 1], by knowing the $[s, u]$ -coordinate of the element $\mathcal{R}^{-1}(\pi^{-1}(q))$.

Remark 5.8. As a consequence of Proposition 5.6, on the discriminant locus $\Sigma = \text{Disc}(\mathcal{C})$ the three OCS's of Corollary 5.7 collapse into two that are independent outside $\hat{\mathbb{C}}_i$ where they coincide with the two constant OCS's $\pm \mathbb{J}_i$ defined as multiplication by $\pm i$.

The non-singular cubic

$$\mathcal{C}' : X_0 X_3^2 + X_3 X_0^2 + X_1 X_2^2 + X_2 X_1^2 = 0$$

studied in [7, 8] is engineered so to have the “twistor geometry [...] as simple as possible” among all the non-singular cubics (see [8]). This is done by maximising the number of twistor lines on a generic non-singular cubic surface $\mathcal{K} \subset \mathbb{C}\mathbb{P}^3$ and the size of the group of symmetries preserving \mathcal{K} and the twistor fibration. Thanks to [7, Theorem 1], a non-singular cubic surface contains at most five twistor lines. Moreover, in the same paper the authors prove the following facts:

- if a smooth cubic contains 5 twistor lines, then the image of these lines under π must all lie on a round 2-sphere in \mathbb{S}^4 ;
- given a line $m \in \mathbb{C}\mathbb{P}^3$ and five points $\{p_1, \dots, p_5\} \subset m$, then, after having identified m with the Riemann sphere, there exists a non-singular cubic surface containing m , $j(m)$ and the five twistor lines joining p_i and $j(p_i)$ if and only if no four of the points p_i lie on a circle under this identification.

Now, in [7, Section 5] the authors show that the the most symmetrical arrangement of 5 points on a 2-sphere such that no four of which lie on a circle, is given by the set of vertices of a triangular bipyramid: two poles and three cubic roots of -1 lying on the equator. The symmetry group of these five points is $\mathbb{Z}_3 \times \mathbb{Z}_2$ and up to conformal transformation of the 2-sphere, it is possible to assume that these points are the elements of G_0 listed in our Proposition 5.2. Then if the 2-sphere $\hat{\mathbb{C}}$ containing G_0 , lifts, by means

of π , to the two lines m and $j(m)$ and we ask for the surface \mathcal{K} to contain both m and $j(m)$, then its symmetry group is given by $\mathbb{Z}_3 \times \mathbb{Z}_2 \times \mathbb{Z}_2$, where the action of the last \mathbb{Z}_2 swaps m and $j(m)$. Up to conformal transformations, the unique non-singular cubic surface having symmetry group $\mathbb{Z}_3 \times \mathbb{Z}_2 \times \mathbb{Z}_2$ is exactly \mathcal{C}' . The discriminant locus of this non-singular surface topologically is $(T_1 \cup T_2)/G_0$, where T_1 and T_2 are two 2-dimensional tori passing through G_0 .

It is interesting to notice that our example carries on part of this geometry. In fact, Theorem 5.2 states that the symmetry group of the cubic scroll \mathcal{C} contains $\mathbb{Z}_3 \times \mathbb{Z}_2$ acting on the twistor lines but not the second \mathbb{Z}_2 part. In fact, while $j(\mathcal{C}') = \mathcal{C}'$, in our case the intersection $j(\mathcal{C}) \cap \mathcal{C}$ is non-trivial.

The construction that we presented and that develops the fascinating interplay between slice regularity and twistor geometry, still has some aspects that need to be better understood. Given a slice-polynomial function P (or more generally a slice regular function), the relation between its singular set (i.e.: the set of points where its differential is not injective) and the discriminant locus of the surface containing the image of its extended twistor lift is not completely clear. In fact, given a slice regular function f , the sets of its wings and degenerate spheres are contained in the singular locus of f but while degenerate spheres correspond to twistor fibres contained in the image of \mathcal{P} , on the other hand the same is not true, in general, for wings. Consider for instance the slice-polynomial function $P(q) = (q + j) * \ell_+ = \ell_+ v + \ell_- j$. As shown in Example 4.6 we have that $P|_{\mathbb{C}_i^-} = j$. Its extended twistor lift is equal to $\mathcal{P}([s, u], v) = [s, u, sv - u, 0]$ and its image lies in the hyperplane $\mathcal{H} := \{X_3 = 0\}$. The hyperplane \mathcal{H} was extensively studied in [2]. In particular $\#(\pi^{-1}(j) \cap \mathcal{H}) = 1$ while $\pi^{-1}(0) \cap \mathcal{H} \simeq \mathbb{CP}^1$. In fact the pre-image of j by $\pi \circ \mathcal{P}$ consists of $([0, 1], v) \subset \mathbb{CP}^1 \times \mathbb{C}_i$, while the pre-image of 0 consists of the surface $([1, v], v) \subset \mathbb{CP}^1 \times \mathbb{C}_i$ for any $v \in \mathbb{C}_i$. From this point of view, j is not a twistor fiber because the projection of its pre-image on the first component is just the point $[0, 1]$, while the projection of the pre-image of 0, $[1, v]$, covers \mathbb{CP}^1 minus a point.

REFERENCES

1. A. Altavilla. Some properties for quaternionic slice regular functions on domains without real points. *Complex Variables and Elliptic Equations*, 60 (1), 59–77, 2014.
2. A. Altavilla. Twistor interpretation of slice regular functions. *Journal of Geometry and Physics*, 123, 184–208, 2018.
3. A. Altavilla. On the real differential of a slice regular function. *Advances in Geometry*, 18 (1), 5–26, 2018.
4. A. Altavilla and E. Ballico. Twistor lines on algebraic surfaces. To appear in *Annals of Global Analysis and Geometry*, DOI: 10.1007/s10455-018-9640-2. See also *arXiv:1802.06697*, 2018.
5. V. Apostolov, P. Gauduchon, and G. Grantcharov. Bihermitian structures on complex surfaces. *Proceedings of the London Mathematical Society*, 79 (2), 414–428, 1999.
6. J. Armstrong. The twistor discriminant locus of the fermat cubic. *New York Journal of Mathematics*, 21, 485–510, 2015.
7. J. Armstrong, M. Povero, and S. Salamon. Twistor lines on cubic surfaces. *Rendiconti del Seminario Matematico Università e Politecnico di Torino*, 71 (3–4), 317–338, 2013.
8. J. Armstrong and S. Salamon. Twistor topology of the fermat cubic. *Symmetry, Integrability and Geometry: Methods and Applications*, Jun 2014.
9. R. L. Bryant. Conformal and minimal immersions of compact surfaces into the 4-sphere. *Journal of Differential Geometry*, 17 (3), 455–473, 1982.
10. F. E. Burstall and J. H. Rawnsley. Twistor theory for riemannian symmetric spaces. *Lecture Notes in Mathematics*, 1990.
11. A. Cayley. On skew surfaces, otherwise scrolls. *Philosophical Transactions of the Royal Society of London*, 153, 453–483, 1863.
12. A. Cayley. A second memoir on skew surfaces, otherwise scrolls. *Philosophical Transactions of the Royal Society of London*, 154, 559–577, 1864.
13. A. Cayley. A memoir on cubic surfaces. *Philosophical Transactions of the Royal Society of London*, 159, 231–326, 1869.
14. A. Fujiki and M. Pontecorvo. Twistors and bi-hermitian surfaces of non-kähler type. *Symmetry, Integrability and Geometry: Methods and Applications*, Apr 2014.

15. G. Gentili, S. Salamon, and C. Stoppato. Twistor transforms of quaternionic functions and orthogonal complex structures. *Journal of the European Mathematical Society*, 16 (11), 2323–2353, 2014.
16. G. Gentili, C. Stoppato, and D. C. Struppa. Regular functions of a quaternionic variable. *Springer Monographs in Mathematics*, 2013.
17. G. Gentili and D.C. Struppa. A new theory of regular functions of a quaternionic variable. *Advances in Mathematics*, 216(1),279–301, 2007.
18. G. Gentili, D.C. Struppa and F. Vlacci. The Fundamental Theorem of Algebra for Hamilton and Cayley numbers. *Mathematische Zeitschrift*, 259, 895–902, 2008.
19. R. Ghiloni and A. Perotti. Slice regular functions on real alternative algebras. *Advances in Mathematics*, 226 (2), 1662–1691, 2011.
20. R. Ghiloni, A. Perotti, and C. Stoppato. Division algebras of slice functions. *arXiv:1711.06604*, 2017.
21. T. Y. Lam. A first course in noncommutative rings. *Graduate Texts in Mathematics*, 2001.
22. C. LeBrun. Orthogonal complex structures on S^6 . *Proceedings of the American Mathematical Society*, 101 (1), 136, 1987.
23. M. Pontecorvo. Complex structures on riemannian four-manifolds. *Mathematische Annalen*, 309 (1), 159–177, 1997.
24. S. Salamon. Orthogonal complex structures. *Note based on a lecture at the conference “Differential Geometry and its Applications” in Brno*, <http://calvino.polito.it/salamon/T/brno.pdf>, 1995.
25. S. Salamon and J. Vialovsky. Orthogonal complex structures on domains in \mathbb{R}^4 . *Mathematische Annalen*, 343 (4), 853–899, 2008.

ALTAVILLA AMEDEO: DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI ROMA "TOR VERGATA", VIA DELLA RICERCA SCIENTIFICA 1, 00133, ROMA, ITALY
E-mail address: `altavilla@mat.uniroma2.it`

GIULIA SARFATTI: DIPARTIMENTO DI MATEMATICA E INFORMATICA "U. DINI", UNIVERSITÀ DI FIRENZE, VIALE MORGAGNI 67/A, 50134 FIRENZE, ITALY
E-mail address: `giulia.sarfatti@unifi.it`