



UNIVERSITÀ POLITECNICA DELLE MARCHE
Repository ISTITUZIONALE

Trends in the quality and hygiene parameters of bulk Italian Mediterranean buffalo (*Bubalus bubalis*) milk: a three year study

This is the peer reviewed version of the following article:

Original

Trends in the quality and hygiene parameters of bulk Italian Mediterranean buffalo (*Bubalus bubalis*) milk: a three year study / Pasquini, M.; Osimani, A.; Tavoletti, S.; Moreno, I.; Clementi, F.; Trombetta, M. F.. - In: ANIMAL SCIENCE JOURNAL. - ISSN 1740-0929. - ELETTRONICO. - 89:1(2018), pp. 176-185. [10.1111/asj.12916]

Availability:

This version is available at: 11566/252482 since: 2022-05-25T17:33:06Z

Publisher:

Published

DOI:10.1111/asj.12916

Terms of use:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. The use of copyrighted works requires the consent of the rights' holder (author or publisher). Works made available under a Creative Commons license or a Publisher's custom-made license can be used according to the terms and conditions contained therein. See editor's website for further information and terms and conditions.

This item was downloaded from IRIS Università Politecnica delle Marche (<https://iris.univpm.it>). When citing, please refer to the published version.

note finali coverpage

(Article begins on next page)



Trends in the quality and hygiene parameters of bulk Italian Mediterranean buffalo (*Bubalus bubalis*) milk: a three year study

Journal:	<i>Animal Science Journal</i>
Manuscript ID	ASJ-2016-0668.R1
Wiley - Manuscript type:	Original Article
Date Submitted by the Author:	n/a
Complete List of Authors:	<p>Pasquini, Marina; Universita Politecnica delle Marche, Dipartimento di Scienze Agrarie, Alimentari ed Ambientali OSIMANI, ANDREA; Universita Politecnica delle Marche, Dipartimento di Scienze Agrarie, Alimentari ed Ambientali Tavoletti, Stefano; Universita Politecnica delle Marche, Dipartimento di Scienze Agrarie, Alimentari ed Ambientali Moreno, Itzhak ; Universita Politecnica delle Marche, Dipartimento di Scienze Agrarie, Alimentari ed Ambientali Clementi, Francesca; Universita Politecnica delle Marche, Dipartimento di Scienze Agrarie, Alimentari ed Ambientali Trombetta, Maria Federica; Universita Politecnica delle Marche, Dipartimento di Scienze Agrarie, Alimentari ed Ambientali</p>
Keywords:	buffalo milk quality, chemical characterization, somatic cell counts, total bacterial counts
Abstract:	<p>Buffalo milk represents an indispensable source of nourishment in many parts of the world and it is the second most consumed milk worldwide. Buffalo milk is actually used for the production of many dairy products such as pasteurized or concentrated milk, butter, yogurt, ice-cream, dehydrated milk products, and cheeses. Due to its high nutritional value and the presence of natural bioactive substances, buffalo milk can also provide health benefits to consumers. In Italy, buffalo milk is used only for cheese making, mainly mozzarella PDO (Protected Designation of Origin), which is a highly valued dairy product. This three-year study, carried out between 2011 and 2013, was aimed at evaluating the quality of bulk Italian Mediterranean buffalo milk by monitoring physico-chemical parameters, somatic cell and total bacterial counts. A total of 51 samples of bulk milk were collected from one herd throughout the monitored period. Analysis of variance, carried out to test month, season, and year main effects, highlighted remarkable seasonal effects for fat, protein, and lactose content, as well as for predicted Mozzarella cheese yield, and somatic cell counts. The calculation of simple correlations allowed the identification of positive correlations between estimated cheese yield and fat and protein content.</p>

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



SCHOLARONE™
Manuscripts

For Review Only

1 Trends in the quality and hygiene parameters of bulk Italian Mediterranean buffalo (*Bubalus*
2 *bubalis*) milk: a three year study

3
4 **Running head:** Quality of Mediterranean buffalo milk

5
6 Marina PASQUINI, Andrea OSIMANI, Stefano TAVOLETTI, Itzhak MORENO, Francesca CLEMENTI
7 and Maria Federica TROMBETTA

8
9 Dipartimento di Scienze Agrarie, Alimentari ed Ambientali, Università Politecnica delle Marche, via
10 Breccie Bianche, 60131, Ancona, Italy

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
Correspondence: Andrea Osimani Ph. D., Dipartimento di Scienze Agrarie, Alimentari ed Ambientali,
Università Politecnica delle Marche, via Breccie Bianche, 60131, Ancona, Italy. (E-mail:
a.osimani@univpm.it).

32 **ABSTRACT**

33 Buffalo milk represents an indispensable source of nourishment in many parts of the world and it is the
34 second most consumed milk worldwide. Buffalo milk is actually used for the production of many dairy
35 products such as pasteurized or concentrated milk, butter, yogurt, ice-cream, dehydrated milk products,
36 and cheeses. Due to its high nutritional value and the presence of natural bioactive substances, buffalo
37 milk can also provide health benefits to consumers. In Italy, buffalo milk is used only for cheese making,
38 mainly mozzarella PDO (Protected Designation of Origin), which is a highly valued dairy product. This
39 three-year study, carried out between 2011 and 2013, was aimed at evaluating the quality of bulk Italian
40 Mediterranean buffalo milk by monitoring physico-chemical parameters, somatic cell and total bacterial
41 counts. A total of 51 samples of bulk milk were collected from one herd throughout the monitored period.
42 Analysis of variance, carried out to test month, season, and year main effects, highlighted remarkable
43 seasonal effects for fat, protein, and lactose content, as well as for predicted Mozzarella cheese yield, and
44 somatic cell counts. The calculation of simple correlations allowed the identification of positive
45 correlations between estimated cheese yield and fat and protein content.

46
47 **Key words:** buffalo milk quality, chemical characterization, somatic cell counts, total bacterial counts.

48

49 INTRODUCTION

50 Buffalo (*Bubalus bubalis*) milk represents an indispensable source of nourishment in many parts of the
51 world. India and Pakistan produce more than 91% of buffalo milk, which is the second most consumed
52 milk worldwide. As recently reviewed by Cazacu *et al.* (2014), the buffalo milk market is still considered
53 an emerging sector. Actually, the dairy industry produces many products based on the use of buffalo
54 milk; among these are pasteurized or concentrated milk, butter, heat-desiccated dairy products, heat-acid
55 coagulated dairy products, yogurt, ice-cream, dehydrated milk products, and cheeses (Cazacu *et al.* 2014).
56 Due to the high nutritional value of buffalo milk, the demand for such products is increasing; however,
57 there is still a shortage of scientific literature on buffalo milk's physico-chemical and hygienic parameters
58 (Zotos & Bampidis 2014).

59 In Europe, the countries that produce the largest quantities of buffalo milk are Italy, Turkey, Bulgaria, and
60 Greece (FAOSTAT 2014). In accordance with a recent report from the European Food Safety Authority
61 (EFSA), Italy is the largest producer (88%) of buffalo milk in the EU (EFSA BIOHAZ Panel 2015); in
62 the period 1990-2012, the Italian production of buffalo milk increased from 43,000 to 192,455 t, allowing
63 in 2012 the production (FAOSTAT 2014) of 51,910 and 37,122 t of mozzarella and mozzarella PDO
64 (Protected Designation of Origin), respectively (INEA 2012; 2013). The rearing of buffalo in Italy
65 constitutes an important reality, not only regarding the number of animals bred but also for the popularity
66 of Italian bubaline dairy products. Buffaloes reared in Italy stand out in the world for their genetics, the
67 applied technologies, the monitoring of pathologies, and the hygiene and quality of the end products
68 (Borghese 2005). In Italy, buffalo milk is used only for cheese making, mainly mozzarella PDO, which is
69 a highly valued product especially in the USA, Germany, France, UK, and Japan (Borghese 2005).

70 Buffalo milk is usually characterized by a rich composition with high content of fat, which constitutes the
71 main fraction, moreover, it is a good source of vitamins A, D, C, and B6; minerals such as Ca and P; and
72 conjugated linoleic acid (Ahmad 2013; Khedkar *et al.* 2016; Simoes *et al.* 2014; Zotos & Bampidis 2014).
73 The presence of trace elements such as boron, cobalt, copper, iron, manganese, sulfur, and zinc has also
74 been ascertained (Ahmad 2013).

75 The consumption of buffalo milk can provide benefits to people who suffer from hypertension, dental
76 decay, dehydration, respiratory problems, obesity, osteoporosis, and some forms of cancer (Ahmad 2013).
77 Furthermore, a recent study carried out by Kapila *et al.* (2013) highlighted that buffalo milk proteins (β -
78 lactoglobulin and casein) are less allergenic than cow milk proteins, which can significantly increase
79 protein-specific IgE sensitization and lymphocyte proliferation index causing allergies in consumers.

1 80 Many parameters can be used to assess the quality of milk, among these are compositional, physical-
2
3 81 chemical, hygienic, and sanitary characteristics. The main traits that are usually considered as milk
4
5 82 quality parameters are defined as proteins, fats, lactose, and solids-not-fat content. All of these parameters,
6
7 83 which can affect the quality of the end products, can vary depending on several factors, including genetic,
8
9 84 age, parity, lactation stage, season, feeding, and geographical area of rearing (Gürler *et al.* 2013).
10
11 85 Moreover, microbiological contamination and the presence of somatic cells can worsen the quality of
12
13 86 milk.

14 87 The aim of this study was to evaluate the quality of bulk Italian Mediterranean buffalo milk through
15
16 88 monitoring physico-chemical parameters, somatic cell counts and total bacterial counts over a three-year
17
18 89 period (2011–2013) in one farm. The effects of month, season, and year on the monitored variables were
19
20 90 also evaluated through statistical analyses.

21 91

22 92 **MATERIALS AND METHODS**

23 93 **Description of the Farm**

24
25
26 94 The monitoring of bulk buffalo milk characteristics was performed on a farm located in the Marche
27
28 95 region (central Italy) over a three-year period (2011-2013). The farm, in the year 2012 produced 184.8 t
29
30 96 of buffalo milk over a total national production of 192,455.3 t; whereas in 2013 the production increased,
31
32 97 reaching 422.7 t of buffalo milk over a national production of 194,892.8 t (Istat, 2014; 2015). The animals
33
34 98 were housed in free stalls with a concrete paddock and a permanent straw bedding area; the paddocks
35
36 99 were equipped with a sprinkler system and fans. The herd was organized in two feeding groups: milking
37
38 100 and dry buffalo cows. During the monitored period, all milking buffalo cows were fed a total mixed ratio
39
40 101 consisting of: corn silage (10.0 kg/head/day), first cut hay (5.0 kg/head/day), alfa-alfa hay (3.5
41
42 102 kg/head/day), maize flour (3.0 kg/head/day), soybean meal (1.0 kg/head/day), cottonseed (1.0
43
44 103 kg/head/day), and mineral-vitamin pre-mix (0.25 kg/head/day).

45
46 104 The milking area was arranged as a double-9 herringbone-milking parlor where buffaloes were placed in
47
48 105 two rows back to back. The herd size of the milking buffalo cows, the average age of lactating animals,
49
50 106 and their average days in milking are reported in Table 1. Buffalo cows were milked twice a day
51
52 107 (morning and afternoon) and the milk daily stored in a refrigerated tank (+4°C). The milk was delivered
53
54 108 within the same day to the dairy plant for cheese making.

55 109

56 110 **Sampling**

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

111 The milk subjected to analyses was obtained by pooling afternoon and morning milkings, stored at +4°C.
112 Before sampling the raw bulk whole milk was mixed in the farm's tank to ensure homogeneity and then
113 five aliquots of 100 mL were manually collected and pooled into a sterilized baker. Approximately 40 mL
114 of bulk buffalo milk was collected from the pooled milk using sterile conical tubes (BD Falcon, Franklin
115 Lakes, NJ, USA). The samples were transferred to the laboratory under refrigerated conditions (+4°C)
116 and subjected to analyses during the same day.
117 A total of 51 samples of bulk milk were collected throughout the monitored period (Table 1).

118

119 **Physico-chemical, Microbiological Analyses and Somatic Cell Counts**

120 All of the analyses were carried out in the same accredited laboratory (ACCREDIA, accreditation No.
121 1239).

122 Milk constituents (fat %, protein %, anhydrous lactose %, expressed as w/w) of bulk buffalo milk samples
123 were quantified via Fourier Transform Infrared (FTIR) spectroscopy using a CombiFoss FT+ composed
124 by Milkoscan FT Plus – 300 and Fossomatic FC (Foss Electric, Hillerød, DK).

125 Somatic cell counts (SCC) were determined in accordance with the ISO 13366:2008 standard by flow
126 cytometry, using a high-capacity somatic cell counter CombiFoss FT+ (Foss Electric). Before analysis,
127 each sample was heated to 40-42°C and subjected to counting within 15 minutes. The results were
128 expressed as number of cells mL⁻¹.

129 Total bacterial counts (TBC) were determined using a fluorimeter (BactoScan FC, Foss Electric) in
130 accordance with Regulation (EC) No 1664/2006, which allows the use of alternative methods for counts
131 at 30 °C when these methods are validated according to the reference horizontal method for the
132 enumeration of microorganisms ISO 4833-1:2013 and the protocol set by the ISO 16140 standards. The
133 results were expressed as colony forming units (cfu) mL⁻¹.

134 The cheese yield formula for estimating the production of mozzarella cheese (kg) was calculated in
135 accordance with the formula proposed by Altiero et al. (1989) as follows: 3.50 (protein %) + 1.23 (fat %)
136 – 0.88. This formula corresponds to a multiple regression equation that attributes 93% of yield variability
137 to fat and protein content as expressly calculated by Altiero et al. (1989) for buffalo mozzarella cheese
138 production. Estimated cheese yields were expressed as kg of mozzarella produced per 100 kg of
139 processed milk (kg 100 kg⁻¹).

140

141 **Statistical Analyses**

1
2 142 Before statistical analysis, the values of SCC and TBC were first log-transformed. Descriptive statistics,
3 143 calculated on the whole data set (51 samples), were carried out on fat, protein, lactose, solids-not-fat,
4 144 cheese yield formula, SCC, and TBC, computing means \pm standard deviation. The values recorded for all
5 145 of the variables were checked for conformance to a normal distribution and then processed by analysis of
6 146 variance (ANOVA) carried out using JMP statistical software version 11.0.0 (SAS Institute Inc., Cary,
7 147 NC, USA). The following main effects were tested: month, season, and year.

8
9 148 The one way ANOVA model was $y_{ij} = \mu + a_i + \epsilon_{ij}$, where μ is the overall mean, a_i is the effect of the i th
10 149 level of factor ($i = \text{January}, \dots, \text{December}$; $i = \text{winter, spring, summer, autumn}$; $i = 2011, 2012, 2013$), and
11 150 ϵ_{ij} is the random error.

12 151 Simple correlation among variables was evaluated by the Pearson product-moment correlation coefficient
13 152 (r).

14 153 Cluster analysis of bulk milk parameters was carried out using the Wards' minimum variance method.

15 154

16 155 **RESULTS AND DISCUSSION**

17 156 The monitoring of bulk buffalo milk composition can reflect the combined effect of environment and
18 157 genetics on the final quality of milk. It is known that the composition of buffalo milk can be influenced
19 158 by breed, age, parity, stage of lactation, seasonality, feeding, udder disorders such as mastitis, and genetic
20 159 polymorphism of milk proteins (Abd El-Salam & El-Shibini 2011). The above-mentioned parameters can
21 160 affect the composition of milk and, in turn, the quality and process yield of dairy products (Gürler *et al.*
22 161 2013).

23 162 Descriptive statistics for quality parameters of bulk buffaloes' milk analyzed between 2011 and 2013 are
24 163 shown in Table 2. The results of ANOVA are shown in Table 3. Multiple comparisons among Least
25 164 Square Means (LSM) using the Tukey HSD test for quality parameters of bulk buffaloes' milk according
26 165 to month, season, and year are shown in Table 4.

27 166 Regarding fat content, the average fat value reported in the present study (7.13%) was slightly lower than
28 167 that reported by Zotos & Bampidis (2014) in Greek buffaloes' milk. The monitored fat values were also
29 168 lower than those reported by Rosati & Van Vleck (2002) for individual milk samples produced by
30 169 buffaloes reared mainly in the South of Italy (average value $8.59 \pm 0.85\%$); additionally, Di Francia *et al.*
31 170 (2007a) found higher fat content in milk produced by Italian buffaloes fed a ration containing extruded
32 171 peas or soybean cake as concentrate components (7.84 and 7.56%, respectively). Finally, the fat values
33 172 we observed in the present study were slightly higher than the values reported by Cunha Neto *et al.*

173 (2005) for Brazilian buffalo milk used for the production of yogurt and by Enb *et al.* (2009) for milk
174 produced by Egyptian buffaloes and used for cheese manufacturing. Moreover, we observed a progressive
175 lowering of measured values during the warmer months with respect to the coldest ones. The highest
176 mean value of fat ($8.54 \pm 0.27\%$) was observed during January, whereas the lowest mean contents were
177 detected during May, June, and July ($6.50 \pm 0.27\%$, $6.27 \pm 0.24\%$, and $6.54 \pm 0.24\%$, respectively) without
178 significant differences among these three months. The data trend reported in the present study for fat
179 content is also in agreement with data reported by other authors in Greek and Italian buffaloes that
180 describe peaks of fat content during December, January, and February (Ahmad *et al.* 2008; Bartocci *et al.*
181 2002; Zotos & Bampidis 2014).

182 The ANOVA according to year did not show any significant differences, while as expected, according to
183 season, winter and autumn showed the highest mean levels of fat ($7.81 \pm 0.20\%$ and $7.76 \pm 0.16\%$,
184 respectively). Han *et al.* (2012) found a similar fat content trend in buffalo milk samples collected in the
185 United States; in more detail, the level of fat was the highest in January (7.63%), then a drop to a
186 minimum was observed in July (6.57%), and a further increase was recorded in November (7.97%).

187 Regarding the seasonal effect, the highest fat values were observed during winter and autumn; these
188 results almost overlap the seasonal trend reported by Gürler *et al.* (2013) for Anatolian buffaloes milk.
189 Interestingly, this latter breed, represents a sub-group of Mediterranean buffaloes as well as Italian
190 Mediterranean ones, and this could explain similar trends in milk compositional parameters.

191 Moreover, Simoes *et al.* (2014) observed that the fat content of Brazilian buffalo milk was lower during
192 warm and humid seasons (rainy season) with a value of 5.53% (w/w) and higher during cold seasons (dry
193 season), reaching a value of 6.74% (w/w). This difference in fat content may be due to greater energy loss
194 by buffaloes for the maintenance of their homeostasis during the warm and humid season associated with
195 their modest physiological sweating due to their low number of sweat glands. It is worth noting that
196 lactation stage, season, and animal diet can also strongly influence lipid synthesis, which in turn can
197 affect the quality of milk and dairy products (Ménard *et al.* 2010; Yadav *et al.* 2015).

198 The physiology of lactation coordinates the production of milk and determines its final composition.
199 However, this mechanism still represents a challenge for the current research. The stage of lactation
200 influences the composition of the milk and the amount and quality of fat. Furthermore, the environmental
201 temperature can also exert an effect on the profile of the fatty acids that compose the fat molecules.
202 Buffalo milk presents the highest fat content with respect to other dairy animals, although the factors that
203 influence its concentration in milk are still under study (Yadav *et al.* 2015; Zotos & Bampidis 2014).

204 Given the high importance of fat in milk's composition, new methods to maximize its content are still
205 under study. To this aim, Shelke *et al.* (2012) showed that buffaloes fed with rumen-protected fat (Ca
206 salts of palm fatty acids) have benefited from greater caloric intake during lactation, leading to a higher
207 yield of cheese and to an increase of unsaturated fatty acids in the fat.

208 As for protein content, the ANOVA according to month and year did not show significant differences,
209 whereas a seasonal effect was discovered. In more detail, we measured the highest mean values
210 ($4.48\pm 0.05\%$ and $4.50\pm 0.05\%$, respectively) during autumn and winter, while we observed the lowest
211 mean value ($4.27\pm 0.05\%$) in samples collected during spring, findings similar to those reported for
212 Anatolian bubaline milk (Gürler *et al.* 2013). As reported by Gürler *et al.* (2013), milk protein, as well as
213 fat, may be inversely related with environment temperature; moreover different level of protein and fat
214 may be ascribed to the lactation period of the majority of milking buffalo cows.

215 The mean values of protein content detected in the present one-farm study were lower than those reported
216 by Bonfatti *et al.* (2013a) for Mediterranean buffaloes reared in the Campania region (Italy), which tested
217 at approximately 6%.

218 It is worth noting that Bonfatti *et al.* (2013b) have recently highlighted that the relative amount of the
219 α S1-casein fraction in buffalo milk can heavily affect its behavior during cheese making, although to
220 clarify the effects of buffalo whey and casein fractions on cheese yield further studies are still needed.

221 Regarding lactose, no significant differences were discovered among months. We observed a progressive
222 lowering of lactose mean value among the years. The mean value dropped from $4.95\pm 0.07\%$ in 2011 to
223 $4.69\pm 0.07\%$ in 2013. We suppose that the decrease of lactose content can be related to the variation of
224 somatic cell counts which reached the highest mean value, although not statistically significant, in 2013
225 when lactose was at minimum. It is worth noting that the increase in SCC lead to a rise of enzymatic
226 activities in mammary tissue, thus reducing lactose synthetic activity (Sharif *et al.* 2007).

227 As for season, we recorded the highest mean value ($4.91\pm 0.06\%$) during summer, whereas we detected
228 the lowest mean value of lactose content during winter ($4.59\pm 0.07\%$). Large differences in lactose mean
229 values are often reported for milks from different parts of the world, although some of these differences
230 could be due to different method of expression (as monohydrate or anhydrous lactose). The recorded
231 values of lactose showed very low variation similar to the monthly variation reported by Han *et al.* (2012)
232 for buffalo species (4.49-4.73%), in a commercial water buffalo dairy farm. Lactose content in buffalo
233 milk and dairy products can be a quality indicator used as a marker of mammary inflammation because its
234 low levels can be the expression of reduced synthetic activity of secretory cells in mammary gland tissue,

1
2 235 thus suggesting mastitis in progress (Sharif *et al.* 2007). Indeed, Tripaldi *et al.* (2010) found that lactose
3 236 content decreased in animals affected by mastitis, this being negatively correlated with somatic cells and
4 237 contaminating bacteria. Hence, though still controversial, a threshold value of 4.7% lactose as a marker of
5 238 mammary inflammation has been proposed by different researchers (Tripaldi *et al.* 2010).
6
7
8
9 239 Solids *not-fat* (SNF) of milk consists of protein, lactose and mineral matter content. The SNF residue,
10 240 after the complete evaporation of water, is a fundamental quality parameter of raw milk that provides
11 241 useful information about the overall suitability of milk for cheese-making (Bassbasi *et al.* 2014).
12
13 242 Regarding SNF content, the ANOVA according to season and year did not show any significant
14 243 differences. However, according to month, we measured the lowest and the highest mean SNF values
15 244 during February ($9.51 \pm 0.07\%$) and September ($10.12 \pm 0.11\%$), respectively. The average values of SNF
16 245 content measured in the present study were slightly lower than those reported by Hussain *et al.* (2012)
17 246 and Zicarelli *et al.* (2007) in the same matrix, which were 10.40% and 10.61-10.43%, respectively. A
18 247 lower mean value of SNF was reported by Gürler *et al.* (2013) for milk produced by Anatolian buffaloes
19 248 and by Simoes *et al.* (2014) for bubaline milk collected during the dry season and used for production of
20 249 Marajó cheeses typical of the homonymous Brazilian island.
21
22 250 Cheese yield represents a parameter with heavy economic repercussions. The theoretical value of cheese
23 251 yield is conditioned by many factors such as the fat and casein content of the milk used, the production
24 252 technology, and moisture and salt content (Melilli *et al.* 2002). Regarding Mozzarella yield prediction, as
25 253 estimated with the formula proposed by Altiero *et al.* (1989), no significant differences were discovered
26 254 according to year, whereas according to month, we recorded the highest and the lowest mean values
27 255 during January (25.84 ± 0.57 kg 100 kg⁻¹) and June (21.80 ± 0.51 kg 100 kg⁻¹), respectively. Cheese yield
28 256 being strictly correlated with both fat and protein content, the trend of the cheese yield formula was
29 257 almost overlapping with that observed for the two parameters. Moreover, we discovered a seasonal effect
30 258 with the lowest average values of theoretical cheese production recorded in spring and summer; this
31 259 finding can be due to the lowest levels of both proteins and fats in milk recorded during these two seasons.
32
33 260 Among the few scientific papers that address the estimation of Mozzarella cheese production according to
34 261 the formula proposed by Altiero *et al.* (1989), Masucci *et al.* (2008) reported higher values of estimated
35 262 Mozzarella production than that observed in the present study. The values observed by Masucci *et al.*
36 263 (2008) of 27.2 and 28.0 kg 100 kg⁻¹ were likely due to the superior characteristics of organic buffalo milk,
37 264 which resulted particularly high in fat and proteins. Our findings for theoretic cheese yield were also
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2 265 lower than those published by Zicarelli *et al.* (2007) who reported an estimation of Mozzarella production
3
4 266 of approximately 25 kg per 100 kg of processed buffalo milk.
5
6 267 While the international literature has mainly provided data on the chemical composition of buffalo milk,
7
8 268 few investigations concerning the trends of the main sanitary (Somatic Cell Count, SCC) and hygienic
9
10 269 (Total Bacterial Counts, TBC) buffalo milk parameters have been carried out to date.
11
12 270 Regarding SCC, we did not discover significant differences among years. We recorded mean SCC values
13
14 271 ranging between 152.84 ± 25.22 and $199.73 \pm 23.43 \times 1,000 \text{ cell mL}^{-1}$ among the three years; these counts,
15
16 272 were always lower than the average value ($314,000 \text{ cell mL}^{-1}$) reported by Tripaldi *et al.* (2010) for
17
18 273 individual buffalo milk samples collected in the Latium region (Italy). Significant differences were
19
20 274 discovered among seasons; in more detail, we recorded the highest mean SCC value in the winter period
21
22 275 ($269.44 \pm 34.54 \times 1,000 \text{ cell mL}^{-1}$), whereas we observed the lowest mean value in samples collected
23
24 276 during summer ($147.00 \pm 27.69 \times 1,000 \text{ cell mL}^{-1}$). Regarding SCC trends across months, we observed the
25
26 277 highest mean value during January ($299.00 \pm 47.34 \times 1,000 \text{ cell mL}^{-1}$) and the lowest one during May
27
28 278 ($87.25 \pm 47.34 \times 1,000 \text{ cell mL}^{-1}$).
29
30 279 Actually, Regulation (EC) No 853/2004 of the European Parliament and of the Council of 29 April 2004,
31
32 280 laying down specific hygiene rules for food of animal origin, set only a threshold SCC value for bovine
33
34 281 raw milk. The threshold SCC value for bovine raw milk is set at $400,000 \text{ cells mL}^{-1}$, while no limits are to
35
36 282 date provided for buffalo raw milk, even though this parameter represents the most reliable indicator of
37
38 283 inflammatory status of mammalian udders, which may also cause a reduction in protein (casein) content
39
40 284 of milk. Somatic cell counts could be used as a quality parameter because, in buffalo milk, they increase
41
42 285 with days in lactation and show the highest values in milk from buffaloes with mastitis (Moroni *et al.*
43
44 286 2006).
45
46 287 The bulk buffalo milk samples under study showed the highest levels of SCC in winter, whereas the
47
48 288 lowest values were recorded during summer. These findings are in agreement with SCC observed by
49
50 289 Simoes *et al.* (2014) for Brazilian milk, which was assessed at 290,000 and 240,000 cells mL^{-1} during the
51
52 290 rainy and dry seasons, respectively. The annual trend of microbiological buffalo milk contamination, as
53
54 291 monitored in the present study, revealed SCC levels always lower than the limit of $200,000 \text{ cell mL}^{-1}$ that
55
56 292 in cows is usually considered a cut-off point for mastitis diagnosis (Tripaldi *et al.* 2010). As reported by
57
58 293 Smith (2002), SCC measurement, which includes polymorphonuclear leukocytes (PMN), macrophages,
59
60 294 and lymphocytes, is commonly used to evaluate the health of mammary glands. Furthermore, Kelly *et al.*
295 (2000) highlighted that the increase of SCC during milk secretion usually reflects an increase in PMN

1
2 296 whose function is the ingestion and destruction of invading microorganisms as well as secretion of
3
4 297 inflammatory regulators. High SCC in milk can exert adverse effects during cheese making, with a direct
5
6 298 negative impact on curd firmness, cheese yield, and sensory characteristics; moreover, the presence of
7
8 299 somatic cells can enhance the quantity of fat loss in whey (More *et al.* 2013). Furthermore, as reported by
9
10 300 Pasquini *et al.* (2003), the lower the number of somatic cells in buffalo milk is, the higher is the quantity
11
12 301 of casein and whey proteins in cheese. Therefore, further investigations on the somatic cell populations in
13
14 302 milk should be carried out to better explain the relationships among them and buffalo milk's aptness for
15
16 303 cheese manufacturing.

17 304 Finally, as for TBC, multiple comparisons among least square means (LSM) showed no significant
18
19 305 differences for the monitored bulk bubaline milk according to month, season, or year.

20 306 TBC are commonly used as indicators to evaluate the hygiene of the entire production process. The limit
21
22 307 values for TBC at 30°C established by Regulation (EC) No 853/2004 are: 100,000 cfu mL⁻¹ for raw cows'
23
24 308 milk; 1,500,000 cfu mL⁻¹ for raw milk from other species; in addition, 500,000 cfu mL⁻¹ limit for raw milk
25
26 309 from species other than cows used for manufacturing cheese products without heat treatments is also
27
28 310 provided.

29 311 TCB counts for the three-year period showed very low levels of raw milk environmental contaminating
30
31 312 microorganisms ranging from 6.46 to 23.41 x 10¹ cfu mL⁻¹. The TBC levels also showed low variability
32
33 313 through months, ranging from 3.66 x 10¹ in June to 9.85 x 10¹ cfu mL⁻¹ in March. Interestingly, the low
34
35 314 microbiological contamination found during the summer months (17.66 x 10¹ cfu mL⁻¹) could reflect
36
37 315 good hygiene practices performed during milking and handling of the raw milk, especially in summer
38
39 316 when warm temperatures could easily increase milk's microorganism content. Although the TBC values
40
41 317 reported in this study are lower than those reported by other authors (Gürler *et al.* 2013; Simoes *et al.*
42
43 318 2014; Tripaldi *et al.* 2010), it is worth noting that sufficiently high TBC could represent a risk for the
44
45 319 consumer because, pathogens can constitute a fraction of this hygiene indicator (Brown *et al.* 2000).
46
47 320 Hence the monitoring of TBC can provide a record of hygiene performance over time.

48 321 Important correlations among variables were identified (Table 5), and only those showing high significant
49
50 322 *P* values (< 0.01) are discussed. Significant positive correlations were found between protein and fat
51
52 323 percentage content ($r=0.64$; $P<0.01$) and between predicted cheese yield and both fat ($r=0.94$; $P<0.01$)
53
54 324 and protein ($r=0.87$; $P<0.01$) contents. Our results showed slightly higher correlation than the results
55
56 325 reported by Napolano *et al.* (2007). Moreover, fat and protein content are the most important parameters
57
58 326 determining cheese yield, which is the most important technological trait in the dairy industry (Bonfatti *et*

1
2 327 *al.* 2013b). Proteins, and in particular the content of casein, represent determining factors in the
3 328 organization of the cheese matrix because they are capable of retaining fat and moisture. Furthermore,
4 329 milk fat determines the amount of whey that drains from the curd. As reported by Mateo *et al.* (2009)
5 330 higher fat milks, with constant protein, resulted in lower yield of whey associated with higher reduction in
6 331 curd moisture content. As expected, SNF (%) was positively correlated with lactose content (Table 5).
7
8 332 The cluster analysis (Figure 1) reflected the different trend of milk compositional parameters among
9 333 seasons, and confirmed the results of ANOVA showing the lack of a significant year effect on
10 334 compositional milk parameters related to cheese yield production.
11
12
13
14
15
16
17

335

336 CONCLUSIONS

337

338 The trends in the compositional and sanitary parameters considered highlighted the high quality of the
339 Italian Mediterranean bulk buffalo milk under study. Due to the remarkable season effects discovered for
340 many of the monitored parameters, we can conclude that a seasonal adjustment of the buffaloes in terms
341 of delivery scheduling should be desirable for maintaining minimum oscillation of milk components
342 throughout the four seasons, thus resulting in bulk milk standardization and constant production of
343 buffalo dairy products. As is well known, milk testing and quality control represent pivotal activities of
344 any milk processing industry; for buffalo milk, such activities can provide important information for the
345 exploitation of this poorly utilized food matrix. Given the richness in composition of buffalo milk, its
346 health properties are supposed to be superior to those of cow milk, thus an increase of knowledge
347 regarding this matrix could be useful to drive the dairy industry toward the development of new buffalo
348 milk-based products with high nutritional properties.

349

350 ACKNOWLEDGMENTS

351

352 The authors wish to thank the *Associazione Regionale Allevatori* of *Marche Region* (ARAM), via
353 Clementina, Falconara Marittima (Italy), for its technical support. The authors also wish to thank the
354 laboratory staff at *Centro Agrochimico Regionale, Qualità delle Produzioni* of *Agenzia per i Servizi nel*
355 *Settore Agroalimentare delle Marche (ASSAM)* for analytical support.

356

357 REFERENCES

358

359

360

- 1 358
2
3 359 Abd El-Salam MH, El-Shibini S. 2011. A comprehensive review on the composition and properties of
4
5 360 buffalo milk. *Dairy Science and Technology* **91**, 663-699.
6
7 361 Ahmad S. 2013. Buffalo milk. Milk and dairy products in human nutrition: Production, Composition and
8
9 362 Health, First Edition. Eds Young W. Park and George F.W. Haenlein. John Wiley & Sons, Ltd. pp.
10 363 519-553.
11
12 364 Ahmad S, Anjum FM, Huma N, Sameen A, Zahor T. 2013. Composition and physico-chemical
13 365 characteristics of buffalo milk with particular emphasis on lipids, proteins, minerals, enzymes and
14
15 366 vitamins. *Journal of Animal and Plant Sciences* **23**, 62-74.
16
17 367 Ahmad S, Gaucher I, Rousseau F, Beaucher E, Piot M, Grongnet JF, *et al.* 2008. Effects of acidification
18 368 on physico-chemical characteristics of buffalo milk: a comparison with cow's milk. *Food*
19 369 *Chemistry* **106**, 11–17.
20
21 370 Altiero V, Moio L, Addeo F. 1989. Prediction of Mozzarella cheese yield on the basis of fat protein
22 371 content of water buffalo milk. *Scienza e Tecnica Lattiero Casearia* **40**,425-433.
23
24 372 Bartocci S, Tripaldi C, Terramoccia S. 2002. Characteristics of foodstuffs and diets, and the quanti-
25 373 qualitative milk parameters of Mediterranean buffaloes bred in Italy using the intensive system. An
26 374 estimate of the nutritional requirements of buffalo herds lactating or dry. *Livestock Production*
27 375 *Science* **77**, 45–58.
28
29 376 Bassbasi M., Platikanov S, Tauler R, Oussama A. 2014. FTIR-ATR determination of solid non fat (SNF)
30 377 in raw milk using PLS and SVM chemometric methods. *Food Chemistry* **146**, 250–254.
31 378 Bonfatti V, Giantin M, Rostellato R, Dacasto M, Carnier P. 2013a. Separation and quantification of water
32 379 buffalo milk protein fractions and genetic variants by RP-HPLC. *Food Chemistry* **136**, 364-367.
33 380 Bonfatti V, Gervaso M, Rostellato R, Coletta A, Carnier P. 2013b. Protein composition affects variation
34 381 in coagulation properties of buffalo milk. *Journal of Dairy Science* **96**, 4182-4190.
35 382 Borghese A. 2005. Buffalo Production And Research. Reu Technical Series 67. Fao Regional Office For
36 383 Europe Inter-Regional Cooperative Research Network On Buffalo (Escorena) Buffalo. Food And
37 384 Agriculture Organization Of The United Nations. Rome, 2005. pp. 312.
38
39 385 Brown MH, Gill CO, Hollingsworth J, Nickelson II R, Seward S, Sheridan JJ, *et al.* 2000. The role of
40 386 microbiological testing in systems for assuring the safety of beef. *International Journal of Food*
41 387 *Microbiology* **62**, 7–16.
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2 388 Cazacu S, Rotsios K, Moshonas G. 2014. Consumers' Purchase Intentions towards Water Buffalo Milk
3 389 Products (WBMPs) in the Greater Area of Thessaloniki, Greece. *Procedia Economics and Finance*
4 390 **9**, 407-416.
- 7 391 Commission Regulation (EC) No 1107/96 of 12 June 1996 on the registration of geographical indications
8 392 and designations of origin under the procedure laid down in Article 17 of Council Regulation
9 393 (EEC) No 2081/92 Available online at: [http://eur-lex.europa.eu/legal-](http://eur-lex.europa.eu/legal-content/EN/TXT/PDF/?uri=CELEX:31996R1107&from=EN)
10 394 [content/EN/TXT/PDF/?uri=CELEX:31996R1107&from=EN](http://eur-lex.europa.eu/legal-content/EN/TXT/PDF/?uri=CELEX:31996R1107&from=EN).
- 14 395 Commission Regulation (EC) No 103/2008 of 4 February 2008 approving non-minor amendments to the
15 396 specification for a name entered in the register of protected designations of origin and protected
16 397 geographical indications — Mozzarella di Bufala Campana (PDO) Available online at: [http://eur-](http://eur-lex.europa.eu/legal-content/EN/TXT/PDF/?uri=CELEX:32008R0103&from=EN)
17 398 [lex.europa.eu/legal-content/EN/TXT/PDF/?uri=CELEX:32008R0103&from=EN](http://eur-lex.europa.eu/legal-content/EN/TXT/PDF/?uri=CELEX:32008R0103&from=EN).
- 22 399 Cunha Neto OC, Oliveira CAF, Hotta RM, Sobral PJA. 2005. Avaliação físico-química e sensorial do
23 400 iogurte natural produzido com leite de búfala contendo diferentes níveis de gordura. *Ciência e*
24 401 *Tecnologia de Alimentos* **25**, 448-453.
- 27 402 Di Francia A, De Rosa G, Masucci F, Romano R, Borriello I, Grassi C. 2007a. Effect of *Pisum Sativum*
28 403 as protein supplement on buffalo milk production. *Italian Journal of Animal Science* **6**, 472-475.
- 31 404 Di Francia A, Masucci F, De Rosa G, Grasso F, Proto V. 2007b. Feeding management and milk
32 405 production in organic and conventional buffalo farms. *Italian Journal of Animal Science* **6**, 571-
33 406 574.
- 37 407 EFSA BIOHAZ Panel (EFSA Panel on Biological Hazards). (2015). Scientific Opinion on the public
38 408 health risks related to the consumption of raw drinking milk. *EFSA Journal* 2015;13(1):3940, 95
39 409 pp. doi:10.2903/j.efsa.2015.3940.
- 42 410 Enb A, Abou Donia MA, Abd-Rabou NS, Abou-Arab AAK, El-Senaiti MH. 2009. Chemical composition
43 411 of raw milk and heavy metals behavior during processing for milk products. *Global Veterinaria* **3**,
44 412 268-275.
- 48 413 FAOSTAT (2014) Food and Agriculture Organization of the United Nations. Available on line:
49 414 <http://faostat.fao.org/site/569/default.aspx#ancor>.
- 52 415 Gürler Z, Kuyucuoğlu Y, Pamuk Ş. 2013. Chemical and microbiological quality of Anatolian Buffalo
53 416 milk. *African Journal of Microbiology Research* **7**, 1512-1517.
- 55 417 Han X, Lee FL, Zhang L, Gou MR. 2012. Chemical composition of water buffalo milk and its low-fat
56 418 symbiotic yogurt development. *Functional Foods in Health and Disease* **2**, 86-106.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
- 419 Hussain I, Grandison AS, Bell AE. 2012. Effects of gelation temperature on Mozzarella-type curd made
420 from buffalo and cows' milk. 1: Rheology and microstructure. *Food Chemistry* **134**, 1500-1508.
- 421 ILAC-G13: 2007. Guidelines for the requirements for the competence of providers of proficiency testing
422 schemes.
- 423 INEA, Istituto Nazionale di Economia Agraria. 2012. Annuario dell'Agricoltura Italiana 2011. LXV,
424 INEA, Rome, pp. 430-431.
- 425 INEA, Istituto Nazionale di Economia Agraria. 2013. Annuario dell'Agricoltura Italiana 2012. LXVI,
426 INEA, Rome, p. 456.
- 427 ISO 4833-1:2013. Microbiology of the food chain - Horizontal method for the enumeration of
428 microorganisms - Part 1: Colony count at 30 degrees C by the pour plate technique.
- 429 ISO 13366-1:2008 (IDF 148-1:2008). Milk - Enumeration of somatic cells -Part 1: Microscopic method
430 (Reference method).
- 431 ISO 5725-2:1994. Accuracy (trueness and precision) of measurement methods and results - part 2.
- 432 ISO 9001:2008. Quality management systems – Requirements.
- 433 ISO/IEC 17043:2010. Conformity assessment – General Requirements for proficiency testing.
- 434 ISTAT, Istituto Italiano di Statistica. 2014. Annuario Statistico Italiano 2014. p. 440. Available online:
435 <http://www.istat.it/it/files/2014/11/Asi-2014.pdf>.
- 436 ISTAT, Istituto Italiano di Statistica. 2015. Annuario Statistico Italiano 2015. p. 462. Available online:
437 <http://www.istat.it/it/files/2015/12/Asi-2015.pdf>.
- 438 Khedkar CD, Kalyankar SD, Deosarkar SS. 2016. Buffalo Milk. Reference Module in Food Science,
439 from *Encyclopedia of Food and Health* 522-528.
- 440 Kelly AL, Tiernan D, O'Sullivan C, Joyce P. 2000. Correlation between bovine milk somatic cell count
441 and polymorphonuclear leukocyte level for sample of bulk milk and milk from individual cows.
442 *Journal of Dairy Science* **83**, 300-304.
- 443 Kapila R, Kavadi KP, Kapila S. 2013. Comparative evaluation of allergic sensitization to milk proteins
444 of cow, buffalo and goat. *Small Ruminant Research* **112**, 191-198.
- 445 Masucci F, Di Francia A, De Rosa G, Romano R, Varricchio ML, Grassi C. 2008. *Pisum sativum* as
446 alternative protein source in diets for buffalo cows in middle and late stage of lactation. 16th
447 IFOAM Organic World Congress, Modena, Italy, June 16-20, pp. 164-167.

- 1
2 448 Mateo MJ, Everard CD, Fagan CC, O'Donnell CP, Castillo M, Payne FA, *et al.* 2009. Effect of milk fat
3 449 concentration and gel firmness on syneresis during curd stirring in cheese-making. *International*
4 450 *Dairy Journal* **19**, 264-268.
- 7 451 Melilli C, Lynch JM, Carpino S, Barbano DM, Licitra G, Cappa A. 2002. An empirical method for
8 452 prediction of cheese yield. *Journal of Dairy Science* **85**, 2699–2704.
- 11 453 Ménard O, Ahmad S, Rousseau F, Briard-Bion V, Gaucheron F, Lopez C. 2010. Buffalo vs. cow milk fat
12 454 globules: Size distribution, zeta-potential, compositions in total fatty acids and in polar lipids from
14 455 the milk fat globule membrane. *Food Chemistry* **120**, 544-551.
- 16 456 More SJ, Clegg TA, Lynch PJ, O'Grady L. 2013. The effect of somatic cell count data adjustment and
17 457 interpretation, as outlined in European Union legislation, on herd eligibility to supply raw milk for
18 458 processing of dairy products. *Journal of Dairy Science* **96**, 3671-3681.
- 22 459 Moroni P, Rossi CS, Pisoni G, Bronzo V, Castiglioni B, Boettcher PJ. 2006. Relationships between
23 460 somatic cell count and intramammary infection in buffaloes. *Journal of Dairy Science* **89**, 998–
25 461 1003.
- 27 462 Napolano R, Potena A, Iovane G, Coletta A, Campanile G, Neglia G, *et al.* 2007. Relationships between
28 463 Italian Mediterranean Buffalo milk compounds. *Italian Journal of Animal Science* **6**, 1088-1091.
- 31 464 Pasquini M, Tommei B, Mattii S. 2003. Buffalo milk: proteins electrophoretic profile and somatic cell
32 465 counts. *Italian Journal of Animal Science* **2**, 299-301.
- 35 466 Pure Appl. Chem. Vol. 78, n°1 pp.145-196, 2006. The International harmonized protocol for the
36 467 proficiency testing of analytical chemistry laboratories.
- 38 468 Regulation (EC) No 853/2004 of the European Parliament and of the Council of 29 April 2004 laying
39 469 down specific hygiene rules for food of animal origin. Official Journal of the European Union.
41 470 Available online: [http://eur-](http://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:L:2004:139:0055:0205:en:PDF)
42 471 [lex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:L:2004:139:0055:0205:en:PDF](http://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:L:2004:139:0055:0205:en:PDF).
- 46 472 Regulation (EC) No 1664/2006 of 6 November 2006 amending Regulation (EC) No 2074/2005 as regards
47 473 implementing measures for certain products of animal origin intended for human consumption and
48 474 repealing certain implementing measures. Official Journal of the European Union. Available
50 475 online: [http://eur-](http://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:L:2006:320:0013:0045:EN:PDF)
51 476 [lex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:L:2006:320:0013:0045:EN:PDF](http://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:L:2006:320:0013:0045:EN:PDF).
- 55
56
57
58
59
60

- 1
2 477 Rosati A, Van Vleck LD. 2002. Estimation of genetic parameters for milk, fat, proteine and mozzarella
3 478 cheese production for the Italian river buffalo *Bubalus bubalis* population. *Livestock Production*
4 479 *Science* **74**, 185-190.
- 7 480 Sharif A, Ahmad T, Bilal MQ, Yousaf A, Muhammad G, Ur-Rehman S, *et al.* 2007. Estimation of milk
8 481 lactose and somatic cells for the diagnosis of sub-clinical mastitis in dairy buffaloes. *International*
9 482 *Journal of Agriculture and Biology* **9**, 267–270.
- 12 483 Shelke SK, Thakur SS, Amrutkar SA. 2012. Effect of feeding protected fat and proteins on milk
14 484 production, composition and nutrient utilization in Murrah buffaloes (*Bubalus bubalis*). *Animal*
15 485 *Feed Science and Technology* **171**, 98-107.
- 18 486 Simoes MG, Portal RE, Rabelo JG, Ferreira CLLF. 2014. Seasonal variations affect the physicochemical
19 487 composition of buffalo milk and artisanal cheeses produced in Marajo island. *Advance Journal of*
20 488 *Food Science and Technology* **6**, 81-91.
- 23 489 Smith KL. 2002. A discussion of normal and abnormal milk based on somatic cell count and clinical
24 490 mastitis. *Bulletin - International Dairy Federation* **372**, 43-45.
- 27 491 Tripaldi C, Palocci G, Miarelli M, Catta M, Orlandini S, Amatiste S, *et al.* 2010. Effects of mastitis on
28 492 buffalo milk quality. *Asian-Australasian Journal of Animal Sciences* **23**, 1319-1324.
- 31 493 Tufarelli V, Dario M, Laudadio V. 2008. Diet composition and milk characteristics of Mediterranean
32 494 water buffaloes reared in Southeastern Italy during spring season. *Livestock Research for Rural*
33 495 *Development* **20**, 10.
- 36 496 Yadav P, Kumar P, Mukesh M, Kataria RS, Yadav A, Mohanty AK, *et al.*, 2015. Kinetics of lipogenic
37 497 genes expression in milk purified mammary epithelial cells (MEC) across lactation and their
38 498 correlation with milk and fat yield in buffalo. *Research in Veterinary Science* **99**, 129-136.
- 41 499 Zicarelli L, Neglia G, Napolano R, Di Rubio M, Caso C, Gasparrini B, *et al.* 2007. Correction of the ratio
42 500 between real cheese yield at 28 hours and protein according to chemical composition of buffalo
43 501 milk. *Italian Journal of Animal Science* **6**, 1108-1111.
- 46 502 Zotos A, Bampidis VA. 2014. Milk fat quality of Greek buffalo (*Bubalus bubalis*). *Journal of Food*
47 503 *Composition and Analysis* **33**, 181-186.
- 51
52
53
54
55
56
57
58
59
60

Table 1 Herd characteristics and sampling plan of bulk buffalo milk (n=51) analyzed in the period 2011-2013.

Herd characteristics	Years		
	2011	2012	2013
	#	#	#
Lactating Cows	74	80	87
Herd average age (years)	5	4.8	5
Milk yield (kg/head)	1,804	1,662	1,745
Days in milking	208	207	216
Daily milk yield (kg/day)	8.67	8.02	8.07
Sampling plan	Years		
	2011	2012	2013
	#	#	#
Winter	2	2	5
Spring	3	5	6
Summer	3	5	6
Autumn	2	7	5
Total	10	19	22

Table 2 Descriptive statistics for quality parameters of bulk buffaloes' milk analyzed between 2011 and 2013.

	Fat (% w/w)	Protein (TN) (% w/w)	Lactose (% w/w)	SNF (%)	CYF (kg 100 kg ⁻¹)	SCC (x 1,000 cells mL ⁻¹)	TBC (x 10 ¹ cfu mL ⁻¹)
Mean	7.13	4.39	4.77	9.86	23.27	175.57	9.91
SD	0.85	0.20	0.24	0.28	1.60	109.87	26.11
Minimum	5.36	4.02	3.75	8.63	20.45	42.00	0.40
Maximum	9.99	4.94	5.12	10.39	27.75	566.00	190.40

TN Total Nitrogen; SNF Solids-not-fat; CYF Cheese yield formula; SCC Somatic cell counts; TBC Total bacterial counts.
cfu colony forming units.
SD standard deviation.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

Table 3 ANOVA results for quality parameters of bulk buffaloes' milk according to month, season and year.

Source of variation	Fat (% w/w)		Protein (TN) (% w/w)		Lactose (% w/w)		SNF (%)		CYF (kg 100 kg ⁻¹)		SCC (x 1,000 cells mL ⁻¹)		TBC (x 10 ¹ cfu mL ⁻¹)	
	df	Variance	df	Variance	df	Variance	df	Variance	df	Variance	df	Variance	df	Variance
Month	11	2.267 ***	11	0.059 n.s.	11	0.107 *	11	0.116 n.s.	11	7.017 ***	11	23,079.5 *	11	47,458.4 n.s.
Error	39	0.285	39	0.037	39	0.042	39	0.067	39	1.312	39	8,965.8	39	74,047.1
Season	3	6.258 ***	3	0.145 *	3	0.189 *	3	0.164 n.s.	3	19.389 ***	3	32,984.8 *	3	41,797.9 n.s.
Error	47	0.368	47	0.035	47	0.048	47	0.072	47	1.493	47	10,735.9	47	69,882.7
Year	2	0.858 n.s.	2	0.075 n.s.	2	0.226 *	2	0.254 *	2	4.085 n.s.	2	11,823.6 n.s.	2	113,462.0 n.s.
Error	48	0.715	48	0.040	48	0.049	48	0.070	48	2.504	48	12,081.1	48	66,312.0

TN Total Nitrogen; SNF Solids-not-fat; CYF Cheese yield formula; SCC Somatic cell counts; TBC Total bacterial counts.
 df degrees of freedom.
 * Significant at $P \leq 0.05$
 ** Significant at $P \leq 0.01$
 *** Significant at $P \leq 0.001$
 n.s. not significant

Table 4 Quality parameters of bulk Mediterranean buffaloes' milk according to month, season and year.

Effect	Fat (% w/w)	Protein (TN) (% w/w)	Lactose (% w/w)	SNF (%)	CYF (kg 100 kg ⁻¹)	SCC (x 1,000 cells mL ⁻¹)	TBC (x 10 ¹ cfu mL ⁻¹)
Month							
January	8.54±0.27 ^a	4.63±0.10 ^a	4.61±0.10 ^a	9.93±0.13 ^{ab}	25.84±0.57 ^a	299.00±47.34 ^a	6.68±13.61 ^a
February	7.02±0.27 ^{bcd}	4.26±0.10 ^a	4.56±0.10 ^a	9.51±0.13 ^b	22.68±0.57 ^{bcd}	274.25±47.34 ^{ab}	7.32±13.61 ^a
March	6.80±0.27 ^{cd}	4.39±0.10 ^a	4.51±0.10 ^a	9.60±0.13 ^{ab}	22.87±0.57 ^{bcd}	127.50±47.34 ^{ab}	9.85±13.61 ^a
April	6.68±0.27 ^{cd}	4.35±0.10 ^a	4.70±0.10 ^a	9.76±0.13 ^{ab}	22.58±0.57 ^{bcd}	293.50±47.34 ^{ab}	3.90±13.61 ^a
May	6.50±0.27 ^d	4.25±0.10 ^a	4.84±0.10 ^a	9.79±0.13 ^{ab}	21.98±0.57 ^{cd}	87.25±47.34 ^b	6.65±13.61 ^a
June	6.27±0.24 ^d	4.28±0.09 ^a	4.94±0.09 ^a	9.91±0.12 ^{ab}	21.80±0.51 ^d	121.80±42.35 ^{ab}	3.66±12.17 ^a
July	6.54±0.24 ^d	4.33±0.09 ^a	4.89±0.09 ^a	9.91±0.12 ^{ab}	22.31±0.51 ^{cd}	125.40±42.35 ^{ab}	4.02±12.17 ^a
August	6.64±0.31 ^{cd}	4.30±0.11 ^a	4.98±0.12 ^a	9.97±0.15 ^{ab}	22.35±0.66 ^{bcd}	116.33±54.67 ^{ab}	5.37±15.71 ^a
September	7.25±0.24 ^{bcd}	4.49±0.09 ^a	4.94±0.09 ^a	10.12±0.12 ^a	23.75±0.51 ^{abcd}	174.00±42.35 ^{ab}	6.74±12.17 ^a
October	7.30±0.24 ^{bcd}	4.42±0.09 ^a	4.84±0.09 ^a	9.95±0.12 ^{ab}	23.56±0.51 ^{abcd}	124.20±42.35 ^{ab}	7.22±12.17 ^a
November	8.04±0.31 ^{abc}	4.50±0.11 ^a	4.71±0.12 ^a	9.90±0.15 ^{ab}	24.77±0.66 ^{abc}	224.33±54.67 ^{ab}	6.83±15.71 ^a
December	8.15±0.24 ^{ab}	4.51±0.09 ^a	4.68±0.09 ^a	9.88±0.12 ^{ab}	24.94±0.51 ^{ab}	175.80±42.35 ^{ab}	8.40±12.17 ^a
Season							
Winter	7.81±0.20 ^a	4.50±0.06 ^a	4.59±0.07 ^b	9.78±0.09 ^a	24.49±0.41 ^a	269.44±34.54 ^a	8.91±8.81 ^a
Spring	6.40±0.16 ^b	4.27±0.05 ^b	4.76±0.06 ^{ab}	9.73±0.07 ^a	21.96±0.33 ^b	153.29±27.69 ^b	5.04±7.06 ^a
Summer	6.79±0.16 ^b	4.36±0.05 ^{ab}	4.91±0.06 ^a	9.95±0.07 ^a	22.73±0.33 ^b	147.00±27.69 ^{ab}	17.66±7.06 ^a
Autumn	7.76±0.16 ^a	4.48±0.05 ^a	4.77±0.06 ^{ab}	9.94±0.07 ^a	24.36±0.33 ^a	166.07±27.69 ^{ab}	7.65±7.06 ^a
Year							
2011	7.09±0.27 ^a	4.33±0.06 ^a	4.95±0.07 ^a	9.97±0.08 ^a	22.99±0.50 ^a	165.60±34.76 ^a	23.41±8.14 ^a
2012	7.36±0.19 ^a	4.46±0.05 ^a	4.78±0.05 ^{ab}	9.93±0.06 ^a	23.79±0.36 ^a	152.84±25.22 ^a	6.46±5.91 ^a
2013	6.95±0.18 ^a	4.37±0.04 ^a	4.69±0.05 ^b	9.74±0.06 ^a	22.95±0.34 ^a	199.73±23.43 ^a	6.74±5.49 ^a

Multiple Comparisons among Least Square Means (LSM) using the Tukey HSD test according to month, season and year (LSM ± mean std. err.).

TN Total Nitrogen; SNF Solids-not-fat; CYF Cheese yield formula; SCC Somatic cell counts; TBC Total bacterial counts.

Within each effect, for each variable, means with different letters are significantly different ($P \leq 0.05$).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49


Table 5 Simple correlation among variables evaluated by the Pearson correlation coefficient (51 bulk milk samples).

	Fat (% w/w)	Protein (TN) (% w/w)	Lactose (% w/w)	SNF (%)	CYF (kg 100 kg ⁻¹)	SCC (x 1,000 cells mL ⁻¹)
Protein (%)	0.6444 **					
Lactose (%)	-0.2777 *	-0.2068 n.s.				
SNF (%)	0.2346 n.s.	0.5514 **	0.7017 **			
CYF (kg 100 kg⁻¹)	0.9398 **	0.8669 **	-0.2735 n.s.	0.3993 **		
SCC (x 1,000 cells mL⁻¹)	0.2684 n.s.	0.3113 *	-0.3579 n.s.	-0.0839 n.s.	0.3141 *	
TBC (x 10¹ cfu mL⁻¹)	-0.0255 n.s.	0.0373 **	0.1775 n.s.	0.1724 n.s.	-0.0005 n.s.	0.0202 n.s.

TN Total Nitrogen; SNF Solids-not-fat; CYF Cheese yield formula; SCC Somatic cell counts; TBC Total bacterial counts.

n.s. not significant
* Significant at $P \leq 0.05$
** Significant at $P \leq 0.01$

Fig. 1 Dendrogram of buffaloes' bulk milk parameters analyzed between 2011 and 2013 according to season resulting from Cluster Analysis carried out using the Wards' minimum variance method.



win winter; spr spring; sum summer; aut autumn
 § mis-assigned records

Review Only