



UNIVERSITÀ POLITECNICA DELLE MARCHE
Repository ISTITUZIONALE

Trends and approaches in the analysis of ecosystem services provided by grazing system: A review

This is the peer reviewed version of the following article:

Original

Trends and approaches in the analysis of ecosystem services provided by grazing system: A review / D'Ottavio, Paride; Francioni, Matteo; Trozzo, Laura; Sedic, Elmir; Budimir, Katarina; Avanzolini, Pietro; Trombetta, Maria Federica; Prqueddu, C.; Santilocchi, Rodolfo; Toderi, Marco. - In: GRASS AND FORAGE SCIENCE. - ISSN 0142-5242. - STAMPA. - 73:(2018), pp. 15-25. [10.1111/gfs12299]

Availability:

This version is available at: 11566/249261 since: 2020-10-09T11:03:39Z

Publisher:

Published

DOI:10.1111/gfs12299

Terms of use:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. The use of copyrighted works requires the consent of the rights' holder (author or publisher). Works made available under a Creative Commons license or a Publisher's custom-made license can be used according to the terms and conditions contained therein. See editor's website for further information and terms and conditions.

This item was downloaded from IRIS Università Politecnica delle Marche (<https://iris.univpm.it>). When citing, please refer to the published version.

note finali coverpage

(Article begins on next page)

1 **Trends and approaches in the analysis of ecosystem services provided by**
2 **grazing systems: a review**

3
4 D'Ottavio P.^{1*}, Francioni M.¹, Trozzo L.¹, Sedić E.¹, Budimir K.¹, Avanzolini P.¹, Trombetta
5 M.F.¹, Porqueddu C.², Santilocchi R.¹, Toderi M.^{1*}

6
7 ¹ Dipartimento di Scienze Agrarie, Alimentari ed Ambientali, Università Politecnica delle
8 Marche, Ancona, Italy

9 ² (CNR) National Research Council, (ISPAAM) Institute for Animal Production System in
10 Mediterranean Environment, Sassari, Italy

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26 ***Correspondence to:** P. D'Ottavio and M. Toderi

27 Dipartimento di Scienze Agrarie, Alimentari ed Ambientali

28 Università Politecnica delle Marche

29 Ancona, Italy

30 E-mails: p.dottavio@univpm.it; m.toderi@univpm.it

31

32

33 **Abstract**

34

35 The ecosystem services (ES) approach is a framework for describing the benefits of nature to
36 human well-being, and this has become a popular instrument for assessment and evaluation of
37 ecosystems and their functions. Grazing lands can provide a wide array of ES that depend on
38 their management practices and intensity. This article reviews the trends and approaches used
39 in the analysis of some relevant ES provided by grazing systems, in line with the framework
40 principles of the Millennium Ecosystem Assessment (MA). The scientific literature provides
41 reports of many studies on ES in general, but the search here focussed on grazing systems,
42 which returned only 62 papers. This review of published papers highlights that: (i) in some
43 papers, the concept of ES as defined by the MA is misunderstood (e.g., lack of anthropocentric
44 vision); (ii) 34% of the papers dealt only with one ES, which neglects the need for the
45 multisectoral approach suggested by the MA; (iii) only a few papers included stakeholder
46 involvement to improve local decision-making processes; (iv) cultural ES have been poorly
47 studied despite being considered the most relevant for local and general stakeholders; and (v)
48 stakeholder awareness of well-being as provided by ES in grazing systems can foster both agri-
49 environmental schemes and the willingness to pay for these services.

50

51

52 **Keywords.** Primary production, habitat services, food, land degradation prevention, water
53 quality regulation, regulation of water flows, climate regulation, moderation of extreme events,
54 natural (landscape) heritage

55

56

57 **1. Introduction**

58

59 Although the first references to the concept of “ecosystem functions, services and values” date
60 back to around the 1960s, the number of scientific papers concerning ecosystem services (ES)
61 has grown exponentially in the last few decades (de Groot et al., 2002). This is particularly the
62 case since the publication of the Millennium Ecosystem Assessment (MA) (Fisher et al., 2009).
63 The MA (Alcamo et al., 2003; Millennium Ecosystem Assessment, 2005) represents one of the
64 most extensive and widely accepted studies on the links between human well-being and the
65 world ecosystems. It defines the ecosystem as “a dynamic complex of plant, animal (including
66 humans), and microorganism communities and the non-living environment interacting as a
67 functional unit”, and ecosystem services as “the benefits people obtain from ecosystems”.
68 According to Alcamo et al. (2003), the goal of MA is to establish the scientific basis for actions
69 that are needed to enhance the contributions of ecosystems to human well-being without
70 undermining their long-term productivity. The MA conceptual framework assumes that there
71 is a dynamic interaction between people and ecosystems that requires a multiscale approach, as
72 this better reflects the multiscale nature of decision making. Effective incorporation of different
73 types of knowledge into ES assessment can both improve the findings and help to increase their
74 adoption by stakeholders. The MA conceptual framework places human well-being as the
75 central focus for assessment.

76 The MA identified four groups of ES: (i) Supporting: services necessary for the
77 production of all other ES (e.g., soil formation, nutrient cycling), where the impact on people
78 is either indirect or occurs over a very long time; (ii) Provisioning: products obtained from
79 ecosystems, such as food and fresh water; (iii) Regulating: benefits obtained from the regulation
80 of ecosystem processes, such as climate and disease control; and (iv) Cultural: non-material
81 benefits people obtain from ecosystems through spiritual enrichment, cognitive development,
82 reflection, recreation, and aesthetic experiences. A second key study concerning ES, The
83 Economics of Ecosystems and Biodiversity (TEEB, 2010), defines ES as “the direct and
84 indirect contributions of ecosystems to human well-being”, and separates the concepts of
85 services and benefits (welfare gains generated by ES), while considering supporting services
86 merely as ecological processes, and not strictly as ES.

87 Although it is recognized that each ecosystem can produce a large number of ES
88 (Alcamo et al., 2003; Millennium Ecosystem Assessment, 2005), ecosystems can also produce
89 ecosystem disservices that are harmful or detrimental to human well-being (von Döhren and
90 Haase, 2015). Thus, the term “ecosystem service” is anthropocentric and is intended to have a

91 positive sense. This vision is one of the recurring critiques of the concept of ES, and according
92 to Schröter et al. (2014), the ES concept is not meant to replace biocentric arguments, but to
93 group together a wide variety of anthropocentric arguments for the protection and sustainable
94 use of ecosystems by humans. Schröter et al. (2014) also counter-argued six other main critiques
95 to the ES concept that were derived from the scientific literature.

96 Ecosystem services are spatial-scale and time-scale dependent, and there is a risk that
97 spatial scale mismatches between ecological processes and decision making will occur. For this
98 reason, the need for an integrated approach that also takes into account the local knowledge of
99 stakeholders is a key requirement in assessing ES (Alcamo et al., 2003; Millennium Ecosystem
100 Assessment, 2005; Reed, 2008).

101 According to Alcamo et al. (2003) and TEEB (2010), ecosystems and biodiversity are
102 closely related concepts, although biodiversity is not strictly considered as an ES, but rather as
103 a source or a regulator of the ecosystem (Harrison et al., 2014). The knowledge gap regarding
104 both the links and the difficulties in understanding the relationships between ES and
105 biodiversity has been highlighted by many authors (e.g., Jax and Heink, 2015; Sircely and
106 Naeem, 2012; Harrison et al., 2014).

107 Livestock systems occupy about a third of the ice-free land surface of the planet, and
108 they represent an important source of income; indeed, they can even be essential for the survival
109 of vulnerable human communities. In these systems, grazing land can provide a large and
110 differentiated number of ES (Porqueddu et al., 2016; Tarrasón et al., 2016). These ES are, in
111 turn, dependent on the different management practices (Fischer et al., 2010; Steiner et al., 2014),
112 such as different grazing regimes (Ford et al., 2012).

113 This article reviews the trends and approaches used in the analysis of some relevant ES
114 provided by grazing systems, in line with the framework principles of the MA. In the context
115 of this review, grazing systems include production systems in which grazing is one of the main
116 management practices adopted across the grazing lands (Allen et al., 2011). This review will
117 analyse: (i) if the papers follow the principles of the MA, and the main reasons behind their
118 missed adoption; (ii) which are the most analysed ES, and which require further investigation
119 within grazing systems; (iii) how different types of knowledge have been incorporated into ES
120 assessment, as requested by the MA; and (iv) how ES concepts have fed the decision-making
121 process. It is intended that the results of this review can be used to derive recommendations for
122 research activities in the analysis of ES.

123

124 **2. Links between biodiversity and ecosystem services**

125

126 Biodiversity is the variability between living organisms, and it includes diversity within and
127 among species and ecosystems. Biodiversity is the source of many goods and services, such as
128 food and genetic resources, and changes in biodiversity can influence the supply of ES (Alcamo
129 et al., 2003). Subsequently the MA (2005) defined biodiversity as a necessary condition for the
130 delivery of all ES, and in most cases, a greater level of biodiversity is associated with a larger
131 or more dependable supply of ES.

132 According to the MA (2005), biodiversity is both a response variable that is affected by
133 the drivers of global change (e.g., climate, change in land use) and a factor that modifies
134 ecosystem processes and ES, and indirectly, human well-being (e.g., health, freedom of choice
135 and action). Changes in human well-being can lead to modifications to management practices,
136 with direct effects on ecosystem processes and biodiversity (Figure 1). Although the MA
137 describes a unilateral relationship between biodiversity and ES, some authors consider
138 biodiversity as a service in its own right; e.g., as the basis of nature-based tourism (van Wilgen
139 et al., 2008). However, others consider that biodiversity can have different roles as a regulator
140 of ecosystem processes, as a service in itself, or as a good (Mace et al., 2012).

141

142 **Insert Fig. 1 about here**

143

144 Habitat provisioning is one of the main ecosystem services that links the effects of
145 livestock grazing to the biodiversity of the host ecosystem (Hoffman et al., 2014). Habitat
146 services arise from the direct interactions of animals with their environments, and are hence
147 related to land-management practices, especially in relation to grazing systems. Unlike the MA
148 (Alcamo et al., 2003; Millennium Ecosystem Assessment, 2005), the TEEB (2010) considers
149 habitat services as a separate category. In agreement with these documents, this review
150 considers habitat services within supporting services, because of their interconnected nature
151 and their shared roles in underpinning the delivery of other services.

152

153 **3. Bibliographic search and analysis criteria**

154

155 This review is based on the ES provided by grazing systems as categorised and defined as
156 prominent by Hoffman et al. (2014) (Table 1). Among these, the ES relevant to the expertise
157 and background of the authors were analysed in detail: primary production (PP), habitat services
158 (HS), food and other livestock-related products (FP), land degradation and soil erosion (LD),

159 water quality regulation/ purification (WQ), regulation of water flows (WF), climate regulation
160 (CR), moderation of extreme events (EE), and natural (landscape) heritage (NH) (Box 1).

161

162 **Insert Box 1**

163

164 Published papers dealing with ES were sampled in January 2016 using the Web of
165 Science™ (WoS). Within the search option of “topic” the basic string *"ecosystem service*" and*
166 *("grassland*" or "rangeland*" or "shrubland*" or "scrubland*") and "grazing"* was used as
167 input in the “field search” (“basic search”), starting from 2004 as the “timespan”. To have a
168 preliminary selection for each analysed ES, specific search terms were added to the basic string
169 according to the keywords (Table 1) included in the Food and Agriculture Organisation report
170 (Hoffman et al., 2014). The additional strings used for the preliminary selection are reported in
171 detail in Table 2.

172 All of the papers extracted with the basic string (155 papers) were analysed to verify the
173 adoption of the MA framework and the attribution of the papers to each ES, which was corrected
174 as necessary. The analysis of the extracted papers allowed the identification which ES were
175 analysed for each paper in the light of the MA, and which did not take the MA into account
176 (i.e., “ecosystem services” and/or “millennium ecosystem assessment” were merely cited in the
177 Introduction or Conclusions).

178 After the analysis of the extracted papers the following manuscripts were excluded from
179 this review: (i) papers dealing with ES that was not analysed (ii) reviews, editorials and meta-
180 analyses; and (iii) papers that did not adopt the MA framework.

181

182 **Insert Table 1**

183

184 **Insert Table 2**

185

186 **4. Trends and approaches in ecosystem services analysis**

187

188 ***4.1. The extracted papers: numbers, exclusion, and reasons for exclusion***

189 The basic string search returned a total of 155 papers (Table 1) with an increasing trend from
190 2010 (Figure 2). The multiple occurrence of different ES within single papers results in a total
191 of 529 findings within the 155 papers. Most papers dealt in particular with supporting (mostly
192 for PP and HS), regulating (in particular, CR and WF) and cultural (NH) ES. Only a few papers

193 dealt with FP, and surprisingly, very few with food itself. The addition of some other terms to
194 the basic string would have resulted in additional papers. For example, by adding *or “good*”*
195 to the basic string, the total number of papers for FP would increase from 12 to 38. This
196 highlights that many authors did not analyse food as an ES according to the MA framework.
197 Similar considerations can be stated for the other ES analysed.

198 The total number of extracted papers is surprisingly low compared to the far more
199 numerous papers that have analysed grazing systems from the economic and/or biophysical
200 perspective, but that did not adopt the MA framework. Indeed, by removing the keyword
201 “*ecosystem service**” from the basic string and maintaining the same time span, the number of
202 papers reached 5,983.

203

204 **Insert Fig. 2**

205

206 According to the review criteria, 29 papers were excluded from this review, as reviews,
207 editorials or meta-analyses, and 64 papers were excluded for only dealing with ES that were
208 not analysed in this review (e.g., fuel, power, pollination; 9 papers) or did not adopt the MA
209 framework (55 papers). With these papers, the term “ecosystem service” was present in the text
210 (e.g., in the Introduction), and for this reason they were extracted.

211 Sixty-two papers (149 findings) were eligible for the present analysis. NH was
212 apparently assessed in 25% of the papers, although it proved to be analysed as a cultural ES in
213 only 6% of the papers (out of 39 publications, 4 were eligible; Table 1). In the papers excluded
214 from the NH ES, the landscape was considered: (i) for the effects that it can have on biodiversity
215 (e.g., Cole et al., 2015; Kearns and Oliveras, 2009; Lindborg et al., 2009; Littlewood et al.,
216 2012; Sanderson et al., 2007); (ii) as support for improving or maintaining other ES, but not as
217 an ES *per se* (e.g., Lavorel et al., 2011, 2015; Schaldach et al., 2013); (iii) as an assessment
218 scale for other ES (e.g., Hussain and Tschirhart, 2013; Medina-Roldán et al., 2012; Peringer et
219 al., 2013; Kimoto et al., 2012); and (iv) for the effects that different drivers had on it without
220 directly analysing the consequences on its cultural value (e.g., Cousins et al., 2015; Lamarque
221 et al., 2014; Schaich et al., 2015). The limited number of papers dealing with the landscape as
222 a cultural ES might be explained by the difficulty for the measurement of this aspect, and to the
223 few currently available indicators (Feld et al., 2009; TEEB, 2010). Rather than being considered
224 as an EE, fire was analysed in some papers as a management tool for the enhancement of other
225 ES (e.g., habitat provisioning, prevention of wildfires), and for this reason these papers were
226 excluded from the EE analysis. For example, Joubert et al. (2014) investigated the effects of

227 annual burning on plant species richness, composition and turnover in three firebreak types,
228 and under different cattle grazing levels. Boughton et al. (2013) conducted an 8-year split-plot
229 experiment to study the effects of the season of burn on the plant composition of a semi-natural
230 grassland in Florida (USA), where in addition to prescribed winter burns, natural historical
231 wildfires occurred on abandoned ranchlands. The response of vegetation disturbance was
232 studied by Hancock and Legg (2012), for prescribed fire management in pine forests and
233 ericaceous heathlands in the UK. These papers were excluded from the NH and EE analyses,
234 but were included in the other ES analysed in this review; e.g., Lavorel et al. (2011) was
235 excluded from NH but was included in the HS, PP and CR analyses. “Landscape” and “fire”
236 were considered as particular cases, as these can have different meanings (e.g., scale of
237 investigation or management tools). The main reasons for the exclusion for the rest of the papers
238 (e.g., Bai et al., 2012; Loucougaray et al., 2015; Zeng et al., 2015) was the lack of adoption of
239 the MA approach or for only mentioning the term “ecosystem service” in the text (e.g., in the
240 Introduction or Abstract). Table 1 summarises these review categories according to the numbers
241 of papers for each ES extracted by the strings, the numbers of papers eligible for the analysis,
242 and the attribution of these papers to each ES.

243

244 ***4.2. The eligible papers: most and least analysed ecosystem services in combinations with*** 245 ***each other***

246 The predominance of papers dealing with PP (63% of the papers), HS (55%) and CR (50%)
247 that emerged in the extracted papers was confirmed for the eligible papers. Although livestock
248 production is clearly related to the forage characteristics of pastures (e.g., yield, quality, species
249 diversity, plant active compounds) (Lieber et al., 2014), only five papers included PP and FP
250 ES in the analyses (Figure 3). From the deep review of the papers, it clearly emerged that PP,
251 CR and HS were often analysed together; i.e., PP was assessed in 80% of the papers dealing
252 with CR (e.g., Medina-Roldán et al., 2012; Oñatibia et al., 2015) and in 60% of the papers
253 dealing with HS (e.g., Duru et al., 2015; Marriot et al., 2010), while HS was analysed in 40%
254 of the papers dealing with PP or CR. At the same time, these three ES were assessed with at
255 least one other ES (e.g., Lamarque et al., 2014; Miller et al., 2011); i.e., PP was analysed in
256 more than 70% of the papers dealing with FP (e.g., Koniak et al., 2011) or LD (e.g., Giese et
257 al., 2013), HS was analysed in 100% of the papers dealing with NH (e.g., Fontana et al., 2014),
258 CR was analysed in about 70% of the papers dealing with FP (e.g., Ford et al., 2012) and in
259 60% of the papers dealing with WQ (e.g., Roche et al., 2014) or with WF (e.g., Fisher et al.,
260 2011) (Figure 3). In the grazing systems, PP and HS were classified as supporting ES, and were

261 thus placed at the base of all of the other ES. This explains the high number of papers that dealt
262 with PP and HS. As a regulating ES, CR is a well-investigated topic, because it is strongly
263 linked to the urgent climate-change issues. Indeed, even if CR was one of the most analysed
264 ES, its analysis was mostly at a global scale, in terms of its role in net sequestration or net
265 emissions of greenhouse gases, while none of the papers analysed how changes in land cover
266 can affect both temperature and precipitation at local levels. The relationships between the
267 supporting ES, PP and HS and the other regulation ES was less analysed; i.e., WQ was assessed
268 only in 3% and 4% of the papers dealing with PP and HS, respectively, while WF was analysed
269 in about 20% of the papers dealing with PP or HS. Also, while 80% of the FP papers analysed
270 the relation with PP and about 67% analysed the relation with HS, only 13% and 11% of the
271 papers that assessed PP or HS included FP. A similar consideration can be derived for the
272 cultural ES NH, where 100% of the papers analysed the NH relationship with HS, and 80%
273 with PP. On the contrary, only 12% and 8% of the papers dealing with HS or PP included the
274 effects of different management options on NH within their study (Figure 3).

275 This analysis highlights that the authors tended to concentrate their research on ES very
276 close to each other in terms of their characteristics and relationships, and that they mostly
277 focussed on the supporting and regulating ES. Indeed, papers that dealt with ES that are distant
278 from each other represented the minority; e.g., between HS and FP or NH. In the next section,
279 the literature was analysed in terms of the advantages that derive from a multisectoral analysis
280 that also includes the provisioning and cultural ES, and how this analysis allows inclusion of
281 different stakeholders in the definition of shared management options or support policies (e.g.,
282 “Payments for ES” or “agri-environmental schemes”).

283

284 **Insert Fig. 3**

285

286 ***4.3. Millennium Ecosystem Assessment principles in the eligible papers***

287 Despite the MA (2005) recommending the implementation of a multisectoral approach to fully
288 evaluate changes in ES, their interactions, and the trade-offs and impact on people, 34% of the
289 62 papers analysed just one ES (i.e., 10 out of 35 papers for HS; 5 out of 31 for CR, and 3 out
290 of 39 for PP), and 23% analysed only two ES (Figure 4). Only 11% of the papers dealt
291 simultaneously with more than five ES.

292

293 **Insert Fig. 4**

294

295 The paper that dealt with one or a few ES turned out to be a very detailed analysis of the
296 single ES, and at the same time, they lost the overview of the system and the potential other
297 effects and trade-offs on the other ES. For example, Kimoto et al. (2012) analysed the effects
298 of different intensities of livestock grazing on native bees, and they concluded that maintaining
299 a heterogeneous landscape with some areas grazed and other not grazed, or with rotation of
300 grazing, might be necessary to support native bee diversity. However, the consequences on FP
301 and NH were not investigated by these authors.

302 In two interesting papers, Cole et al. (2012, 2015) analysed the effects of the main
303 physical and botanical attributes and of the different management options of riparian field
304 margins on ground beetle and pollinator diversity, and they concluded that wide riparian
305 margins strategically placed within the landscape can enhance taxonomic and functional
306 diversity. Nevertheless, this study did not analyse the effects on the landscape as cultural ES
307 (i.e., the aesthetic value) generated by the different management options, and so they missed
308 the opportunity to highlight further positive effects or trade-offs.

309 Another example is provided by Peringer et al. (2013), who analysed silvopastoral
310 systems as traditional components of the landscape in the Swiss Jura Mountains, for the
311 prevention of the loss of species-rich open grasslands and forest-grassland ecotones. In this
312 paper, the landscape was an assessment scale for the other ES (i.e., HS), and so it was not an
313 ES.

314 Other authors enlarged their analyses to other ES, to highlight potential trade-offs or
315 existing relationships; e.g., between different management options on FP or on the aesthetic
316 value of the landscape to produce income from tourism. In this vision, Fontana et al. (2014)
317 analysed the effects of management changes of larch grasslands in the Italian Alps
318 (abandonment and intensification *vs.* traditional management) on PP, HS and pollination, and
319 also on valuable cultural ES (i.e., scenic beauty, traditional healing plants). They conducted a
320 phyto-sociological study on plots that were randomly selected using geographic information
321 systems. For each plant species recorded, three out of eight plant traits were chosen explicitly
322 for their relevance for ES provision: flower colour, high diversity of pollination agents, and the
323 occurrence of edible or healing value for traditional meals and medicines. The provision of
324 scenic beauty and other ES was associated with specific management systems to be addressed
325 when planning future subsidies, and with specific financial support for a traditional agroforestry
326 system.

327 Other authors analysed the effects of several scenarios (e.g., climate change, policies,
328 management) on FP and on other ES for a more holistic analysis; e.g., Koniak et al. (2011)

329 addressed issues related to honey production, and developed a mathematical model that
330 predicted the dynamics of multiple services in response to management scenarios (grazing, fire,
331 and their combination) mediated by vegetation changes. These authors combined the potential
332 contribution to honey production with other ES from different groups into one “ES basket”
333 (e.g., carbon retention for CR, forage production for PP, density of geophytes for HS), despite
334 their different natures, which can help land managers to evaluate the effects and trade-offs of
335 alternative management scenarios. Another example of a holistic approach is provided by Dong
336 et al. (2012a, 2012b; 2014), who used the emergy¹ approach to calculate the performance of
337 several ES (i.e., CR, EE, FP, WR, PP) under different systems and scenarios, to support local
338 resource management and larger-scale environmental resource decision making. Ford et al.
339 (2012) used a wide range of ES for each of the MA category of ES to test the hypothesis that
340 changes in grazing intensity of semi-natural grassland differentially affect individual services
341 and alter the balance of supporting, provisioning, regulating and cultural ES provision. This
342 holistic approach underlined that in addition to biodiversity measures of “success” in
343 conservation, ES measures and trade-offs need to be taken into account when choosing an
344 appropriate grassland management scheme. Reed et al. (2015) analysed a combination of many
345 ES to produce tools and frameworks to support the stakeholder decision-making processes for
346 land management. These authors identified new economic instruments (e.g., payments for ES)
347 to enhance the flow of ES provided by grazing systems.

348

349 ***4.4. Ecosystem services, and different types of knowledge and decision making in the eligible*** 350 ***papers***

351 A further approach to the analysis of ES provided by grazing systems emerged from some
352 papers that included the involvement of stakeholders in different phases of the evaluation
353 process and with different aims. Some other authors applied a holistic approach that combined
354 the ES analysis with stakeholder involvement to explore the relationships between land
355 management and ES. This approach was intended to influence the decision-making processes,
356 to increase the stakeholder ES knowledge and awareness of the consequences of their activity.
357 Lamarque et al. (2014) applied a role game, in which farmers were faced with changes in ES
358 (i.e., PP, HS, WQ, CR) under climatic and socio-economic scenarios, and prompted to plan for
359 the future and to take land-management decisions as deemed necessary. The results

¹ Emergy was defined as the amount of available energy of one type (usually solar) that is directly or indirectly required to provide a given flow or storage of energy or matter (Odum, 1996).

360 demonstrated that the farmers were not aware, e.g., of the potential effects of their activities on
361 nitrate leaching, and that feedback loops between ES and land-management decisions can
362 favour more sustainable ES management. A global-scale study was performed by Petz et al.
363 (2014a) in South African rangelands that were affected by historical issues of land conservation
364 and degradation due to overgrazing (e.g., vegetation cover, species diversity, soil erosion,
365 carbon stock, water quality). These authors used the combined approach of a literature review,
366 collected data, and models (i.e., “IMAGE-USLE”) to study the interactions between input data,
367 livestock density, and ES, to strengthen and optimise the choices of local stakeholders for the
368 future management of the area in three different land-management scenarios. A further example
369 of the effectiveness of the use of this approach to identify the best land-management options
370 was provided by Fisher et al. (2011). These authors explored the variations in ES delivery that
371 resulted from different management practices in UK wetlands. In particular, the role of species-
372 led (both animals and plants) management on biodiversity was investigated. In a following step,
373 consultation with stakeholders and experts was carried out through workshops and meetings, to
374 elaborate specific details of the management impact on CR, WQ and WR, linked to the range
375 of management practices. These results are particularly relevant for the drafting of management
376 plans that need to carefully balance the effects of management practices. One example in this
377 sense was provided by Van Horn et al. (2012), who suggested taking into account grazing-
378 related effects on some ES, such as water-quality parameters like turbidity and temperature.

379 Other authors used different approaches for the analysis of ES, with the integration of
380 scientific knowledge with local knowledge, to create “hybrid knowledge”. In this vision, for a
381 pastoral system of a semi-arid region of northern Nicaragua, Tarrasón et al. (2016) highlighted
382 the importance of engaging relevant and interested stakeholders in dialogue with each other and
383 with the researchers, and encouraging the participation of local stakeholders in the decision-
384 making processes. They applied a participatory methodological framework to identify features
385 of LD and links with other ES provisions. The study designed a four-step methodological
386 framework to integrate local and scientific knowledge within a participatory assessment of land
387 degradation. Field visits and in-depth interviews with key informants and farmers produced
388 information that was integrated with the scientific knowledge that was validated by focus
389 groups, and then used in a state-and-transition conceptual model. Field data on the cover
390 vegetation and the plot life forms were used in thematic working groups with different
391 stakeholders to discuss the results of the previous phases and to develop adaptive management
392 options to maintain or improve ES.

393 The increase in awareness of local and general stakeholders (e.g., citizens, inhabitants,
394 tourists) of the flow of ES provided by grazing systems was considered by some authors as a
395 key element. The increased awareness of these stakeholders favours the acceptance of new
396 economic instruments (e.g., Payments for ES), which increased their “willingness to pay” for
397 ES. An example emerges from the analysis of Bernués et al. (2014), who attempted to determine
398 the socio-cultural and economic value of some ES delivered by mountain agroecosystems in
399 northeast Spain (e.g., forest fires, habitats for species, aesthetic and recreational values of the
400 landscape, product quality linked to the territory), by identifying stakeholder willingness to pay
401 for their provision. Focus groups and survey-based stated preference methods were combined
402 to identify the effects on ES of three different scenarios that were derived from contrasting
403 policies, and to test the willingness to pay for ES. Cultural ES were demonstrated to be a useful
404 tool to engage with stakeholders to support grazing system policies. From this analysis, it
405 emerged that the farmers were more interested in supporting ES, the local and general
406 stakeholders were more interested in cultural ES, and the local stakeholders were more
407 interested in the landscape than the general stakeholders. In any case, the willingness to pay for
408 ES was higher compared to the current level of EU agri-environmental support.

409

410 **5. Concluding remarks**

411

412 The extraction criteria used for this bibliographic review resulted in a relatively small number
413 of papers. The keyword “ecosystem service” was the dividing term between a vast literature
414 that deals with biophysical and socio-economic features of the grazing systems and the minimal
415 results of papers in this analysis that used the ES concept.

416 Although the MA has been the most widely accepted ES assessment framework since
417 2003, the analysis of these extracted papers has highlighted misunderstandings concerning the
418 concept of ES. One clear example is the confusion concerning biodiversity, which contrary to
419 the MA, was considered in several papers as an ES *per se* (e.g., Lindborg et al., 2009; Mace et
420 al., 2012). Also, not all of the analysed papers understood or accepted the anthropocentric vision
421 of the ES framework; e.g., some authors proposed biocentric solutions to reverse the inner
422 dynamics of systems without taking into account stakeholder opinions or needs (e.g., Cole et
423 al., 2015).

424 The need to examine the supply and condition of each ES, as well as the trade-offs (e.g.,
425 Marriot et al., 2010; Oñatibia et al., 2015) and interactions between them (as requested by the
426 MA), was applied in a number of these analysed papers (e.g., Koniak et al., 2011; Petz et al.,

2014a). Management and development options should take into account the internal dynamics of systems and the biophysical components, and also the socio-economic, socio-cultural and institutional features (Caballero and Fernández-Santos, 2009). Despite this, only a few authors integrated a multi-stakeholder approach into their analysis of ES and the interactions between these (e.g., Bernués et al., 2014; Petz et al., 2014b; Tarrasón et al., 2016). The need for stakeholder involvement emerged in some papers that underlined how the ES concept was not familiar to stakeholders, and was often confused, e.g., with the responsibility of humans to preserve nature (e.g., Bernués et al., 2014; Tarrasón et al., 2016). The use of ES as a basis for discussion might favour more sustainable practices, to increase the awareness of the effects of different management options on stakeholder well-being (e.g., Lamarque et al., 2014).

Other authors emphasised how the stakeholders and their knowledge inclusion is needed to improve the effectiveness of local decision-making processes (e.g., Lindborg et al., 2009; Tarrasón et al., 2016). The integration of local and scientific knowledge generates hybrid knowledge, thereby encouraging the participation of local stakeholders in the decision-making processes. This allowed the identification of adaptive strategies for key services to be maintained into the future (Lamarque et al., 2014; Francioni et al., 2014); e.g., through the implementation of *in-situ* experiments on native pasture management (Tarrasón et al., 2016). Many tools that are commonly used in scientific activities, such as mathematical models, future scenarios, indicators and biophysical data, were adopted by these authors to engage the stakeholders or to facilitate discussion with and between them.

In the analysed literature, cultural ES were poorly studied, despite these being considered the most relevant for local and general stakeholders (Bernués et al., 2014). This thus limited the ES framework to agriculture-related aspects. Better stakeholder awareness of the well-being provided by ES in grazing systems might foster agri-environmental schemes and the willingness to pay for these services. Many papers analysed and proposed different management options to improve the provision of ES (e.g., Cole et al., 2015), but did not analyse the effects on the natural heritage (e.g., the landscape aesthetic value), which can be relevant in policy-making processes (Bulte et al., 2008) and, for instance, in the definition of Payments for ES. Compensation and market-related policies have gained prominence to encourage farmers, policy makers and land managers to change their behaviour, and these might represent a mechanism to align potentially opposing interests; e.g., in the areas of wildlife management and biodiversity conservation.

459

460

461 **Acknowledgements**

462 The study was carried out with the support of the project MACSUR (D.M. 2660/7303/2012 -
463 www.MACSUR.eu), funded for the Italian partnership by the Italian Ministry of Agricultural,
464 Food and Forestry Policies. An earlier version of this study was presented at the 15th Meeting
465 of the Mediterranean Sub-Network of the Food and Agriculture Organisation–International
466 Centre for Advanced Mediterranean Agronomic Studies (FAO-CHIEAM), International
467 Network for the Research and Development of Pastures and Fodder Crops, “Ecosystem services
468 and socio-economic benefits of Mediterranean grasslands”, Orestiada (Greece), 12-14 April
469 2016.

470

471

472 **References**

- 473 Alcamo, J., Ash, N. J., Butler, C. D., Callicot, J. B., Capistrano, D., & Carpenter, S. R.
474 (2003). *Ecosystems and human well-being: a framework for assessment*. Washington,
475 DC: Island Press.
- 476 Allen, V. G., Batello, C., Berretta, E. J., Hodgson, J., Kothmann, M., Li, X., Mcivor, J.,
477 Milne, J., Morris, C., Peeters, A., & Sanderson, M. (2011). An international terminology
478 for grazing lands and grazing animals. *Grass and Forage Science*, 66, 2–28.
- 479 Bai, Y., Wu, J., Clark, C. M., Pan, Q., Zhang, L., Chen, S., Wang, Q., & Han, X. (2012).
480 Grazing alters ecosystem functioning and C: N: P stoichiometry of grasslands along a
481 regional precipitation gradient. *Journal of Applied Ecology*, 49, 1204–1215.
- 482 Bernués, A., Rodríguez-Ortega, T., Ripoll-Bosch, R., & Alfnes, F. (2014). Socio-cultural and
483 economic valuation of ecosystem services provided by Mediterranean mountain
484 agroecosystems. *PloS one*, 9, e102479.
- 485 Boughton, E. H., Bohlen, P. J., & Steele C. (2013). Season of fire and nutrient enrichment
486 affect plant community dynamics in subtropical semi-natural grasslands released from
487 agriculture. *Biological Conservation*, 158, 239–247.
- 488 Bulte, E., Boone, R. B., Stringer, R., & Thornton, P. K. (2008). Elephants or onions? Paying
489 for nature in Amboseli, Kenya. *Environment and Development Economics*, 13, 395–414.
- 490 Caballero, R., & Fernández-Santos, X. (2009). Grazing institutions in Castilla-La Mancha,
491 dynamic or downward trend in the Spanish cereal-sheep system. *Agricultural Systems*,
492 101, 69–79.
- 493 Cole, L. J., Brocklehurst, S., Elston, D. A., & McCracken, D. I. (2012). Riparian field
494 margins: can they enhance the functional structure of ground beetle (Coleoptera:
495 Carabidae) assemblages in intensively managed grassland landscapes? *Journal of*
496 *Applied Ecology*, 49, 1384–1395.
- 497 Cole, L. J., Brocklehurst, S., Robertson, D., Harrison, W., & McCracken, D. I. (2015).
498 Riparian buffer strips: Their role in the conservation of insect pollinators in intensive
499 grassland systems. *Agriculture, Ecosystems and Environment*, 211, 207–220.
- 500 Cousins, S. A., Auffret, A. G., Lindgren, J., & Tränk, L. (2015). Regional-scale land-cover
501 change during the 20th century and its consequences for biodiversity. *Ambio*, 44, 17–27.
- 502 de Groot, R. S., Wilson, M. A., & Boumans, R. M. (2002). A typology for the classification,
503 description and valuation of ecosystem functions, goods and services. *Ecological*
504 *Economics*, 41, 393–408.
- 505 Dong, X., Brown, M. T., Pfahler, D., Ingwersen, W. W., Kang, M., Jin, Y., Yu, B., Zhang, X.,
506 & Ulgiati, S. (2012a). Carbon modeling and emergy evaluation of grassland management
507 schemes in Inner Mongolia. *Agriculture, Ecosystems and Environment*, 158, 49–57.
- 508 Dong, X., Yang, W., Ulgiati, S., Yan, M., & Zhang, X. (2012b). The impact of human
509 activities on natural capital and ecosystem services of natural pastures in North Xinjiang,
510 China. *Ecological Modelling*, 225, 28–39.
- 511 Dong, X. B., Yu, B.H., Brown, M. T., Zhang, Y. S., Kang, M. Y., Jin, Y., Zhang, X. S., &
512 Ulgiati, S. (2014). Environmental and economic consequences of the overexploitation of
513 natural capital and ecosystem services in Xilinguole League, China. *Energy Policy*, 67,

- 514 767–780.
- 515 Duru, M., Jouany, C., Theau, J. P., Granger, S., & Cruz, P. (2015). A plant-functional-type
516 approach tailored for stakeholders involved in field studies to predict forage services and
517 plant biodiversity provided by grasslands. *Grass and Forage Science*, 70, 2–18.
- 518 Hoffman, I., From, T., & Boerma, D. (2014). Ecosystem services provided by livestock
519 species and breeds, with special consideration to the contributions of small-scale
520 livestock keepers and pastoralists. FAO Commission on genetic resources for food and
521 agriculture. Background study paper 66.
- 522 Feld, C. K., Martins da Silva, P., Paulo Sousa, J., De Bello, F., Bugter, R., Grandin, U.,
523 Hering, D., Lavorel, S., Mountford, O., Pardo, I., & Pärtel, M. I. (2009). Indicators of
524 biodiversity and ecosystem services: a synthesis across ecosystems and spatial scales.
525 *Oikos*, 118, 1862–1871.
- 526 Fischer, M., Bossdorf, O., Gockel, S., Hänsel, F., Hemp, A., Hessenmöller, D., Korte, G.,
527 Nieschulze, J., Pfeiffer, S., Prati, D., Renner, S., Schöning, I., Schumacher, U., Wells,
528 K., Buscot, F., Kalko, E. K. V., Linsenmair, K. E., Shulze, E.-D., & Weisser, W. W.
529 (2010). Implementing large-scale and long-term functional biodiversity research: The
530 Biodiversity Exploratories. *Basic and Applied Ecology*, 11, 473–485.
- 531 Fisher, B., Bradbury, R. B., Andrews, J. E., Ausden, M., Bentham-Green, S., White, S. M., &
532 Gill, J. A. (2011). Impacts of species-led conservation on ecosystem services of
533 wetlands: understanding co-benefits and tradeoffs. *Biodiversity and Conservation*, 20,
534 2461–2481.
- 535 Fisher, B., Turner, R. K., & Morling, P. (2009). Defining and classifying ecosystem services
536 for decision making. *Ecological Economics*, 68, 643–653.
- 537 Fontana, V., Radtke, A., Walde, J., Tasser, E., Wilhelm, T., Zerbe, S., & Tappeiner, U.
538 (2014). What plant traits tell us: Consequences of land-use change of a traditional agro-
539 forest system on biodiversity and ecosystem service provision. *Agriculture, Ecosystems
540 and Environment*, 186, 44–53.
- 541 Ford, H., Garbutt, A., Jones, D. L., & Jones, L. (2012). Impacts of grazing abandonment on
542 ecosystem service provision: Coastal grassland as a model system. *Agriculture,
543 Ecosystems and Environment*, 162, 108–115.
- 544 Francioni, M., Toderi, M., Catorci, A., Pancotto, D., & D'Ottavio, P. (2014). Agri-
545 environmental measures for the conservation of semi-natural grassland: a case of study
546 in Natura 2000 sites in Marche Region (Italy). *Options Méditerranéennes A*, 109, 655–
547 659.
- 548 Giese, M., Brueck, H., Gao, Y. Z., Lin, S., Steffens, M., Kögel-Knabner, I., Glindemann, T.,
549 Susenbeth, A., Taube, F., Butterbach-Bahl, K., Zheng, X. H., Hoffmann, C., Bai, Y. F.,
550 & Han, X. G. (2013). N balance and cycling of Inner Mongolia typical steppe: a
551 comprehensive case study of grazing effects. *Ecological Monographs*, 83, 195–219.
- 552 Hancock, M. H., & Legg, C. J. (2012). Diversity and stability of ericaceous shrub cover
553 during two disturbance experiments: one on heathland and one in forest. *Plant Ecology
554 and Diversity*, 5, 275–287.
- 555 Harrison, P. A., Berry, P. M., Simpson, G., Haslett, J. R., Blicharska, M., Bucur, M., Dunford,
556 R., Egoh, B., Garcia-Llorente, M., Geamăna, N., Geertsema, W., Lommelen, E.,

- 557 Meiresonne, L., & Turkelboom, F. (2014). Linkages between biodiversity attributes and
558 ecosystem services: a systematic review. *Ecosystem Services*, 9, 191–203.
- 559 Hussain, A. T., & Tschirhart, J. (2013). Economic/ecological tradeoffs among ecosystem
560 services and biodiversity conservation. *Ecological Economics*, 93, 116–127.
- 561 Jax, K., & Heink, U. (2015). Searching for the place of biodiversity in the ecosystem services
562 discourse. *Biological Conservation*, 191, 198–205.
- 563 Joubert, L., Pryke, J. S., & Samways, M. J. (2014). Annual burning drives plant communities
564 in remnant grassland ecological networks in an afforested landscape. *South African
565 Journal of Botany*, 92, 126–133.
- 566 Kearns, C. A., & Oliveras, D. M. (2009). Environmental factors affecting bee diversity in
567 urban and remote grassland plots in Boulder, Colorado. *Journal of Insect
568 Conservation*, 13, 655–665.
- 569 Kimoto, C., DeBano, S. J., Thorp, R. W., Taylor, R. V., Schmalz, H., DelCurto, T., Johnson,
570 T., Kennedy, P. L., & Rao, S. (2012). Short-term responses of native bees to livestock
571 and implications for managing ecosystem services in grasslands. *Ecosphere*, 3, 1–19.
- 572 Koniak, G., Noy-Meir, I., & Perevolotsky, A. (2011). Modelling dynamics of ecosystem
573 services basket in Mediterranean landscapes: a tool for rational management. *Landscape
574 Ecology*, 26, 109–124.
- 575 Lamarque, P., Meyfroidt, P., Nettièr, B., & Lavorel, S. (2014). How ecosystem services
576 knowledge and values influence farmers' decision-making. *PloS one*, 9, e107572.
- 577 Lavorel, S., Colloff, M. J., McIntyre, S., Doherty, M. D., Murphy, H. T., Metcalfe, D. J.,
578 Dunlop, M., Williams, R. J., Wise, R. M., & Williams, K. J. (2015). Ecological
579 mechanisms underpinning climate adaptation services. *Global Change Biology*, 21, 12–
580 31.
- 581 Lavorel, S., Grigulis, K., Lamarque, P., Colace, M. P., Garden, D., Girel, J., Pellet, G., &
582 Douzet, R. (2011). Using plant functional traits to understand the landscape distribution
583 of multiple ecosystem services. *Journal of Ecology*, 99, 135–147.
- 584 Leiber, F., Jouven, M., Martin, B., Priolo, A., Coppa, M., Prache, S., Heckendorn, F., &
585 Baumont, R. (2014). Potentials and challenges for sustainable grassland utilisation in
586 animal production. *Options Méditerranéennes A*, 109, 33–47.
- 587 Lindborg, R., Stenseke, M., Cousins, S. A., Bengtsson, J., Berg, Å., Gustafsson, T., Sjödin, N.
588 E., & Eriksson, O. (2009). Investigating biodiversity trajectories using scenarios-Lessons
589 from two contrasting agricultural landscapes. *Journal of Environmental Management*, 91,
590 499–508.
- 591 Littlewood, N. A., Stewart, A. J., & Woodcock, B. A. (2012). Science into practice—how can
592 fundamental science contribute to better management of grasslands for invertebrates?
593 *Insect Conservation and Diversity*, 5, 1–8.
- 594 Loucogouray, G., Dobremez, L., Gos, P., Pauthenet, Y., Nettièr, B., & Lavorel, S. (2015).
595 Assessing the effects of grassland management on forage production and environmental
596 quality to identify paths to ecological intensification in mountain grasslands.
597 *Environmental Management*, 56, 1039–1052.
- 598 Mace, G. M., Norris, K., & Fitter, A. H. (2012). Biodiversity and ecosystem services: a

- 599 multilayered relationship. *Trends in Ecology and Evolution*, 27, 19–26.
- 600 Marriott, C. A., Fisher, J. M., Hood, K., & Pakeman, R. J. (2010). Impacts of extensive
601 grazing and abandonment on grassland soils and productivity. *Agriculture, Ecosystems
602 and Environment*, 139, 476–482.
- 603 Medina-Roldán, E., Paz-Ferreiro, J., & Bardgett, R. D. (2012). Grazing exclusion affects soil
604 and plant communities, but has no impact on soil carbon storage in an upland grassland.
605 *Agriculture, Ecosystems and Environment*, 149, 118–123.
- 606 Millennium Ecosystem Assessment (2005). *Ecosystems and human well-being: synthesis*.
607 Washington, DC: Island Press.
- 608 Miller, M. E., Belote, R. T., Bowker, M. A., & Garman, S. L. (2011). Alternative states of a
609 semiarid grassland ecosystem: implications for ecosystem services. *Ecosphere*, 2, 1–18.
- 610 Odum, H. T. (1996). *Environmental Accounting: Emergy and Environmental Decision
611 Making*. New York: Wiley.
- 612 Oñatibia, G. R., Aguiar, M. R., & Semmartin, M. (2015). Are there any trade-offs between
613 forage provision and the ecosystem service of C and N storage in arid rangelands?
614 *Ecological Engineering*, 77, 26–32.
- 615 Peringer, A., Siehoff, S., Chételat, J., Spiegelberger, T., Buttler, A., & Gillet, F. (2013). Past
616 and future landscape dynamics in pasture-woodlands of the Swiss Jura Mountains under
617 climate change. *Ecology and Society*, 18, 11.
- 618 Petz, K., Alkemade, R., Bakkenes, M., Schulp, C. J., van der Velde, M., & Leemans, R.
619 (2014a). Mapping and modelling trade-offs and synergies between grazing intensity and
620 ecosystem services in rangelands using global-scale datasets and models. *Global
621 Environmental Change*, 29, 223–234.
- 622 Petz, K., Glenday, J., & Alkemade, R. (2014b). Land management implications for ecosystem
623 services in a South African rangeland. *Ecological Indicators*, 45, 692–703.
- 624 Porqueddu, C., Ates, S., Louhaichi, M., Kyriazopoulos, A. P., Moreno, G., del Pozo, A.,
625 Ovalle, C., Ewing M. A., & Nichols, P. G. H. (2016). Grasslands in ‘Old World’ and
626 ‘New World’ Mediterranean-climate zones: past trends, current status and future
627 research priorities. *Grass and Forage Science*, 71, 1–35.
- 628 Reed, M. S. (2008). Stakeholder participation for environmental management: a literature
629 review. *Biological Conservation*, 141, 2417–2431.
- 630 Reed, M. S., Stringer, L. C., Dougill, A. J., Perkins, J. S., Atlhopheng, J. R., Mulale, K., &
631 Favretto, N. (2015). Reorienting land degradation towards sustainable land management:
632 Linking sustainable livelihoods with ecosystem services in rangeland systems, *Journal of
633 Environmental Management*, 151, 472–485.
- 634 Roche, L. M., O'Geen, A. T., Latimer, A. M., & Eastburn, D. J. (2014). Montane meadow
635 hydrology, plant community, and herbivore dynamics. *Ecosphere*, 5, 1–16.
- 636 Sanderson, M. A., Goslee, S. C., Soder, K. J., Skinner, R. H., Tracy, B. F., & Deak, A.
637 (2007). Plant species diversity, ecosystem function, and pasture management - a
638 perspective. *Canadian Journal of Plant Science*, 87, 479–487.
- 639 Schaich, H., Kizos, T., Schneider, S., & Plieninger, T. (2015). Land change in eastern
640 Mediterranean wood-pasture landscapes: the case of deciduous oak woodlands in Lesvos

641 (Greece). *Environmental Management*, 56, 110–126.

642 Schaldach, R., Wimmer, F., Koch, J., Volland, J., Geißler, K., & Köchy, M. (2013). Model-
643 based analysis of the environmental impacts of grazing management on eastern
644 Mediterranean ecosystems in Jordan. *Journal of Environmental Management*, 127, S84–
645 S95.

646 Schröter, M., van der Zanden, E. H., van Oudenhoven, A. P. E., Remme, R. P., Serna-Chavez,
647 H. M., de Groot, R. S., & Opdam, P. (2014). Ecosystem Services as a contested concept:
648 a synthesis of critique and counter-arguments. *Conservation Letters*, 7, 514–523.

649 Sircely, J., & Naeem, S. (2012). Biodiversity and ecosystem multi-functionality: observed
650 relationships in smallholder fallows in Western Kenya. *PloS one*, 7, e50152.

651 Steiner, J. L., Engle, D. M., Xiao, X., Saleh, A., Tomlinson, P., Rice, C. W., Cole, N. A.,
652 Coleman, S. W., Osei, E., Basara, J., Middendorf, G., Gowda, P., Todd, R., Moffet, C.,
653 Anandhi, A., Starks, P. J., Ocshner, T., Reuter, R., & Devlin, D. (2014). Knowledge and
654 tools to enhance resilience of beef grazing systems for sustainable animal protein
655 production. *Annals of the New York Academy of Science*, 1328, 10–17.

656 Tarrasón, D., Ravera, F., Reed, M. S., Dougill, A. J., & Gonzalez, L. (2016). Land
657 degradation assessment through an ecosystem services lens: Integrating knowledge and
658 methods in pastoral semi-arid systems. *Journal of Arid Environments*, 124, 205–213.

659 TEEB (2010). *The economics of ecosystems and biodiversity. Ecological and economic*
660 *foundations*. Edited by Kumar, P. London and Washington: Earthscan.


661 Van Horn, D. J., White, C. S., Martinez, E. A., Hernandez, C., Merrill, J. P., Parmenter, R. R.,
662 & Dahm, C. N. (2012). Linkages between riparian characteristics, ungulate grazing, and
663 geomorphology and nutrient cycling in montane grassland streams. *Rangeland Ecology*
664 *and Management*, 65, 475–485.


665 van Wilgen, B. W., Reyers, B., Le Maitre, D. C., Richardson, D. M., & Schonegevel, L.
666 (2008). A biome-scale assessment of the impact of invasive alien plants on ecosystem
667 services in South Africa. *Journal of Environmental Management*, 89, 336–349.










668 von Döhren, P., & Haase, D. (2015). Ecosystem disservices research: A review of the state of
669 the art with a focus on cities. *Ecological Indicators*, 52, 490–497.

670 Zeng, C., Wu, J., & Zhang, X. (2015). Effects of grazing on above-vs. below-ground biomass
671 allocation of alpine grasslands on the Northern Tibetan Plateau. *PloS one*, 10, e0135173.

672





Primary production**Habitat services****Climate regulation****Extreme events****Food****Natural (landscape) heritage****Land degradation****Water flows****Water quality**




Table 1. Papers dealing with ecosystem services provided by grazing systems returned by the basic string from the Web of Science™ and after selection according to the review criteria. Each paper can deal with more than one ecosystem service.

| Ecosystem services group | Ecosystem service | Description | Papers | | | |
|--------------------------|--|--|-------------------------------|---|------|--|
| | | | Extracted ¹ (n) | Satisfying analysis criteria ² | | |
| | | | | (n) | (%) | |
| Supporting | Maintenance of soil structure and fertility | Nutrient cycling on farms and across landscapes; soil formation | 12 | n.a. | n.a. | |
| | Primary production | Improving vegetation growth/ cover | 72 | 39 | 63 | |
| | <i>Habitat services (as part of supporting services)</i> | | | | | |
| | Maintenance of life cycles of species | Habitat for species, especially migratory species | 78 | 35 | 56 | |
| | Habitat connectivity | Seed dispersal in guts and coats | 2 | n.a. | n.a. | |
| | Maintenance of genetic diversity | Gene pool protection and conservation | 0 | 0 | 0 | |
| Provisioning | Food | Meat, milk, eggs, honey, wool, leather, hides, skins, etc. | 12 | 6 | 10 | |
| | Fertiliser | Manure and urine for fertiliser | 9 | n.a. | n.a. | |
| | Fuel | Manure and CH ₄ for energy, manure biogas, etc. | 11 | n.a. | n.a. | |
| | Power | Draught animal power | 0 | 0 | 0 | |
| | Genetic resources | Basis for breed improvement and medicinal purposes | 10 | n.a. | n.a. | |
| | Biotechnical/ medicinal resources | Laboratory animals, test organisms, biochemical products | 0 | 0 | 0 | |
| Regulating | Waste recycling and conversion of non-human edible feed | Recycling of crop residues, household waste, swill, primary vegetation consumption | 1 | n.a. | n.a. | |
| | Land degradation and erosion prevention | Maintenance of vegetation cover | 26 | 10 | 16 | |
| | Water quality regulation/ purification | Water purification/ filtering in soils | 8 | 5 | 8 | |
| | Regulation of water flows | Natural drainage and drought prevention, influence of vegetation on rainfall, timing/ magnitude of run-off/ flooding | 44 | 15 | 24 | |
| | Climate regulation | Soil carbon sequestration, greenhouse gas mitigation | 60 | 31 | 50 | |

| | | | | | |
|-----------------|--|--|----|------|------|
| | Moderation of extreme events | Avalanche and fire control | 19 | 4 | 6 |
| | Pollination | Yield/ seed quality of crops and natural vegetation; genetic diversity | 17 | n.a. | n.a. |
| | Biological control and animal/ human disease control | Destruction of habitats of pest and disease vectors; yields | 3 | 0 | 0 |
| Cultural | Opportunities for recreation | Eco/ agro-tourism, sports, shows and other recreational activities involving specific animal breeds | 50 | n.a. | n.a. |
| | Knowledge systems and educational values | Traditional and formal knowledge about breeds, grazing and socio-cultural systems of the area | 23 | n.a. | n.a. |
| | Cultural and historic heritage | Presence of the breed in the area helps to maintain elements of the local culture that are valued as part of the local heritage; cultural identity | 21 | n.a. | n.a. |
| | Inspiration for culture, art and design | Traditional art/ handicraft; fashion; cultural, intellectual and spiritual enrichment and inspiration; pet animals, advertising | 12 | n.a. | n.a. |
| | Natural (landscape) heritage | Values associated with landscape as shaped by animals themselves or as a part of landscape; e.g., aesthetic values, sense of place, inspiration | 39 | 4 | 6 |
| | Spiritual and religious experience | Values related to religious rituals and the human life-cycle, such as religious ceremonies, funerals or weddings | 0 | 0 | 0 |

¹, 155 papers extracted from the Web of ScienceTM, for a total of 529 findings;

², 62 papers, for 149 findings, satisfying the analysis criteria.

n.a., not analysed

Table 2. Basic and additional strings used for the extraction of the papers, according to the keywords included in the Food and Agriculture Organisation report (Hoffman et al., 2014).

| Ecosystem service analysed | Extraction string |
|---|---|
| Ecosystem services (basic string) | <i>"ecosystem service*" and ("grassland*" or "rangeland*" or "shrubland*" or "scrubland*") and "grazing"</i> |
| Primary production | <i>("primary production" or "vegetation growth" or "vegetation cover" or "vegetation" or "NPP" or "net primary production")</i> |
| Habitat services | <i>("species" or "habitat" or "life cycle")</i> |
| Food and other livestock related products | <i>("meat" or "milk" or "honey" or "wool" or "leather" or "hide" or "skin" or "wax")</i> |
| Land degradation and soil erosion | <i>("land degradation" or "erosion" or "cover crop*" or "vegetation cover")</i> |
| Water quality regulation/ purification | <i>("water quality" or "water regulation" or "water purification" or "water filtering in soil")</i> |
| Regulation of water flows | <i>("water" or "natural drainage" or "drought prevention" or "runoff" or "rainfall" or "flooding")</i> |
| Climate regulation | <i>("climate" or "soil carbon" or "greenhouse gas*" or "GHG" or "CO₂" or "CH₄" or "N₂O")</i> |
| Moderation of extreme events | <i>("avalanche*" or "fire" or "extreme event*")</i> |
| Natural (landscape) heritage | <i>("landscape" or "aesthetic" or "inspiration")</i> |