

UNIVERSITÀ POLITECNICA DELLE MARCHE Repository ISTITUZIONALE

Cold brewing of rooibos tea affects its sensory profile and physicochemical properties compared to regular hot, and boiled brewing

This is the peer reviewd version of the followng article:

Original

Cold brewing of rooibos tea affects its sensory profile and physicochemical properties compared to regular hot, and boiled brewing / Muller, Magdalena; De Beer, Dalene; Truzzi, Cristina; Annibaldi, Anna; Carloni, Patricia; Girolametti, Federico; Damiani, Elisabetta; Joubert, Elizabeth. - In: LEBENSMITTEL-WISSENSCHAFT + TECHNOLOGIE. - ISSN 0023-6438. - ELETTRONICO. - 132:(2020). [10.1016/j.lwt.2020.109919]

Availability: This version is available at: 11566/283772 since: 2024-03-28T13:02:41Z

Publisher:

Published DOI:10.1016/j.lwt.2020.109919

Terms of use:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. The use of copyrighted works requires the consent of the rights' holder (author or publisher). Works made available under a Creative Commons license or a Publisher's custom-made license can be used according to the terms and conditions contained therein. See editor's website for further information and terms and conditions. This item was downloaded from IRIS Università Politecnica delle Marche (https://iris.univpm.it). When citing, please refer to the

This item was downloaded from IRIS Università Politecnica delle Marche (https://iris.univpm.it). When citing, please refer to the published version.

Cold brewing of rooibos tea affects its sensory profile and physicochemical properties compared to regular hot-, and boiled-brewing

Magdalena Muller^a, Dalene De Beer^{a,b}, Cristina Truzzi^c, Anna Annibaldi^c, Patricia Carloni^d, Federico Girolametti^c, Elisabetta Damiani^c, Elizabeth Joubert^{a,b,*}

^a Department of Food Science, Stellenbosch University, Private Bag X1, Matieland, Stellenbosch 7602, South Africa

^b Plant Bioactives Group, Post-Harvest and Agro-Processing Technologies, Agricultural Research Council (ARC) Infruitec-Nietvoorbij, Private Bag X5026, Stellenbosch 7599, South Africa

^c Department of Life and Environmental Sciences, Università Politecnica delle Marche, Via Brecce Bianche, I-60131 Ancona, Italy

^d Department of Agricultural, Food and Environmental Sciences, Università Politecnica delle Marche, Via Brecce Bianche, I-60131 Ancona, Italy

 * Correspondence to: Elizabeth Joubert, Plant Bioactives Group, Post-Harvest and Agro-Processing Technologies, Agricultural Research Council (ARC) Infruitec-Nietvoorbij, Private Bag X5026, Stellenbosch 7599, South Africa. Tel: +27218093416, E-mail: joubertL@arc.agric.za

1 ABSTRACT

2 Cold-brewing of rooibos tea has gained popularity, in particular in Japan, one of its major markets, due to the convenience of preparation. The sensory profile of traditional 3 4 ('fermented') rooibos, prepared by regular brewing (5 min infused in freshly-boiled 5 water) and served hot, is well described. However, no sensory profiling of cold-brewed 6 rooibos tea, rooibos tea served at ambient temperature or green rooibos tea has been 7 performed to date. In this study, the sensory profile of both fermented and green rooibos prepared by cold brewing (8 h at \leq 5 °C), regular brewing and boiling (5 min), and 8 consumed at ambient temperature (21 °C), were investigated. 'Rooibos woody' and 9 'fynbos-floral' aroma notes of fermented rooibos were unaffected by brewing 10 procedure, but its cold brew was sweeter and less astringent than the other brews. The 11 12 sensory profile of green rooibos, dominated by a 'hay/dried grass' aroma, bitter taste 13 and astringency, was affected to a greater extent by brewing procedure, but mostly, cold and regular brews had similar aroma attribute intensities. Higher levels of flavonoids 14 may explain the trends for astringency. Fe and Al were present in the highest levels in 15 boiled brews, although at levels too low to impact on health. 16

17

18 Keywords

19 Rooibos; cold brew; sensory profiling; flavonoids; aspalathin; Fe and Al content

20

21 **1.** Introduction

Sugar-sweetened beverages, including ready-to-drink teas, come at a price, not 22 23 only in terms of 'sugar-tax', but also the socio-economic burden of non-communicable 24 diseases. These beverages make a substantial contribution to sugar intake and have 25 consequently been linked to the onset of obesity, diabetes, and heart diseases (Benade & Essop, 2017). In response, the beverage industry has reformulated beverages with lower 26 27 sugar content and/or replaced sugar with non-nutritive sweeteners. A recent report predicted huge growth in the global sugar-free tea market by 2025 with the increasing 28 consumer awareness towards healthy drinks and a growing acceptance of natural herbs 29 30 identified as some of the market drivers (https://www.openpr.com/news/2015719/sugarfree-tea-market-to-see-huge-growth-by-2025-unilever). 31

32 Another market trend is the emergence of cold-brewed tea, which could boost the acceptance of sugar-free tea. Cold brewing delivers a beverage with less caffeine 33 and subsequently a reduced bitter taste as demonstrated for green tea (Lin, Liu, & Mau, 34 35 2008; Lin et al., 2014). A caffeine-free herbal tea that has grown in popularity on the global market is rooibos (Aspalathus linearis) (Joubert & De Beer, 2011). The main 36 product is 'fermented' (oxidized) rooibos with its natural, slightly sweet taste and mild 37 38 astringency. The characteristic sweet-associated aroma notes, 'caramel' and 'honey', of hot-brewed fermented rooibos (Jolley, Van der Rijst, Joubert, & Muller, 2017) could 39 40 even enhance the perception of sweetness through cross-modal aroma-taste interactions (Alcaire, Antúnez, Vidal, Giménez, & Ares, 2017). In contrast to fermented rooibos, 41 green (unoxidized) rooibos is a relatively new product, developed to capitalize on the 42 43 market opportunities for antioxidant-rich beverages. High levels of aspalathin, a potent

44 *C*-glucosyl dihydrochalcone antioxidant, can be found in the unoxidized plant material (Joubert & De Beer, 2011). Mounting evidence of aspalathin and green rooibos extracts 45 46 in the prevention of the metabolic syndrome (Johnson et al., 2018; Muller et al., 2018) merits the development of green rooibos as a functional beverage. However, a ready-to-47 drink beverage containing aspalathin-enriched green rooibos extract as an ingredient 48 49 was perceived as having an overt 'plant-like' character, disliked by consumers (Viljoen, Muller, De Beer, & Joubert, 2017). Studies on functional foods have demonstrated that 50 51 consumers are not inclined to compromise on flavor and taste for health (Verbeke, 2006). Changing the brewing procedure of green rooibos may, therefore, be merited to 52 improve flavor and taste. Brewing temperature and duration of green tea were found to 53 54 affect flavor (Lee & Chambers, 2009) and the extraction of health-promoting substances (Pastoriza, Pérez-Burillo, & Rufián-Henares, 2017). Cold-brewing of fermented rooibos 55 during summer has already gained acceptance in Japan, one of its main markets, due to 56 convenience. Adding a rooibos tea bag to water and infusing it for a prolonged period at 57 ambient temperature or in a refrigerator is a convenient alternative to steeping in hot 58 59 water or boiling on a stove. The latter practice was traditionally used to prepare rooibos tea (Joubert, Gelderblom, Louw, & De Beer, 2008). 60

Against this background, the aim of this study was to determine the effect of brewing procedure on the sensory profile and composition of both fermented rooibos and green rooibos. The sensory profile, color, and turbidity of cold-brewed rooibos were therefore compared with that of regular- and boiled-brewed rooibos. Color and turbidity are important parameters for visual appeal. The chemical composition of the beverages was quantified in terms of their *Z*-2-(β -D-glucopyranosyloxy)-3-phenylpropenoic acid

67 (PPAG), flavonoid, Al, and Fe content. PPAG was previously demonstrated to elicit a68 bitter taste (Joubert et al., 2013).

69

70 2. Experimental

71 2.1. Chemicals

72 Reagent-grade chemicals were obtained from Sigma-Aldrich (St. Louis, MO, 73 USA) or Merck (Darmstadt, Germany). Ultrapure water for HPLC and metal analysis 74 was prepared by a two-stage water purification system (Elix and Milli-Q) (Merck). Authentic standards (purity \geq 95%) were purchased from Extrasynthese (Genay, France; 75 76 isoorientin, orientin, 7-β-D-glucopyranosyleriodictyol (miscanthoside), isovitexin and hyperoside), Carl Roth (Karlsruhe, Germany; vitexin), Sigma-Aldrich (isoquercitrin, 77 78 ferulic acid), Transmit (Gießen, Germany; rutin), Phytolab (Vestenbergsgreuth, 79 Germany; aspalathin (91% purity)) and SAMRC (Cape Town, South Africa; nothofagin). PPAG was from our compound library (Plant Bioactives Group, ARC 80 Infruitec-Nietvoorbij). Stock solutions of the standards (ca. 1 g/L) were prepared in 81 DMSO and diluted with water as required. The diluted standard mixtures contained ca. 82 10 g/L ascorbic acid (Sigma-Aldrich) and were filtered through 0.45 µm hydrophilic 83 PVDF filters (Merck). 84 Working standard solutions for metal analysis were prepared by appropriate 85 86 dilution of inorganic atomic absorption standard solutions (1000 mg/L) of Fe (Carlo Erba, Milan, Italy) and Al (Sigma-Aldrich) and stored in a refrigerator at ca. 4 °C 87 protected from light. Ultrapure nitric acid (670-690 g/kg) was purchased from Carlo 88

89 Erba.

90 2.2. Rooibos samples

Samples from production batches (n=12) of both fermented rooibos and green
rooibos were sourced from Bokkeveld Rooibos (Nieuwoudtville, Northern Cape, South
Africa) and Rooibos Ltd. (Clanwilliam, Western Cape, South Africa), respectively.
Samples were mechanically sieved to remove dust (<0.42 mm) and coarse material
(>1.68 mm) (Jolley et al., 2017) and then stored at 21 °C in sealed glass jars.

96 2.3. Preparation of infusions

A cold brew (C), regular brew (R), and boiled brew (B) were prepared from each 97 batch of fermented and green rooibos by adding 12.5 g plant material to 1 L of distilled 98 99 water (equivalent to 'cup-of-tea' strength) as outlined in Fig. 1. Cold brewing entailed 100 adding water (21 °C) to the plant material in a wide-necked Schott bottle and steeping 101 for 8 h in a refrigerator (0-5 °C). For regular brew, freshly-boiled water was added to 102 the plant material in a glass jug followed by stirring for 5 s and steeping for 5 min (Koch, Muller, Joubert, Van der Rijst, & Næs, 2012). The boiled brew was prepared by 103 104 adding boiling water to the plant material in a glass jug followed by microwave heating 105 (1000 W) for 5 min with stirring every minute to prevent boiling over. All brews were strained through a fine-meshed strainer into 1 L Schott bottles directly after brewing and 106 107 stored overnight in a refrigerator (12 h). The brews were equilibrated to 21 °C prior to sensory analysis. 108

109 2.4. Descriptive sensory analysis

Panel training and testing were performed separately for fermented and green rooibos. Eleven assessors, with extensive experience in DSA of fermented rooibos, were trained according to a basic protocol (Koch et al., 2012) to generate appropriate aroma, flavor, taste, and mouthfeel attributes for green and fermented rooibos (supplementary material, Table A.1). Attribute intensities were scored, using an unstructured line scale (0 = not detected; 100 = high intensity), and the scores were captured electronically (Compusense®, Guelph, Canada). Three batches of plant material of each of the brew variants were analyzed per day in three consecutive sessions. The sample order was randomized per assessor per session. The blind-coded samples were served in black wine tasting glasses (ca. 70 mL per glass) covered with plastic lids. Analyses were conducted in individual tasting booths under standard lighting and controlled temperature (21 °C).

122 2.5. Physicochemical analyses

Turbidity, objective color, and soluble solids (SS) content of the brews were
determined in triplicate as described by Sishi, Muller, De Beer, Van der Rijst, & Joubert
(2019).

126 The concentration of the major flavonoids and PPAG in the brews was 127 determined in duplicate by high performance liquid chromatography with diode array detection (HPLC-DAD) using an Agilent 1200 HPLC system (Agilent Technologies, 128 129 Santa Clara, CA, USA). All compounds, except aspalathin and ferulic acid, were quantified using the validated method described by Walters, De Villiers, Joubert, & De 130 131 Beer (2017). Aspalathin and ferulic acid were quantified after separation on a Poroshell 120 EC-C₁₈ column (50×3 mm, 2.7 µm particle size; Agilent Technologies) at 25 °C. 132 The mobile phases (0.1% aqueous formic acid (A) and acetonitrile (B)) were run at 0.43 133 mL/min using the following gradient: 0–5 min, 12.4–14.5% B; 5–5.5 min, 14.5–80% B; 134 5.5-6.5 min, 80% B; 6.5-7 min, 80-12.4% B; 7-11 min, 12.4% B. UV-Vis spectra were 135 recorded between 200 and 450 nm with selected wavelength monitoring at 288 nm for 136 137 aspalathin and 320 nm for ferulic acid and the compounds quantified, using eight-point calibration curves. Peak identity was verified by performing the same separation on a 138

Waters Acquity UPLC coupled to a Synapt G2 quadrupole-time-of-flight (Q-TOF) MS
detector, equipped with an electrospray ionization (ESI) source (Waters, Milford, USA).
The mass spectrometer was operated in negative MS^E mode with a collision energy
ramp from 20–60 V. All other MS parameters were as described by Walters et al.
(2017).

144 2.6. Quantification of Fe and Al

Samples were prepared and analyzed in triplicate in a clean room laboratory ISO 145 146 14644–1 Class 6, with areas at ISO Class 5 under laminar flow. All the laboratory 147 materials were acid-cleaned as described by Illuminati, Annibaldi, Truzzi, & Scarponi (2014). An aliquot of each sample (5 mL) was put into Teflon vessels of a Microwave 148 Accelerated Reaction System, MARS-5, 1500 W (CEM, Mathews, NC, USA) and 149 150 digested (program reported in supplementary material, Table A.2) without any 151 pretreatment with 3 mL of HNO₃ (Annibaldi et al., 2019; Łozak, Sołtyk, Ostapczuk, & Fijałek, 2002). An HP-500 control vessel containing the same matrix of samples was 152 used to control the temperature and pressure during the process. Water (2 mL) was used 153 154 to clean up the vessels at the end of the mineralization process and added to the 155 samples, to reach a final volume of 10 mL. The quantitative determinations of Al and Fe 156 were carried out using a graphite furnace atomic absorption spectrophotometer (DUO 157 240FS AA-GTA120 Graphite Tube Atomizer, Agilent, Santa Clara, CA, USA), equipped with Zeeman background correction with argon (purity = 99.998%) as carrier 158 159 gas and multi-element hollow cathode lamps as light source. The instrumental parameters used for each element are reported in the supplementary material (Table 160 A.3). The quantitative values were calculated from calibration curves generated using 161 162 diluted standard solutions (20–80 μ g/L for Al and 10–50 μ g/L for Fe).

163	For assessment of the accuracy of the data, Al and Fe were determined using a
164	certified reference material (dogfish muscle DORM-2, NRCC; Ottawa, ON, Canada).
165	Certified mean values and experimental mean values obtained from the analysis of
166	DORM-2, expressed in mg/kg dry weight (d.w.), are as follows: Fe, Certified value =
167	142 ± 10 , Analytical result found = 132 ± 1 ; Al, Certified value = 10.9 ± 1.7 , Analytical
168	result found = 10.2 ± 0.8 . No statistically significant differences (P ≥ 0.05 , Student's T-
169	test) between certified and measured values were detected, and results were in good
170	agreement with the certified values, proving good accuracy of the analytical
171	methodology.
172	2.7. Statistical procedures

173 Data obtained for fermented and green rooibos were analyzed separately. DSA 174 data were subjected to various statistical techniques to confirm panel reliability (Næs, Brockhoff, & Tomic, 2010) and normality. Subsequent statistical analyses were 175 conducted on means over assessors. DSA and composition data were subjected to 176 177 analysis of variance (ANOVA) to test for treatment differences using SAS software (Version 9.2, SAS Institute Inc., Cary, USA). Fisher's least significant difference (LSD) 178 was calculated (5% level) to compare treatment means. P values < 0.05 were considered 179 180 significant.

Principal component analysis (PCA), using the correlation matrix, was performed, using XLStat (Version 2018, Addinsoft, Paris, France). Using the combined fermented and green rooibos data sets, the Pearson correlation coefficient (r) was calculated to determine the relationship between individual compound content and bitterness and astringency, respectively.

186 **3.** Results and discussion

187 Twelve batches of both fermented and green rooibos were subjected to three 188 different brewing procedures, i.e. cold (C), regular (R), and boiled (B), to determine the 189 impact on their sensory profiles. As the mean intensity scores for the aroma and flavor 190 attributes followed similar trends with flavor attributes mostly perceived at lower 191 intensities, the focus will fall on the aroma.

192 *3.1* Sensory profile

193 The PCA bi-plot for fermented rooibos (Fig. 2a) shows that only 52.7% of the 194 variance is explained. The regular- and boiled-brewed samples tend to associate on the 195 left of PC1, while the cold-brewed samples were positioned on the right, indicating that 196 the application of heat played a role in discriminating between treatments. On PC2, 197 samples largely separated between the boiled-brewed samples at the top and the cold-198 and regular-brewed samples at the bottom. The intensities of the primary aroma attributes ('rooibos-woody', 'fynbos-floral' and 'honey') of regular-brewed fermented 199 rooibos consumed hot (Jolley et al., 2017), as well as 'cooked apple', a secondary 200 201 aroma attribute of fermented rooibos, were not significantly ($P \ge 0.05$) affected by 202 treatment (Fig. 3a). The boiled-brewed samples associated with 'raisin' and 'caramel' (Fig. 2a), which were perceived at significantly higher (P<0.05) intensities in these 203 204 brews (Fig. 3a). The cold-brewed samples associated with 'tobacco' and sweet taste, but 205 not with astringency (Fig. 2a). The cold-brewed samples were significantly sweeter and less astringent (P<0.05) than the regular- and boiled-brewed samples (Fig. 3a), which 206 207 could contribute to consumer acceptance of cold-brewed rooibos. In view of the need to reduce the sugar content of beverages to limit sugar intake, the inherent slightly sweeter 208 taste of cold-brewed rooibos is beneficial. The 'tobacco' aroma note is generally 209

associated with the dry 'fermented' leaf product. Furthermore, 'seaweed', a negative
aroma note, was not perceptible in the cold- and boiled-brewed samples, and just
perceptible in the regular brews (P<0.05).

213 For green rooibos, PC1 and PC2 explained 77.6% of the variance and a clear distinction between the samples of the three brewing procedures is evident (Fig. 2b). On 214 215 PC1, the boiled-brewed samples clustered on the right, whereas the cold- and regular-216 brewed samples formed separate clusters on the left. On PC2, the separation between 217 samples was driven by brewing temperature with cold-brewed samples positioned at the 218 top and both regular- and boiled-brewed samples at the bottom. The cold brews had 219 significantly higher intensities (P<0.05) of the aroma notes 'fresh apple' and 'tobacco', 220 as well as sweet taste, compared to those of the other brew types (Fig. 3b). In contrast, 221 the boiled brews had significantly higher intensities of the attributes 'oats 222 porridge/cooked grains', 'caramel', 'stewed fruit', bitter taste and astringency (P<0.05) than the other two treatments, whereas the negative attributes, 'green grass' and 'putty', 223 were barely perceptible in the boiled brews (P<0.05) (Fig. 3b). The 'seaweed' aroma 224 225 note was prominent in the regular brews (intensity >15), barely perceptible in the cold brews (<5), and practically not perceptible in the boiled brews. The trend is the same as 226 227 for fermented rooibos, suggesting that the short heat exposure is required to form 228 volatile compound(s) responsible for this negative aroma note, while a more severe 229 application of heat resulted in sufficient loss of the formed volatile compound(s).

The fermented and green rooibos samples did not originate from the same batches of fresh plant material. However, the sensory profile of green rooibos was consistently and vastly different to that of fermented rooibos, especially in that the vegetal ('green grass', 'cooked vegetables') and cereal-like ('oats porridge/cooked grains') aroma attributes

234 were present in moderate to high intensities in green rooibos brews, depending on the 235 brewing procedure. While these aroma notes seem integral to the sensory profile of 236 green rooibos, they are considered to be negative for fermented rooibos (Jolley et al., 2017). The 'hay/dried grass' intensity was not affected by the brewing procedure. A 237 short steam treatment of the dry fermented rooibos leaves also did not affect the 'hay-238 239 like' intensity of rooibos infusions (regular-brewed; consumed hot) (Koch, Muller, De 240 Beer, Næs, & Joubert, 2013). Fermented rooibos brews had a sweeter taste, while the 241 green rooibos brews were more astringent with a prominent bitter taste.

242 3.2 SS content, color and turbidity

243 The SS content and turbidity of the fermented rooibos brews were significantly 244 (P<0.05) affected by brewing procedure, increasing progressively from cold to regular 245 to boiled brew (Table 1). The CIELab color parameters, lightness (L*), and hue (h) decreased and redness (a*) increased significantly (P<0.05) (Table 1) with the increase 246 in the soluble solids content of the brew. The yellowness parameter, b*, and 247 248 subsequently also chroma of the cold brews were significantly (P<0.05) lower than those for the regular and boiled brews. Whilst the color parameters indicate differences 249 250 between the brews, the values are difficult to relate to the actual visual color in a cup or 251 glass due to dichroism, i.e. the hue or color depends on the degree of dilution or container size (Francis & Clydesdale, 1975). 252

The different procedures used for the brewing of green rooibos had similar effects to fermented rooibos on the soluble solids content, turbidity, L*, a* and hue of the brews (P<0.05) (Table 2). For all these parameters, except a*, higher values were obtained for the green rooibos brews than for the fermented rooibos brews (Tables 1 and 257 2). The lower a* values of green rooibos brews (1.0-14.3 vs 16.01-30.9) were to be
258 expected as the plant material is not oxidized.

259 3.3 PPAG and flavonoid content

260 The PPAG concentration in the fermented rooibos brews was slightly increased 261 (P<0.05) by boiling compared to cold and regular brewing (Table 1; representative 262 chromatograms shown in Fig. A.1). The concentration of most of the flavonoids increased significantly (P<0.05) when heat was applied during the brewing of fermented 263 264 rooibos, with the boiling procedure being most effective in extracting the compounds 265 from the plant material (Table 1). The aspalathin, bioquercetin, hyperoside, and isoquercitrin concentration of the regular and boiled brews did not differ significantly 266 267 (P≥0.05).

268 For the green rooibos brews, most phenolic compounds showed a significantly 269 (P<0.05) higher concentration in the boiled brew compared to the other brews (Table 2; 270 representative chromatograms shown in Fig. A.2). Some exceptions were PPAG, 271 bioquercetin, and rutin with the lowest (P < 0.05) concentration in the regular brew. The PPAG concentration of the cold and boiled brews was not significantly (P≥0.05) 272 273 different, but the bioquercetin and rutin concentrations were significantly (P<0.05) higher in the boiled brew than in the cold brew. Our previous study on the same sample 274 275 set provided more insight into the phenolic profile of the brews. A large number of compounds were putatively annotated (Damiani et al., 2019). Of 187 putatively 276 277 annotated compounds, 32 and 35 polyphenols were considered as markers that 278 discriminate between cold and boiled brews, and cold and regular brews, respectively.

The investigation of the association between the flavonoid content of rooibos infusions and the intensities of the taste modalities and astringency (when served hot) is 281 limited (Koch et al., 2013). The association between PPAG and sweet taste was notable, 282 however, when tested at 21 °C as a single compound, dissolved in water, it was bitter at 283 a concentration similar to that of regular-brewed rooibos. The PPAG concentration of 284 the brews analyzed for the present study (Tables 1 and 2) was much higher than its 285 detection threshold in water (0.4 mg/L) (Joubert et al., 2013). PPAG could, therefore, be 286 expected to contribute to the bitter taste. Considering that the green rooibos brews were 287 substantially more bitter than the fermented rooibos brews, while their respective PPAG 288 content did not differ to the same extent, other compounds or interaction with other compounds (i.e. masking and/or enhancing effects) contributed to bitterness. A 289 290 moderate correlation was found between the content of the individual flavonoids and the 291 bitter intensity of the brews ($r \ge 0.5$; $P \le 0.002$). The eriodictyol glucosides had the highest 292 correlation with bitter intensity (r>0.65; P<0.0001).

It is well-known that phenolic compounds contribute to the astringency of food, wine, and tea (Bordenave, Hamaker, & Ferruzzi, 2014). Despite the green rooibos brews having a substantially higher flavonoid content than the fermented rooibos brews, the green rooibos brews were not substantially more astringent (Tables 1 and 2; Fig. 3). A moderate, but significant correlation was found between the individual flavonoid content of the brews and astringency ($r \ge 0.41$; P<0.01). PPAG did not correlate significantly with astringency (r = 0.325; P=0.053).

300 3.4 Al and Fe content

Al and Fe were present in significantly higher (P<0.05) levels in the boiled brews of both fermented and green rooibos (Tables 1 and 2), while no significant differences (P \ge 0.05) between the cold and regular brews were found, except in the case of the Fe content of fermented rooibos brews. In this case, the Fe content increased
significantly (P<0.05) from cold to regular to boiled brew.

306 The Fe and Al content values were similar to those previously reported for hot water infusions of fermented rooibos (Joubert et al., 2008; Malik, Szakova, Drabek, 307 Balik, & Kokoska, 2008). Consumption of one cup of the brews would contribute 308 309 <0.2% of the nutrient reference value (NRV) for adults, i.e. 18 mg/day, according to 310 South African food labeling regulations. Chronic Al intake, on the other hand, has 311 negative health implications (Pérez-Granados & Vaquero, 2002). A tolerable weekly 312 intake (TWI) of Al is 1 mg/kg body weight/week (Aguilar et al., 2008), i.e. 10 mg/day 313 by a 70 kg adult. One cup of rooibos per day would contribute <0.3% of this intake.

314 3.5 "To cold-brew or not to cold-brew"

315 Boiling increased the flavonoid content of rooibos tea, irrespective of the type. 316 Similarly, in our previous study boiling also delivered brews with the highest 317 antioxidant capacities (ORAC and DPPH) (Damiani et al., 2019). However, it is not clear whether these increases would translate into a specific health benefit. No 318 recommended daily intake values have been established for flavonoids, as this group of 319 320 phytochemicals differs vastly in terms of bioavailability and bioactivity, while the value of antioxidant capacity in the assessment of health promotion is the subject of debate 321 322 (De Camargo et al., 2019; Williamson, Kay, & Crozier, 2018).

The flavor of beverage products is regarded as one of the main drivers of consumer preference. The sensory profile associated with the respective rooibos brews would be the deciding factor if convenience is not considered important. Cold brewing of fermented rooibos delivered a similar primary and secondary sensory profile as regular brewing, thus cold brewing offers no advantage or disadvantage in terms of its 328 sensory profile. The slightly higher intensities of 'raisin' and 'caramel' aroma notes of 329 boiled-brewed rooibos may be one of the reasons why rooibos was traditionally brewed 330 on a hot stove for extended periods. The 'caramel' note was also more prominent in 331 boiled-brewed green rooibos. The intensity of the 'oats porridge/cooked grains' note, not present in fermented rooibos, was substantially increased with the boiling of green 332 333 rooibos. While the brewing procedure did not impact on astringency, boiling of green 334 rooibos increased the bitterness of the tea, linked to increased extraction of flavonoids. 335 To date, no study has been conducted to determine whether rooibos consumers are 336 willing to forfeit taste for health. Even the Fe and Al content of a cup of rooibos tea, 337 irrespective of type or brewing procedure, was found to be too low to impact, 338 respectively, positively and negatively on health and thus would not drive choice.

339 4. Conclusions

Cold brewing of both fermented rooibos and green rooibos did not offer many advantages in terms of flavonoid content and sensory profile of the brew compared to regular brewing. Notable was a less bitter and/or sweeter taste of the cold brew compared to the boiled brew.

344 Acknowledgments

345 Clinton Gass, Rooibos Marketing Ltd, Japan, and Prof. Yutaka Miura, Tokyo

346 University of Agriculture and Technology, Fuchu, Japan, are thanked for information

347 on the preparation of rooibos tea by Japanese consumers. This work was supported by

the National Research Foundation of South Africa (NRF grants 119193 and 112136,

- awarded to EJ and DdB, respectively) and the Northern Cape Department of
- 350 Agriculture, Land Reform and Rural Development (Contract NC/DALC/0679). The
- 351 funding bodies had no involvement in: study design; collection, analysis, and

- interpretation of data; writing of the manuscript; or decision to publish the work. ED,
- 353 PC, CT, AA, and FG acknowledge funding from the Polytechnic University of Marche.

354 Conflict of interest

355 The authors declare no conflict of interest.

356 **References**

- 357 Aguilar, F., Autrup, H., Barlow, S., Castle, L., Crebelli, R., Dekant, W., Gontard, N.,
- 358 Gott, D., Grilli, S., Leclercq, C., Pratt, I., Rietjens, I., Tobback, P., & Aids, P.
- 359 (2008). Safety of aluminium from dietary intake Scientific Opinion of the Panel
- 360 on Food Additives, Flavourings, Processing Aids and Food Contact Materials
- 361 (AFC). *The EFSA Journal*, *754*, 1–34.
- Alcaire, F., Antúnez, L., Vidal, L., Giménez, A., & Ares, G. (2017). Aroma-related

363 cross-modal interactions for sugar reduction in milk desserts: Influence on
364 consumer perception. *Food Research International*, *97*, 45–50.

- 365 Annibaldi, A., Truzzi, C., Carnevali, O., Pignalosa, P., Api, M., Scarponi, G., &
- 366 Illuminati, S. (2019). Determination of Hg in farmed and wild Atlantic Bluefin
- tuna (*Thunnus thynnus* L.) muscle. *Molecules*, 24, 1273.
- Benade, J., & Essop, M. F. (2017). Introduction of "Sugar Tax" in South Africa:
- Placebo or panacea to curb the onset of cardio-metabolic diseases? *SAHeart*, *14*,
 148–153.
- Bordenave, N., Hamaker, B. R., & Ferruzzi, M. G. (2014). Nature and consequences of
 non-covalent interactions between flavonoids and macronutrients in foods. *Food and Function*, *5*, 18–34.

374	Damiani, E., Carloni, P., Rocchetti, G., Senizza, B., Tiano, L., Joubert, E., De Beer, D.,			
375	& Lucini, L. (2019). Impact of cold versus hot brewing on the phenolic profile and			
376	antioxidant capacity of rooibos (Aspalathus linearis) herbal tea. Antioxidants, 8,			
377	499.			
378	De Camargo, A. C., Biasoto, A. C. T., Schwember, A. R., Granato, D., Rasera, G. B.,			
379	Franchin, M., Rosalen, P. L., Alencar, S. M., & Shahidi, F. (2019). Should we ban			
380	total phenolics and antioxidant screening methods? The link between antioxidant			
381	potential and activation of NF-KB using phenolic compounds from grape by-			
382	products. Food Chemistry, 290, 229-238.			
383	Francis, F. J., & Clydesdale, F. M. (1975). Food Colorimetry: Theory and Applications.			
384	Westport, USA: Avi Publishing Company.			
385	Illuminati, S., Annibaldi, A., Truzzi, C., & Scarponi, G. (2014). Recent temporal			
386	variations of trace metal content in an Italian white wine. Food Chemistry, 159,			
387	493–497.			
388	Johnson, R., De Beer, D., Dludla, P., Ferreira, D., Muller, C. J. F., & Joubert, E. (2018).			
389	Aspalathin from rooibos (Aspalathus linearis): A bioactive C-glucosyl			
390	dihydrochalcone with potential to target the metabolic syndrome. Planta Medica,			
391	84, 568–583.			
392	Jolley, B., Van der Rijst, M., Joubert, E., & Muller, M. (2017). Sensory profile of			
393	rooibos originating from the Western and Northern Cape governed by production			
394	year and development of rooibos aroma wheel. South African Journal of Botany,			
395	110, 161–166.			
396	Joubert, E., & De Beer, D. (2011). Rooibos (Aspalathus linearis) beyond the farm gate:			

- From herbal tea to potential phytopharmaceutical. *South African Journal of Botany*, 77, 869–886.
- Joubert, E., De Beer, D., Malherbe, C. J., Muller, N., Bonnet, S. L., Van der
- 400 Westhuizen, J. H., & Ferreira, D. (2013). Occurrence and sensory perception of Z-
- 401 2-(β-D-glucopyranosyloxy)-3-phenylpropenoic acid in rooibos (*Aspalathus*
- 402 *linearis*). *Food Chemistry*, *136*, 1078–1085.
- 403 Joubert, E., Gelderblom, W. C. A., Louw, A., & De Beer, D. (2008). South African
- 404 herbal teas: *Aspalathus linearis*, *Cyclopia* spp. and *Athrixia phylicoides*—A
 405 review. *Journal of Ethnopharmacology*, *119*, 376–412.
- 406 Koch, I. S., Muller, M., Joubert, E., Van der Rijst, M., & Næs, T. (2012). Sensory
- 407 characterization of rooibos tea and the development of a rooibos sensory wheel and
 408 lexicon. *Food Research International*, 46, 217–228.
- Koch, I. S., Muller, N., De Beer, D., Næs, T., & Joubert, E. (2013). Impact of steam
- 410 pasteurization on the sensory profile and phenolic composition of rooibos
- 411 (Aspalathus linearis) herbal tea infusions. Food Research International, 53, 704–
- 412 712.
- Lee, J., & Chambers, D. H. (2009). Sensory descriptive evaluation: brewing methods
 affect flavour of green tea. *Asian Journal of Food and Agro-Industry*, 2, 427–439.
- Lin, S.-D., Liu, E.-H., & Mau, J.-L. (2008). Effect of different brewing methods on
- antioxidant properties of steaming green tea. *LWT Food Science and Technology*,
 417 41, 1616–1623.
- Lin, S.-D., Yang, J.-H., Hsieh, Y.-J., Liu, E.-H., & Mau, J.-L. (2014). Effect of different

- 419 brewing methods on quality of green tea. *Journal of Food Processing and*
- 420 *Preservation*, *38*, 1234–1243.
- 421 Łozak, A., Sołtyk, K., Ostapczuk, P., & Fijałek, Z. (2002). Determination of selected
- trace elements in herbs and their infusions. *Science of the Total Environment*, 289,
 33–40.
- Malik, J., Szakova, J., Drabek, O., Balik, J., & Kokoska, L. (2008). Determination of
 certain micro and macroelements in plant stimulants and their infusions. *Food Chemistry*, *111*, 520–525.
- 427 Muller, C. J. F., Malherbe, C. J., Chellan, N., Yagasaki, K., Miura, Y., & Joubert, E.
- 428 (2018). Potential of rooibos, its major C-glucosyl flavonoids, and Z-2-(β -D-
- 429 glucopyranosyloxy)-3-phenylpropenoic acid in prevention of metabolic syndrome.

430 *Critical Reviews in Food Science and Nutrition*, 58, 227–246.

- 431 Næs, T., Brockhoff, P., & Tomic, O. (2010). *Statistics for Sensory and Consumer*432 *Science*. Wiley: New York, USA.
- 433 Pastoriza, S., Pérez-Burillo, S., & Rufián-Henares, J. Á. (2017). How brewing
- 434 parameters affect the healthy profile of tea. *Current Opinion in Food Science*, *14*,
 435 7–12.
- 436 Pérez-Granados, A. M., & Vaquero, M. P. (2002). Silicon, aluminium, arsenic and
- 437 lithium: Essentiality and human health implications. *Journal of Nutrition, Health*438 *and Aging*, *6*, 154–162.
- 439 Sishi, M., Muller, M., De Beer, D., Van der Rijst, M., & Joubert, E. (2019). Rooibos
 440 agro-processing waste as herbal tea products: optimisation of soluble solids

441	extraction from dust and application to improve sensory profile, colour and
442	flavonoid content of stem infusions. Journal of the Science of Food and
443	Agriculture, 99, 3653–3661.
444	Verbeke, W. (2006). Functional foods: Consumer willingness to compromise on taste
445	for health? Food Quality and Preference, 17, 126–131.
446	Viljoen, M., Muller, M., De Beer, D., & Joubert, E. (2017). Identification of broad-
447	based sensory attributes driving consumer preference of ready-to-drink rooibos
448	iced tea with increased aspalathin content. South African Journal of Botany, 110,
449	177–183.
450	Walters, N. A., De Villiers, A., Joubert, E., & De Beer, D. (2017). Improved HPLC
451	method for rooibos phenolics targeting changes due to fermentation. Journal of
452	Food Composition and Analysis, 55, 20–29.
453	Williamson, G., Kay, C. D., & Crozier, A. (2018). The bioavailability, transport, and
454	bioactivity of dietary flavonoids: A review from a historical perspective.
455	Comprehensive Reviews in Food Science and Food Safety, 17, 1054–1112.

458

459 **Figure captions**

- 460 Fig. 1. Basic procedure used for the preparation of cold-, regular-, and boiled-brewed
- 461 fermented and green rooibos herbal tea.
- 462 Fig. 2. PCA bi-plot of the sensory profile of cold-, regular-, and boiled-brewed
- 463 fermented (a) and green (b) rooibos herbal tea with brewing procedure indicated by C,
- 464 R, and B, respectively. The numbers (1 to 12) refer to the individual samples,
- 465 representing independent replicates.
- 466 Fig. 3. ANOVA results of selected sensory attributes of cold-, regular-, and boiled-
- 467 brewed fermented (a) and green (b) rooibos herbal tea. Bars colored white, grey and
- 468 black represent cold-, regular-, and boiled-brewed rooibos, indicating mean values and
- 469 error bars (standard deviation).
- 470





Figure 2



Figure 3

Table 1

Parameter	Cold brew	Regular brew	Boiled brew	LSD
L*	81 ± 2 a	72 ± 2 b	$64 \pm 3 c$	1
a*	$16 \pm 3 c$	$26 \pm 2 b$	31 ± 2 a	1
b*	93 ± 5 b	97 ± 3 a	95 ± 3 a	2
С	94 ± 5 b	101 ± 3 a	100 ± 3 a	2
h	80 ± 1 a	75 ± 1 b	$72 \pm 1 c$	1
Turbidity	$5 \pm 1 c$	22 ± 7 b	42 ± 14 a	6
Soluble solids	$1.0 \pm 0.1 \text{ c}$	$1.5 \pm 0.2 \text{ b}$	1.9 ± 0.2 a	0.1
SE6G ^a	$7.3 \pm 0.7 \text{ c}$	$10.8 \pm 1.0 \text{ b}$	14.0 ± 1.4 a	0.5
RE6G ^b	$6.9 \pm 0.8 \text{ c}$	$10.5 \pm 1.0 \text{ b}$	13.8 ± 1.5 a	0.5
SE8G ^c	2.0 ± 0.3 c	3.0 ± 0.4 b	4.3 ± 0.7 a	0.2
RE8G ^d	2.5 ± 0.2 c	3.4 ± 0.3 b	4.7 ± 0.5 a	0.2
Aspalathin	3 ± 2 b	5 ± 4 a	5 ± 3 a	1
PPAG ^e	9 ± 1 b	8 ± 1 b	10 ± 2 a	0.4
Nothofagin	$0.3 \pm 0.2 \text{ c}$	$0.6 \pm 0.3 \text{ b}$	0.9 ± 0.4 a	0.1
Isoorientin	$4.4 \pm 0.6 \text{ c}$	11.3 ± 1.4 b	15.9 ± 1.3 a	0.6
Orientin	8 ± 1 c	$18 \pm 2 b$	25 ± 2 a	1
Ferulic acid	1.2 ± 0.4 c	1.3 ± 0.3 b	1.6 ± 0.4 a	0.1
Bioquercetin	5 ± 3 b	9 ± 4 a	$10 \pm 3 a$	1
Vitexin	$1.3 \pm 0.1 \text{ c}$	3.1 ± 0.3 b	4.1 ± 0.3 a	0.2
Hyperoside	$0.9 \pm 0.9 \text{ b}$	2.4 ± 1.8 a	2.4 ± 0.9 a	0.5
Rutin	0.8 ± 0.6 c	1.5 ± 1.0 b	2.1 ± 1.4 a	0.5
Isovitexin	$1.3 \pm 0.1 \text{ c}$	3.2 ± 0.3 b	4.4 ± 0.3 a	0.2
Isoquercitrin	$0.3 \pm 0.4 \text{ b}$	1.8 ± 1.4 a	1.9 ± 1.0 a	0.6
Luteoleoside	0.02 ± 0.04 c	0.51 ± 0.18 b	0.94 ± 0.22 a	0.12
Sum of	45	85	111	-
flavonoids				_
Fe	$105 \pm 13 c$	$119 \pm 14 \text{ b}$	$142 \pm 5 a$	7
Al	$42 \pm 22 \text{ b}$	57 ± 21 b	108 ± 35 a	20

Colour, turbidity (NTU), soluble solids content (SS, g/L) and chemical composition (mg/L, except Fe and Al, μ g/L) of cold brew, regular brew and boiled brew fermented rooibos.

^a (*S*)-6-β-D-glucopyranosyleriodictyol.

^b (*R*)-6- β -D-glucopyranosyleriodictyol.

^c (*S*)-8-β-D-glucopyranosyleriodictyol.

^d (*R*)-8- β -D-glucopyranosyleriodictyol.

^e Z-2-(β-D-glucopyranosyloxy)-3-phenylpropenoic acid.

Table 2

Parameter	Cold brew	Regular brew	Boiled brew	LSD
L*	93 ± 1 a	88 ± 3 b	80 ± 2 c	1
a*	1 ± 1 c	$6 \pm 2 b$	$14 \pm 2 a$	1
b*	$40 \pm 7 b$	$35 \pm 6 c$	$46 \pm 5 a$	3
С	$40 \pm 7 b$	$35 \pm 7 c$	49 ± 5 a	3
h	89 ± 1 a	$80 \pm 2 b$	$73 \pm 1 c$	1
Turbidity	9 ± 3 c	$33 \pm 10 \text{ b}$	60 ± 9 a	5
Soluble solids	$1.9 \pm 0.2 \text{ b}$	$1.7 \pm 0.4 \text{ b}$	2.5 ± 0.4 a	0.2
Aspalathin	189 ± 27 b	176 ± 48 b	267 ± 40 a	17
PPAG ^a	13 ± 2 a	9 ± 2 b	13 ± 1 a	1
Nothofagin	19 ± 4 b	18 ± 6 b	28 ± 6 a	2
Isoorientin	$13 \pm 2 b$	$13 \pm 3 b$	21 ± 3 a	1
Orientin	$19 \pm 2 \mathrm{b}$	$19 \pm 4 b$	30 ± 4 a	2
Bioquercetin	18 ± 5 b	$14 \pm 5 c$	21 ± 6 a	1
Vitexin	3.0 ± 0.4 b	$2.8 \pm 0.7 \text{ b}$	4.5 ± 0.6 a	0.3
Hyperoside	5 ± 2 b	$5 \pm 2 b$	7 ± 3 a	1
Rutin	8 ± 2 b	$6 \pm 2 c$	10 ± 2 a	1
Isovitexin	3.6 ± 0.4 b	$3.7 \pm 0.9 \text{ b}$	5.7 ± 0.8 a	0.3
Isoquercitrin	8 ± 3 b	8 ± 4 b	13 ± 5 a	1
Luteoleoside	$0.7 \pm 0.3 c$	$1.1 \pm 0.4 \text{ b}$	1.9 ± 0.6 a	0.2
Sum of	286	267	409	-
flavonoids				
Fe	21±8 b	21 ± 7 b	53 ± 14 a	6
Al	82 ± 42 b	81 ± 38 b	149 ± 45 a	32

Colour, turbidity (NTU), soluble solids content (SS, g/L) and chemical composition (mg/L, except Fe and Al, μ g/L) of cold brew, regular brew and boiled brew green rooibos.

^a Z-2-(β -D-glucopyranosyloxy)-3-phenylpropenoic acid.

