



UNIVERSITÀ POLITECNICA DELLE MARCHE
Repository ISTITUZIONALE

Corrigendum to the paper: A way to model stochastic perturbations in population dynamics models with bounded realizations. *Commun Nonlinear Sci Numer Simulat*, 77 (2019), 239–257

This is the peer reviewed version of the following article:

Original

Corrigendum to the paper: A way to model stochastic perturbations in population dynamics models with bounded realizations. *Commun Nonlinear Sci Numer Simulat*, 77 (2019), 239–257 / Caraballo, T.; Colucci, R.; Lopez-de-la-Cruz, J.; Rapaport, A.. - In: COMMUNICATIONS IN NONLINEAR SCIENCE & NUMERICAL SIMULATION. - ISSN 1007-5704. - STAMPA. - 96:(2021). [10.1016/j.cnsns.2020.105681]

Availability:

This version is available at: 11566/290151 since: 2024-07-09T08:50:13Z

Publisher:

Published

DOI:10.1016/j.cnsns.2020.105681

Terms of use:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. The use of copyrighted works requires the consent of the rights' holder (author or publisher). Works made available under a Creative Commons license or a Publisher's custom-made license can be used according to the terms and conditions contained therein. See editor's website for further information and terms and conditions.

This item was downloaded from IRIS Università Politecnica delle Marche (<https://iris.univpm.it>). When citing, please refer to the published version.

note finali coverpage

(Article begins on next page)

Corrigendum to the paper: A way to model stochastic perturbations in population dynamics models with bounded realizations

Commun Nonlinear Sci Numer Simulat, **77** (2019), 239–257

TOMÁS CARABALLO

Departamento de Ecuaciones Diferenciales y Análisis Numérico
C/ Tarfia s/n, Facultad de Matemáticas, Universidad de Sevilla,
41012 Sevilla, Spain

RENATO COLUCCI

Dipartimento di Ingegneria Industriale e Scienze Matematiche,
Università Politecnica delle Marche,
Via Breccie Bianche 12, 60131, Ancona, Italy.

JAVIER LÓPEZ-DE-LA-CRUZ*

Departamento de Ecuaciones Diferenciales y Análisis Numérico
C/ Tarfia s/n, Facultad de Matemáticas, Universidad de Sevilla
41012 Sevilla, Spain

AND ALAIN RAPAPORT

MISTEA, Univ. Montpellier, Inra, Montpellier SupAgro
2, place Pierre Viala
34060 Montpellier, France

ABSTRACT. In this corrigendum we correct an error in our paper [T. Caraballo, R. Colucci, J. López-de-la-Cruz and A. Rapaport. A way to model stochastic perturbations in population dynamics models with bounded realizations, *Commun Nonlinear Sci Numer Simulat*, **77** (2019) 239–257]. We present a correct way to model real noisy perturbations by considering a slightly different stochastic process based, as in the original paper, on the Ornstein-Uhlenbeck process. Namely, we correct the formulae that generates the noisy realizations to ensure the boundedness property to be satisfied with probability one (which turns out not to be true in our original paper even though it was observed in all the simulations). Once this modification is done, every result and every example in the initial work remain valid, in fact, the same goal is achieved.

1. INTRODUCTION

In our paper [3] we presented a new way to model real noisy perturbations by making use of the well-known Ornstein-Uhlenbeck (OU) process. In addition, we illustrated this idea with different examples coming from population dynamics and several numerical simulations.

Let us describe firstly the explanations given in [3] in order to let the reader understand the error in [3] and the changes needed to preserve the spirit and the

2010 *Mathematics Subject Classification*. Primary 92D25, 34F05, Secondary 34C60.

Key words and phrases. random perturbation, population dynamics, logistic model, prey-predator model, observer.

*Corresponding author: jlopez78@us.es

We thank the anonymous referees of another work, who pointed out a problem when we applied a result from [1], helping us to find the solution.

goal of our original work.

It is well known that most real phenomena are subject to suffer the effects of some kind of noise. Even though there are many works in the literature leading with stochastic models, they mainly focus on making use of the standard Wiener process, in spite of the fact that it is unbounded with probability one. This last property is completely unrealistic from the point of view of applications, since random perturbations in real life are bounded. In order to understand better the drawback caused by the standard Wiener process when modeling noise, we refer the readers to [4, 5, 6], where the authors investigate chemostat models by means of the Brownian motion.

Motivated by this fact, we presented in [3] a way to model bounded real perturbations involving the OU process, which has proved to fit in a very loyal way the random disturbances observed in real life.

More precisely, we explained that we could consider perturbations of some parameter in a random system of the following kind

$$(1) \quad \dot{x} = f(x, z_{\beta, \gamma}^*(\theta_t \omega)),$$

where $z_{\beta, \gamma}^*(\theta_t \omega)$ denotes the OU process. This type of systems arises when replacing, for instance, some deterministic parameter, namely a , by $a + \alpha z_{\beta, \gamma}^*(\theta_t \omega)$, where $\alpha > 0$ denotes the intensity of the noise.

In order to deal with system (1), we introduced in [3] the OU process which is defined by the random variable

$$(2) \quad z_{\beta, \gamma}^*(\theta_t \omega) = -\beta\gamma \int_{-\infty}^0 e^{\beta s} \theta_t \omega(s) ds, \quad t \in \mathbb{R}, \omega \in \Omega, \beta, \gamma > 0,$$

and solves the Langevin equation

$$(3) \quad dz + \beta z dt = \gamma d\omega,$$

and then we provided some properties, as the ones in Proposition 2 in [3] that we recall below.

Proposition 1.1. *There exists a θ_t -invariant set $\tilde{\Omega} \in \mathcal{F}$ of Ω of full \mathbb{P} -measure such that for $\omega \in \tilde{\Omega}$ and $\beta, \gamma > 0$, we have*

- (i) *the random variable $|z_{\beta, \gamma}^*(\omega)|$ is tempered.*
- (ii) *the mapping*

$$(t, \omega) \rightarrow z_{\beta, \gamma}^*(\theta_t \omega) = -\beta\gamma \int_{-\infty}^0 e^{\beta s} \omega(t+s) ds + \omega(t)$$

is a stationary solution of (2) with continuous trajectories;

(iii) for any $\omega \in \tilde{\Omega}$ one has

$$\begin{aligned} \lim_{t \rightarrow \pm\infty} \frac{|z_{\beta,\gamma}^*(\theta_t\omega)|}{t} &= 0; \\ \lim_{t \rightarrow \pm\infty} \frac{1}{t} \int_0^t z_{\beta,\gamma}^*(\theta_s\omega) ds &= 0; \\ \lim_{t \rightarrow \pm\infty} \frac{1}{t} \int_0^t |z_{\beta,\gamma}^*(\theta_s\omega)| ds &= \mathbb{E}[z_{\beta,\gamma}^*] < \infty; \end{aligned}$$

(iv) finally, for any $\omega \in \tilde{\Omega}$,

$$\lim_{\beta \rightarrow \infty} z_{\beta,\gamma}^*(\theta_t\omega) = 0, \quad \text{for all } t \in \mathbb{R}.$$

Hence, the idea was the following: (1) we are given an interval, namely $[a_1, a_2]$, (2) we fix any event $\omega \in \tilde{\Omega}$ and (3) for every fixed $\omega \in \tilde{\Omega}$, thanks to Proposition 1.1 (iv), which can be found in [1], Lemma 4.1, we can choose β large enough such that $a + \alpha z_{\beta,\gamma}^*(\theta_t\omega) \in [a_1, a_2]$ for all $t \in \mathbb{R}$.

Then, Proposition 1.1 (iv) was the key of our paper since it allowed us to ensure that the OU process $z_{\beta,\gamma}^*(\theta_t\omega)$ was bounded in some desired interval (typically fixed by practitioners) for every fixed event $\omega \in \tilde{\Omega}$ and β large enough.

As we explained in [3], since β could depend on ω , the resulting random system does not generate a random dynamical system (RDS) and then the theory of RDSs and pullback attractors could not be applied. But this was not a problem at all, since the random system could be analyzed for every fixed event ω . In fact, this allowed us to prove every result of the paper forwards in time, which is a more natural way than the pullback one.

Nevertheless, we noticed that property (iv) in Proposition 1.1 (Proposition 2 (iv) in [3]) is not true. By having a deeper look at the proof of the result in [1] we realized that such a result cannot be proved uniformly in time (as we first understood in [1]), but for every fixed t .

After this, we found a slightly different way to model bounded noises which allows us to prove every result and present every example in the original paper [3] in a very similar manner. Let us present now the new correct idea.

Assume that we want to consider some random perturbations in a certain parameter a of a deterministic differential system. In addition, as in real life, those disturbances must be bounded (for all the time) in some interval around a , namely $[a - d, a + d]$, where $d > 0$.

Instead of replacing a by $a + \alpha z_{\beta,\gamma}^*(\theta_t\omega)$, as we did in [3], we replace a by $a + \Phi(z^*(\theta_t\omega))$, where $\Phi(z) = \frac{2d}{\pi} \arctan(z)$ and $z^*(\theta_t\omega)$ denotes the OU process (2). In this case, β and γ are fixed through the whole work. Hence, instead of having to deal with system (1), the resulting random system is of the kind

$$(4) \quad \dot{x} = f(x, \Phi(z^*(\theta_t\omega))),$$

and it will be investigated for every fixed event $\omega \in \Omega$, as in the original work [3].

We notice that, in this case, as β and γ are fixed, the resulting random system (4) generates a RDS and then the theory of RDSs and pullback attractors could

be applied (which was not the case in the original paper). Nevertheless, instead of using that theory and pullback attraction, we study the random system (4) for every fixed $\omega \in \Omega$ since it allows us to prove every result forwards in time (which is, as explained before, more natural than the pullback one).

It is easy to check that the random perturbations $a + \Phi(z^*(\theta_t\omega))$ are bounded for every $t \in \mathbb{R}$ and any $\omega \in \Omega$, since Φ is bounded. In fact, for every $\omega \in \Omega$, we have

$$a - d \leq a + \Phi(z^*(\theta_t\omega)) \leq a + d, \quad \text{for all } t \in \mathbb{R}.$$

In addition, we can prove the following property.

Proposition 1.2. *Let $\Phi(z) = \frac{2d}{\pi} \arctan(z)$. Then*

$$(5) \quad \lim_{t \rightarrow +\infty} \frac{1}{t} \int_0^t \Phi(z^*(\theta_s\omega)) ds = 0, \quad \text{a.s. in } \Omega.$$

Proof. Since $\Phi(z) = \frac{2d}{\pi} \arctan(z)$, then we have

$$\int_{\Omega} \left| \frac{2d}{\pi} \arctan(z^*(\omega)) \right| d\mathbb{P}(\omega) \leq d|\Omega| = d$$

whence $\Phi(z^*(\cdot)) \in L^1(\Omega, \mathcal{F}, \mathbb{P})$. Then, since \mathbb{P} is invariant by θ (see [2, 7]), from the Birkhoff ergodic theorem we deduce that

$$\lim_{t \rightarrow +\infty} \frac{1}{t} \int_0^t \Phi(z^*(\theta_s\omega)) ds = \mathbb{E}[\Phi(z^*(\omega))], \quad \text{a.s. in } \Omega.$$

Hence, it is enough to show that $\mathbb{E}[\Phi(z^*(\omega))] = 0$. In fact, we have

$$\mathbb{E}[\Phi(z^*(\omega))] = \int_{\mathbb{R}} \Phi(x) f_{OU}(x) dx = 0,$$

where f_{OU} denotes the density function of the random variable $z^*(\cdot)$, which is an even function since it is Gaussian, and Φ is an odd function. \square

Hence, the stochastic process $\Phi(z^*(\omega))$ satisfies the desired ergodic property (which was false in Proposition 1.1 (iv)) and it is bounded. Then, every result in our original work [3] is true when considering this little modification and every example there can be given to illustrate this new way to model bounded random perturbations.

Concerning the numerical simulations, they can be done again by considering the new function $\Phi(z) = \frac{2d}{\pi} \arctan(z)$ and they are analogous to the ones in the original work [3].

To finish, we would like to remark that we could also use any odd measurable function Φ such that

$$\lim_{z \rightarrow +\infty} \Phi(z) = d < +\infty.$$

REFERENCES

- [1] S. Al-Azzawi, J. Liu and X. Liu, Convergence rate of synchronization of systems with additive noise, *Discrete Contin. Dyn. Syst. Ser. B*, **22** (2017), 227–245.
- [2] L. Arnold, *Random Dynamical Systems*, Springer Berlin Heidelberg, 1998.
- [3] T. Caraballo, R. Colucci, J. López-de-la-Cruz and A. Rapaport, A way to model stochastic perturbations in population dynamics models with bounded realizations, *Communications in Nonlinear Science and Numerical Simulation*.
- [4] T. Caraballo, M. J. Garrido-Atienza and J. López-de-la-Cruz, *Some Aspects Concerning the Dynamics of Stochastic Chemostats*, vol. 69, chapter 11, 227–246, Springer International Publishing, Cham, 2016.

- [5] T. Caraballo, M. J. Garrido-Atienza and J. López-de-la-Cruz, Dynamics of some stochastic chemostat models with multiplicative noise, *Communications on Pure and Applied Analysis*, **16** (2017), 1893–1914.
- [6] T. Caraballo, M. J. Garrido-Atienza, J. López-de-la-Cruz and A. Rapaport, Modeling and analysis of random and stochastic input flows in the chemostat model, *Discrete & Continuous Dynamical Systems - Series B*.
- [7] T. Caraballo and X. Han, *Applied Nonautonomous and Random Dynamical Systems, Applied Dynamical Systems*, Springer International Publishing, 2016.

E-mail address: caraball@us.es

E-mail address: renatocolucci@hotmail.com

E-mail address: jlopez78@us.es

E-mail address: alain.rapaport@inra.fr